

Title

D. Zack Garza

Monday 27th July, 2020

Contents

1	Spring 2017	1
1.1	1	1
1.2	2	2
	1.2.1 a	2
	1.2.2 b	3
1.3	3	3
	1.3.1 a	3
	1.3.2 b	4
1.4	4	4
1.5	5	4

1 Spring 2017

1.1 1

Concepts used:

- Definition: A is *nowhere dense* \iff every interval I contains a subinterval $S \subseteq A^c$.

Solution

- Claim: K is compact.
 - It suffices to show that $K^c := [0, 1] \setminus K$ is open; Then K will be a closed and bounded subset of \mathbb{R} and thus compact by Heine-Borel.
 - Strategy: write K^c as the union of open intervals (since these form a basis for the Euclidean topology on \mathbb{R}).
 - Identify K^c as the set of real numbers in $[0, 1]$ whose decimal expansion **does** use a 4.
 - Let $x \in K^c$, suppose a 4 occurs as the k th digit, and write

$$x = 0.d_1d_2 \cdots d_{k-1} 4 d_{k+1} \cdots = \left(\sum_{j=1}^k d_j 10^{-j} \right) + (4 \cdot 10^{-k}) + \left(\sum_{j=k+1}^{\infty} d_j 10^{-j} \right).$$

- Set $r < 10^{-k}$ and pick any $y \in [0, 1]$ such that $y \in B_r(x)$, i.e. $|x - y| < 10^{-k}$.

- If we write $y = \sum_{j=1}^{\infty} c_j 10^{-j}$, this means that for all $j \leq k$ we have $d_j = c_j$, and in particular $d_k = 4 = c_k$, so y has a 4 in its decimal expansion.
- But then $K^c = \bigcup_x B_r(x)$ is a union of open sets and thus open.

• **Claim: K is nowhere dense and $m(K) = 0$:**

Since K is closed, we'll show that K can not properly contain any interval, so $(\overline{K})^\circ = \emptyset$.

As in the construction of the Cantor set, let

- K_1 denote $[0, 1]$ with 1 interval $[0.4, 0.5]$ of length $\frac{1}{10}$ deleted
- K_2 denote K_1 with 9 intervals $[0.04, 0.05], [0.14, 0.15], \dots [0.94, 0.95]$ length $\frac{1}{100}$ deleted
- K_n denote K_{n-1} with 9^{n-1} such intervals of length 10^{-n} deleted.

Then $K = \bigcap K_n$, and

$$m(K) = 1 - m(K^c) = 1 - \sum_{j=0}^{\infty} \frac{9^j}{10^{j+1}} = 1 - \frac{1}{10} \left(\frac{1}{1 - \frac{9}{10}} \right) = 0,$$

and since any interval has strictly positive measure, K can not contain any interval.

• **Claim: K has no isolated points:**

A point $x \in K$ is isolated iff there is an open ball $B_r(x)$ containing x such that $B_r(x) \cap K = \{x\}$, so every point in this ball has a 4 in its decimal expansion.

Note that $m(K_n) = \left(\frac{9}{10}\right)^n \rightarrow 0$ and that the endpoints of intervals are never removed and are thus elements of K . Then for every ε , we can choose n such that $\left(\frac{9}{10}\right)^n < \varepsilon$; then there is an endpoint of a removed interval e_n satisfying $|x - e_n| \leq \left(\frac{9}{10}\right)^n < \varepsilon$.

So every ball containing x contains some endpoint of a removed interval, and thus an element of K . ■

1.2 2

$$\lambda \ll \mu \iff E \in \mathcal{M}, \mu(E) = 0 \implies \lambda(E) = 0.$$

1.2.1 a

By Radon-Nikodym, if $\lambda \ll \mu$ then $d\lambda = f d\mu$, which would yield

$$\int g d\lambda = \int g f d\mu.$$

So let E be measurable and suppose $\mu(E) = 0$. Then

$$\lambda(E) := \int_E f \, d\mu = \lim_n \left\{ \varphi_n := \sum_j c_j \mu(E_j) \right\},$$

where we take a sequence of simple functions increasing to f .

But since each $E_j \subseteq E$, we must have $\mu(E_j) = 0$ for any such E_j , so every such φ_n must be zero and thus $\lambda(E) = 0$.

1.2.2 b

By Radon-Nikodym, there exists a positive f such that

$$\int g \, dm = \int gf \, d\mu,$$

where we can take $g(x) = x^2$, then the LHS is zero by assumption and thus so is the RHS.

Note that gf is positive.

Define $A_k = \left\{ x \in X \mid gf\chi_E > \frac{1}{k} \right\}$, then by Chebyshev

$$\mu(A_k) \leq k \int_E gf \, d\mu = 0,$$

which holds for every k .

Then noting that $A_k \searrow A := \{x \in E \mid x^2 > 0\}$, and gf is positive, we have

$$x \in E \iff gf\chi_E(x) > 0 \iff x \in A,$$

so $E = A$ and $\mu(E) = \mu(A)$.

But since $m \ll \mu$ by construction, we can conclude that $m(E) = 0$. ■

1.3 3

1.3.1 a

Letting $x_n := \frac{1}{n}$, we have

$$\sum_{k=1}^{\infty} |f_k(x)| \geq |f_n(x_n)| = \left| ae^{-ax} - be^{-bx} \right| := M.$$

In particular, $\sup_x |f_n(x)| \not\rightarrow 0$, so the terms do not go to zero and the sum can not converge.

1.3.2 b

?

1.4 4

Switching to polar coordinates and integrating over a half-circle contained in I^2 , we have

$$\int_{I^2} f \geq \int_0^\pi \int_0^1 \frac{\cos(\theta) \sin(\theta)}{r^2} dr d\theta = \infty,$$

so f is not integrable.

1.5 5

See <https://math.stackexchange.com/questions/507263/prove-that-c1a-b-with-the-c1-norm-is-a-banach-space>

This is clearly a norm, which we'll write $\|\cdot\|_u$

Let f_n be a Cauchy sequence and define a candidate limit $f(x) = \lim_n f_n(x)$.

Then noting that $\|f_n\|_\infty, \|f'_n\|_\infty \leq \|f_n\|_u < \infty$, both f_n, f'_n are Cauchy sequences in $C^0([a, b], \|\cdot\|_\infty)$, which is a Banach space.

So $f_n \rightarrow f$ uniformly, and $f'_n \rightarrow g$ uniformly for some g , and moreover $f, g \in C^0([a, b])$.

We thus have

$$\begin{aligned} f_n(x) - f_n(a) &\xrightarrow{u} f(x) - f(a) \\ \int_a^x f'_n &\xrightarrow{u} \int_a^x g, \end{aligned}$$

and by the FTC, the left-hand sides are equal, and by uniqueness of limits so are the right-hand sides, so $f' = g$.

Since $f, f' \in C^0([a, b])$, they are bounded, and so $\|f\|_u < \infty$. This means that $\|f_n - f\|_u \rightarrow 0$, so f_n converges to f , which is in the same space. ■