

Topology Qualifying Exam Notes

D. Zack Garza

Friday 29th May, 2020

Contents

1	Definitions	1
2	Examples	2
2.1	Common Spaces and Operations	2
2.2	Alternative Topologies	5
3	Theorems	5
3.1	Point-Set	5
3.2	Algebraic	7

1 Definitions

- Topology: Closed under arbitrary unions and finite intersections.
- Basis: A subset $\{B_i\}$ is a basis iff
 - $x \in X \implies x \in B_i$ for some i .
 - $x \in B_i \cap B_j \implies x \in B_k \subset B_i \cap B_j$.
 - Topology generated by this basis: $x \in N_x \implies x \in B_i \subset N_x$ for some i .
- Dense: A subset $Q \subset X$ is dense iff $y \in N_y \subset X \implies N_y \cap Q \neq \emptyset$ iff $\bar{Q} = X$.
- Neighborhood: A neighborhood of a point x is any open set containing x .
- Hausdorff
- Second Countable: admits a countable basis.
- Closed (several characterizations)
- Closure in a subspace: $Y \subset X \implies \text{cl}_Y(A) := \text{cl}_X(A) \cap Y$.
- Bounded
- Compact: A topological space (X, τ) is **compact** if every open cover has a *finite* subcover.

That is, if $\{U_j \mid j \in J\} \subset \tau$ is a collection of open sets such that $X \subseteq \bigcup_{j \in J} U_j$, then there exists a *finite* subset $J' \subset J$ such that $X \subseteq \bigcup_{j \in J'} U_j$.

- Locally compact For every $x \in X$, there exists a $K_x \ni x$ such that K_x is compact.
- Connected: There does not exist a disconnecting set $X = A \amalg B$ such that $\emptyset \neq A, B \subsetneq X$, i.e. X is the union of two proper disjoint nonempty sets.

Equivalently, X contains no proper nonempty clopen sets.

– Additional condition for a subspace $Y \subset X$: $\text{cl}_Y(A) \cap V = A \cap \text{cl}_Y(B) = \emptyset$.

- Locally connected: A space is locally connected at a point x iff $\forall N_x \ni x$, there exists a $U \subset N_x$ containing x that is connected.
- Retract: A subspace $A \subset X$ is a *retract* of X iff there exists a continuous map $f : X \rightarrow A$ such that $f|_A = \text{id}_A$. Equivalently it is a *left* inverse to the inclusion.
- Uniform Continuity: For $f : (X, d_x) \rightarrow (Y, d_Y)$ metric spaces,

$$\forall \varepsilon > 0, \exists \delta > 0 \text{ such that } d_X(x_1, x_2) < \delta \implies d_Y(f(x_1), f(x_2)) < \varepsilon.$$

- Lebesgue number: For (X, d) a compact metric space and $\{U_\alpha\} \rightrightarrows X$, there exist $\delta_L > 0$ such that

$$A \subset X, \text{diam}(A) < \delta_L \implies A \subseteq U_\alpha \text{ for some } \alpha.$$

- Paracompact
- Components: Set $x \sim y$ iff there exists a connected set $U \ni x, y$ and take equivalence classes.
- Path Components: Set $x \sim y$ iff there exists a path-connected set $U \ni x, y$ and take equivalence classes.
- Separable: Contains a countable dense subset.
- Limit Point: For $A \subset X$, x is a limit point of A if every punctured neighborhood P_x of x satisfies $P_x \cap A \neq \emptyset$, i.e. every neighborhood of x intersects A in some point other than x itself.

Equivalently, x is a limit point of A iff $x \in \text{cl}_X(A \setminus \{x\})$.

2 Examples

2.1 Common Spaces and Operations

Point-Set:

- Finite discrete sets with the discrete topology
- Subspaces of \mathbb{R} : $(a, b), (a, b], (a, \infty)$, etc.

$$- \{0\} \cup \left\{ \frac{1}{n} \mid n \in \mathbb{Z}^{\geq 1} \right\}$$

- \mathbb{Q}
- The topologist's sine curve
- One-point compactifications
- \mathbb{R}^ω


Non-Hausdorff spaces:

- The cofinite topology on any infinite set.
- \mathbb{R}/\mathbb{Q}
- The line with two origins.

General Spaces:

$$S^n, \mathbb{D}^n, T^n \mathbb{R}P^n, \mathbb{C}P^n, M, K, \Sigma_g, \mathbb{R}P^\infty, \mathbb{C}P^\infty.$$

- Knot complements in S^3
- Lens spaces
- Prism spaces
- $\mathbb{H}P^n$
- Dunce Cap
- Matrix groups
- Pair of pants
- Covering spaces (hyperbolic geometry)
- Seifert surfaces
- Surgery
- Hawaiian earring
- Horned sphere
- Cantor set
- Simplicial Complexes
 - Nice minimal example:



Operations

- Cartesian product $A \times B$
- Wedge product $A \vee B$

- Connect Sum $A \# B$
- Quotienting A/B
- Puncturing $A \setminus \{a_i\}$
- Smash product
- Join
- Cones
- Suspension
- Loop space
- Identifying a finite number of points

2.2 Alternative Topologies

- Discrete
- Cofinite
- Discrete and Indiscrete
- Uniform

The cofinite topology:

- Non-Hausdorff
- Compact

The discrete topology:

- Discrete iff points are open
- Always Hausdorff
- Compact iff finite
- Totally disconnected
- If the domain, every map is continuous

The indiscrete topology:

- Only open sets are \emptyset, X
- Non-Hausdorff
- If the codomain, every map is continuous
- Compact

3 Theorems

3.1 Point-Set

- Closed subsets of Hausdorff spaces are compact? (check)
- Cantor's intersection theorem?
- Tube lemma
- Properties pushed forward through continuous maps:
 - Compactness?
 - Connectedness (when surjective)
 - Separability

- Density **only when** f is surjective
- **Not** openness
- **Not** closedness
- A retract of a Hausdorff/connected/compact space is closed/connected/compact respectively.

Proposition 3.1.

A continuous function on a compact set is uniformly continuous.

Proof.

Take $\{B_{\frac{\varepsilon}{2}}(y) \mid y \in Y\} \rightrightarrows Y$, pull back to an open cover of X , has Lebesgue number $\delta_L > 0$, then $x' \in B_{\delta_L}(x) \implies f(x), f(x') \in B_{\frac{\varepsilon}{2}}(y)$ for some y . ■

- Lipschitz continuity implies uniform continuity (take $\delta = \varepsilon/C$)
 - Counterexample to converse: $f(x) = \sqrt{x}$ on $[0, 1]$ has unbounded derivative.
- Extreme Value Theorem: for $f : X \rightarrow Y$ continuous with X compact and Y ordered in the order topology, there exist points $c, d \in X$ such that $f(x) \in [f(c), f(d)]$ for every x .

Theorem 3.2.

Points are closed in T_1 spaces.

Theorem 3.3.

A metric space X is sequentially compact iff it is complete and totally bounded.

Theorem 3.4.

A metric space is totally bounded iff every sequence has a Cauchy subsequence.

Theorem 3.5.

A metric space is compact iff it is complete and totally bounded.

Theorem 3.6 (Baire).

If X is a complete metric space, then the intersection of countably many dense open sets is dense in X .

Theorem 3.7.

A continuous bijective open map is a homeomorphism.

Theorem 3.8.

A closed subset A of a compact set B is compact.

Proof.

- Let $\{A_i\} \rightrightarrows A$ be a covering of A by sets open in A .
- Each $A_i = B_i \cap A$ for some B_i open in B (definition of subspace topology)
- Define $V = \{B_i\}$, then $V \rightrightarrows A$ is an open cover.

- Since A is closed, $W := B \setminus A$ is open
- Then $V \cup W$ is an open cover of B , and has a finite subcover $\{V_i\}$
- Then $\{V_i \cap A\}$ is a finite open cover of A .

■

Theorem 3.9.

The continuous image of a compact set is compact.

Theorem 3.10.

A closed subset of a Hausdorff space is compact.

3.2 Algebraic

Todo: Merge the two van Kampen theorems.

Theorem 3.11 (Van Kampen).

The pushout is the northwest colimit of the following diagram

$$\begin{array}{ccc} A \amalg_Z B & \longleftarrow & A \\ & & \uparrow \iota_A \\ B & \xleftarrow{\iota_B} & Z \end{array}$$

For groups, the pushout is given by the amalgamated free product: if $A = \langle G_A \mid R_A \rangle$, $B = \langle G_B \mid R_B \rangle$, then $A *_Z B = \langle G_A, G_B \mid R_A, R_B, T \rangle$ where T is a set of relations given by $T = \{ \iota_A(z) \iota_B(z)^{-1} \mid z \in Z \}$.

Example: $A = \mathbb{Z}/4\mathbb{Z} = \langle x \mid x^4 \rangle$, $B = \mathbb{Z}/6\mathbb{Z} = \langle y \mid y^6 \rangle$, $Z = \mathbb{Z}/2\mathbb{Z} = \langle z \mid z^2 \rangle$. Then we can identify Z as a subgroup of A, B using $\iota_A(z) = x^2$ and $\iota_B(z) = y^3$. So

$$A *_Z B = \langle x, y \mid x^4, y^6, x^2 y^{-3} \rangle$$

Suppose $X = U_1 \cup U_2$ such that $U_1 \cap U_2 \neq \emptyset$ is path connected. Then taking $x_0 \in U := U_1 \cap U_2$ yields a pushout of fundamental groups

$$\pi_1(X; x_0) = \pi_1(U_1; x_0) *_{\pi_1(U; x_0)} \pi_1(U_2; x_0).$$

Theorem 3.12 (Van Kampen).

If $X = U \cup V$ where $U, V, U \cap V$ are all path-connected then

$$\pi_1(X) = \pi_1 U *_{\pi_1(U \cap V)} \pi_1 V,$$

where the amalgamated product can be computed as follows: If we have presentations

$$\begin{aligned}\pi_1(U, w) &= \langle u_1, \dots, u_k \mid \alpha_1, \dots, \alpha_l \rangle \\ \pi_1(V, w) &= \langle v_1, \dots, v_m \mid \beta_1, \dots, \beta_n \rangle \\ \pi_1(U \cap V, w) &= \langle w_1, \dots, w_p \mid \gamma_1, \dots, \gamma_q \rangle\end{aligned}$$

then

$$\begin{aligned}\pi_1(X, w) &= \langle u_1, \dots, u_k, v_1, \dots, v_m \rangle \\ &\quad \text{mod } \langle \alpha_1, \dots, \alpha_l, \beta_1, \dots, \beta_n, I(w_1)J(w_1)^{-1}, \dots, I(w_p)J(w_p)^{-1} \rangle \\ &= \frac{\pi_1(U) * \pi_1(B)}{\langle \{I(w_i)J(w_i)^{-1} \mid 1 \leq i \leq p\} \rangle}\end{aligned}$$

where

$$\begin{aligned}I &: \pi_1(U \cap V, w) \rightarrow \pi_1(U, w) \\ J &: \pi_1(U \cap V, w) \rightarrow \pi_1(V, w).\end{aligned}$$