

# UGA Algebra Qualifying Exam Questions

D. Zack Garza

Monday 8<sup>th</sup> June, 2020

## Contents

<b>1</b>	<b>Spring 2020</b>	<b>2</b>
1.1	1 . . . . .	2
1.2	2 . . . . .	2
1.3	3 . . . . .	2
1.4	4 . . . . .	3
1.5	5 . . . . .	3
1.6	6 . . . . .	3
1.7	7 . . . . .	3
1.8	8 . . . . .	4
<b>2</b>	<b>Spring 2019</b>	<b>4</b>
2.1	1 . . . . .	4
2.2	2 . . . . .	4
2.3	3 . . . . .	4
2.4	4 . . . . .	4
2.5	5 . . . . .	5
2.6	6 . . . . .	5
2.7	7 . . . . .	5
2.8	8 . . . . .	5
<b>3</b>	<b>Fall 2019</b>	<b>6</b>
3.1	1 . . . . .	6
3.2	2 . . . . .	6
3.3	3 . . . . .	6
3.4	4 . . . . .	6
3.5	5 . . . . .	6
3.6	6 . . . . .	7
3.7	7 . . . . .	7
3.8	8 . . . . .	7
<b>4</b>	<b>2019 Course Exams</b>	<b>8</b>
4.1	Midterm . . . . .	8
4.2	Final . . . . .	8

---

<b>5</b>	<b>Spring 2018</b>	<b>9</b>
5.1	1. . . . .	9
5.2	2. . . . .	9
5.3	3. . . . .	9
5.4	4. . . . .	10
5.5	5. . . . .	10
5.6	6. . . . .	10
5.7	7. . . . .	11
5.8	8. . . . .	11
<b>6</b>	<b>Fall 2018</b>	<b>11</b>
6.1	1. . . . .	11
6.2	2. . . . .	11
6.3	3. . . . .	12
6.4	4. . . . .	12
6.5	5. . . . .	12
6.6	6. . . . .	12
6.7	7. . . . .	12
<b>7</b>	<b>Fall 2017</b>	<b>13</b>
7.1	1. . . . .	13
7.2	2. . . . .	13
7.3	3. . . . .	14
7.4	4. . . . .	14
7.5	5. . . . .	14
7.6	6. . . . .	14
7.7	7. . . . .	14

## 1 Spring 2020

### 1.1 1

- Show that any group of order 2020 is solvable.
- Give (without proof) a classification of all abelian groups of order 2020.
- Describe one nonabelian group of order 2020.

### 1.2 2

Let  $H$  be a normal subgroup of a finite group  $G$  where the order of  $H$  and the index of  $H$  in  $G$  are relatively prime. Prove that no other subgroup of  $G$  has the same order as  $H$ .

### 1.3 3

Let  $E$  be an extension field of  $F$  and  $\alpha \in E$  be algebraic of odd degree over  $F$ .

- Show that  $F(\alpha) = F(\alpha^2)$ .

- b. Prove that  $\alpha^{2020}$  is algebraic of odd degree over  $F$ .

**1.4 4**

Let  $f(x) = x^4 - 2 \in \mathbb{Q}[x]$ .

- Define what it means for a finite extension field  $E$  of a field  $F$  to be a Galois extension.
- Determine the Galois group  $\text{Gal}(E/\mathbb{Q})$  for the polynomial  $f(x)$ , and justify your answer carefully.
- Exhibit a subfield  $K$  in (b) such that  $\mathbb{Q} \leq K \leq E$  with  $K$  not a Galois extension over  $\mathbb{Q}$ . Explain.

**1.5 5**

Let  $R$  be a ring and  $f : M \rightarrow N$  and  $g : N \rightarrow M$  be  $R$ -module homomorphisms such that  $g \circ f = \text{id}_M$ . Show that  $N \cong \text{im } f \oplus \ker g$ .

**1.6 6**

Let  $R$  be a ring with unity.

- Give a definition for a free module over  $R$ .
- Define what it means for an  $R$ -module to be torsion free.
- Prove that if  $F$  is a free module, then any short exact sequence of  $R$ -modules of the following form splits:

$$0 \rightarrow N \rightarrow M \rightarrow F \rightarrow 0.$$

- Let  $R$  be a PID. Show that any finitely generated  $R$ -module  $M$  can be expressed as a direct sum of a torsion module and a free module. You may assume that a finitely generated torsion module over a PID is free.

**1.7 7**

Let

$$A = \begin{bmatrix} 2 & 0 & 0 \\ 4 & 6 & 1 \\ -16 & -16 & -2 \end{bmatrix} \in M_3(\mathbb{C}).$$

- Find the Jordan canonical form  $J$  of  $A$ .
- Find an invertible matrix  $P$  such that  $P^{-1}AP = J$ . You should not need to compute  $P^{-1}$ .
- Write down the minimal polynomial of  $A$ .

**1.8 8**

Let  $T : V \rightarrow V$  be a linear transformation where  $V$  is a finite-dimensional vector space over  $\mathbb{C}$ . Prove the Cayley-Hamilton theorem: if  $p(x)$  is the characteristic polynomial of  $T$ , then  $p(T) = 0$ . You may use canonical forms.

**2 Spring 2019****2.1 1.**

Let  $A$  be a square matrix over the complex numbers. Suppose that  $A$  is nonsingular and that  $A^{2019}$  is diagonalizable over  $\mathbb{C}$ .

Show that  $A$  is also diagonalizable over  $\mathbb{C}$ .

**2.2 2.**

Let  $F = \mathbb{F}_p$ , where  $p$  is a prime number.

- (a) Show that if  $\pi(x) \in F[x]$  is irreducible of degree  $d$ , then  $\pi(x)$  divides  $x^{p^d} - x$ .
- (b) Show that if  $\pi(x) \in F[x]$  is an irreducible polynomial that divides  $x^{p^n} - x$ , then  $\deg \pi(x)$  divides  $n$ .

**2.3 3.**

How many isomorphism classes are there of groups of order 45?

Describe a representative from each class.

**2.4 4.**

For a finite group  $G$ , let  $c(G)$  denote the number of conjugacy classes of  $G$ .

- (a) Prove that if two elements of  $G$  are chosen uniformly at random, then the probability they commute is precisely

$$\frac{c(G)}{|G|}.$$

- (b) State the class equation for a finite group.
- (c) Using the class equation (or otherwise) show that the probability in part (a) is at most

$$\frac{1}{2} + \frac{1}{2[G : Z(G)]}.$$

Here, as usual,  $Z(G)$  denotes the center of  $G$ .

**2.5 5.**

Let  $R$  be an integral domain. Recall that if  $M$  is an  $R$ -module, the *rank* of  $M$  is defined to be the maximum number of  $R$ -linearly independent elements of  $M$ .

- (a) Prove that for any  $R$ -module  $M$ , the rank of  $\text{Tor}(M)$  is 0.
- (b) Prove that the rank of  $M$  is equal to the rank of  $M/\text{Tor}(M)$ .
- (c) Suppose that  $M$  is a non-principal ideal of  $R$ .

Prove that  $M$  is torsion-free of rank 1 but not free.

**2.6 6.**

Let  $R$  be a commutative ring with 1.

Recall that  $x \in R$  is nilpotent iff  $x^n = 0$  for some positive integer  $n$ .

- (a) Show that every proper ideal of  $R$  is contained within a maximal ideal.
- (b) Let  $J(R)$  denote the intersection of all maximal ideals of  $R$ .  
Show that  $x \in J(R) \iff 1 + rx$  is a unit for all  $r \in R$ .
- (c) Suppose now that  $R$  is finite. Show that in this case  $J(R)$  consists precisely of the nilpotent elements in  $R$ .

**2.7 7.**

Let  $p$  be a prime number. Let  $A$  be a  $p \times p$  matrix over a field  $F$  with 1 in all entries except 0 on the main diagonal.

Determine the Jordan canonical form (JCF) of  $A$

- (a) When  $F = \mathbb{Q}$ ,
- (b) When  $F = \mathbb{F}_p$ .

Hint: In both cases, all eigenvalues lie in the ground field. In each case find a matrix  $P$  such that  $P^{-1}AP$  is in JCF.

**2.8 8.**

Let  $\zeta = e^{2\pi i/8}$ .

- (a) What is the degree of  $\mathbb{Q}(\zeta)/\mathbb{Q}$ ?
- (b) How many quadratic subfields of  $\mathbb{Q}(\zeta)$  are there?
- (c) What is the degree of  $\mathbb{Q}(\zeta, \sqrt[4]{2})$  over  $\mathbb{Q}$ ?

---

### 3 Fall 2019

#### 3.1 1

Let  $G$  be a finite group with  $n$  distinct conjugacy classes. Let  $g_1 \cdots g_n$  be representatives of the conjugacy classes of  $G$ .

Prove that if  $g_i g_j = g_j g_i$  for all  $i, j$  then  $G$  is abelian.

#### 3.2 2

Let  $G$  be a group of order 105 and let  $P, Q, R$  be Sylow 3, 5, 7 subgroups respectively.

- (a) Prove that at least one of  $Q$  and  $R$  is normal in  $G$ .
- (b) Prove that  $G$  has a cyclic subgroup of order 35.
- (c) Prove that both  $Q$  and  $R$  are normal in  $G$ .
- (d) Prove that if  $P$  is normal in  $G$  then  $G$  is cyclic.

#### 3.3 3

Let  $R$  be a ring with the property that for every  $a \in R$ ,  $a^2 = a$ .

- (a) Prove that  $R$  has characteristic 2.
- (b) Prove that  $R$  is commutative.

#### 3.4 4

Let  $F$  be a finite field with  $q$  elements.

Let  $n$  be a positive integer relatively prime to  $q$  and let  $\omega$  be a primitive  $n$ th root of unity in an extension field of  $F$ .

Let  $E = F[\omega]$  and let  $k = [E : F]$ .

- (a) Prove that  $n$  divides  $q^k - 1$ .
- (b) Let  $m$  be the order of  $q$  in  $\mathbb{Z}/n\mathbb{Z}$ . Prove that  $m$  divides  $k$ .
- (c) Prove that  $m = k$ .

#### 3.5 5

Let  $R$  be a ring and  $M$  an  $R$ -module.

Recall that the set of torsion elements in  $M$  is defined by

$$\text{Tor}(M) = \{m \in M \mid \exists r \in R, r \neq 0, rm = 0\}.$$

- (a) Prove that if  $R$  is an integral domain, then  $\text{Tor}(M)$  is a submodule of  $M$ .

- (b) Give an example where  $\text{Tor}(M)$  is not a submodule of  $M$ .
- (c) If  $R$  has zero-divisors, prove that every non-zero  $R$ -module has non-zero torsion elements.

### 3.6 6

Let  $R$  be a commutative ring with multiplicative identity. Assume Zorn's Lemma.

- (a) Show that

$$N = \{r \in R \mid r^n = 0 \text{ for some } n > 0\}$$

is an ideal which is contained in any prime ideal.

- (b) Let  $r$  be an element of  $R$  not in  $N$ . Let  $S$  be the collection of all proper ideals of  $R$  not containing any positive power of  $r$ . Use Zorn's Lemma to prove that there is a prime ideal in  $S$ .
- (c) Suppose that  $R$  has exactly one prime ideal  $P$ . Prove that every element  $r$  of  $R$  is either nilpotent or a unit.

### 3.7 7

Let  $\zeta_n$  denote a primitive  $n$ th root of  $1 \in \mathbb{Q}$ . You may assume the roots of the minimal polynomial  $p_n(x)$  of  $\zeta_n$  are exactly the primitive  $n$ th roots of 1.

Show that the field extension  $\mathbb{Q}(\zeta_n)$  over  $\mathbb{Q}$  is Galois and prove its Galois group is  $(\mathbb{Z}/n\mathbb{Z})^\times$ .

How many subfields are there of  $\mathbb{Q}(\zeta_{20})$ ?

### 3.8 8

Let  $\{e_1, \dots, e_n\}$  be a basis of a real vector space  $V$  and let

$$\Lambda := \left\{ \sum r_i e_i \mid r_i \in \mathbb{Z} \right\}$$

Let  $\cdot$  be a non-degenerate ( $v \cdot w = 0$  for all  $w \in V \iff v = 0$ ) symmetric bilinear form on  $V$  such that the Gram matrix  $M = (e_i \cdot e_j)$  has integer entries.

Define the dual of  $\Lambda$  to be

$$\Lambda^\vee := \{v \in V \mid v \cdot x \in \mathbb{Z} \text{ for all } x \in \Lambda\}.$$

- (a) Show that  $\Lambda \subset \Lambda^\vee$ .
- (b) Prove that  $\det M \neq 0$  and that the rows of  $M^{-1}$  span  $\Lambda^\vee$ .
- (c) Prove that  $\det M = |\Lambda^\vee / \Lambda|$ .

---

## 4 2019 Course Exams

### 4.1 Midterm

1. Let  $G$  be a group of order  $p^2q$  for  $p, q$  prime. Show that  $G$  has a nontrivial normal subgroup.
2. Let  $G$  be a finite group and let  $P$  be a Sylow  $p$ -subgroup for  $p$  prime. Show that  $N(N(P)) = N(P)$  where  $N$  is the normalizer in  $G$ .
3. Show that there exist no simple groups of order 148.
4. Let  $p$  be a prime. Show that  $S_p = \langle \tau, \sigma \rangle$  where  $\tau$  is a transposition and  $\sigma$  is a  $p$ -cycle.
5. Let  $G$  be a nonabelian group of order  $p^3$  for  $p$  prime. Show that  $Z(G) = [G, G]$ .
6. Compute the Galois group of  $f(x) = x^3 - 3x - 3 \in \mathbb{Q}[x]/\mathbb{Q}$ .
7. Show that a field  $k$  of characteristic  $p \neq 0$  is perfect  $\iff$  for every  $x \in k$  there exists a  $y \in k$  such that  $y^p = x$ .
8. Let  $k$  be a field of characteristic  $p \neq 0$  and  $f \in k[x]$  irreducible. Show that  $f(x) = g(x^{p^d})$  where  $g(x) \in k[x]$  is irreducible and separable. Conclude that every root of  $f$  has the same multiplicity  $p^d$  in the splitting field of  $f$  over  $k$ .
9. Let  $n \geq 3$  and  $\zeta_n$  be a primitive  $n$ th root of unity. Show that  $[\mathbb{Q}(\zeta_n + \zeta_n^{-1}) : \mathbb{Q}] = \varphi(n)/2$  for  $\varphi$  the totient function.
10. Let  $L/K$  be a finite normal extension
  - Show that if  $L/K$  is cyclic and  $E/K$  is normal with  $L/E/K$  then  $L/E$  and  $E/K$  are cyclic.
  - Show that if  $L/K$  is cyclic then there exists exactly one extension  $E/K$  of degree  $n$  with  $L/E/K$  for each divisor  $n$  of  $[L : K]$ .

### 4.2 Final

1. Let  $A$  be an abelian group, and show  $A$  is a  $\mathbb{Z}$ -module in a unique way.
2. Consider the  $\mathbb{Z}$ -submodule  $N$  of  $\mathbb{Z}^3$  spanned by  $f_1 = [-1, 0, 1]$ ,  $f_2 = [2, -3, 1]$ ,  $f_3 = [0, 3, 1]$ ,  $f_4 = [3, 1, 5]$ . Find a basis for  $N$  and describe  $\mathbb{Z}^3/N$ .
3. Let  $R = k[x]$  for  $k$  a field and let  $M$  be the  $R$ -module given by

$$M = \frac{k[x]}{(x-1)^3} \oplus \frac{k[x]}{(x^2+1)^2} \oplus \frac{k[x]}{(x-1)(x^2+1)^4} \oplus \frac{k[x]}{(x+2)(x^2+1)^2}.$$

Describe the elementary divisors and invariant factors of  $M$ .

4. Let  $I = (2, x)$  be an ideal in  $R = \mathbb{Z}[x]$ , and show that  $I$  is not a direct sum of nontrivial cyclic  $R$ -modules.
5. Let  $R$  be a PID.
  - Classify irreducible  $R$ -modules up to isomorphism.
  - Classify indecomposable  $R$ -modules up to isomorphism.
6. Let  $V$  be a finite-dimensional  $k$ -vector space and  $T : V \rightarrow V$  a non-invertible  $k$ -linear map. Show that there exists a  $k$ -linear map  $S : V \rightarrow V$  with  $T \circ S = 0$  but  $S \circ T \neq 0$ .
7. Let  $A \in M_n(\mathbb{C})$  with  $A^2 = A$ . Show that  $A$  is similar to a diagonal matrix, and exhibit an explicit diagonal matrix similar to  $A$ .
8. Exhibit the rational canonical form for
  - $A \in M_6(\mathbb{Q})$  with minimal polynomial  $(x-1)(x^2+1)^2$ .
  - $A \in M_{10}(\mathbb{Q})$  with minimal polynomial  $(x^2+1)^2(x^3+1)$ .



---

9. Exhibit the rational and Jordan canonical forms for the following matrix  $A \in M_4(\mathbb{C})$ :

$$A = \begin{pmatrix} 2 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ -2 & -2 & 0 & 1 \\ -2 & 0 & -1 & -2 \end{pmatrix}.$$

10. Show that the eigenvalues of a Hermitian matrix  $A$  are real and that  $A = PDP^{-1}$  where  $P$  is an invertible matrix with orthogonal columns.

## 5 Spring 2018

### 5.1 1.

- (a) Use the Class Equation (equivalently, the conjugation action of a group on itself) to prove that any  $p$ -group (a group whose order is a positive power of a prime integer  $p$ ) has a nontrivial center.
- (b) Prove that any group of order  $p^2$  (where  $p$  is prime) is abelian.
- (c) Prove that any group of order  $5^2 \cdot 7^2$  is abelian.
- (d) Write down exactly one representative in each isomorphism class of groups of order  $5^2 \cdot 7^2$ .

### 5.2 2.

Let  $f(x) = x^4 - 4x^2 + 2 \in \mathbb{Q}[x]$ .

- (a) Find the splitting field  $K$  of  $f$ , and compute  $[K : \mathbb{Q}]$ .
- (b) Find the Galois group  $G$  of  $f$ , both as an explicit group of automorphisms, and as a familiar abstract group to which it is isomorphic.
- (c) Exhibit explicitly the correspondence between subgroups of  $G$  and intermediate fields between  $\mathbb{Q}$  and  $K$ .

### 5.3 3.

Let  $K$  be a Galois extension of  $\mathbb{Q}$  with Galois group  $G$ , and let  $E_1, E_2$  be intermediate fields of  $K$  which are the splitting fields of irreducible  $f_i(x) \in \mathbb{Q}[x]$ .

Let  $E = E_1E_2 \subset K$ .

Let  $H_i = \text{Gal}(K/E_i)$  and  $H = \text{Gal}(K/E)$ .

- (a) Show that  $H = H_1 \cap H_2$ .
- (b) Show that  $H_1H_2$  is a subgroup of  $G$ .
- (c) Show that

$$\text{Gal}(K/(E_1 \cap E_2)) = H_1H_2.$$

**5.4 4.**

Let

$$A = \begin{bmatrix} 0 & 1 & -2 \\ 1 & 1 & -3 \\ 1 & 2 & -4 \end{bmatrix} \in M_3(\mathbb{C})$$

- (a) Find the Jordan canonical form  $J$  of  $A$ .
- (b) Find an invertible matrix  $P$  such that  $P^{-1}AP = J$ .

You should not need to compute  $P^{-1}$ .

**5.5 5.**

Let

$$M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \quad \text{and} \quad N = \begin{pmatrix} x & u \\ -y & -v \end{pmatrix}$$

over a commutative ring  $R$ , where  $b$  and  $x$  are units of  $R$ . Prove that

$$MN = \begin{pmatrix} 0 & 0 \\ 0 & * \end{pmatrix} \implies MN = 0.$$

**5.6 6.**

Let

$$M = \{(w, x, y, z) \in \mathbb{Z}^4 \mid w + x + y + z \in 2\mathbb{Z}\},$$

and

$$N = \{(w, x, y, z) \in \mathbb{Z}^4 \mid 4 \mid (w - x), 4 \mid (x - y), 4 \mid (y - z)\}.$$

- (a) Show that  $N$  is a  $\mathbb{Z}$ -submodule of  $M$ .
- (b) Find vectors  $u_1, u_2, u_3, u_4 \in \mathbb{Z}^4$  and integers  $d_1, d_2, d_3, d_4$  such that

$$\{u_1, u_2, u_3, u_4\}$$

is a free basis for  $M$ , and

$$\{d_1u_1, d_2u_2, d_3u_3, d_4u_4\}$$

is a free basis for  $N$ .

- (c) Use the previous part to describe  $M/N$  as a direct sum of cyclic  $\mathbb{Z}$ -modules.

**5.7 7.**

Let  $R$  be a PID and  $M$  be an  $R$ -module. Let  $p$  be a prime element of  $R$ . The module  $M$  is called  $\langle p \rangle$ -primary if for every  $m \in M$  there exists  $k > 0$  such that  $p^k m = 0$ .

- (a) Suppose  $M$  is  $\langle p \rangle$ -primary. Show that if  $m \in M$  and  $t \in R$ ,  $t \notin \langle p \rangle$ , then there exists  $a \in R$  such that  $atm = m$ .
- (b) A submodule  $S$  of  $M$  is said to be *pure* if  $S \cap rM = rS$  for all  $r \in R$ . Show that if  $M$  is  $\langle p \rangle$ -primary, then  $S$  is pure if and only if  $S \cap p^k M = p^k S$  for all  $k \geq 0$ .

**5.8 8.**

Let  $R = C[0, 1]$  be the ring of continuous real-valued functions on the interval  $[0, 1]$ . Let  $I$  be an ideal of  $R$ .

- (a) Show that if  $f \in I$ ,  $a \in [0, 1]$  are such that  $f(a) \neq 0$ , then there exists  $g \in I$  such that  $g(x) \geq 0$  for all  $x \in [0, 1]$ , and  $g(x) > 0$  for all  $x$  in some open neighborhood of  $a$ .
- (b) If  $I \neq R$ , show that the set  $Z(I) = \{x \in [0, 1] \mid f(x) = 0 \text{ for all } f \in I\}$  is nonempty.
- (c) Show that if  $I$  is maximal, then there exists  $x_0 \in [0, 1]$  such that  $I = \{f \in R \mid f(x_0) = 0\}$ .

**6 Fall 2018****6.1 1.**

Let  $G$  be a finite group whose order is divisible by a prime number  $p$ . Let  $P$  be a normal  $p$ -subgroup of  $G$  (so  $|P| = p^c$  for some  $c$ ).

- (a) Show that  $P$  is contained in every Sylow  $p$ -subgroup of  $G$ .
- (b) Let  $M$  be a maximal proper subgroup of  $G$ . Show that either  $P \subseteq M$  or  $|G/M| = p^b$  for some  $b \leq c$ .

**6.2 2.**

- (a) Suppose the group  $G$  acts on the set  $X$ . Show that the stabilizers of elements in the same orbit are conjugate.
- (b) Let  $G$  be a finite group and let  $H$  be a proper subgroup. Show that the union of the conjugates of  $H$  is strictly smaller than  $G$ , i.e.

$$\bigcup_{g \in G} gHg^{-1} \subsetneq G$$

- (c) Suppose  $G$  is a finite group acting transitively on a set  $S$  with at least 2 elements. Show that there is an element of  $G$  with no fixed points in  $S$ .

**6.3 3.**

Let  $F \subset K \subset L$  be finite degree field extensions. For each of the following assertions, give a proof or a counterexample.

- (a) If  $L/F$  is Galois, then so is  $K/F$ .
- (b) If  $L/F$  is Galois, then so is  $L/K$ .
- (c) If  $K/F$  and  $L/K$  are both Galois, then so is  $L/F$ .

**6.4 4.**

Let  $V$  be a finite dimensional vector space over a field (the field is not necessarily algebraically closed).

Let  $\varphi : V \rightarrow V$  be a linear transformation. Prove that there exists a decomposition of  $V$  as  $V = U \oplus W$ , where  $U$  and  $W$  are  $\varphi$ -invariant subspaces of  $V$ ,  $\varphi|_U$  is nilpotent, and  $\varphi|_W$  is nonsingular.

**6.5 5.**

Let  $A$  be an  $n \times n$  matrix.

- (a) Suppose that  $v$  is a column vector such that the set  $\{v, Av, \dots, A^{n-1}v\}$  is linearly independent. Show that any matrix  $B$  that commutes with  $A$  is a polynomial in  $A$ .
- (b) Show that there exists a column vector  $v$  such that the set  $\{v, Av, \dots, A^{n-1}v\}$  is linearly independent  $\iff$  the characteristic polynomial of  $A$  equals the minimal polynomial of  $A$ .

**6.6 6.**

Let  $R$  be a commutative ring, and let  $M$  be an  $R$ -module. An  $R$ -submodule  $N$  of  $M$  is maximal if there is no  $R$ -module  $P$  with  $N \subsetneq P \subsetneq M$ .

- (a) Show that an  $R$ -submodule  $N$  of  $M$  is maximal  $\iff M/N$  is a simple  $R$ -module: i.e.,  $M/N$  is nonzero and has no proper, nonzero  $R$ -submodules.
- (b) Let  $M$  be a  $\mathbb{Z}$ -module. Show that a  $\mathbb{Z}$ -submodule  $N$  of  $M$  is maximal  $\iff \#M/N$  is a prime number.
- (c) Let  $M$  be the  $\mathbb{Z}$ -module of all roots of unity in  $\mathbb{C}$  under multiplication. Show that there is no maximal  $\mathbb{Z}$ -submodule of  $M$ .

**6.7 7.**

Let  $R$  be a commutative ring.

- (a) Let  $r \in R$ . Show that the map

$$\begin{aligned} r\bullet : R &\longrightarrow R \\ x &\mapsto rx. \end{aligned}$$

is an  $R$ -module endomorphism of  $R$ .

- (b) We say that  $r$  is a **zero-divisor** if  $\mathbf{r} \bullet$  is not injective. Show that if  $r$  is a zero-divisor and  $r \neq 0$ , then the kernel and image of  $R$  each consist of zero-divisors.
- (c) Let  $n \geq 2$  be an integer. Show: if  $R$  has exactly  $n$  zero-divisors, then  $\#R \leq n^2$ .
- (d) Show that up to isomorphism there are exactly two commutative rings  $R$  with precisely 2 zero-divisors.

You may use without proof the following fact: every ring of order 4 is isomorphic to exactly one of the following:

$$\frac{\mathbb{Z}}{4\mathbb{Z}}, \quad \frac{\frac{\mathbb{Z}}{2\mathbb{Z}}[t]}{(t^2 + t + 1)}, \quad \frac{\frac{\mathbb{Z}}{2\mathbb{Z}}[t]}{(t^2 - t)}, \quad \frac{\frac{\mathbb{Z}}{2\mathbb{Z}}[t]}{(t^2)}.$$

## 7 Fall 2017

### 7.1 1.

Suppose the group  $G$  acts on the set  $A$ . Assume this action is faithful (recall that this means that the kernel of the homomorphism from  $G$  to  $\text{Sym}(A)$  which gives the action is trivial) and transitive (for all  $a, b$  in  $A$ , there exists  $g$  in  $G$  such that  $g \cdot a = b$ .)

- (a) For  $a \in A$ , let  $G_a$  denote the stabilizer of  $a$  in  $G$ . Prove that for any  $a \in A$ ,

$$\bigcap_{\sigma \in G} \sigma G_a \sigma^{-1} = \{1\}.$$

- (b) Suppose that  $G$  is abelian. Prove that  $|G| = |A|$ . Deduce that every abelian transitive subgroup of  $S_n$  has order  $n$ .

### 7.2 2.

- (a) Classify the abelian groups of order 36.

For the rest of the problem, assume that  $G$  is a non-abelian group of order 36.

You may assume that the only subgroup of order 12 in  $S_4$  is  $A_4$  and that  $A_4$  has no subgroup of order 6.

- (b) Prove that if the 2-Sylow subgroup of  $G$  is normal,  $G$  has a normal subgroup  $N$  such that  $G/N$  is isomorphic to  $A_4$ .
- (c) Show that if  $G$  has a normal subgroup  $N$  such that  $G/N$  is isomorphic to  $A_4$  and a subgroup  $H$  isomorphic to  $A_4$  it must be the direct product of  $N$  and  $H$ .
- (d) Show that the dihedral group of order 36 is a non-abelian group of order 36 whose Sylow-2 subgroup is not normal.

**7.3 3.**

Let  $F$  be a field. Let  $f(x)$  be an irreducible polynomial in  $F[x]$  of degree  $n$  and let  $g(x)$  be any polynomial in  $F[x]$ . Let  $p(x)$  be an irreducible factor (of degree  $m$ ) of the polynomial  $f(g(x))$ .

Prove that  $n$  divides  $m$ . Use this to prove that if  $r$  is an integer which is not a perfect square, and  $n$  is a positive integer then every irreducible factor of  $x^{2n} - r$  over  $\mathbb{Q}[x]$  has even degree.

**7.4 4.**

- (a) Let  $f(x)$  be an irreducible polynomial of degree 4 in  $\mathbb{Q}[x]$  whose splitting field  $K$  over  $\mathbb{Q}$  has Galois group  $G = S_4$ .

Let  $\theta$  be a root of  $f(x)$ . Prove that  $\mathbb{Q}[\theta]$  is an extension of  $\mathbb{Q}$  of degree 4 and that there are no intermediate fields between  $\mathbb{Q}$  and  $\mathbb{Q}[\theta]$ .

- (b) Prove that if  $K$  is a Galois extension of  $\mathbb{Q}$  of degree 4, then there is an intermediate subfield between  $K$  and  $\mathbb{Q}$ .

**7.5 5.**

A ring  $R$  is called *simple* if its only two-sided ideals are  $0$  and  $R$ .

- (a) Suppose  $R$  is a commutative ring with  $1$ . Prove  $R$  is simple if and only if  $R$  is a field.  
(b) Let  $k$  be a field. Show the ring  $M_n(k)$ ,  $n \times n$  matrices with entries in  $k$ , is a simple ring.

**7.6 6.**

For a ring  $R$ , let  $U(R)$  denote the multiplicative group of units in  $R$ . Recall that in an integral domain  $R$ ,  $r \in R$  is called *irreducible* if  $r$  is not a unit in  $R$ , and the only divisors of  $r$  have the form  $ru$  with  $u$  a unit in  $R$ .

We call a non-zero, non-unit  $r \in R$  *prime* in  $R$  if  $r \mid ab \implies r \mid a$  or  $r \mid b$ . Consider the ring  $R = \{a + b\sqrt{-5} \mid a, b \in \mathbb{Z}\}$ .

- (a) Prove  $R$  is an integral domain.  
(b) Show  $U(R) = \{\pm 1\}$ .  
(c) Show  $3$ ,  $2 + \sqrt{-5}$ , and  $2 - \sqrt{-5}$  are irreducible in  $R$ .  
(d) Show  $3$  is not prime in  $R$ .  
(e) Conclude  $R$  is not a PID.

**7.7 7.**

Let  $F$  be a field and let  $V$  and  $W$  be vector spaces over  $F$ .

Make  $V$  and  $W$  into  $F[x]$ -modules via linear operators  $T$  on  $V$  and  $S$  on  $W$  by defining  $X \cdot v = T(v)$  for all  $v \in V$  and  $X \cdot w = S(w)$  for all  $w \in W$ .

Denote the resulting  $F[x]$ -modules by  $V_T$  and  $W_S$  respectively.

- (a) Show that an  $F[x]$ -module homomorphism from  $V_T$  to  $W_S$  consists of an  $F$ -linear transformation  $R : V \rightarrow W$  such that  $RT = SR$ .
- (b) Show that  $VT \cong WS$  as  $F[x]$ -modules  $\iff$  there is an  $F$ -linear isomorphism  $P : V \rightarrow W$  such that  $T = P^{-1}SP$ .
- (c) Recall that a module  $M$  is *simple* if  $M \neq 0$  and any proper submodule of  $M$  must be zero. Suppose that  $V$  has dimension 2. Give an example of  $F, T$  with  $V_T$  simple.
- (d) Assume  $F$  is algebraically closed. Prove that if  $V$  has dimension 2, then any  $V_T$  is not simple.