

Topology Qualifying Exam Solutions

D. Zack Garza

Tuesday 26th May, 2020

Contents

1	Definitions	1
2	Theorems	2
3	General Topology	2
3.1	2	2
3.1.1	Alternate Proof of (ii)	2
3.2	4	3
3.3	6	4
3.3.1	Proof 1 (DZG)	4
3.4	8	4
3.4.1	Proof 1 (DZG)	4
4	10	6
5	12	6
6	14	7
7	16	7

1 Definitions

- Closed (several characterizations)
- Bounded
- Compact
- Connectedness: There does not exist a disconnecting set $X = A \amalg B$ such that $\emptyset \neq A, B \subsetneq X$, i.e. X is the union of two proper disjoint nonempty sets. Equivalently, X contains no proper nonempty clopen sets.
- Subspace topology
- Retract: A subspace $A \subset X$ is a *retract* of X iff there exists a continuous map $f : X \rightarrow A$ such that $f|_A = \text{id}_A$. Equivalently it is a *left inverse* to the inclusion.

2 Theorems

- Closed subsets of Hausdorff spaces are compact? (check)
- Cantor's intersection theorem?
- Tube lemma
- Properties pushed forward through continuous maps:
 - Compactness?
 - Connectedness (when surjective)
 - Separability
 - Density **only when** f is surjective
 - **Not** openness
 - **Not** closedness
- Results that only work for metric spaces
 - ?
- A retract of a Hausdorff/connected/compact space is closed/connected/compact respectively.

3 General Topology

3.1 2

Statement: state the definition of compactness, determine if the sets $\{0\} \cup \left\{\frac{1}{n}\right\}, (0, 1]$ are compact.

- i. A topological space (X, τ) is **compact** if every open cover has a *finite* subcover. That is, if $\{U_j \mid j \in J\} \subset \tau$ is a collection of open sets such that $X \subseteq \bigcup_{j \in J} U_j$, then there exists a *finite* subset $J' \subset J$ such that $X \subseteq \bigcup_{j \in J'} U_j$.

- ii. Use Heine-Borel theorem: a set $U \subset \mathbb{R}^n$ is compact $\iff U$ is *closed* and *bounded*.

- X is closed in \mathbb{R} , since we can write its complement as an arbitrary union of open intervals:

$$X^c = (-\infty, 0) \cup \left(\bigcup_{n \in \mathbb{Z}^+} \left(\frac{1}{n}, \frac{1}{n+1} \right) \right) \cup (1, \infty)$$

- X is *bounded*, since we can pick $r = 1$, then $x, y \in X \implies d(x, y) \leq r = 1$.

- iii. Use Heine-Borel again: X is not closed because it does not contain all of its limit points, e.g. the sequence $\left\{x_n := \frac{1}{n} \mid n \in \mathbb{Z}^{\geq 1}\right\} \subset X$ but $x_n \xrightarrow{n \rightarrow \infty} 0 \in X^c$. Thus X is **not** compact.

3.1.1 Alternate Proof of (ii)

See Munkres p.164

- Let $\{U_i \mid i \in J\} \Rightarrow X$; then $0 \in U_j$ for some $j \in J$.
- In the subspace topology, U_i is given by some $V \in \tau(\mathbb{R})$ such that $V \cap X = U_i$


- A basis for the subspace topology on \mathbb{R} is open intervals, so write V as a union of open intervals $V = \bigcup_{k \in K} I_k$.
- Since $0 \in U_j$, $0 \in I_k$ for some k .
- Since I_k is an interval, it contains infinitely many points of the form $x_n = \frac{1}{n} \in X$
- Then $I_k \cap X \subset U_j$ contains infinitely many such points.
- So there are only *finitely* many points in $X \setminus U_j$, each of which is in $U_{j(n)}$ for some $j(n) \in J$ depending on n .
- So U_j and the *finitely* many $U_{j(n)}$ form a finite subcover of X . ■

3.2 4

Statement: show that the *Lebesgue number* is well-defined for compact metric spaces.

Note: this is a question about the *Lebesgue Number*. See Wikipedia for detailed proof.

- Write $U = \{U_i \mid i \in I\}$, then $X \subseteq \bigcup_{i \in I} U_i$. Need to construct a $\delta > 0$.
- By compactness of X , choose a finite subcover U_1, \dots, U_n .
- Define the distance between a point x and a set $Y \subset X$: $d(x, Y) = \inf_{y \in Y} d(x, y)$.
 - **Claim:** the function $d(\cdot, Y) : X \rightarrow \mathbb{R}$ is continuous for a fixed set.
 - Proof: Todo, not obvious.



- Define a function

$$f : X \longrightarrow \mathbb{R}$$

$$x \mapsto \frac{1}{n} \sum_{i=1}^n d(x, X \setminus U_i).$$

- Note this is a sum of continuous functions and thus continuous.

- **Claim:**

$$\delta := \inf_{x \in X} f(x) = \min_{x \in X} f(x) = f(x_{\min}) > 0$$

suffices.

- That the infimum is a minimum: f is a continuous function on a compact set, apply the extreme value theorem: it attains its minimum.
- That $\delta > 0$: otherwise, $\delta = 0 \implies \exists x_0$ such that $d(x_0, X \setminus U_i) = 0$ for all i .
 - * Forces $x_0 \in X \setminus U_i$ for all i , but $X \setminus \bigcup U_i = \emptyset$ since the U_i cover X .
- That it satisfies the Lebesgue condition:

$$\forall x \in X, \exists i \text{ such that } B_\delta(x) \subset U_i$$

- * Let $B_\delta(x) \ni x$; then by minimality $f(x) \geq \delta$.
- * Thus it can *not* be the case that $d(x, X \setminus U_i) < \delta$ for *every* i , otherwise

$$f(x) \leq \frac{1}{n}(\delta + \dots + \delta) = \frac{n\delta}{n} = \delta$$

- * So there is some particular i such that $d(x, X \setminus U_i) \geq \delta$.
- * But then $B_\delta \subseteq U_i$ as desired.

3.3 6

Statement: prove that $[0, 1] \subset \mathbb{R}$ is compact.

3.3.1 Proof 1 (DZG)

Todo: find a direct proof.

3.4 8

Topic: proof of the tube lemma.

Statement: show $X, Y \in \text{Top}_{\text{compact}} \iff X \times Y \in \text{Top}_{\text{compact}}$

3.4.1 Proof 1 (DZG)

\Leftarrow :

- By universal properties, the product $X \times Y$ is equipped with continuous projections
- The continuous image of a compact set is compact, and $\pi_1(X \times Y) = X, \pi_2(X \times Y) = Y$
- So X, Y are compact.

\Rightarrow :

Proof of Tube Lemma:

- Let $\{U_j \times V_j \mid j \in J\} \Rightarrow X \times Y$.
- Fix a point $x_0 \in X$, then $\{x_0\} \times Y \subset N$ for some open set N .
- By the tube lemma, there is a $U^x \subset X$ such that the tube $U^x \times Y \subset N$.
- Since $\{x_0\} \times Y \cong Y$ which is compact, there is a finite subcover $\{U_j \times V_j \mid j \leq n\} \Rightarrow \{x_0\} \times Y$.
- “Integrate the X ”: write

$$W = \bigcap_{j=1}^n U_j,$$

then $x_0 \in W$ and W is a finite intersection of open sets and thus open.

- Claim: $\{U_j \times V_j \mid j \leq n\} \Rightarrow W \times Y$
 - Let $(x, y) \in W \times Y$; want to show $(x, y) \in U_j \times V_j$ for some $j \leq n$.
 - Then $(x_0, y) \in \{x_0\} \times Y$ is on the same horizontal line
 - $(x_0, y) \in U_j \times V_j$ for some j by construction
 - So $y \in V_j$ for this j
 - Since $x \in W$, $x \in U_j$ for *every* j , thus $x \in U_j$.
 - So $(x, y) \in U_j \times V_j$

Actual Proof:

- Let $\{U_j \mid j \in J\} \Rightarrow X \times Y$.
- Fix $x_0 \in X$, the slice $\{x_0\} \times Y$ is compact and can be covered by finitely many elements $\{U_j \mid j \leq m\} \Rightarrow \{x_0\} \times Y$.
 - Sum: write $N = \bigcup_{j=1}^m U_j$; then $\{x_0\} \times Y \subset N$.
 - Apply the tube lemma to N : produce $\{x_0\} \times Y \in W \times Y \subset N$; then $\{U_j \mid j \leq m\} \Rightarrow W \times Y$.
- Now let $x \in X$ vary: for each $x \in X$, produce $W_x \times Y$ as above, then $\{W_x \times Y \mid x \in X\} \Rightarrow X$.
 - By above argument, every tube $W_x \times Y$ can be covered by *finitely* many U_j .
- Since $\{W_x \mid x \in X\} \Rightarrow X$ and X is compact, produce a finite subset $\{W_k \mid k \leq m'\} \Rightarrow X$.
- Then $\{W_k \times Y \mid k \leq m'\} \Rightarrow X \times Y$; the claim is that it is a finite cover.
 - Finitely many k
 - For each k , the tube $W_k \times Y$ is covered by finitely by U_j
 - And finite \times finite = finite. ■

Shorter mnemonic:

19.U It is sufficient to consider a cover consisting of elementary sets. Since Y is compact, each fiber $x \times Y$ has a finite subcovering $\{U_i^x \times V_i^x\}$. Put $W^x = \cap U_i^x$. Since X is compact, the cover $\{W^x\}_{x \in X}$ has a finite subcovering W^{x_j} . Then $\{U_i^{x_j} \times V_i^{x_j}\}$ is the required finite subcovering.

4 10

X is connected:

- Write $X = L \amalg G$ where $L = \{0\} \times [-1, 1]$ and $G = \{\Gamma(\sin(x)) \mid x \in (0, 1]\}$ is the graph of $\sin(x)$.
- $L \cong [0, 1]$ which is connected
 - Claim: Every interval is connected (todo)
- Claim: G is connected
 - The function

$$\begin{aligned} f : (0, 1] &\longrightarrow [-1, 1] \\ x &\mapsto \sin(x) \end{aligned}$$

is continuous (how to prove?)

- Claim: The diagonal map $\Delta : Y \longrightarrow Y \times Y$ where $\Delta(t) = (t, t)$ is continuous for any Y since $\Delta = (\text{id}, \text{id})$
- The composition of continuous function is continuous
- So the composition is continuous:

$$\begin{aligned} F : (0, 1] &\xrightarrow{\Delta} (0, 1]^2 \xrightarrow{(\text{id}, f)} (0, 1] \times [-1, 1] \\ t &\mapsto (t, t) \mapsto (t, f(t)) \end{aligned}$$

- Then $G = F((0, 1])$ is the continuous image of a connected set and thus connected.
- Claim: X is connected
 - Suppose there is a disconnecting cover $X = A \amalg B$ such that $\bar{A} \cap B = A \cap \bar{B} = \emptyset$ and $A, B \neq \emptyset$.
 - WLOG suppose $(x, \sin(x)) \in B$ for $x > 0$.
 - Claim: $B = G$
 - * It can't be the case that A intersects G : otherwise $X = A \amalg B \implies G = (A \cap G) \amalg (B \cap G)$ disconnects G . So $A \cap G = \emptyset$, forcing $A \subseteq L$
 - * Similarly L can not be disconnected, so $B \cap L = \emptyset$ forcing $B \subset G$
 - * So $A \subset L$ and $B \subset G$, and since $X = A \amalg B$, this forces $A = L$ and $B = G$.
 - But any open set U in the subspace topology $L \subset \mathbb{R}^2$ (generated by open balls) containing $(0, 0) \in L$ is the restriction of a ball $V \subset \mathbb{R}^2$ of positive radius $r > 0$, i.e. $U = V \cap X$.
 - * But any such ball contains points of G : namely take n large enough such that $\frac{1}{n\pi} < r$.
 - * So $U \cap L \cap G \neq \emptyset$, contradicting $L \cap G = \emptyset$.

5 12

- Using the fact that $[0, \infty) \subset \mathbb{R}$ is Hausdorff, any retract must be closed, so any closed interval $[\varepsilon, N]$ for $0 \leq \varepsilon \leq N \leq \infty$.
 - Note that $\varepsilon = N$ yields all one point sets $\{x_0\}$ for $x_0 \geq 0$.
- No finite discrete sets occur, since the retract of a connected set is connected.
- ?

6 14

- Take two connected sets X, Y ; then there exists $p \in X \cap Y$.
- Write $X \cup Y = A \amalg B$ with both $A, B \subset A \amalg B$ open.
- Since $p \in X \cup Y = A \amalg B$, WLOG $p \in A$. We will show B must be empty.
- Claim: $A \cap X$ is clopen in X .
 - $A \cap X$ is open in X : ?
 - $A \cap X$ is closed in X : ?
- The only clopen sets of a connected set are empty or the entire thing, and since $p \in A$, we must have $A \cap X = X$.
- By the same argument, $A \cap Y = Y$.
- So $A \cap (X \cup Y) = (A \cap X) \cup (A \cap Y) = X \cup Y$
- Since $A \subset X \cup Y$, $A \cap (X \cup Y) = A$
- Thus $A = X \cup Y$, forcing $B = \emptyset$.

7 16

- $S \subset X$ is **not** connected iff there exist two sets $A, B \subset X$ such that
 - $A, B \neq \emptyset$
 - $A \cap B = \emptyset$
 - $S = A \amalg B$
 - $\bar{A} \cap B = A \cap \bar{B} = \emptyset$.