

# Algebra Qualifying Exam Notes

D. Zack Garza

Friday 12<sup>th</sup> June, 2020

## Contents

<b>1</b>	<b>Study Guide for Algebra Qualifying Exam</b>	<b>3</b>
1.1	Group Theory . . . . .	4
1.2	Linear Algebra . . . . .	4
1.3	Rings and Modules . . . . .	4
1.4	Field Theory . . . . .	5
<b>2</b>	<b>Remarks</b>	<b>5</b>
2.1	Group theory: . . . . .	5
2.2	Rings: . . . . .	6
2.3	Field Theory / Galois Theory. . . . .	7
<b>3</b>	<b>Group Theory</b>	<b>7</b>
3.1	Random References . . . . .	7
3.2	Big List of Notation . . . . .	7
3.3	Basics . . . . .	7
3.4	Finitely Generated Abelian Groups . . . . .	9
3.5	The Symmetric Group . . . . .	10
3.6	Counting Theorems . . . . .	11
3.6.1	Examples of Orbit-Stabilizer . . . . .	11
3.6.2	Sylow Theorems . . . . .	12
3.6.3	Sylow 1 (Cauchy for Prime Powers) . . . . .	12
3.6.4	Sylow 2 (Sylows are Conjugate) . . . . .	13
3.6.5	Sylow 3 (Numerical Constraints) . . . . .	13
3.7	Products . . . . .	13
3.8	Isomorphism Theorems . . . . .	14
3.9	Special Classes of Groups . . . . .	15
3.10	Series of Groups . . . . .	16
3.11	Classification of Groups . . . . .	17
<b>4</b>	<b>Rings</b>	<b>17</b>
4.1	Definitions . . . . .	17
4.2	Nontrivial Properties . . . . .	19
4.3	Ideals . . . . .	19
4.3.1	Maximal and Prime Ideals . . . . .	19
4.3.2	Nilradical and Jacobson Radical . . . . .	19

4.3.3	Zorn's Lemma . . . . .	20
<b>5</b>	<b>Fields</b>	<b>20</b>
5.1	Finite Fields . . . . .	21
5.2	Galois Theory . . . . .	23
5.2.1	Lemmas About Towers . . . . .	23
5.2.2	Examples . . . . .	25
5.3	Cyclotomic Polynomials . . . . .	26
<b>6</b>	<b>Modules</b>	<b>27</b>
6.1	General Modules . . . . .	27
6.2	Classification of Modules over a PID . . . . .	28
6.3	Minimal / Characteristic Polynomials . . . . .	28
6.4	Canonical Forms . . . . .	29
6.4.1	Rational Canonical Form . . . . .	29
6.4.2	Jordan Canonical Form . . . . .	29
6.5	Using Canonical Forms . . . . .	30
6.6	Diagonalizability . . . . .	30
6.7	Matrix Counterexamples . . . . .	31
6.8	Miscellaneous . . . . .	32
<b>7</b>	<b>Extra Problems</b>	<b>33</b>
7.1	Group Theory . . . . .	33
7.1.1	Basic Structure . . . . .	33
7.1.2	Primes in Group Theory . . . . .	33
7.1.3	Classification . . . . .	33
7.1.4	Series . . . . .	33
7.2	Ring Theory . . . . .	34
7.3	Field Theory . . . . .	34
7.4	Modules and Linear Algebra . . . . .	35
7.5	Commutative Algebra . . . . .	35
<b>8</b>	<b>List of Topics</b>	<b>35</b>
<b>9</b>	<b>Groups</b>	<b>37</b>
9.1	Definitions . . . . .	37
9.1.1	<b>Subgroup Generated</b> by a set $A$ . . . . .	37
9.1.2	<b>Free Group</b> on a set $X$ . . . . .	37
9.1.3	<b>Centralizer</b> of an element or a subgroup . . . . .	37
9.1.4	<b>Center</b> of a group . . . . .	37
9.1.5	<b>Normalizer</b> of a subgroup . . . . .	38
9.1.6	<b>Normal Core</b> of a subgroup . . . . .	38
9.1.7	<b>Normal Closure</b> of a subgroup . . . . .	38
9.1.8	<b>Group Action</b> of a group on a set . . . . .	39
9.1.9	Transitive group actions . . . . .	40
9.1.10	Orbit of an Element of a Set . . . . .	40
9.1.11	Stabilizer of an Element of a Set . . . . .	40
9.1.12	<b>Automorphisms</b> of a group . . . . .	40

---

9.1.13	<b>Inner Automorphisms</b> of a group . . . . .	40
9.1.14	<b>Outer Automorphisms</b> of a group . . . . .	40
9.1.15	<b>Conjugacy Class</b> of an element . . . . .	41
9.1.16	<b>Characteristic</b> subgroup . . . . .	41
9.1.17	<b>Simple</b> group . . . . .	41
9.1.18	<b>Commutator</b> of an element, or of subgroups . . . . .	41
9.2	Structural Results . . . . .	41
9.2.1	Isomorphisms Theorems . . . . .	41
9.3	Misc Results . . . . .	42
9.4	Numeric Results . . . . .	43
9.4.1	Cauchy's Theorem . . . . .	43
9.4.2	Sylow Theorems: $ G  = p^k m$ where $p \nmid m$ . . . . .	43
9.4.3	Orbit-stabilizer Theorem . . . . .	43
9.4.4	Burnside's Lemma . . . . .	43
9.4.5	The class equation . . . . .	44
9.4.6	General facts . . . . .	44
9.5	Common Groups . . . . .	45
9.5.1	$S_3$ . . . . .	45
9.5.2	$S_n$ . . . . .	45
9.5.3	$A_n$ . . . . .	45
9.5.4	$D_n$ . . . . .	45
<b>10</b>	<b>Rings</b> . . . . .	<b>46</b>
10.1	Facts about ideals: . . . . .	46
10.2	Maximal ideals . . . . .	46
10.3	Prime ideals . . . . .	46
10.4	Radicals . . . . .	46
10.5	Other ideals . . . . .	46
10.6	Orders less than 16: . . . . .	47

## 1 Study Guide for Algebra Qualifying Exam

### References:

- [1]. David Dummit and Richard Foote, Abstract Algebra, Wiley, 2003.
- [2]. Kenneth Hoffman and Ray Kunze, Linear Algebra, Prentice-Hall, 1971.
- [3]. Thomas W. Hungerford, Algebra, Springer, 1974.
- [4]. Roy Smith, Algebra Course Notes (843-1 through 845-3), <http://www.math.uga.edu/~roy/>,

As a general rule, students are responsible for knowing both the theory (proofs) and practical applications (e.g. **how to find the Jordan or rational canonical form** of a given matrix, **or the Galois group of a given polynomial**) of the topics mentioned.  
 A supplement to this study guide is available at:  
<http://www.math.uga.edu/sites/default/files/PDFs/Graduate/QualsStudyGuides/AlgebraPhDqualremarks.pdf>

### 1.1 Group Theory

- Subgroups and quotient groups
- Lagrange's Theorem
- Fundamental homomorphism theorems
- Group actions with applications to the structure of groups such as
  - The Sylow Theorems
- Group constructions such as:
  - Direct and semi-direct products
- Structures of special types of groups such as:
  - p-groups
  - Dihedral,
  - Symmetric and Alternating groups
    - \* Cycle decompositions
- The simplicity of  $A_n$ , for  $n \geq 5$
- Free groups, generators and relations
- Solvable groups

References: [1,3,4]

### 1.2 Linear Algebra

- Determinants
- Eigenvalues and eigenvectors
- Cayley-Hamilton Theorem
- Canonical forms for matrices
- Linear groups ( $GL_n, SL_n, O_n, U_n$ )
- Duality
  - Dual spaces,
  - Dual bases,
  - Induced dual map,
  - Double duals
- Finite-dimensional spectral theorem

References: [1,2,4]

### 1.3 Rings and Modules

- Zorn's Lemma

- Every vector space has a basis
- Maximal ideals exist
- Properties of ideals and quotient rings
- Fundamental homomorphism theorems for rings and modules
- Characterizations and properties of special domains such as:
  - Euclidean  $\implies$  PID  $\implies$  UFD
- Classification of finitely generated modules over PIDs (*with emphasis on Euclidean Domains*)
- Applications to the structure of:
  - Finitely generated abelian groups
  - Canonical forms of matrices

References: [1,3,4]

## 1.4 Field Theory

- Algebraic extensions of fields
- Fundamental theorem of Galois theory
- Properties of finite fields
- Separable extensions
- Computations of Galois groups of polynomials of small degree and cyclotomic
- Polynomials
- Solvability of polynomials by radicals

References: [1,3,4]

## 2 Remarks

Adapted from remark written by Roy Smith, August 2006

### 2.1 Group theory:

The first 6 chapters (220 pages) of DF are excellent.

All the definitions and proofs of these theorems on groups are given in Smith's web based lecture notes for math 843 part 1.

#### Key topics:

- Sylow theorems
- Simplicity of  $A_n$  for  $n > 4$ .
- The first isomorphism theorem,

## 2.2 Rings:

---

- The Jordan Holder theorem,

The last two (one easy, one hard) are left as exercises.

**The proof JH is seldom tested on the qual**, but proofs are always of interest.

- Fundamental theorem of finite abelian groups

*DF Exercises 12.1.16-19*

- The simple groups of order between 60 and 168 have prime order

## 2.2 Rings:

- DF Chapters 7,8,9.
- Gauss's important theorem on unique factorization of polynomials:
  - $\mathbb{Z}[x]$  is a UFD
  - $R[x]$  is a UFD when  $R$  is a UFD
- The fundamental isomorphism theorems for rings (easy and useful exercise)
- How to use Zorn's lemma
  - To find maximal ideals
  - Construct algebraic field closures
  - Why it is unnecessary in countable or noetherian rings.

Smith discusses extensively in 844-1.

- Results about PIDs  
(DF Section 8.2)
  - Example of a PID that is not a Euclidean domain  
(*DF p.277*)
  - Proof that a Euclidean domain is a PID and hence a UFD
  - Proof that  $\mathbb{Z}$  and  $k[x]$  are UFDs  
(*p.289 Smith, p.300 DF*)
- A polynomial ring in infinitely many variables over a UFD is still a ufd  
(*Easy, DF, p.305*)
- Eisenstein's criterion  
(*DF p.309*)
  - Stated only for monic polynomials – proof of general case identical.
  - See Smith's notes for the full version.
- Cyclic product structure of  $(\mathbb{Z}/n\mathbb{Z})^\times$   
(*exercise in DF, Smith 844-2, section 18*)

- Grobner bases and division algorithms for polynomials in several variables  
(DF 9.6.)

- Modules over pid's and Canonical forms of matrices.

DF sections 10.1, 10.2, 10.3, and 12.1, 12.2, 12.3.

- Constructive proof of decomposition: DF Exercises 12.1.16-19
- Smith 845-1 and 845-2: Detailed discussion of the constructive proof.

## 2.3 Field Theory / Galois Theory.

- DF chapters 13,14 (about 145 pages).
- Smith:
  - 843-2, sections 11,12, and 16-21 (39 pages)
  - 844-1, sections 7-9 (20 pages)
  - 844-2, sections 10-16, (37 pages)

## 3 Group Theory

### 3.1 Random References

### 3.2 Big List of Notation

$C_G(x) =$	$\{g \in G \mid gxg^{-1} = x\}$	$\subseteq G$	Centralizer
$? =$	$\{ghg^{-1} \mid g \in G\}$	$\subseteq G$	Conjugacy Class
$\mathcal{O}_x, G \cdot x =$	$\{g.x \mid x \in X\}$	$\subseteq X$	Orbit
$\text{Stab}_G(x), G_x =$	$\{g \in G \mid g.x = x\}$	$\subseteq G$	Stabilizer
$X^g =$	$\{x \in X \mid \forall g \in G, g.x = x\}$	$\subseteq X$	Fixed Points
$Z(G) =$	$\{x \in G \mid \forall g \in G, gxg^{-1} = x\}$	$\subseteq G$	Center
$N_G(H) =$	$\{g \in G \mid gHg^{-1} = H\}$	$\subseteq G$	Normalizer
$\text{Inn}(G) =$	$\{\varphi_g(x) = gxg^{-1}\}$	$\subseteq \text{Aut}(G)$	Inner Aut.
$\text{Out}(G) =$	$\text{Aut}(G)/\text{Inn}(G)$	$\hookrightarrow \text{Aut}(G)$	Outer Aut.

### 3.3 Basics

**Definition (Centralizer):**

$$C_G(H) = \{g \in G \mid ghg^{-1} = h \forall h \in H\}$$

**Definition (Normalizer):**

$$N_G(H) = \{g \in G \mid gHg^{-1} = H\}$$

**Lemma:**  $C_G(H) \trianglelefteq N_G(H)$

**Lemma:** The size of the conjugacy class of  $H$  is the index of its centralizer, i.e.

$$\left| \{gHg^{-1} \mid g \in G\} \right| = [G : C_G(H)].$$

Proof: Orbit-stabilizer.

**Lemma (“The Fundamental Theorem of Cosets”):**

$$aH = bH \iff a^{-1}b \in H \text{ or } aH \cap bH = \emptyset$$

**Definition:**  $[x, y] = x^{-1}y^{-1}xy$  is the **commutator**, and  $[G, G] := \{[x, y] \mid x, y \in G\}$  is the **commutator subgroup**.

**Lemma:**

$$[G, G] \leq H \text{ and } H \trianglelefteq G \implies G/H \text{ is abelian.}$$

**Lemmas:**

- Every subgroup of a cyclic group is itself cyclic.
- Intersections of subgroups are still subgroups
  - Intersections of distinct coprime-order subgroups are trivial
  - Intersections of subgroups of the same prime order are either trivial or equality
- The Quaternion group has only one element of order 2, namely  $-1$ .
  - They also have the presentation

$$\begin{aligned} Q &= \langle x, y, z \mid x^2 = y^2 = z^2 = xyz = -1 \rangle \\ &= \langle x, y \mid x^4 = y^4 = e, x^2 = y^2, yxy^{-1} = x^{-1} \rangle. \end{aligned}$$

- A dihedral group always has a presentation of the form

$$D_n = \langle x, y \mid x^n = y^2 = (xy)^2 = e \rangle,$$

yielding at least 2 distinct elements of order 2.



### 3.4 Finitely Generated Abelian Groups

Invariant factor decomposition:

$$G \cong \mathbb{Z}^r \times \prod_{j=1}^m \mathbb{Z}/(n_j) \quad \text{where } n_1 \mid \cdots \mid n_m.$$

**Going from invariant divisors to elementary divisors:**

- Take prime factorization of each factor
- Split into coprime pieces

*Example:*

$$\begin{aligned} & \mathbb{Z}/(2) \oplus \mathbb{Z}/(2) \oplus \mathbb{Z}/(2^3 \cdot 5^2 \cdot 7) \\ & \cong \mathbb{Z}/(2) \oplus \mathbb{Z}/(2) \oplus \mathbb{Z}/(2^3) \oplus \mathbb{Z}/(5^2) \oplus \mathbb{Z}/(7) \end{aligned}$$

**Going from elementary divisors to invariant factors:**

- Bin up by primes occurring (keeping exponents)
- Take highest power from each prime as *last* invariant factor
- Take highest power from all remaining primes as next, etc

*Example:* Given the invariant factor decomposition

$$G = \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_3 \times \mathbb{Z}_{25},$$

$p = 2$	$p = 3$	$p = 5$
$2, 2, 2$	$3, 3$	$5^2$

$$\implies n_m = 5^2 \cdot 3 \cdot 2$$

$p = 2$	$p = 3$	$p = 5$
$2, 2$	$3$	$\emptyset$

$$\implies n_{m-1} = 3 \cdot 2$$

$p = 2$	$p = 3$	$p = 5$
$2$	$\emptyset$	$\emptyset$

$$\implies n_{m-2} = 2$$

and thus

$$G \cong \mathbb{Z}/(2) \oplus \mathbb{Z}/(3 \cdot 2) \oplus \mathbb{Z}/(5^2 \cdot 3 \cdot 2).$$

### Classifying Abelian Groups of a Given Order:

Let  $p(x)$  be the integer partition function.

Example:  $p(6) = 11$ , given by  $6, 5 + 1, 4 + 2, \dots$ .

Write  $G = p_1^{k_1} p_2^{k_2} \dots$ ; then there are  $p(k_1)p(k_2) \dots$  choices, each yielding a distinct group.

## 3.5 The Symmetric Group

### Definitions:

- A cycle is **even**  $\iff$  product of an *even* number of transpositions.
  - A cycle of even *length* is **odd**
  - A cycle of odd *length* is **even**

**Definition** The **alternating group** is the subgroup of **even** permutations, i.e.  $A_n := \{\sigma \in S_n \mid \text{sign}(\sigma) = 1\}$  where  $\text{sign}(\sigma) = (-1)^m$  where  $m$  is the number of cycles of even length.

*Corollary:* Every  $\sigma \in A_n$  has an even number of *odd* cycles (i.e. an even number of *even-length* cycles).

*Example:*

$$\begin{aligned} A_4 = \{ & \text{id}, \\ & (1, 3)(2, 4), (1, 2)(3, 4), (1, 4)(2, 3), \\ & (1, 2, 3), (1, 3, 2), \\ & (1, 2, 4), (1, 4, 2), \\ & (1, 3, 4), (1, 4, 3), \\ & (2, 3, 4), (2, 4, 3) \}. \end{aligned}$$

### Lemmas:

- The transitive subgroups of  $S_3$  are  $S_3, A_3$
- The transitive subgroups of  $S_4$  are  $S_4, A_4, D_4, \mathbb{Z}_2^2, \mathbb{Z}_4$ .
- $S_4$  has two normal subgroups:  $A_4, \mathbb{Z}_2^2$ .
- $S_{n \geq 5}$  has one normal subgroup:  $A_n$ .
- $Z(S_n) = 1$  for  $n \geq 3$
- $Z(A_n) = 1$  for  $n \geq 4$
- $[S_n, S_n] = A_n$
- $[A_4, A_4] \cong \mathbb{Z}_2^2$
- $[A_n, A_n] = A_n$  for  $n \geq 5$ , so  $A_{n \geq 5}$  is nonabelian.
- $A_{n \geq 5}$  is *simple*.

### 3.6 Counting Theorems

**Lagrange's Theorem:**

$$H \leq G \implies |H| \mid |G|.$$

*Corollary:* The order of every element divides the size of  $G$ , i.e.

$$g \in G \implies o(g) \mid o(G) \implies g^{|G|} = e.$$

**Warning:** There does **not** necessarily exist  $H \leq G$  with  $|H| = n$  for every  $n \mid |G|$ .  
Counterexample:  $|A_4| = 12$  but has no subgroup of order 6.

**Cauchy's Theorem:**

For every prime  $p$  dividing  $|G|$ , there is an element (and thus a subgroup) of order  $p$ .

This is a partial converse to Lagrange's theorem, and strengthened by Sylow's theorem.

**Notation:** For a group  $G$  acting on a set  $X$ ,

- $G \cdot x = \{g \curvearrowright x \mid g \in G\} \subseteq X$  is the orbit
- $G_x = \{g \in G \mid g \curvearrowright x = x\} \subseteq G$  is the stabilizer
- $X/G \subset \mathcal{P}(X)$  is the set of orbits
- $X^g = \{x \in X \mid g \curvearrowright x = x\} \subseteq X$  are the fixed points

**Orbit-Stabilizer:**

$$|G \cdot x| = [G : G_x] = |G|/|G_x| \quad \text{if } G \text{ is finite}$$

Mnemonic:  $G/G_x \cong G \cdot x$ .

#### 3.6.1 Examples of Orbit-Stabilizer

1. Let  $G$  act on itself by conjugation.

- $G \cdot x$  is the **conjugacy class** of  $x$
- $G_x = Z(x) := C_G(x) = \{g \mid [g, x] = e\}$ , the **centralizer** of  $x$ .
- $G^g$  (the fixed points) is the **center**  $Z(G)$ .

*Corollary:* The number of conjugates of an element (i.e. the size of its conjugacy class) is the index of its centralizer,  $[G : C_G(x)]$ .

*Corollary:* the **Class Equation**:

$$|G| = |Z(G)| + \sum_{\substack{\text{One } x_i \text{ from} \\ \text{each conjugacy} \\ \text{class}}} [G : Z(x_i)]$$

1. Let  $G$  act on  $S$ , its set of *subgroups*, by conjugation.
  - $G \cdot H = \{gHg^{-1}\}$  is the **set of conjugate subgroups** of  $H$
  - $G_H = N_G(H)$  is the **normalizer** of in  $G$  of  $H$
  - $S^G$  is the set of **normal subgroups** of  $G$

Corollary: Given  $H \leq G$ , the number of conjugate subgroups is  $[G : N_G(H)]$ .

1. For a fixed proper subgroup  $H < G$ , let  $G$  act on its cosets  $G/H = \{gH \mid g \in G\}$  by left-multiplication.
  - $G \cdot gH = G/H$ , i.e. this is a *transitive* action.
  - $G_{gH} = gHg^{-1}$  is a *conjugate subgroup* of  $H$
  - $(G/H)^G = \emptyset$

*Application:* If  $G$  is simple,  $H < G$  proper, and  $[G : H] = n$ , then there exists an injective map  $\varphi : G \hookrightarrow S_n$ .

*Proof:* This action induces  $\varphi$ ; it is nontrivial since  $gH = H$  for all  $g$  implies  $H = G$ ;  $\ker \varphi \trianglelefteq G$  and  $G$  simple implies  $\ker \varphi = 1$ .

**Burnside's Formula:**

$$|X/G| = \frac{1}{|G|} \sum_{g \in G} |X^g|.$$

### 3.6.2 Sylow Theorems

**Notation:** For any  $p$ , let  $\text{Syl}_p(G)$  be the set of Sylow- $p$  subgroups of  $G$ .

Write

- $|G| = p^n m$  where  $(m, p) = 1$ ,
- $S_p$  a Sylow- $p$  subgroup, and
- $n_p$  the number of Sylow- $p$  subgroups.

**Definition:** A  $p$ -group is a group  $G$  such that every element is order  $p^k$  for some  $k$ . If  $G$  is a finite  $p$ -group, then  $|G| = p^j$  for some  $j$ .

**Lemma:**  $p$ -groups have nontrivial centers.

Some useful facts:

- Coprime order subgroups are disjoint, or more generally  $\mathbb{Z}_p, \mathbb{Z}_q \subset G \implies \mathbb{Z}_p \cap \mathbb{Z}_q = \mathbb{Z}_{(p,q)}$ .
- The Chinese Remainder theorem:  $(p, q) = 1 \implies \mathbb{Z}_p \times \mathbb{Z}_q \cong \mathbb{Z}_{pq}$

### 3.6.3 Sylow 1 (Cauchy for Prime Powers)

$\forall p^n$  dividing  $|G|$  there exists a subgroup of size  $p^n$ .

If  $|G| = \prod p_i^{\alpha_i}$ , then there exist subgroups of order  $p_i^{\beta_i}$  for every  $i$  and every  $0 \leq \beta_i \leq \alpha_i$ . In particular, Sylow  $p$ -subgroups always exist.

### 3.6.4 Sylow 2 (Sylows are Conjugate)

All sylow- $p$  subgroups  $S_p$  are conjugate, i.e.

$$S_p^1, S_p^2 \in \text{Syl}_p(G) \implies \exists g \text{ such that } gS_p^1g^{-1} = S_p^2.$$

**Corollary:**  $n_p = 1 \iff S_p \trianglelefteq G$

### 3.6.5 Sylow 3 (Numerical Constraints)

1.  $n_p \mid m$  (in particular,  $n_p \leq m$ ),
2.  $n_p \equiv 1 \pmod{p}$ ,
3.  $n_p = [G : N_G(S_p)]$  where  $N_G$  is the normalizer.

**Corollary:**  $p$  does not divide  $n_p$ .

**Lemma:** Every  $p$ -subgroup of  $G$  is contained in a Sylow  $p$ -subgroup.

*Proof:* Let  $H \leq G$  be a  $p$ -subgroup. If  $H$  is not *properly* contained in any other  $p$ -subgroup, it is a Sylow  $p$ -subgroup by definition. Otherwise, it is contained in some  $p$ -subgroup  $H^1$ . Inductively this yields a chain  $H \subsetneq H^1 \subsetneq \dots$ , and by Zorn's lemma  $H := \bigcup_i H^i$  is maximal and thus a Sylow  $p$ -subgroup.

**Fratini's Argument:** If  $H \trianglelefteq G$  and  $P \in \text{Syl}_p(G)$ , then  $HN_G(P) = G$  and  $[G : H]$  divides  $|N_G(P)|$ .

## 3.7 Products

**Characterizing direct products:**  $G \cong H \times K$  when

- $G = HK = \{hk \mid h \in H, k \in K\}$
- $H \cap K = \{e\} \subset G$
- $H, K \trianglelefteq G$

Can relax to only  $H \trianglelefteq G$  to get a semidirect product instead

**Characterizing semidirect products:**  $G = N \rtimes_{\psi} H$  when

- $G = NH$
- $N \trianglelefteq G$
- $H \curvearrowright N$  by conjugation via a map

$$\begin{aligned} \psi : H &\longrightarrow \text{Aut}(N) \\ h &\mapsto h(\cdot)h^{-1}. \end{aligned}$$

### Useful Facts

- If  $\sigma \in \text{Aut}(H)$ , then  $N \rtimes_{\psi} H \cong N \rtimes_{\psi \circ \sigma} H$ .
- $\text{Aut}(\mathbb{Z}/(p)^n) \cong \text{GL}(n, \mathbb{F}_p)$ , which has size  $|\text{Aut}(\mathbb{Z}/(p)^n)| = (p^n - 1)(p^n - p) \cdots (p^n - p^{n-1})$ .

- If this occurs in a semidirect product, it suffices to consider similarity classes of matrices (i.e. just use canonical forms)
- $\text{Aut}(\mathbb{Z}/(n)) \cong \mathbb{Z}/(n)^\times \cong \mathbb{Z}/(\varphi(n))$  where  $\varphi$  is the totient function.
  - $\varphi(p^k) = p^{k-1}(p-1)$
- If  $G, H$  have coprime order then  $\text{Aut}(G \oplus H) \cong \text{Aut}(G) \oplus \text{Aut}(H)$ .

### 3.8 Isomorphism Theorems


**Lemma:** If  $H, K \leq G$  and  $H \leq N_G(K)$  (or  $K \trianglelefteq G$ ) then  $HK \leq G$  is a subgroup.

Note that this implies that  $HK$  is not always a subgroup.

**Diamond Theorem / 2nd Isomorphism Theorem:**

If  $S \leq G$  and  $N \trianglelefteq G$ , then

$$\frac{SN}{N} \cong \frac{S}{S \cap N} \quad \text{and} \quad |SN| = \frac{|S||N|}{|S \cap N|}$$



Mnemonic:

Note: for this to make sense, we also have

- $SN \leq G$ ,
- $S \cap N \trianglelefteq S$ ,

**Cancellation / 3rd Isomorphism Theorem**

If  $H, K \trianglelefteq G$  with  $H \trianglelefteq K$ , then

$$\frac{G/H}{G/K} \cong \frac{G}{K}$$

Note: for this to make sense, we also have  $G/K \trianglelefteq G/H$ .

**The Correspondence Theorem / 4th Isomorphism Theorem:** Suppose  $N \trianglelefteq G$ , then there exists a correspondence:

$$\begin{aligned} \left\{ H < G \mid N \subseteq H \right\} &\Longleftrightarrow \left\{ H \mid H < \frac{G}{N} \right\} \\ \left\{ \begin{array}{c} \text{Subgroups of } G \\ \text{containing } N \end{array} \right\} &\Longleftrightarrow \left\{ \begin{array}{c} \text{Subgroups of the} \\ \text{quotient } G/N \end{array} \right\}. \end{aligned}$$

In words, subgroups of  $G$  containing  $N$  correspond to subgroups of the quotient group  $G/N$ . This is given by the map  $H \mapsto H/N$ .

Note:  $N \trianglelefteq G$  and  $N \subseteq H < G \implies N \trianglelefteq H$ .

### 3.9 Special Classes of Groups

**Definition:** The “**2 out of 3 property**” is satisfied by a class of groups  $\mathcal{C}$  iff whenever  $G \in \mathcal{C}$ , then  $N, G/N \in \mathcal{C}$  for any  $N \trianglelefteq G$ .

**Definition:** If  $|G| = p^k$ , then  $G$  is a **p-group**.

**Facts about p-groups:**

- If  $k = 1$  then  $G$  is cyclic
- If  $k = 2$ , then  $G \cong \mathbb{Z}/(p)^2$  or  $\mathbb{Z}/(p^2)$ .
- p-groups have nontrivial centers
  - Proof: Use class equation.
- Every normal subgroup is contained in the center
- Normalizers grow
- Every maximal is normal
- Every maximal has index  $p$
- p-groups are *nilpotent*
- p-groups are *solvable*

**Facts about other special order groups:**

General strategy: find a normal subgroup (usually a Sylow) and use recognition of semidirect products.

- $|G| = pq$ : Two possibilities. By cases:
  - If  $p$  divides  $q - 1$ , two cases:
    - \*  $G \cong \mathbb{Z}/(pq)$  or  $\mathbb{Z}(p) \times \mathbb{Z}/(q)$
  - Otherwise,  $G \cong \mathbb{Z}/(pq)$

Proof: Sylow theorems. Note: Such groups are never simple.

- $|G| = p^2q$ :
  - $q \mid p^2 - 1$ : Two abelian possibilities,  $\mathbb{Z}/(p) \times \mathbb{Z}/(q^2)$ , or  $\mathbb{Z}/(pq) \times \mathbb{Z}/(q)$ .

- Otherwise, the sylow-q subgroup  $H$  is normal and order  $q^2$ , so either  $\mathbb{Z}/(q)^2$  or  $\mathbb{Z}/(q^2)$ .
  - \* Case 2:  $|\text{Aut}(\mathbb{Z}/(q)^2)| = q(q-1)$ , so only trivial action
  - \* Case 1:  $|\text{Aut}(\mathbb{Z}/(q^2))| = q(q-1)^2(q+1)$ 
    - If  $p$  doesn't divide  $q+1$ , noting new
    - Otherwise, a nontrivial semidirect product.

**Definition:** A group  $G$  is **simple** iff  $H \trianglelefteq G \implies H = \{e\}, G$ , i.e. it has no non-trivial proper subgroups.

**Lemma:** If  $G$  is *not* simple, then for any  $N \trianglelefteq G$ , it is the case that  $G \cong E$  for an extension of the form  $N \longrightarrow E \longrightarrow G/N$ .  $>$

**Definition:** A group  $G$  is **solvable** iff  $G$  has a terminating normal series with abelian factors, i.e.

$$G \longrightarrow G^1 \longrightarrow \cdots \longrightarrow \{e\} \text{ with } G^i/G^{i+1} \text{ abelian for all } i.$$

**Lemmas:**

- $G$  is solvable iff  $G$  has a terminating *derived series*.
- Solvable groups satisfy the 2 out of 3 property
- Abelian  $\implies$  solvable
- Every group of order less than 60 is solvable.

**Definition:** A group  $G$  is **nilpotent** iff  $G$  has a terminating central series, upper central series, or lower central series.

Moral: the adjoint map is nilpotent.

**Lemma:** For  $G$  a finite group, TFAE:

- $G$  is nilpotent
- Normalizers grow (i.e.  $H < N_G(H)$  whenever  $H$  is proper)
- Every Sylow-p subgroup is normal
- $G$  is the direct product of its Sylow p-subgroups
- Every maximal subgroup is normal
- $G$  has a terminating *Lower Central Series*
- $G$  has a terminating *Upper Central Series*

**Lemmas:**

- $G$  nilpotent  $\implies G$  solvable
- Nilpotent groups satisfy the 2 out of 3 property.
- $G$  has normal subgroups of order  $d$  for *every*  $d$  dividing  $|G|$
- $G$  nilpotent  $\implies Z(G) \neq 0$
- Abelian  $\implies$  nilpotent
- p-groups  $\implies$  nilpotent

### 3.10 Series of Groups

**Definition:** A **normal series** of a group  $G$  is a sequence  $G \longrightarrow G^1 \longrightarrow G^2 \longrightarrow \cdots$  such that  $G^{i+1} \trianglelefteq G_i$  for every  $i$ .



**Definition** A **composition series** of a group  $G$  is a finite normal series such that  $G^{i+1}$  is a *maximal proper* normal subgroup of  $G^i$ .

**Theorem (Jordan-Holder):** Any two composition series of a group have the same length and isomorphic factors (up to permutation).<sup>1</sup>

**Definition** A **derived series** of a group  $G$  is a normal series  $G \rightarrow G^1 \rightarrow G^2 \rightarrow \dots$  where  $G^{i+1} = [G^i, G^i]$  is the commutator subgroup.

The derived series terminates iff  $G$  is *solvable*.

**Definition:** A **central series** for a group  $G$  is a terminating normal series  $G \rightarrow G^1 \rightarrow \dots \rightarrow \{e\}$  such that each quotient is **central**, i.e.  $[G, G^i] \leq G^{i-1}$  for all  $i$ .

**Definition:** A **lower central series** is a terminating normal series  $G \rightarrow G^1 \rightarrow \dots \rightarrow \{e\}$  such that  $G^{i+1} = [G^i, G]$

Moral: Iterate the adjoint map  $[\cdot, G]$ .

$G$  is nilpotent  $\iff$  the LCS terminates.

**Definition:** An **upper central series** is a terminating normal series  $G \rightarrow G^1 \rightarrow \dots \rightarrow \{e\}$  such that  $G^1 = Z(G)$  and  $G^{i+1}$  is defined such that  $G^{i+1}/G^i = Z(G^i)$ .

Moral: Iterate taking “higher centers”.

## 3.11 Classification of Groups

- Keith Conrad: Classifying Groups of Order 12
- Order  $p$ : cyclic.
- Order  $pq$ : ?
- Order  $p^2q$ : ?

## 4 Rings

### 4.1 Definitions

**Lemma:** Intersections, products, and sums (but not necessarily unions) of ideals are ideals.

**Theorem (Krull):** Every ring has proper maximal ideals, and any proper ideal is contained in a maximal ideal.

**Definition:** A ring  $R$  is **simple** iff every ideal  $I \trianglelefteq R$  is either 0 or  $R$ .

**Definition:** An element  $r \in R$  is **irreducible** iff  $r = ab \implies a$  is a unit or  $b$  is a unit.

**Definition:** An element  $r \in R$  is **prime** iff  $ab \mid r \implies a \mid r$  or  $b \mid r$  whenever  $a, b$  are nonzero and not units.

**Definition:**  $\mathfrak{p}$  is a **prime ideal**  $\iff ab \in \mathfrak{p} \implies a \in \mathfrak{p}$  or  $b \in \mathfrak{p}$ .

**Definition:**  $\text{Spec}(R) = \{\mathfrak{p} \trianglelefteq R \mid \mathfrak{p} \text{ is prime}\}$  is the **spectrum** of  $R$ .

**Definition:**  $\mathfrak{m}$  is **maximal**  $\iff I \triangleleft R \implies I \subseteq \mathfrak{m}$ .

Example: Maximal ideals of  $R[x]$  are of the form  $I = (x - a_i)$  for some  $a_i \in R$ .

**Definition:**  $\text{Spec}_{\max}(R) = \{\mathfrak{m} \trianglelefteq R \mid \mathfrak{m} \text{ is maximal}\}$  is the **max-spectrum** of  $R$ .

Note: nonstandard notation / definition.

**Lemmas (Quotients of Rings):**

- $R/I$  is a domain  $\iff I$  is prime,
- $R/I$  is a field  $\iff I$  is maximal.
- For  $R$  a PID,  $I$  is prime  $\iff I$  is maximal.

**Lemma (Characterizations of Rings):**

- $R$  a commutative division ring  $\implies R$  is a field
- $R$  a finite integral domain  $\implies R$  is a field.
- $\mathbb{F}$  a field  $\implies \mathbb{F}[x]$  is a Euclidean domain.
- $\mathbb{F}$  a field  $\implies \mathbb{F}[x]$  is a PID.
- $\mathbb{F}$  is a field  $\iff \mathbb{F}$  is a commutative simple ring.
- $R$  is a UFD  $\iff R[x]$  is a UFD.
- $R$  a PID  $\implies R[x]$  is a UFD
- $R$  a PID  $\implies R$  Noetherian
- $R[x]$  a PID  $\implies R$  is a field.

**Lemma:** Fields  $\subset$  Euclidean domains  $\subset$  PIDs  $\subset$  UFDs  $\subset$  Integral Domains  $\subset$  Rings

- A Euclidean Domain that is not a field:  $\mathbb{F}[x]$  for  $\mathbb{F}$  a field  
– *Proof:* Use previous lemma, and  $x$  is not invertible
- A PID that is not a Euclidean Domain:  $\mathbb{Z}\left[\frac{1 + \sqrt{-19}}{2}\right]$ .  
– *Proof:* complicated.
- A UFD that is not a PID:  $\mathbb{F}[x, y]$ .  
– *Proof:*  $\langle x, y \rangle$  is not principal
- An integral domain that is not a UFD:  $\mathbb{Z}[\sqrt{-5}]$   
– *Proof:*  $(2 + \sqrt{-5})(2 - \sqrt{-5}) = 9 = 3 \cdot 3$ , where all factors are irreducible (check norm).
- A ring that is not an integral domain:  $\mathbb{Z}/(4)$   
– *Proof:*  $2 \bmod 4$  is a zero divisor.

**Lemma:** In  $R$  a UFD, an element  $r \in R$  is prime  $\iff r$  is irreducible.

Note: For  $R$  an integral domain, prime  $\implies$  irreducible, but generally not the converse.

Example of a prime that is not irreducible:  $x^2 \bmod (x^2 + x) \in \mathbb{Q}[x]/(x^2 + x)$ . Check that  $x$  is prime directly, but  $x = x \cdot x$  and  $x$  is not a unit.

Example of an irreducible that is not prime:  $3 \in \mathbb{Z}[\sqrt{-5}]$ . Check norm to see irreducibility, but  $3 \mid 9 = (2 + \sqrt{-5})(2 - \sqrt{-5})$  and doesn't divide either factor.

**Lemma:** If  $R$  is a PID, then every element in  $R$  has a unique prime factorization.

**Definition:** A nonzero unital ring  $R$  is **semisimple** iff  $R \cong \bigoplus_{i=1}^n M_i$  with each  $M_i$  a simple module.

**Theorem (Artin-Wedderburn):** If  $R$  is a nonzero, unital, semisimple ring then  $R \cong \bigoplus_{i=1}^m \text{Mat}(n_i, D_i)$ ,

a finite sum of matrix rings over division rings.

*Corollary:* If  $M$  is a simple ring over  $R$  a division ring, the  $M$  is isomorphic to a matrix ring.

## 4.2 Nontrivial Properties

**Lemma:** Every  $a \in R$  for a finite ring is either a unit or a zero divisor.

*Proof:* Let  $a \in R$  and define  $\varphi(x) = ax$ . If  $\varphi$  is injective, then it is surjective, so  $1 = ax$  for some  $x \implies x^{-1} = a$ . Otherwise,  $ax_1 = ax_2$  with  $x_1 \neq x_2 \implies a(x_1 - x_2) = 0$  and  $x_1 - x_2 \neq 0$ , so  $a$  is a zero divisor.

## 4.3 Ideals

### 4.3.1 Maximal and Prime Ideals

**Lemma:** Maximal  $\implies$  prime, but generally not the converse.

*Counterexample:*  $(0) \in \mathbb{Z}$  is prime since  $\mathbb{Z}$  is a domain, but not maximal since it is properly contained in any other ideal.

*Proof:* Suppose  $\mathfrak{m}$  is maximal,  $ab \in \mathfrak{m}$ , and  $b \notin \mathfrak{m}$ . Then there is a containment of ideals  $\mathfrak{m} \subsetneq \mathfrak{m} + (b) \implies \mathfrak{m} + (b) = R$ .  
So

$$1 = m + rb \implies a = am + r(ab),$$

but  $am \in \mathfrak{m}$  and  $ab \in \mathfrak{m} \implies a \in \mathfrak{m}$ . ■

**Lemma:** If  $x$  is not a unit, then  $x$  is contained in some maximal ideal  $\mathfrak{m}$ .

*Proof:* Zorn's lemma.

**Lemma:**  $R/\mathfrak{m}$  is a field  $\iff \mathfrak{m}$  is maximal.

**Lemma:**  $R/\mathfrak{p}$  is an integral domain  $\iff \mathfrak{p}$  is prime.

### 4.3.2 Nilradical and Jacobson Radical

**Definition:**  $\mathfrak{N} := \{x \in R \mid x^n = 0 \text{ for some } n\}$  is the **nilradical** of  $R$ .

**Lemma:** The nilradical is the intersection of all **prime** ideals, i.e.

$$\mathfrak{N}(R) = \bigcap_{\mathfrak{p} \in \text{Spec}(R)} \mathfrak{p}$$

*Proof:*

$\mathfrak{N} \subseteq \bigcap \mathfrak{p}$ :  $x \in \mathfrak{N} \implies x^n = 0 \in \mathfrak{p} \implies x \in \mathfrak{p}$  or  $x^{n-1} \in \mathfrak{p}$ .

$\mathfrak{N}^c \subseteq \bigcup \mathfrak{p}^c$ : Define  $S = \{I \trianglelefteq R \mid a^n \notin I \text{ for any } n\}$ . Then apply Zorn's lemma to get a maximal ideal  $\mathfrak{m}$ , and maximal  $\implies$  prime.

**Lemma:**  $R/\mathfrak{N}(R)$  has no nonzero nilpotent elements.

*Proof:*

$$\begin{aligned}
a + \mathfrak{N}(R) \text{ nilpotent} &\implies (a + \mathfrak{N}(R))^n := a^n + \mathfrak{N}(R) = \mathfrak{N}(R) \\
&\implies a^n \in \mathfrak{N}(R) \\
&\implies \exists \ell \text{ such that } (a^n)^\ell = 0 \\
&\implies a \in \mathfrak{N}(R).
\end{aligned}$$

**Definition:** The **Jacobson radical** is the intersection of all **maximal** ideals, i.e.

$$J(R) = \bigcap_{\mathfrak{m} \in \text{Spec}_{\max}} \mathfrak{m}$$

**Lemma:**  $\mathfrak{N}(R) \subseteq J(R)$ .

*Proof:* Maximal  $\implies$  prime, and so if  $x$  is in every prime ideal, it is necessarily in every maximal ideal as well.

### 4.3.3 Zorn's Lemma

**Lemma:** A field has no nontrivial proper ideals.

**Lemma:** If  $I \trianglelefteq R$  is a proper ideal  $\iff I$  contains no units.

$$\text{Proof: } r \in R^\times \cap I \implies r^{-1}r \in I \implies 1 \in I \implies x \cdot 1 \in I \quad \forall x \in R.$$

**Lemma:** If  $I_1 \subseteq I_2 \subseteq \dots$  are ideals then  $\bigcup_j I_j$  is an ideal.

**Example Application of Zorn's Lemma:** Every proper ideal is contained in a maximal ideal.

*Proof:* Let  $0 < I < R$  be a proper ideal, and consider the set

$$S = \left\{ J \mid I \subseteq J < R \right\}.$$

Note  $I \in S$ , so  $S$  is nonempty. The claim is that  $S$  contains a maximal element  $M$ .  $S$  is a poset, ordered by set inclusion, so if we can show that every chain has an upper bound, we can apply Zorn's lemma to produce  $M$ .

Let  $C \subseteq S$  be a chain in  $S$ , so  $C = \{C_1 \subseteq C_2 \subseteq \dots\}$  and define  $\widehat{C} = \bigcup_i C_i$ .

**$\widehat{C}$  is an upper bound for  $C$ :**

This follows because every  $C_i \subseteq \widehat{C}$ .

**$\widehat{C}$  is in  $S$ :**

Use the fact that  $I \subseteq C_i < R$  for every  $C_i$  and since no  $C_i$  contains a unit,  $\widehat{C}$  doesn't contain a unit, and is thus proper. ■

## 5 Fields

Let  $k$  denote a field.

**Lemmas:**

- The characteristic of any field  $k$  is either 0 or  $p$  a prime.
- All fields are simple rings (no proper nontrivial ideals).
- If  $L/k$  is algebraic, then  $\min(\alpha, L)$  divides  $\min(\alpha, k)$ .
- Every field morphism is either zero or injective.

**Theorem 5.1.**

Every finite extension is algebraic.

*Proof .*

Todo? ■

**Theorem 5.2 (Gauss' Lemma).**

Let  $R$  be a UFD and  $F$  its field of fractions. Then a primitive  $p \in R[x]$  is irreducible in  $R[x] \iff p$  is irreducible in  $F[x]$ .

**Corollary 5.3.**

A primitive polynomial  $p \in \mathbb{Q}[x]$  is irreducible  $\iff p$  is irreducible in  $\mathbb{Z}[x]$ .

**Theorem 5.4 (Eisenstein's Criterion).**

If  $f(x) = \sum_{i=0}^n \alpha_i x^i \in \mathbb{Q}[x]$  and  $\exists p$  such that

- $p$  divides every coefficient *except*  $a_n$  and
- $p^2$  does not divide  $a_0$ ,

then  $f$  is irreducible over  $\mathbb{Q}[x]$ , and by Gauss' lemma, over  $\mathbb{Z}[x]$ .

**Definition 5.4.1 (Primitive).**

For  $R$  a UFD, a polynomial  $p \in R[x]$  is **primitive** iff the greatest common divisors of its coefficients is a unit.

## 5.1 Finite Fields

**Definition 5.4.2.**

The **prime subfield** of a field  $F$  is the subfield generated by 1.

**Lemma 5.5 (Characterization of Prime Subfields).**

The prime subfield of any field is isomorphic to either  $\mathbb{Q}$  or  $\mathbb{F}_p$  for some  $p$ .

**Proposition 5.6 (Freshman's Dream).**

If  $\text{char } k = p$  then  $(a + b)^p = a^p + b^p$  and  $(ab)^p = a^p b^p$ .

*Proof .*

Todo ■

**Theorem 5.7 (Construction of Finite Fields).**

$\mathbb{GF}(p^n) \cong \frac{\mathbb{F}_p}{(f)}$  where  $f \in \mathbb{F}_p[x]$  is any irreducible of degree  $n$ , and  $\mathbb{GF}(p^n) \cong \mathbb{F}[\alpha] \cong \text{span}_{\mathbb{F}} \{1, \alpha, \dots, \alpha^{n-1}\}$  for any root  $\alpha$  of  $f$ .

**Lemma 5.8 (Prime Subfields of Finite Fields).**

Every finite field  $F$  is isomorphic to a unique field of the form  $\mathbb{GF}(p^n)$  and if  $\text{char } F = p$ , it has prime subfield  $\mathbb{F}_p$ .

**Lemma 5.9 (Containment of Finite Fields).**

$\mathbb{GF}(p^\ell) \leq \mathbb{GF}(p^k) \iff \ell \text{ divides } k$ .

**Lemma 5.10 (Identification of Finite Fields as Splitting Fields).**

$\mathbb{GF}(p^n)$  is the splitting field of  $\rho(x) = x^{p^n} - x$ , and the elements are exactly the roots of  $\rho$ .

*Proof.*

Todo. Every element is a root by Cauchy's theorem, and the  $p^n$  roots are distinct since its derivative is identically  $-1$ . ■

**Lemma 5.11 (Splits Product of Irreducibles).**

Let  $\rho_n := x^{p^n} - x$ . Then  $f(x) \mid \rho_n(x) \iff \deg f \mid n$  and  $f$  is irreducible.

**Corollary 5.12.**

$x^{p^n} - x = \prod f_i(x)$  over all irreducible monic  $f_i \in \mathbb{F}_p[x]$  of degree  $d$  dividing  $n$ .

*Proof.*

$\Leftarrow$  : Suppose  $f$  is irreducible of degree  $d$ . Then  $f \mid x^{p^d} - x$  (consider  $F[x]/\langle f \rangle$ ) and  $x^{p^d} - x \mid x^{p^n} - x \iff d \mid n$ .

$\Rightarrow$  :

- $\alpha \in \mathbb{GF}(p^n) \iff \alpha^{p^n} - \alpha = 0$ , so every element is a root of  $\varphi_n$  and  $\deg \min(\alpha, \mathbb{F}_p) \mid n$  since  $\mathbb{F}_p(\alpha)$  is an intermediate extension.
- So if  $f$  is an irreducible factor of  $\varphi_n$ ,  $f$  is the minimal polynomial of some root  $\alpha$  of  $\varphi_n$ , so  $\deg f \mid n$ .  $\varphi'_n(x) = p^n x^{p^n-1} \neq 0$ , so  $\varphi_n$  has distinct roots and thus no repeated factors. So  $\varphi_n$  is the product of all such irreducible  $f$ . ■

**Lemma 5.13.**

No finite field is algebraically closed.

*Proof .*  
 Todo?

■

## 5.2 Galois Theory

### Definition 5.13.1.

A field extension  $L/k$  is **algebraic** iff every  $\alpha \in L$  is the root of some polynomial  $f \in k[x]$ .

### Definition 5.13.2.

Let  $L/k$  be a finite extension. Then TFAE:

- $L/k$  is **normal**.
- Every irreducible  $f \in k[x]$  that has one root in  $L$  has *all* of its roots in  $L$   
 – i.e. every polynomial splits into linear factors
- Every embedding  $\sigma : L \hookrightarrow \bar{k}$  that is a lift of the identity on  $k$  satisfies  $\sigma(L) = L$ .
- If  $L$  is separable:  $L$  is the splitting field of some irreducible  $f \in k[x]$ .

### Definition 5.13.3.

Let  $L/k$  be a field extension,  $\alpha \in L$  be arbitrary, and  $f(x) := \min(\alpha, k)$ . TFAE:

- $L/k$  is **separable**
- $f$  has no repeated factors/roots
- $\gcd(f, f') = 1$ , i.e.  $f$  is coprime to its derivative
- $f' \neq 0$

### Lemma 5.14.

If  $\text{char } k = 0$  or  $k$  is finite, then every *algebraic* extension  $L/k$  is separable.

### Definition 5.14.1.

$\text{Aut}(L/k) = \left\{ \sigma : L \longrightarrow L \mid \sigma|_k = \text{id}_k \right\}$ .

### Lemma 5.15.

If  $L/k$  is algebraic, then  $\text{Aut}(L/k)$  permutes the roots of irreducible polynomials.

### Lemma 5.16.

$|\text{Aut}(L/k)| \leq [L : k]$  with equality precisely when  $L/k$  is normal.

### Definition 5.16.1.


If  $L/k$  is Galois, we define  $\text{Gal}(L/k) := \text{Aut}(L/k)$ .

### 5.2.1 Lemmas About Towers

Let  $L/F/k$  be a finite tower of field extensions

- Multiplicativity:  $[L : k] = [L : F][F : k]$

- $L/k$  normal/algebraic/Galois  $\implies L/F$  normal/algebraic/Galois.
  - *Proof (normal)*:  $\min(\alpha, F) \mid \min(\alpha, k)$ , so if the latter splits in  $L$  then so does the former.
  - *Corollary*:  $\alpha \in L$  algebraic over  $k \implies \alpha$  algebraic over  $F$ .
  - *Corollary*:  $E_1/k$  normal and  $E_2/k$  normal  $\implies E_1E_2/k$  normal and  $E_1 \cap E_2/k$  normal.



- $F/k$  algebraic and  $L/F$  algebraic  $\implies L/k$  algebraic.
- If  $L/k$  is algebraic, then  $F/k$  separable and  $L/F$  separable  $\iff L/k$  separable



- $F/k$  Galois and  $L/F$  Galois  $\implies F/k$  Galois **only if**  $\text{Gal}(L/F) \trianglelefteq \text{Gal}(L/k)$ 
  - $\implies \text{Gal}(F/k) \cong \frac{\text{Gal}(L/k)}{\text{Gal}(L/F)}$



### Common Counterexamples:

- $\mathbb{Q}(\zeta_3, 2^{1/3})$  is normal but  $\mathbb{Q}(2^{1/3})$  is not since the irreducible polynomial  $x^3 - 2$  has only one root in it.

### Definition 5.16.2 (Characterizations of Galois Extensions).

Let  $L/k$  be a finite field extension. TFAE:

- $L/k$  is **Galois**
- $L/k$  is finite, normal, and separable.
- $L/k$  is the splitting field of a separable polynomial
- $|\text{Aut}(L/k)| = [L : k]$
- The fixed field of  $\text{Aut}(L/k)$  is exactly  $k$ .

### Theorem 5.17 (Fundamental Theorem of Galois Theory).



Let  $L/k$  be a Galois extension, then there is a correspondence:

$$\begin{aligned} \{\text{Subgroups } H \leq \text{Gal}(L/k)\} &\iff \left\{ \begin{array}{l} \text{Fields } F \text{ such} \\ \text{that } L/F/k \end{array} \right\} \\ H &\rightarrow \{E^H := \text{The fixed field of } H\} \\ \left\{ \text{Gal}(L/F) := \left\{ \sigma \in \text{Gal}(L/k) \mid \sigma(F) = F \right\} \right\} &\leftarrow F. \end{aligned}$$

- This is contravariant with respect to subgroups/subfields.
- $[F : k] = [G : H]$ , so degrees of extensions over the base field correspond to indices of subgroups.
- $[K : F] = |H|$
- $L/F$  is Galois and  $\text{Gal}(K/F) = H$
- $F/k$  is Galois  $\iff H$  is normal, and  $\text{Gal}(F/k) = \text{Gal}(L/k)/H$ .
- The compositum  $F_1 F_2$  corresponds to  $H_1 \cap H_2$ .
- The subfield  $F_1 \cap F_2$  corresponds to  $H_1 H_2$ .

### 5.2.2 Examples

1.  $\text{Gal}(\mathbb{Q}(\zeta_n)/\mathbb{Q}) \cong \mathbb{Z}/(n)^\times$  and is generated by maps of the form  $\zeta_n \mapsto \zeta_n^j$  where  $(j, n) = 1$ .  
I.e., the following map is an isomorphism:

$$\begin{aligned} \mathbb{Z}/(n)^\times &\longrightarrow \text{Gal}(\mathbb{Q}(\zeta_n), \mathbb{Q}) \\ r \pmod n &\mapsto (\varphi_r : \zeta_n \mapsto \zeta_n^r). \end{aligned}$$

2.  $\text{Gal}(\mathbb{GF}(p^n)/\mathbb{F}_p) \cong \mathbb{Z}/(n)$ , a cyclic group generated by powers of the Frobenius automorphism:

$$\begin{aligned} \varphi_p : \mathbb{GF}(p^n) &\longrightarrow \mathbb{GF}(p^n) \\ x &\mapsto x^p. \end{aligned}$$

#### Lemma 5.18.

Every quadratic extension is Galois.

#### Lemma 5.19.

If  $K$  is the splitting field of an irreducible polynomial of degree  $n$ , then  $\text{Gal}(K/\mathbb{Q}) \leq S_n$  is a transitive subgroup.

#### Corollary 5.20.

$n$  divides the order  $|\text{Gal}(K/\mathbb{Q})|$ .

#### Definition 5.20.1.

TFAE:

- $k$  is a **perfect** field.
- Every irreducible polynomial  $p \in k[x]$  is separable
- Every finite extension  $F/k$  is separable.

- If  $\text{char } k > 0$ , the Frobenius is an automorphism of  $k$ .

**Theorem 5.21.**

- If  $\text{char } k = 0$  or  $k$  is finite, then  $k$  is perfect.
- $k = \mathbb{Q}, \mathbb{F}_p$  are perfect, and any finite normal extension is Galois.
- Every splitting field of a polynomial over a perfect field is Galois.

**Proposition 5.22 (Composite Extensions).**

If  $F/k$  is finite and Galois and  $L/k$  is arbitrary, then  $FL/L$  is Galois and

$$\text{Gal}(FL/L) = \text{Gal}(F/F \cap L) \subset \text{Gal}(F/k).$$

### 5.3 Cyclotomic Polynomials

**Definition 5.22.1** (Cyclotomic Polynomials).

Let  $\zeta_n = e^{2\pi i/n}$ , then the  $n$ th cyclotomic polynomial is given by

$$\Phi_n(x) = \prod_{\substack{k=1 \\ (k,n)=1}}^n (x - \zeta_n^k),$$

which is a product over primitive roots of unity. It is the unique irreducible polynomial which is a divisor of  $x^n - 1$  but *not* a divisor of  $x^k - 1$  for any  $k < n$ .

**Proposition 5.23.**

$\deg \Phi_n(x) = \varphi(n)$  for  $\varphi$  the totient function.

*Proof.*

$\deg \Phi_n(x)$  is the number of  $n$ th primitive roots, which is the number of numbers less than and coprime to  $n$ . ■

**Computing  $\Phi_n$ :**

1.

$$\Phi_n(z) = \prod_{d|n, d>0} (z^d - 1)^{\mu(\frac{n}{d})}$$

where

$$\mu(n) \equiv \begin{cases} 0 & \text{if } n \text{ has one or more repeated prime factors} \\ 1 & \text{if } n = 1 \\ (-1)^k & \text{if } n \text{ is a product of } k \text{ distinct primes,} \end{cases}$$

2.

$$x^n - 1 = \prod_{d|n} \Phi_d(x) \implies \Phi_n(x) = \frac{x^n - 1}{\prod_{\substack{d|n \\ d < n}} \Phi_d(x)},$$

so just use polynomial long division.

**Lemma 5.24.**

$$\begin{aligned}\Phi_p(x) &= x^{p-1} + x^{p-2} + \cdots + x + 1 \\ \Phi_{2p}(x) &= x^{p-1} - x^{p-2} + \cdots - x + 1.\end{aligned}$$

**Lemma 5.25.**

$$k \mid n \implies \Phi_{nk}(x) = \Phi_n(x^k)$$

**Definition 5.25.1.**

An extension  $F/k$  is **simple** if  $F = k[\alpha]$  for a single element  $\alpha$ .

**Theorem 5.26 (Primitive Element).**

Every finite separable extension is simple.

**Corollary 5.27.**

$\mathbb{GF}(p^n)$  is a simple extension over  $\mathbb{F}_p$ .

## 6 Modules

### 6.1 General Modules

**Definition:** A module is **simple** iff it has no nontrivial proper submodules.

**Definition:** A **free** module is a module with a basis (i.e. a spanning, linearly independent set).

*Example:*  $\mathbb{Z}/(6)$  is a  $\mathbb{Z}$ -module that is *not* free.

**Definition:** A module  $M$  is **projective** iff  $M$  is a direct summand of a free module  $F = M \oplus \cdots$ .

Free implies projective, but not the converse.

**Definition:** A sequence of homomorphisms  $0 \xrightarrow{d_1} A \xrightarrow{d_2} B \xrightarrow{d_3} C \longrightarrow 0$  is *exact* iff  $\text{im } d_i = \ker d_{i+1}$ .

**Lemma:** If  $0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$  is a short exact sequence, then

- $C$  free  $\implies$  the sequence splits
- $C$  projective  $\implies$  the sequence splits

- $A$  injective  $\implies$  the sequence splits

Moreover, if this sequence splits, then  $B \cong A \oplus C$ .

## 6.2 Classification of Modules over a PID

Let  $M$  be a finitely generated modules over a PID  $R$ . Then there is an invariant factor decomposition

$$M \cong F \oplus R/(r_i) \quad \text{where } r_1 \mid r_2 \mid \cdots,$$

and similarly an elementary divisor decomposition.

## 6.3 Minimal / Characteristic Polynomials

Fix some notation:

$\min_A(x)$  : The minimal polynomial of  $A$

$\chi_A(x)$  : The characteristic polynomial of  $A$ .

**Definition:** The minimal polynomial is the unique polynomial  $\min_A(x)$  of minimal degree such that  $\min_A(A) = 0$ .

**Definition:** The **characteristic polynomial** of  $A$  is given by

$$\chi_A(x) = \det(A - xI) = \det(SNF(A - xI)).$$

*Useful lemma:* If  $A$  is upper triangular, then  $\det(A) = \prod_i a_{ii}$

**Theorem (Cayley-Hamilton):** The minimal polynomial divides the characteristic polynomial, and in particular  $\chi_A(A) = 0$ .

**Lemma:** Writing

$$\begin{aligned} \min_A(x) &= \prod (x - \lambda_i)^{a_i} \\ \chi_A(x) &= \prod (x - \lambda_i)^{b_i} \end{aligned}$$

- $a_i \leq b_i$
- The roots both polynomials are precisely the eigenvalues of  $A$ .

*Proof:* By Cayley-Hamilton,  $\min$  divides  $\chi_A$ . Every  $\lambda_i$  is a root of  $\mu_M$ :  
Let  $(\mathbf{v}_i, \lambda_i)$  be a nontrivial eigenpair. Then by linearity,

$$\min_A(\lambda_i)\mathbf{v}_i = \min_A(A)\mathbf{v}_i = \mathbf{0},$$

which forces  $\min_A(\lambda_i) = 0$ .

**Definition:** Two matrices  $A, B$  are **similar** (i.e.  $A = PBP^{-1}$ )  $\iff A, B$  have the same Jordan Canonical Form (JCF).

**Definition:** Two matrices  $A, B$  are **equivalent** (i.e.  $A = PBQ$ )  $\iff$

- They have the same rank,
- They have the same invariant factors, *and*
- They have the same (JCF)

### Finding the minimal polynomial:

Let  $m(x)$  denote the minimal polynomial  $A$ .

1. Find the characteristic polynomial  $\chi(x)$ ; this annihilates  $A$  by Cayley-Hamilton. Then  $m(x) \mid \chi(x)$ , so just test the finitely many products of irreducible factors.
2. Pick any  $\mathbf{v}$  and compute  $T\mathbf{v}, T^2\mathbf{v}, \dots, T^k\mathbf{v}$  until a linear dependence is introduced. Write this as  $p(T) = 0$ ; then  $\min_A(x) \mid p(x)$ .

**Definition:** Given a monic  $p(x) = a_0 + a_1x + a_2x^2 + \dots + a_{n-1}x^{n-1} + x^n$ , the **companion matrix** of  $p$  is given by

$$C_p := \begin{bmatrix} 0 & 0 & \dots & 0 & -a_0 \\ 1 & 0 & \dots & 0 & -a_1 \\ 0 & 1 & \dots & 0 & -a_2 \\ \vdots & & \ddots & & \vdots \\ 0 & 0 & \dots & 1 & -a_{n-1} \end{bmatrix}.$$

## 6.4 Canonical Forms

### 6.4.1 Rational Canonical Form

Corresponds to the **Invariant Factor Decomposition** of  $T$ .

**Lemma:**  $RCF(A)$  is a block matrix where each block is the companion matrix of an invariant factor of  $A$ .

#### Derivation:

- Let  $k[x] \curvearrowright V$  using  $T$ , take invariant factors  $a_i$ ,
- Note that  $T \curvearrowright V$  by multiplication by  $x$
- Write  $\bar{x} = \pi(x)$  where  $F[x] \xrightarrow{\pi} F[x]/(a_i)$ ; then  $\text{span}\{\bar{x}\} = F[x]/(a_i)$ .
- Write  $a_i(x) = \sum b_i x^i$ , note that  $V \longrightarrow F[x]$  pushes  $T \curvearrowright V$  to  $T \curvearrowright k[x]$  by multiplication by  $\bar{x}$
- WRT the basis  $\bar{x}$ ,  $T$  then acts via the companion matrix on this summand.
- Each invariant factor corresponds to a block of the RCF.

### 6.4.2 Jordan Canonical Form

Corresponds to the **Elementary Divisor Decomposition** of  $T$ .

**Lemma:** The elementary divisors of  $A$  are the minimal polynomials of the Jordan blocks.

**Lemma:** Writing

$$\begin{aligned}\min_A(x) &= \prod (x - \lambda_i)^{a_i} \\ \chi_A(x) &= \prod (x - \lambda_i)^{b_i}\end{aligned}$$

- $a_i \leq b_i$
- $a_i$  tells you the size of the **largest** Jordan block associated to  $\lambda_i$ ,
- $b_i$  is the **sum of sizes** of all Jordan blocks associated to  $\lambda_i$
- $\dim E_{\lambda_i}$  is the **number of Jordan blocks** associated to  $\lambda_i$

## 6.5 Using Canonical Forms

**Lemma:** The characteristic polynomial is the *product of the invariant factors*, i.e.

$$\chi_A(x) = \prod_{j=1}^n f_j(x).$$

**Lemma:** The minimal polynomial of  $A$  is the *invariant factor of highest degree*, i.e.

$$\min_A(x) = f_n(x).$$

**Lemma:** For a linear operator on a vector space of nonzero finite dimension, TFAE:

- The minimal polynomial is equal to the characteristic polynomial.
- The list of invariant factors has length one.
- The Rational Canonical Form has a single block.
- The operator has a matrix similar to a companion matrix.
- There exists a *cyclic vector*  $\mathbf{v}$  such that  $\text{span}_k \{T^j \mathbf{v} \mid j = 1, 2, \dots\} = V$ .
- $T$  has  $\dim V$  distinct eigenvalues

## 6.6 Diagonalizability

*Notation:*  $A^*$  denotes the conjugate transpose of  $A$ .

**Lemma:** Let  $V$  be a vector space over  $k$  an algebraically closed and  $A \in \text{End}(V)$ . Then if  $W \subseteq V$  is an invariant subspace, so  $A(W) \subseteq W$ , the  $A$  has an eigenvector in  $W$ .

**Theorem (The Spectral Theorem):**

1. Hermitian matrices (i.e.  $A^* = A$ ) are diagonalizable over  $\mathbb{C}$ .
2. Symmetric matrices (i.e.  $A^t = A$ ) are diagonalizable over  $\mathbb{R}$ .

*Proof:* Suppose  $A$  is Hermitian. Since  $V$  itself is an invariant subspace,  $A$  has an eigenvector  $\mathbf{v}_1 \in V$ . Let  $W_1 = \text{span}_k \{\mathbf{v}_1\}^\perp$ . Then for any  $\mathbf{w}_1 \in W_1$ ,

$$\langle \mathbf{v}_1, A\mathbf{w}_1 \rangle = \langle A\mathbf{v}_1, \mathbf{w}_1 \rangle = \lambda \langle \mathbf{v}_1, \mathbf{w}_1 \rangle = 0,$$

so  $A(W_1) \subseteq W_1$  is an invariant subspace, etc.

Suppose now that  $A$  is symmetric. Then there is an eigenvector of norm 1,  $\mathbf{v} \in V$ .

$$\lambda = \lambda \langle \mathbf{v}, \mathbf{v} \rangle = \langle A\mathbf{v}, \mathbf{v} \rangle = \langle \mathbf{v}, A\mathbf{v} \rangle = \bar{\lambda} \implies \lambda \in \mathbb{R}.$$

**Lemma:**  $\{A_i\}$  pairwise commute  $\iff$  they are all simultaneously diagonalizable.

*Proof:* By induction on number of operators

- $A_n$  is diagonalizable, so  $V = \bigoplus E_i$  a sum of eigenspaces
- Restrict all  $n - 1$  operators  $A$  to  $E_n$ .
- The commute in  $V$  so they commute in  $E_n$
- **(Lemma)** They were diagonalizable in  $V$ , so they're diagonalizable in  $E_n$
- So they're simultaneously diagonalizable by I.H.
- But these eigenvectors for the  $A_i$  are all in  $E_n$ , so they're eigenvectors for  $A_n$  too.
- Can do this for each eigenspace. ■

Full details here

### Theorem (Characterizations of Diagonalizability)

$M$  is diagonalizable over  $\mathbb{F} \iff \min_M(x, \mathbb{F})$  splits into distinct linear factors over  $\mathbb{F}$ , or equivalently iff all of the roots of  $\min_M$  lie in  $\mathbb{F}$ .

*Proof:*  $\implies$  : If  $\min_A$  factors into linear factors, so does each invariant factor, so every elementary divisor is linear and  $JCF(A)$  is diagonal.

$\impliedby$  : If  $A$  is diagonalizable, every elementary divisor is linear, so every invariant factor factors into linear pieces. But the minimal polynomial is just the largest invariant factor.

## 6.7 Matrix Counterexamples

1. A matrix that is:

- Not diagonalizable over  $\mathbb{R}$  but diagonalizable over  $\mathbb{C}$
- No eigenvalues in  $\mathbb{R}$  but distinct eigenvalues over  $\mathbb{C}$
- $\min_M(x) = \chi_M(x) = x^2 + 1$

$$M = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix} \sim \left[ \begin{array}{c|c} -1\sqrt{-1} & 0 \\ \hline 0 & 1\sqrt{-1} \end{array} \right].$$

2.

$$M = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}.$$

- Not diagonalizable over  $\mathbb{C}$

- Eigenvalues  $[1, 1]$  (repeated, multiplicity 2)
  - $\min_M(x) = \chi_M(x) = x^2 - 2x + 1$
3. Non-similar matrices with the same characteristic polynomial

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \text{ and } \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

4. A full-rank matrix that is not diagonalizable:

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}.$$

5. Matrix roots of unity:

$$\sqrt{I_2} = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}.$$

$$\sqrt{-I_2} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}.$$

## 6.8 Miscellaneous

**Lemma:**  $I \trianglelefteq R$  is a free  $R$ -module iff  $I$  is a principal ideal.

Proof:  $\implies$  :

Suppose  $I$  is free as an  $R$ -module, and let  $B = \{\mathbf{m}_j\}_{j \in J} \subseteq I$  be a basis so we can write  $M = \langle B \rangle$ .

Suppose that  $|B| \geq 2$ , so we can pick at least 2 basis elements  $\mathbf{m}_1 \neq \mathbf{m}_2$ , and consider

$$\mathbf{c} = \mathbf{m}_1\mathbf{m}_2 - \mathbf{m}_2\mathbf{m}_1,$$

which is also an element of  $M$ .

Since  $R$  is an integral domain,  $R$  is commutative, and so

$$\mathbf{c} = \mathbf{m}_1\mathbf{m}_2 - \mathbf{m}_2\mathbf{m}_1 = \mathbf{m}_1\mathbf{m}_2 - \mathbf{m}_1\mathbf{m}_2 = \mathbf{0}_M$$

However, this exhibits a linear dependence between  $\mathbf{m}_1$  and  $\mathbf{m}_2$ , namely that there exist  $\alpha_1, \alpha_2 \neq 0_R$  such that  $\alpha_1\mathbf{m}_1 + \alpha_2\mathbf{m}_2 = \mathbf{0}_M$ ; this follows because  $M \subset R$  means that we can take  $\alpha_1 = -m_2, \alpha_2 = m_1$ . This contradicts the assumption that  $B$  was a basis, so we must have  $|B| = 1$  and so  $B = \{\mathbf{m}\}$  for some  $\mathbf{m} \in I$ . But then  $M = \langle B \rangle = \langle \mathbf{m} \rangle$  is generated by a single element, so  $M$  is principal.

$\impliedby$  :

Suppose  $M \trianglelefteq R$  is principal, so  $M = \langle \mathbf{m} \rangle$  for some  $\mathbf{m} \neq \mathbf{0}_M \in M \subset R$ .

Then  $x \in M \implies x = \alpha\mathbf{m}$  for some element  $\alpha \in R$  and we just need to show that  $\alpha\mathbf{m} = \mathbf{0}_M \implies \alpha = 0_R$  in order for  $\{\mathbf{m}\}$  to be a basis for  $M$ , making  $M$  a free  $R$ -module.

But since  $M \subset R$ , we have  $\alpha, m \in R$  and  $\mathbf{0}_M = 0_R$ , and since  $R$  is an integral domain, we have  $\alpha m = 0_R \implies \alpha = 0_R$  or  $m = 0_R$ .

Since  $m \neq 0_R$ , this forces  $\alpha = 0_R$ , which allows  $\{\mathbf{m}\}$  to be a linearly independent set and thus a basis for  $M$  as an  $R$ -module. ■



---

## 7 Extra Problems

### 7.1 Group Theory

#### 7.1.1 Basic Structure

15. Show that if  $G/Z(G)$  is cyclic then  $G$  is abelian.
16. Show that the intersection of two subgroups is again a subgroup.
17. Show that if  $G \curvearrowright X$  is a group action, then the stabilizer  $G_x$  of a point is a subgroup.
18. Show that  $G = H \times K$  iff the conditions for recognizing direct products hold.
19. Show that if  $H, K \trianglelefteq G$  and  $H \cap K = \emptyset$ , then  $hk = kh$  for all  $h \in H, k \in K$ .
20. Show that every normal subgroup of  $G$  is contained in  $Z(G)$ .
21. Show that  $|G|/|H| = [G : H]$ .
22. Show that the order of any element in a group divides the order of the group.
23. Show that  $\varphi(n) = n \prod p \mid n \left(1 - \frac{1}{p}\right)$ .
24. Show that  $Z(G) \subseteq C_G(H) \subseteq N_G(H)$ .

#### 7.1.2 Primes in Group Theory

14. Show that any group of prime order is cyclic and simple.
15. Analyze groups of order  $p^2$ .
16. Analyze groups of order  $pq$ .
17. Show that a group of order  $pq$  with  $q < p$  and  $q$  not dividing  $p - 1$  is cyclic of order  $pq$ .
18. Analyze groups of order  $p^2q$ .
19. Show that no group of order  $p^2q^2$  is simple for  $p < q$  primes.
20. Show that a group of order  $p^2q^2$  has a normal Sylow subgroup.
21. Show that a group of order  $p^2q^2$  where  $q$  does not divide  $p^2 - 1$  and  $p$  does not divide  $q^2 - 1$  is abelian.
22. Show that every group of order  $pqr$  with  $p < q < r$  primes contains a normal Sylow subgroup.
23. Show that every  $p$ -group is nilpotent.
24. Show that every  $p$ -group is solvable.
25. Show that  $p$ -groups have nontrivial centers.
26. Show that any normal  $p$ -subgroup is contained in every Sylow  $p$ -subgroup of  $G$ .

#### 7.1.3 Classification

10. Show that no group of order 36 is simple.
11. Show that no group of order 90 is simple.
12. Show that all groups of order 45 are abelian.

#### 7.1.4 Series


6. Show that  $A_n$  is simple for  $n \geq 5$
7. Give a necessary and sufficient condition for a cyclic group to be solvable.
8. Prove that every simple abelian group is cyclic.

## 7.2 Ring Theory

1. Show that if  $x \in R$  a PID, then  $x$  is irreducible  $\iff \langle x \rangle \trianglelefteq R$  is maximal.

## 7.3 Field Theory

1. What is  $[\mathbb{Q}(\sqrt{2} + \sqrt{3}) : \mathbb{Q}]$ ?
2. What is  $[\mathbb{Q}(2^{\frac{3}{2}}) : \mathbb{Q}]$ ?
3. Show that every field is simple.
4. Show that any field morphism is either 0 or injective.
5. Show that if  $p \in \mathbb{Q}[x]$  and  $r \in \mathbb{Q}$  is a rational root, then in fact  $r \in \mathbb{Z}$ .
6. If  $\{\alpha_i\}_{i=1}^n \subset F$  are algebraic over  $K$ , show that  $K[\alpha_1, \dots, \alpha_n] = K(\alpha_1, \dots, \alpha_n)$ .
7. Show that the Galois group of  $x^n - 2$  is  $D_n$ , the dihedral group on  $n$  vertices.
8. Compute all intermediate field extensions of  $\mathbb{Q}(\sqrt{2}, \sqrt{3})$ , show it is equal to  $\mathbb{Q}(\sqrt{2} + \sqrt{3})$ , and find a corresponding minimal polynomial.



9. Compute all intermediate field extensions of  $\mathbb{Q}(2^{\frac{1}{4}}, \zeta_8)$ .
10. Show that  $\mathbb{Q}(2^{\frac{1}{3}})$  and  $\mathbb{Q}(\zeta_3 2^{\frac{1}{3}})$
11. Show that if  $L/K$  is separable, then  $L$  is normal  $\iff$  there exists a polynomial  $p(x) = \prod_{i=1}^n (x - \alpha_i) \in K[x]$  such that  $L = K(\alpha_1, \dots, \alpha_n)$  (so  $L$  is the splitting field of  $p$ ).
12. Is  $\mathbb{Q}(2^{\frac{1}{3}})/\mathbb{Q}$  normal?
13. Show that any finite integral domain is a field.
14. Prove that if  $R$  is an integral domain, then  $R[t]$  is again an integral domain.
15. Show that  $f(R[t]) = f(R)(t)$ .
16. Prove that  $x^{p^n} - x$  is the product of all monic irreducible polynomials in  $\mathbb{F}_p[x]$  with degree dividing  $n$ .
17. Prove that an irreducible  $\pi(x) \in \mathbb{F}_p[x]$  divides  $x^{p^n} - x \iff \deg \pi(x)$  divides  $n$ .
18. Show that a field with  $p^n$  elements has exactly one subfield of size  $p^d$  for every  $d$  dividing  $n$ .
19. Show that  $\mathbb{GF}(p^n)$  is the splitting field of  $x^{p^n} - x \in \mathbb{F}_p[x]$ .
20. Show that  $x^{p^d} - x \mid x^{p^n} - x \iff d \mid n$
21. Show that  $\mathbb{GF}(p^d) \leq \mathbb{GF}(p^n) \iff d \mid n$
22. Show that  $x^{p^n} - x = \prod f_i(x)$  over all irreducible monic  $f_i$  of degree  $d$  dividing  $n$ .
23. Compute the Galois group of  $x^n - 1 \in \mathbb{Q}[x]$  as a function of  $n$ .

24. Identify all of the elements of the Galois group of  $x^p - 2$  for  $p$  an odd prime (note: this has a complicated presentation).
25. Classify the five groups of order 12.
26. Classify the four groups of order 28.
27. Show if  $G$  is finite, then  $G$  is solvable  $\iff$  all of its composition factors are of prime order.
28. Show that if  $N$  and  $G/N$  are solvable, then  $G$  is solvable.
29. Show that  $S_n$  is generated by disjoint cycles.
30. Show that  $S_n$  is generated by transpositions.
31. Show that an  $m$ -cycle is an odd permutation iff  $m$  is an even number.
32. Show that a permutation is odd iff it has an odd number of even cycles.
33. Prove Burnside's theorem.
- 34.

## 7.4 Modules and Linear Algebra

1. Prove the Cayley-Hamilton theorem.
2. Prove that the minimal polynomial divides the characteristic polynomial.
3. Prove that the cokernel of  $A \in \text{Mat}(n \times n, \mathbb{Z})$  is finite  $\iff \det A \neq 0$ , and show that in this case  $|\text{coker}(A)| = |\det(A)|$ .
4. Show that a nilpotent operator is diagonalizable.
5. Show that if  $A, B$  are diagonalizable and  $[A, B] = 0$  then  $A, B$  are simultaneously diagonalizable.
6. Does diagonalizable imply invertible? The converse?

## 7.5 Commutative Algebra

1. Show that a finitely generated module over a Noetherian local ring is flat iff it is free using Nakayama and Tor.

## 8 List of Topics

Chapters 1-9 of Dummit and Foote

- Left and right cosets
- Lagrange's theorem
- Isomorphism theorems
- Group generated by a subset
- Structure of cyclic groups
- Composite groups
  - $HK$  is a subgroup iff  $HK = KH$
- Normalizer
  - $HK \leq H$  if  $H \leq N_G(K)$
- Symmetric groups
  - Conjugacy classes are determined by cycle types
- Group actions
  - Actions of  $G$  on  $X$  are equivalent to homomorphisms from  $G$  into  $\text{Sym}(X)$
- Cayley's theorem

- 
- Orbits of an action
  - Orbit stabilizer theorem
  - Orbits act on left cosets of subgroups
  - Subgroups of index  $p$ , the smallest prime dividing  $|G|$ , are normal
  - Action of  $G$  on itself by conjugation
  - Class equation
  - $p$ -groups
    - Have non trivial center
  - $p^2$  groups are abelian
  - Automorphisms, the automorphism group
    - Inner automorphisms
    - $\text{Inn}(G) \cong Z/Z(G)$
    - $\text{Aut}(S_n) = \text{Inn}(S_n)$  unless  $n = 6$
    - $\text{Aut}(G)$  for cyclic groups
    - $G \cong Z_p^n$ , then  $\text{Aut}(G) \cong GL_n(Z_p)$
  - Proof of Sylow theorems
  - $A_n$  is simple for  $n \geq 5$
  - Recognition of internal direct product
  - Recognition of semi-direct product
  - Classifications:
    - $pq$
  - Free group & presentations
  - Commutator subgroup
  - Solvable groups
    - $S_n$  is solvable for  $n \leq 4$
  - Derived series
    - Solvable iff derived series reaches  $e$
  - Nilpotent groups
    - Nilpotent iff all sylow- $p$  subgroups are normal
    - Nilpotent iff all maximal subgroups are normal
  - Upper central series
    - Nilpotent iff series reaches  $G$
  - Lower central series
    - Nilpotent iff series reaches  $e$
  - Fratini's argument
  - Rings
    - $I$  maximal iff  $R/I$  is a field
    - Zorn's lemma
    - Chinese remainder theorem
    - Localization of a domain
    - Field of fractions
    - Factorization in domains
    - Euclidean algorithm
    - Gaussian integers
    - Primes and irreducibles
    - Domains
      - \* Primes are irreducible
    - UFDs

- 
- \* Have GCDs
  - \* Sometimes PIDs
  - PIDs
    - \* Noetherian
    - \* Irreducibles are prime
    - \* Are UFDs
    - \* Have GCDs
  - Euclidean domains
    - \* Are PIDs
  - Factorization in  $\mathbb{Z}[i]$
  - Polynomial rings
  - Gauss' lemma
  - Remainder and factor theorem
  - Polynomials
  - Reducibility
  - Rational root test
  - Eisenstein's criterion

## 9 Groups

### 9.1 Definitions

#### 9.1.1 Subgroup Generated by a set $A$

- $\langle A \rangle = \{a_1^{\pm 1}, a_2^{\pm 1}, \dots, a_n^{\pm 1} : a_i \in A, n \in \mathbb{N}\}$
- Equivalently, the intersection of all  $H$  such that  $A \subseteq H \leq G$

#### 9.1.2 Free Group on a set $X$

- Equivalently, words over the alphabet  $X$  made into a group via concatenation

#### 9.1.3 Centralizer of an element or a subgroup

- $C_G(a) = \{g \in G : ga = ag\}$
- 

$$C_G(H) = \{g \in G : \forall h \in H, gh = hg\} = \bigcap_{h \in H} C_G(h)$$

- Note - requires the same  $g$  on both sides!

- Facts:
  - $C_G(H) \leq G$
  - $C_G(H) \trianglelefteq N_G(H)$
  - $C_G(G) = Z(G)$
  - $C_H(a) = H \cap C_G(a)$

#### 9.1.4 Center of a group

- $Z(G) = \{g \in G : \forall x \in G, gx = xg\}$

- Facts

—

$$Z(G) = \bigcap_{a \in G} C_G(a)$$

### 9.1.5 Normalizer of a subgroup

- 

$$N_G(H) = \{g \in G : gHg^{-1} = H\}$$

- Equivalently,  $\bigcup \{K : H \trianglelefteq K \leq G\}$  (the largest  $K \leq G$  for which  $H \trianglelefteq K$ )
- Equivalently, the stabilizer of  $H$  under  $G$  acting on its subgroups via conjugation
- Differs from centralizer; can have  $gh = h'g$
- Facts:
  - $C_G(H) \subseteq N_G(H) \leq G$
  - $N_G(H)/C_G(H) \cong A \leq \text{Aut}(H)$
  - Given  $H \subseteq G$ , let

$$S(H) = \bigcup_{g \in G} gHg^{-1}$$

, so  $|S(H)|$  is the number of conjugates to  $H$ . Then

$$|S(H)| = [G : N_G(H)]$$

\* i.e. the number of subgroups conjugate to  $H$  equals the index of the normalizer of  $H$ .

### 9.1.6 Normal Core of a subgroup

- 

$$H_G = \bigcap_{g \in G} gHg^{-1}$$

- Equivalently,  $H_G = \langle N : N \trianglelefteq G \text{ \& } N \leq H \rangle$ 
  - Largest normal subgroup that contains  $H$
- Equivalently,  $H_G = \ker \psi$  where  $\psi : G \rightarrow \text{Sym}(G/H)$ ;  $g \sim (xH) = (gx)H$
- Facts:
  - $H_G \trianglelefteq G$  and is an idempotent operation

### 9.1.7 Normal Closure of a subgroup

- $H^G = \{gHg^{-1} : g \in G\}$
- Equivalently,

$$H^G = \bigcap \{N : H \leq N \trianglelefteq G\}$$

- (The smallest normal subgroup of  $G$  containing  $H$ )

**9.1.8 Group Action of a group on a set**

- Given as a function

$$\begin{aligned}\varphi : G \times X &\rightarrow X \\ (g, x) &\mapsto g \sim x\end{aligned}$$

which gives rise to a function

$$\begin{aligned}\varphi_g : X &\rightarrow X \\ x &\mapsto g \sim x\end{aligned}$$

(which is a bijection) where  $\sim$  denotes a group element acting on a set element, and  $\forall x \in X$ ,

- $e \sim x = x$
- $(gh) \sim x = g \sim (h \sim x)$

- Equivalently, a function

$$\begin{aligned}\psi : G &\rightarrow \text{Sym}(X) \\ g &\mapsto \varphi_g\end{aligned}$$

–

$$\ker \psi = \bigcap_{x \in X} G_x$$

(intersection of all stabilizers)

- Interesting actions:
  - Left multiplication of  $G$  on  $G$ :

$$\begin{aligned}\varphi : G &\rightarrow G \rightarrow G \\ g &\mapsto \varphi_g : G \rightarrow G \\ h &\mapsto gh\end{aligned}$$

\*  $\mathcal{O}_x = G$  (transitive)

\*  $G_x = e$

- $G$  acting via conjugation on itself:

$$\begin{aligned}\varphi : G &\rightarrow G \rightarrow G \\ g &\mapsto \psi_g : G \rightarrow G \\ h &\mapsto ghg^{-1}\end{aligned}$$

\* A common notation is  $x^g = g^{-1}xg$  which obeys  $(x^g)^h = x^{gh}$

\*  $\mathcal{O}_x = [x]$  (Conjugacy classes, so not generally transitive)

\*  $G_x = \{g \in G : gxg^{-1} = g\} = C_G(x)$

- $G$  acting on  $S = \{H : H \leq G\}$  via conjugation:

\*

$$\begin{aligned}\varphi : G &\rightarrow S \rightarrow S \\ g &\mapsto \psi_g : S \rightarrow S \\ H &\mapsto gHg^{-1}\end{aligned}$$

\*  $\mathcal{O}_H = [H] = \{gHg^{-1} : g \in G\}$ , conjugate subgroups of  $H$

\*  $G_x = N_G(H) = \{g \in G : gHg^{-1} = H\}$

**9.1.9 Transitive group actions**

- $\forall x, y \in X, \exists g \in G : g \cdot x = y$
- Equivalent, actions with a single orbit

**9.1.10 Orbit of an Element of a Set**

$$\mathcal{O}_x = \{g \cdot x \mid x \in X\} = \bigcup_{g \in G} \{g \cdot x\}.$$

- The set of all orbits is denoted  $X/G = \{\mathcal{O}_x \mid x \in X\}$
- Partitions  $X$  according to the equivalence relation  $x \cong y \iff \exists g \in G : g \cdot x = y$
- $G$  acts transitively on  $X$  if restricted to any single orbit

**9.1.11 Stabilizer of an Element of a Set**

- $G_x = \{g \in G : g \cdot x = x\}$
- Facts:
  - $G_x \leq G$ , not usually normal
  - $x, y \in \mathcal{O}_x \Rightarrow G_x$  is conjugate to  $G_y$

**9.1.12 Automorphisms of a group**

- $\text{Aut}(G) = \{\varphi : G \rightarrow G : \varphi \text{ is an isomorphism}\}$

**9.1.13 Inner Automorphisms of a group**

- $\text{Inn}(G) = \{\varphi_g \in \text{Aut}(G) : \varphi_g(x) = gxg^{-1}\}$
- Also consider the map

$$\begin{aligned}\psi : G &\rightarrow \text{Aut}(G) \\ g &\mapsto (\lambda : x \mapsto gxg^{-1})\end{aligned}$$

Then  $\text{im}\psi = \text{Inn}(G)$ ,  $\ker\psi = Z(G)$

- Facts:
  - $\text{Inn}(G) \trianglelefteq \text{Aut}(G)$
  - $\text{Inn}(G) \cong G/Z(G)$

**9.1.14 Outer Automorphisms of a group**

- $\text{Out}(G) = \text{Aut}(G)/\text{Inn}(G)$



**9.1.15 Conjugacy Class of an element**

- 

$$[a] = \{gag^{-1} : g \in G\} = \bigcup_{g \in G} \{gag^{-1}\}$$

- Equivalently,  $[a] = \mathcal{O}_a$  under  $G$  acting on itself via conjugation
- Facts:
  - Equivalence relation, partitions the group
  - $|[a]|$  divides  $|G|$
  - $a \in Z(G) \Rightarrow [a] = \{a\}$

**9.1.16 Characteristic subgroup**

- $H \text{ char } G \iff \forall \varphi \in \text{Aut}(G), \varphi(H) = H$ 
  - i.e.,  $H$  is fixed by all automorphisms of  $G$ .

**9.1.17 Simple group**

- $G$  is simple  $\iff H \trianglelefteq G \Rightarrow H = e \text{ or } G$ 
  - No non-trivial normal subgroups

**9.1.18 Commutator of an element, or of subgroups**

- $[g, h] = ghg^{-1}h^{-1}$
- $[G, H] = \langle [g, h] : g \in G, h \in H \rangle$  (Subgroup generated by commutators)

**9.2 Structural Results**

- Cyclic  $\Rightarrow$  abelian
- $G/Z(G)$  cyclic  $\Rightarrow G$  is abelian
- Intersections of subgroups are also subgroups

**9.2.1 Isomorphisms Theorems**

-\*First Isomorphism Theorem\*\*

- Conditions:
  - $\varphi : G \rightarrow G'$  is a homomorphism.
- Result:
  - $\ker \varphi \trianglelefteq G$
  - $\text{im } \varphi \leq G'$
  - $G/\ker \varphi \cong \text{im } \varphi$ .
- Corollaries:
  - $\ker \varphi = e \Rightarrow G \cong G'$

-\*Second Isomorphism Theorem\*\*

- Conditions:

- $N \trianglelefteq G, H \leq G$
- Results:
  - $HN \leq G$
  - $N \cap H \trianglelefteq H$
  -

$$\frac{H}{H \cap N} \cong \frac{HN}{N}$$

- Corrolaries:
  - (Weaker) Relaxing  $N \trianglelefteq G$  to  $H \subseteq N(N)$  yields
    - \*  $N \cap H \subseteq G$  (Not normal)
    - \*  $N \cap H \trianglelefteq H$

-\*Third Isomorphism Theorem\*\*

- Conditions:
  - $N \trianglelefteq G, N \leq A \leq G$
- Results:
  - $A/N \leq G/N$
  - \* Every subgroup of  $G/N$  is of this form for *some* such  $A$

$$\frac{G/N}{A/N} \cong \frac{G}{A}$$

- \* Cancel the  $N$ !
- Corrolaries:
  - $A \trianglelefteq G \Rightarrow A/N \trianglelefteq G/N$
  - \* All normal subgroups of  $G/N$  are of this form for some  $A$ .

### 9.3 Misc Results

- $G/N$  is abelian  $\iff [G, G] \leq N$
- $HK$  is not always a subgroup - see conditions in 2nd Isomorphism theorem'
- $H \trianglelefteq G, K \trianglelefteq G$  and  $H \cap K = e \Rightarrow hk = kh \forall h \in H, k \in K$ 
  - Normal subgroups with trivial intersection commute
- $H \text{ char } G \Rightarrow H \trianglelefteq G$ 
  - Characteristic is a strictly stronger condition than normality
- $H \text{ char } K \text{ char } G \Rightarrow H \text{ char } G$ 
  - Characteristic is transitive
- $H \leq G, K \trianglelefteq G, H \text{ char } K \Rightarrow H \trianglelefteq G$ 
  - i.e., normality is **not** transitive, strengthening normality to char gives “weak transitivity”
- Recognizing (Internal) Direct Products:  $H \leq G, K \leq G$ 
  - $H \cap K = e$
  - $\forall g \in G, \exists h \in H, k \in K : g = hk$
  - $H \trianglelefteq G, K \trianglelefteq G$ 
    - \* **OR** Every element in  $H$  commutes with every element in  $K$
- P Groups
  - $\bigcap P = O_P(G) \text{ char } G$ . And  $O_P(G) \trianglelefteq G$  as well.

- $N \trianglelefteq G$  implies that  $P_N \leq N$  are of the form  $N \bigcap P_G$
- $P \bigcap Q = e$

## 9.4 Numeric Results

### 9.4.1 Cauchy's Theorem

- For any  $p$  dividing  $|G|$ , there is a subgroup of order  $p$ .

### 9.4.2 Sylow Theorems: $|G| = p^k m$ where $p \nmid m$

- At least one Sylow- $p$  subgroup always exists:  $\exists P \leq G$  with  $|P| = p^k$
- All such subgroups are conjugate:  $\forall P, P', \exists g \in G : gPg^{-1} = P'$
- $n_p$  satisfies:
  - $n_p$  divides  $m = [G : P]$
  - $n_p \equiv 1 \pmod{p}$
  - $n_p = [G : N_G(P)]$  (Not as useful)
- Every  $p$ -subgroup of  $G$  is a  $p$ -subgroup of  $P$  (i.e.  $P$  is maximal and contains all subgroups of order  $p^l$  with  $l \leq k$ )

### 9.4.3 Orbit-stabilizer Theorem

- Given a group action,  $G/G_x \cong \mathcal{O}_x$
- Gives the numeric result  $|\mathcal{O}_x| = |G/G_x| = [G : G_x] = \frac{|G|}{|G_x|}$
- Also useful in the form  $|G| = |\mathcal{O}_x| |G_x|$
- Proof:
  - Use the map

$$\begin{aligned}\varphi : G &\rightarrow X \\ g &\mapsto g \sim x\end{aligned}$$

Where  $\text{im}\varphi = \mathcal{O}_x$  and  $\ker\varphi = G_x$ .

### 9.4.4 Burnside's Lemma

•

$$|X_G| = \frac{1}{|G|} \sum_{g \in G} |X^g|$$

- $|X_G|$  is the number of orbits
- $X^g = \{x \in X : g \sim x = x\}$

### 9.4.5 The class equation

- 

$$|G| = |Z(G)| + \sum_{a \in A} [G : C_G(a)]$$

- Where  $A = \{a_1, a_2, \dots, a_n : a_1 \in [a_1], a_2 \in [a_2], \dots\}$  is a set containing one element from each conjugacy class
- $[G : C_G(a)]$  is the number of elements in  $[a]$
- Each element in  $Z(G)$  has a singleton conjugacy class

### 9.4.6 General facts

- $|G| = p \Rightarrow G$  is cyclic
- $|G| = p^e \Rightarrow Z(G) \neq e$
- $|G| = p^e$  (P-groups)
  - $Z(G) \neq \{e\}$  (Use class equation)
- $|G| = p$ 
  - Always cyclic
    - \* Proof: Any nontrivial cyclic subgroup's order is  $> 1$  and divides  $p$ , so equals  $p$ .
- $|G| = p^2$ 
  - Always abelian
    - \* Proof:  $|G/Z(G)| = 1, p$ . If  $p$ , it's cyclic, and  $G$  is abelian. Otherwise it's 1, so  $G = Z(G)$ .
  - Two possibilities:
    - \*  $Z_{p^2}$  (cyclic)
    - \*  $Z_p \times Z_p$
- $|G| = pq$ 
  - $p \nmid q-1$  ( $q \not\equiv 1 \pmod p$ ):
    - \* One possibility:
      - $G \cong Z_{pq}$  (cyclic)
    - \* Facts:
      - $\exists P \trianglelefteq G$  (A Sylow- $P$  subgroup)
  - $p$  divides  $q-1$  ( $q \equiv 1 \pmod p$ ):
    - \* Two possibilities:
      - $G \cong Z_{pq}$  (cyclic)
      - $G \cong Z_q \rtimes Z_p$
  - Never simple
- $|G| = p^2q$ 
  - $\exists P \trianglelefteq G$  (A Sylow- $P$  subgroup)
- $|G| = p_1p_2p_3$  (distinct)

- Not simple

## 9.5 Common Groups

### 9.5.1 $S_3$

$$S_3 = \langle (12), (23), (13) \rangle$$

- $Z(S_3) = e$
- $\text{Aut}(S_3) = \text{Inn}(S_3)$ , since

$$\begin{aligned} Z(G) &= e = \ker \psi \\ \Rightarrow \text{Out}(S_3) &= \text{Inn}(S_3) \\ \Rightarrow \text{Aut}(S_3) &\cong S_3 \end{aligned}$$

### 9.5.2 $S_n$

$$S_n, n \geq 4$$

- $Z(S_n) = e$ 
  - Let  $\sigma(a) = b$ , choose  $\tau = (bc)$  so  $\tau\sigma(a) = \tau(b) = c \neq b = \sigma(a)$
- Conjugacy classes are determined entirely by cycle structure
  - There are exactly  $p(n)$  of them (partition function)
- Disjoint cycles commute
- $\sigma \circ (a_1 \cdots a_k) \circ \sigma^{-1} = (\sigma(a_1), \cdots \sigma(a_k))$
- Every element is a product of disjoint cycles
- Every element is a product of transpositions
  - A cycle of length  $k$  can be written as  $k - 1$  transpositions
  - Parity of the cycle equals the parity of  $k - 1$ .
- The order of an element is the lcm of the size of the cycles.

### 9.5.3 $A_n$

- Simple for  $n \geq 5$
- Index 2 in  $S_n$ , so  $A_n \trianglelefteq S_n$

### 9.5.4 $D_n$

- $\langle a, b \mid a^n = b^2 = 1, bab^{-1} = a^{-1} \rangle \cong \langle r, s \rangle$
- $D_n/N$  is always another dihedral group for any  $N \trianglelefteq D_n$
- All subgroups:
  - $\langle r^d \rangle \cong Z_{n/d}$  where  $d$  divides  $n$  (index  $2d$ )
  - $\langle r^d, r^i s \rangle \cong D_{\frac{n}{d}}$  where  $d$  divides  $n$  and  $0 \leq i \leq d - 1$  (index  $d$ )
  - \* All dihedral

---

## 10 Rings

### 10.1 Facts about ideals:

- Intersections, products, and sums of ideals are ideals
- Not necessarily unions
- Every ring has proper maximal ideals
- Apply Z.L. to  $\{I \trianglelefteq R : I \neq R\}$
- Every proper ideal is contained in a maximal ideal

### 10.2 Maximal ideals

$I \trianglelefteq R$  *maximal* if  $\nexists J \trianglelefteq R : I \subset J \subset R$

- Every nonzero ring has a maximal ideal (Krull's Theorem)
- $R$  commutative  $\implies R/I$  a field
- Union of maximal ideals  $= R - R^\times$
- $(X - a) \trianglelefteq R[X]$  is maximal for  $a \in R$

### 10.3 Prime ideals

$I \trianglelefteq R$  *prime* when  $pq \in I \implies p \in I \vee q \in I$

- $I$  prime  $\iff R/I$  an integral domain,
- (maximal  $\implies$  prime)
- $\text{rad}(I^n) = I$

### 10.4 Radicals

$I \trianglelefteq R$  *radical* when  $\forall a \in R, a^n \in I \implies a \in I$

- The nilradical:  $\text{nilrad}(I) = \bigcap P$  such that  $P \trianglelefteq R$  is prime
- $\text{rad}(I) = \{x \in R \mid \exists n : x^n \in I\}$
- $\text{rad}(0) = \text{nilrad}(R)$
- $\text{rad}(IJ) = \text{rad}(I) \cap \text{rad}(J)$
- $\text{rad}(I) = \bigcap J$  such that  $I \subset J, J$  prime (i.e. intersection of all prime ideals containing  $I$ )

### 10.5 Other ideals

- $I \trianglelefteq R$  *primary* when  $pq \in I \implies a \in I \vee \exists n \in \mathbb{N} : b^n \in I$
- Prime  $\implies$  primary
- $I \trianglelefteq R$  *principal* when  $\exists a \in R : I = \langle a \rangle$
- $I \trianglelefteq R$  *irreducible* when  $\nexists \{J \trianglelefteq R : I \subset J\} : I = \bigcap J$
- $I \subset R \iff 1, u \notin I (u \in R^\times)$

- $\{I : I \trianglelefteq R\}$  is a poset
- Zorn's lemma can be applied to  $\{I \trianglelefteq R : 1 \notin I\}$
- Every proper ideal is contained in a maximal ideal.
- Facts about units
- $R^\times$  is closed under multiplication, but *not* under addition.
- $R - R^\times$  an additive group  $\iff R$  is a local ring
- Integral Domain
- Principal Ideal Domain
- (Prime  $\implies$  maximal)  $\implies$  UFD
- Unique Factorization Domain
- Field
- When  $(0)$  is the only proper ideal
- $R/M$  a field  $\iff M$  maximal
- Localization
- Zorn's Lemma: For every poset  $P$ , every chain in  $P$  has an upper bound  $\implies P$  has a maximal element.
- Noetherian: Every ideal is finitely generated
- iff the ascending chain condition for ideals holds

## 10.6 Orders less than 16:

(Normal: Diamond, grouped by conjugacy class)

- 1 (The trivial group)
  - $Z_1 = \{e\}$
- 2 (One group)
  - $Z_2 \cong Z_3^\times \cong Z_4^\times \cong Z_6^\times$
  - $= \{e, a\}$
  - \* Cyclic
  - \* One element of order 2
- 3 (One group)
  - $Z_3 \cong A_3$
  - $\cong \{(), (123), (132)\}$
  - \* Cyclic
  - \* One element of order 3
- 4 (Two groups, both abelian)



Figure 1: img

- $Z_4 \cong Z_5^\times \cong Z_8^\times \cong Z_{10}^\times \cong Z_{12}^\times$ 
  - \* Cyclic
  - \* One element of order 4
- $Z_2 \times Z_2 \cong V_4 \cong D_2 \cong Z_8^\times$ , which are all isomorphic to  $\langle a, b \mid a^2 = b^2 = (ab)^2 = e \rangle \cong \langle (12)(34), (13)(24), (14)(23) \rangle$ 
  - \* Not cyclic, but abelian
  - \* All elements have order 2
  - \*  $V_4 \trianglelefteq A_4 \leq S_4$
- 5 (One group)
  - $Z_5$ 
    - \* Cyclic, one element of order 5
- 6 (Two groups)
  - $Z_6 \cong Z_7^\times \cong Z_9^\times \cong Z_{14}^\times$ 
    - \* Cyclic, one element of order 6
  - $S_3 \cong D_6$






Figure 2: File:S3latticeofsubgroups.png

$$\cong \langle a, b, c \mid a^2 = b^2 = c^3 = abc = e \rangle$$

\* Non-abelian (smallest one)

- 7 (One group)
  - $Z_7$ 
    - \* Cyclic, one element of order 7
- 8 (**Five groups**)



- $Z_8 \cong Z_{15}^\times \cong Z_{16}^\times$  (cyclic)
- $Z_2 \times Z_4$







Figure 3: img




- \* Abelian, one element of order 4
- $Z_2 \times Z_2 \times Z_2$



- \* Abelian, every element has order 2
- $D_8 \cong \langle r, s \mid r^4 = s^2 = e, srs^{-1} = r^{-1} \rangle$



- $\cong \{(), (1234), (13)(24), (1432), (13)(24), (14)(23), (12)(34)\} \leq S_4$
- $Q_8 \cong \langle i, j, k \mid i^2 = j^2 = k^2 = ijk \rangle$
- $\cong \langle a, b, c \mid a^4 = b^4 = e, a^2 = b^2, ba = a^3b \rangle$
- \* Every element has order 4



- \* All subgroups are normal, but not abelian

- 9 (*Two groups*)
  - $Z_9$
  - $Z_3 \times Z_3$
- 10 (*Two groups*)
  - $Z_{10} \cong Z_{11}^\times$
  - $D_{10}$
- 11 (One group)
  - $Z_{11}$
- 12 (**Five groups**)
  - $Z_{12} \cong Z_{13}^\times$
- 13 (One group)
  - $Z_{13}$
- 14 (*Two groups*)
  - $Z_{14}$
- 15 (One group)
  - $Z_{15}$
- 16 (***Fourteen groups!***)