UGA Algebra Qualifying Exam Questions (Spring 2011 – Spring 2021)

D. Zack Garza

Table of Contents

Contents

Table of Contents 2

1 Group Theory: General 11
- 1.1 Spring 2020 #2 11
- 1.2 Spring 2019 #4 11
  - 1.2.1 a 11
  - 1.2.2 b 11
  - 1.2.3 c 11
- 1.3 Spring 2012 #2 12
  - 1.3.1 a 12
  - 1.3.2 b 12
- 1.4 Spring 2017 #1 12
- 1.5 Fall 2016 #1 12
- 1.6 Fall 2015 #1 13
- 1.7 Spring 2015 #1 13
- 1.8 Fall 2014 #6 13
- 1.9 Spring 2013 #3 13
- 1.10 Fall 2019 Midterm #1 13
- 1.11 Fall 2019 Midterm #4 13
- 1.12 Fall 2019 Midterm #5 14
- 1.13 Spring 2021 #2 14

2 Groups: Sylow Theory 14
- 2.1 Fall 2019 #1 14
- 2.2 Fall 2019 Midterm #2 14
- 2.3 Fall 2013 #2 14
  - 2.3.1 a 15
  - 2.3.2 b 15
  - 2.3.3 c 15
  - 2.3.4 d 15
- 2.4 Spring 2014 #2 15
  - 2.4.1 a 15
  - 2.4.2 b 15
  - 2.4.3 c 15
  - 2.4.4 d 16
- 2.5 Fall 2014 #2 16
  - 2.5.1 a 16
  - 2.5.2 b 16
- 2.6 Spring 2016 #3 16
  - 2.6.1 a 16
  - 2.6.2 b 16

2.6.3 c . . . 16
2.7 Spring 2017 #2 . . . 17
2.7.1 a . . . 17
2.7.2 b . . . 17
2.7.3 c . . . 17
2.8 Fall 2017 #2 . . . 17
2.8.1 a . . . 17
2.8.2 b . . . 17
2.8.3 c . . . 18
2.8.4 d . . . 18
2.9 Fall 2012 #2 . . . 18
2.9.1 a . . . 18
2.9.2 b . . . 18
2.9.3 c . . . 18
2.10 Fall 2018 #1 . . . 18
2.10.1 a . . . 18
2.10.2 b . . . 19
2.11 Fall 2019 #2 . . . 19
2.11.1 a . . . 19
2.11.2 b . . . 19
2.11.3 c . . . 19
2.11.4 d . . . 19
2.12 Spring 2021 #3 . . . 19
2.13 Fall 2020 #1 . . . 20
2.14 Fall 2020 #2 . . . 20

3 Groups: Group Actions **20**
3.1 Fall 2012 #1 . . . 20
3.1.1 a . . . 20
3.1.2 b . . . 20
3.2 Fall 2015 #2 . . . 21
3.3 Spring 2016 #5 . . . 21
3.3.1 a . . . 21
3.3.2 b . . . 21
3.4 Fall 2017 #1 . . . 21
3.4.1 a . . . 21
3.4.2 b . . . 22
3.5 Fall 2018 #2 . . . 22
3.5.1 a . . . 22
3.5.2 b . . . 22
3.5.3 c . . . 22

4 Groups: Classification **22**
4.1 Spring 2020 #1 . . . 22
4.1.1 a . . . 22
4.1.2 a . . . 23
4.1.3 c . . . 23
4.2 Spring 2019 #3 . . . 23

4.3 Spring 2012 #3 . . . 23
4.3.1 a . . . 23
4.3.2 a . . . 23
4.4 Fall 2016 #3 . . . 23
4.5 Spring 2018 #1 . . . 24
4.5.1 a . . . 24
4.5.2 b . . . 24
4.5.3 c . . . 24
4.5.4 d . . . 24

5 Groups: Simple and Solvable **24**
5.1 ⋆ Fall 2016 #7 . . . 24
5.1.1 a . . . 24
5.1.2 a . . . 24
5.2 Spring 2015 #4 . . . 25
5.2.1 a . . . 25
5.2.2 a . . . 25
5.2.3 c . . . 25
5.3 Spring 2014 #1 . . . 25
5.4 Fall 2013 #1 . . . 25
5.4.1 a . . . 26
5.4.2 a . . . 26
5.5 Spring 2013 #4 . . . 26
5.6 Fall 2019 Midterm #3 . . . 26

6 Commutative Algebra **26**
6.1 Spring 2020 #5 . . . 26
6.2 Fall 2019 #3 . . . 26
6.2.1 a . . . 27
6.2.2 b . . . 27
6.3 Fall 2019 #6 . . . 27
6.3.1 a . . . 27
6.3.2 b . . . 27
6.3.3 c . . . 27
6.4 Spring 2019 #6 . . . 27
6.4.1 a . . . 28
6.4.2 b . . . 28
6.4.3 c . . . 28
6.5 Fall 2018 #7 . . . 28
6.5.1 a . . . 28
6.5.2 b . . . 28
6.5.3 c . . . 29
6.5.4 d . . . 29
6.6 Spring 2018 #5 . . . 29
6.7 Spring 2018 #8 . . . 29
6.7.1 a . . . 29
6.7.2 b . . . 30
6.7.3 c . . . 30

6.8 Fall 2017 #5 . . . 30
6.8.1 a . . . 30
6.8.2 b . . . 30
6.9 Fall 2017 #6 . . . 30
6.9.1 a . . . 30
6.9.2 b . . . 31
6.9.3 c . . . 31
6.9.4 d . . . 31
6.9.5 e . . . 31
6.10 Spring 2017 #3 . . . 31
6.10.1 a . . . 31
6.10.2 b . . . 31
6.11 Spring 2017 #4 . . . 32
6.11.1 a . . . 32
6.11.2 b . . . 32
6.12 Spring 2016 #8 . . . 32
6.12.1 a . . . 32
6.12.2 b . . . 32
6.13 Fall 2015 #3 . . . 33
6.14 Fall 2015 #4 . . . 33
6.15 Spring 2015 #7 . . . 33
6.16 Fall 2014 #7 . . . 33
6.17 Fall 2014 #8 . . . 34
6.18 Spring 2014 #5 . . . 34
6.19 Spring 2014 #6 . . . 34
6.19.1 a . . . 34
6.19.2 b . . . 34
6.20 Fall 2013 #3 . . . 34
6.20.1 a . . . 34
6.20.2 b . . . 35
6.21 Fall 2013 #4 . . . 35
6.21.1 a . . . 35
6.21.2 b . . . 35
6.21.3 c . . . 35
6.21.4 d . . . 35
6.22 Spring 2013 #1 . . . 36
6.22.1 a . . . 36
6.22.2 b . . . 36
6.22.3 c . . . 36
6.23 Spring 2013 #2 . . . 36
6.23.1 a . . . 36
6.23.2 b . . . 36
6.23.3 c . . . 37
6.23.4 d . . . 37

7 Fields and Galois Theory **37**
7.1 ⋆ Fall 2016 #5 . . . 37

7.2 ⋆ Fall 2013 #7 . . . 37
7.2.1 a . . . 37
7.2.2 b . . . 37
7.3 Fall 2019 #4 . . . 37
7.3.1 a . . . 38
7.3.2 b . . . 38
7.3.3 c . . . 38
7.4 Fall 2019 #7 . . . 38
7.5 Spring 2019 #2 . . . 38
7.5.1 a . . . 39
7.5.2 b . . . 39
7.6 Spring 2019 #8 . . . 39
7.6.1 a . . . 39
7.6.2 b . . . 39
7.6.3 c . . . 39
7.7 Fall 2018 #3 . . . 39
7.7.1 a . . . 40
7.7.2 b . . . 40
7.7.3 c . . . 40
7.8 Spring 2018 #2 . . . 40
7.8.1 a . . . 40
7.8.2 b . . . 40
7.8.3 c . . . 40
7.9 Spring 2018 #3 . . . 41
7.9.1 a . . . 41
7.9.2 b . . . 41
7.9.3 c . . . 41
7.10 Spring 2020 #4 . . . 41
7.10.1 a . . . 41
7.10.2 b . . . 41
7.10.3 c . . . 42
7.11 Spring 2020 #3 . . . 42
7.11.1 a . . . 42
7.11.2 b . . . 42
7.12 Fall 2017 #4 . . . 42
7.12.1 a . . . 42
7.12.2 b . . . 42
7.13 Fall 2017 #3 . . . 43
7.14 Spring 2017 #7 . . . 43
7.14.1 a . . . 43
7.14.2 b . . . 43
7.14.3 c . . . 43
7.15 Spring 2017 #8 . . . 43
7.15.1 a . . . 43
7.15.2 b . . . 44
7.16 Fall 2016 #4 . . . 44
7.16.1 a . . . 44
7.16.2 b . . . 44

7.16.3 c . . . 44
7.17 Spring 2016 #2 . . . 44
7.17.1 a . . . 44
7.17.2 b . . . 44
7.17.3 c . . . 45
7.18 Spring 2016 #6 . . . 45
7.19 Fall 2015 #5 . . . 45
7.19.1 a . . . 45
7.19.2 b . . . 45
7.19.3 c . . . 45
7.20 Fall 2015 #6 . . . 45
7.20.1 a . . . 45
7.20.2 b . . . 46
7.20.3 c . . . 46
7.21 Spring 2015 #2 . . . 46
7.21.1 a . . . 46
7.21.2 b . . . 46
7.21.3 c . . . 46
7.22 Spring 2015 #5 . . . 46
7.22.1 a . . . 47
7.22.2 b . . . 47
7.23 Fall 2014 #1 . . . 47
7.23.1 a . . . 47
7.23.2 b . . . 47
7.24 Fall 2014 #3 . . . 47
7.24.1 a . . . 47
7.24.2 b . . . 47
7.25 Spring 2014 #3 . . . 48
7.25.1 a . . . 48
7.25.2 b . . . 48
7.26 Spring 2014 #4 . . . 48
7.26.1 a . . . 48
7.26.2 b . . . 48
7.26.3 c . . . 48
7.27 Fall 2013 #5 . . . 49
7.27.1 a . . . 49
7.27.2 b . . . 49
7.27.3 c . . . 49
7.28 Fall 2013 #6 . . . 49
7.28.1 a . . . 49
7.28.2 b . . . 49
7.28.3 c . . . 50
7.28.4 d . . . 50
7.29 Spring 2013 #7 . . . 50
7.29.1 a . . . 50
7.29.2 b . . . 50
7.29.3 c . . . 50
7.29.4 d . . . 50

7.29.5 e . . . 50
7.30 Spring 2013 #8 . . . 51
7.30.1 a . . . 51
7.30.2 b . . . 51
7.31 Fall 2012 #3 . . . 51
7.32 Fall 2012 #4 . . . 51
7.33 Spring 2012 #1 . . . 51
7.33.1 a . . . 51
7.33.2 b . . . 52
7.34 Spring 2012 #4 . . . 52
7.34.1 a . . . 52
7.34.2 b . . . 52
7.35 Fall 2019 Midterm #6 . . . 52
7.36 Fall 2019 Midterm #7 . . . 52
7.37 Fall 2019 Midterm #8 . . . 52
7.38 Fall 2019 Midterm #9 . . . 53
7.39 Spring 2021 #4 . . . 53
7.40 Spring 2021 #7 . . . 53
7.41 Fall 2020 #3 . . . 53
7.42 Fall 2020 #4 . . . 54

8 Modules **54**
8.1 General Questions . . . 54
8.1.1 Fall 2018 #6 ✨ . . . 54
8.1.2 Fall 2019 Final #2 . . . 54
8.1.3 Spring 2018 #6 . . . 55
8.1.4 Spring 2018 #7 . . . 55
8.1.5 Fall 2016 #6 . . . 55
8.1.6 Spring 2016 #4 . . . 56
8.1.7 Spring 2015 #8 . . . 56
8.1.8 Fall 2012 #6 . . . 56
8.1.9 Fall 2019 Final #1 . . . 56
8.1.10 Fall 2020 #6 . . . 57
8.2 Torsion and the Structure Theorem . . . 57
8.2.1 ⋆ Fall 2019 #5 ✨ . . . 57
8.2.2 ⋆ Spring 2019 #5 ✨ . . . 57
8.2.3 ⋆ Spring 2020 #6 ✨ . . . 58
8.2.4 Spring 2012 #5 . . . 58
8.2.5 Spring 2017 #5 . . . 59
8.2.6 Fall 2019 Final #3 . . . 59
8.2.7 Fall 2019 Final #4 . . . 59
8.2.8 Fall 2019 Final #5 . . . 59
8.2.9 Fall 2019 Final #6 . . . 60
8.2.10 Fall 2019 Final #7 . . . 60
8.2.11 Fall 2019 Final #10 . . . 60
8.2.12 Fall 2020 #7 . . . 60

9 Ring Theory **60**
9.1 Spring 2021 #5 . . . 60
9.2 Spring 2021 #6 . . . 61

10 Linear Algebra: Diagonalizability **61**
10.1 Fall 2017 #7 . . . 61
10.1.1 a . . . 61
10.1.2 b . . . 61
10.1.3 c . . . 62
10.1.4 d . . . 62
10.2 Spring 2015 #3 . . . 62
10.3 Fall 2016 #2 . . . 62
10.4 Spring 2019 #1 . . . 62

11 Linear Algebra: Misc **62**
11.1 ⋆ Spring 2012 #6 . . . 63
11.1.1 a . . . 63
11.1.2 b . . . 63
11.2 ⋆ Spring 2014 #7 . . . 63
11.2.1 a . . . 63
11.2.2 b . . . 63
11.3 Fall 2012 #7 . . . 64
11.4 Fall 2012 #8 . . . 64
11.5 Fall 2012 #5 . . . 64
11.6 Fall 2015 #7 . . . 64
11.6.1 a . . . 64
11.6.2 b . . . 65
11.7 Fall 2014 #4 . . . 65
11.8 Fall 2015 #8 . . . 65
11.8.1 a . . . 65
11.8.2 b . . . 65
11.9 Fall 2018 #4 . . . 66
11.10 Fall 2018 #5 . . . 66
11.10.1 a . . . 66
11.10.2 b . . . 66
11.11 Fall 2019 #8 . . . 66
11.11.1 a . . . 67
11.11.2 b . . . 67
11.11.3 c . . . 67
11.12 Spring 2013 #6 . . . 67
11.12.1 a . . . 67
11.12.2 b . . . 67
11.13 Fall 2020 #8 . . . 68

12 Linear Algebra: Canonical Forms **68**
12.1 ⋆ Spring 2012 #8 . . . 68
12.1.1 a . . . 68
12.1.2 b . . . 68

12.1.3 c . . . 68
12.2 ⋆ Spring 2020 #8 . . . 69
12.3 ⋆ Spring 2012 #7 . . . 69
12.3.1 a . . . 69
12.3.2 b . . . 69
12.3.3 c . . . 69
12.4 Fall 2019 Final #8 . . . 69
12.5 Fall 2019 Final #9 . . . 70
12.6 Spring 2016 #7 . . . 70
12.7 Spring 2020 #7 . . . 70
12.7.1 a . . . 70
12.7.2 b . . . 70
12.7.3 c . . . 70
12.8 Spring 2019 #7 . . . 71
12.8.1 a . . . 71
12.8.2 b . . . 71
12.9 Spring 2018 #4 . . . 71
12.9.1 a . . . 71
12.9.2 b . . . 71
12.10 Spring 2017 #6 . . . 72
12.10.1 a . . . 72
12.10.2 b . . . 72
12.11 Spring 2016 #1 . . . 72
12.11.1 a . . . 72
12.11.2 b . . . 72
12.12 Spring 2015 #6 . . . 73
12.13 Fall 2014 #5 . . . 73
12.14 Spring 2013 #5 . . . 73
12.14.1 a . . . 73
12.14.2 b . . . 73
12.15 Spring 2021 #1 . . . 74
12.16 Fall 2020 #5 . . . 74

1 | Group Theory: General

1.1 Spring 2020 #2

Let H be a normal subgroup of a finite group G where the order of H and the index of H in G are relatively prime. Prove that no other subgroup of G has the same order as H.

Work this problem.

1.2 Spring 2019 #4

For a finite group G, let $c(G)$ denote the number of conjugacy classes of G.

1.2.1 a

Prove that if two elements of G are chosen uniformly at random,then the probability they commute is precisely

$$\frac{c(G)}{|G|}.$$

1.2.2 b

State the class equation for a finite group.

1.2.3 c

Using the class equation (or otherwise) show that the probability in part (a) is at most

$$\frac{1}{2} + \frac{1}{2[G : Z(G)]}.$$

Here, as usual, $Z(G)$ denotes the center of G.

1.3 Spring 2012 #2

Let G be a finite group and p a prime number such that there is a normal subgroup $H \trianglelefteq G$ with $|H| = p^i > 1$.

1.3.1 a

Show that H is a subgroup of any Sylow p-subgroup of G.

1.3.2 b

Show that G contains a nonzero abelian normal subgroup of order divisible by p.

1.4 Spring 2017 #1

Let G be a finite group and $\pi : G \to \mathrm{Sym}(G)$ the Cayley representation.

(Recall that this means that for an element $x \in G$, $\pi(x)$ acts by left translation on G.)

Prove that $\pi(x)$ is an odd permutation $\iff$ the order $|\pi(x)|$ of $\pi(x)$ is even and $|G|/|\pi(x)|$ is odd.

1.5 Fall 2016 #1

Let G be a finite group and $s, t \in G$ be two distinct elements of order 2. Show that subgroup of G generated by s and t is a dihedral group.

Recall that the dihedral groups of order $2m$ for $m \geq 2$ are of the form

$$D_{2m} = \left\langle \sigma, \tau \;\middle|\; \sigma^m = 1 = \tau^2, \tau\sigma = \sigma^{-1}\tau \right\rangle.$$

1.6 Fall 2015 #1

Let G be a group containing a subgroup H not equal to G of finite index. Prove that G has a normal subgroup which is contained in every conjugate of H which is of finite index.

1.7 Spring 2015 #1

For a prime p, let G be a finite p-group and let N be a normal subgroup of G of order p. Prove that N is contained in the center of G.

1.8 Fall 2014 #6

Let G be a group and $H, K < G$ be subgroups of finite index. Show that

$$[G : H \cap K] \leq [G : H]\ [G : K].$$

1.9 Spring 2013 #3

Let P be a finite p-group. Prove that every nontrivial normal subgroup of P intersects the center of P nontrivially.

1.10 Fall 2019 Midterm #1

Let G be a group of order p^2q for p, q prime. Show that G has a nontrivial normal subgroup.

1.11 Fall 2019 Midterm #4

Let p be a prime. Show that $S_p = \langle \tau, \sigma \rangle$ where τ is a transposition and σ is a p-cycle.

1.12 Fall 2019 Midterm #5

Let G be a nonabelian group of order p^3 for p prime. Show that $Z(G) = [G, G]$

1.13 Spring 2021 #2

Let $H \trianglelefteq G$ be a normal subgroup of a finite group G, where the order of H is the smallest prime p dividing $|G|$. Prove that H is contained in the center of G.

2 | Groups: Sylow Theory

2.1 Fall 2019 #1

Let G be a finite group with n distinct conjugacy classes. Let $g_1 \cdots g_n$ be representatives of the conjugacy classes of G.

Prove that if $g_i g_j = g_j g_i$ for all i, j then G is abelian.

2.2 Fall 2019 Midterm #2

Let G be a finite group and let P be a sylow p-subgroup for p prime. Show that $N(N(P)) = N(P)$ where N is the normalizer in G.

2.3 Fall 2013 #2

Let G be a group of order 30.

2.3.1 a

Show that G has a subgroup of order 15.

2.3.2 b

Show that every group of order 15 is cyclic.

2.3.3 c

Show that G is isomorphic to some semidirect product $\mathbb{Z}_{15} \rtimes \mathbb{Z}_2$.

2.3.4 d

Exhibit three nonisomorphic groups of order 30 and prove that they are not isomorphic. You are not required to use your answer to (c).

2.4 Spring 2014 #2

Let $G \subset S_9$ be a Sylow-3 subgroup of the symmetric group on 9 letters.

2.4.1 a

Show that G contains a subgroup H isomorphic to $\mathbb{Z}_3 \times \mathbb{Z}_3 \times \mathbb{Z}_3$ by exhibiting an appropriate set of cycles.

2.4.2 b

Show that H is normal in G.

2.4.3 c

Give generators and relations for G as an abstract group, such that all generators have order 3.

Also exhibit elements of S_9 in cycle notation corresponding to these generators.

2.4.4 d

Without appealing to the previous parts of the problem, show that G contains an element of order 9.

2.5 Fall 2014 #2

Let G be a group of order 96.

2.5.1 a

Show that G has either one or three 2-Sylow subgroups.

2.5.2 b

Show that either G has a normal subgroup of order 32, or a normal subgroup of order 16.

2.6 Spring 2016 #3

2.6.1 a

State the three Sylow theorems.

2.6.2 b

Prove that any group of order 1225 is abelian.

2.6.3 c

Write down exactly one representative in each isomorphism class of abelian groups of order 1225.

2.7 Spring 2017 #2

2.7.1 a

How many isomorphism classes of abelian groups of order 56 are there? Give a representative for one of each class.

2.7.2 b

Prove that if G is a group of order 56, then either the Sylow-2 subgroup or the Sylow-7 subgroup is normal.

2.7.3 c

Give two non-isomorphic groups of order 56 where the Sylow-7 subgroup is normal and the Sylow-2 subgroup is *not* normal. Justify that these two groups are not isomorphic.

2.8 Fall 2017 #2

2.8.1 a

Classify the abelian groups of order 36.

For the rest of the problem, assume that G is a non-abelian group of order 36.

You may assume that the only subgroup of order 12 in S_4 is A_4 and that A_4 has no subgroup of order 6.

2.8.2 b

Prove that if the 2-Sylow subgroup of G is normal, G has a normal subgroup N such that G/N is isomorphic to A_4.

2.8.3 c

Show that if G has a normal subgroup N such that G/N is isomorphic to A_4 and a subgroup H isomorphic to A_4 it must be the direct product of N and H.

2.8.4 d

Show that the dihedral group of order 36 is a non-abelian group of order 36 whose Sylow-2 subgroup is not normal.

2.9 Fall 2012 #2

Let G be a group of order 30.

2.9.1 a

Show that G contains normal subgroups of orders 3, 5, and 15.

2.9.2 b

Give all possible presentations and relations for G.

2.9.3 c

Determine how many groups of order 30 there are up to isomorphism.

2.10 Fall 2018 #1

Let G be a finite group whose order is divisible by a prime number p. Let P be a normal p-subgroup of G (so $|P| = p^c$ for some c).

2.10.1 a

Show that P is contained in every Sylow p-subgroup of G.

2.10.2 b

Let M be a maximal proper subgroup of G. Show that either $P \subseteq M$ or $|G/M| = p^b$ for some $b \leq c$.

2.11 Fall 2019 #2 ✨

Let G be a group of order 105 and let P, Q, R be Sylow 3, 5, 7 subgroups respectively.

2.11.1 a

Prove that at least one of Q and R is normal in G.

2.11.2 b

Prove that G has a cyclic subgroup of order 35.

2.11.3 c

Prove that both Q and R are normal in G.

2.11.4 d

Prove that if P is normal in G then G is cyclic.

2.12 Spring 2021 #3 🚩

a. Show that every group of order p^2 with p prime is abelian.

b. State the 3 Sylow theorems.

c. Show that any group of order $4225 = 5^2 13^2$ is abelian.

d. Write down one representative from each isomorphism class of abelian groups of order 4225.

2.13 Fall 2020 #1

a. Using Sylow theory, show that every group of order $2p$ where p is prime is not simple.

b. Classify all groups of order $2p$ and justify your answer. For the nonabelian group(s), give a presentation by generators and relations.

2.14 Fall 2020 #2

Let G be a group of order 60 whose Sylow 3-subgroup is normal.

a. Prove that G is solvable.

b. Prove that the Sylow 5-subgroup is also normal.

3 | Groups: Group Actions

3.1 Fall 2012 #1

Let G be a finite group and X a set on which G acts.

3.1.1 a

Let $x \in X$ and $G_x \coloneqq \left\{ g \in G \;\middle|\; g \cdot x = x \right\}$. Show that G_x is a subgroup of G.

3.1.2 b

Let $x \in X$ and $G \cdot x \coloneqq \left\{ g \cdot x \;\middle|\; g \in G \right\}$. Prove that there is a bijection between elements in $G \cdot x$ and the left cosets of G_x in G.

3.2 Fall 2015 #2

Let G be a finite group, H a p-subgroup, and P a sylow p-subgroup for p a prime. Let H act on the left cosets of P in G by left translation.

Prove that this is an orbit under this action of length 1.

Prove that xP is an orbit of length 1 $\iff$ H is contained in xPx^{-1}.

3.3 Spring 2016 #5

Let G be a finite group acting on a set X. For $x \in X$, let G_x be the stabilizer of x and $G \cdot x$ be the orbit of x.

3.3.1 a

Prove that there is a bijection between the left cosets G/G_x and $G \cdot x$.

3.3.2 b

Prove that the center of every finite p-group G is nontrivial by considering that action of G on $X = G$ by conjugation.

3.4 Fall 2017 #1

Suppose the group G acts on the set A. Assume this action is faithful (recall that this means that the kernel of the homomorphism from G to $\mathrm{Sym}(A)$ which gives the action is trivial) and transitive (for all a, b in A, there exists g in G such that $g \cdot a = b$.)

3.4.1 a

For $a \in A$, let G_a denote the stabilizer of a in G. Prove that for any $a \in A$,

$$\bigcap_{\sigma \in G} \sigma G_a \sigma^{-1} = \{1\}\,.$$

3.4.2 b

Suppose that G is abelian. Prove that $|G| = |A|$. Deduce that every abelian transitive subgroup of S_n has order n.

3.5 Fall 2018 #2 ✨

3.5.1 a

Suppose the group G acts on the set X . Show that the stabilizers of elements in the same orbit are conjugate.

3.5.2 b

Let G be a finite group and let H be a proper subgroup. Show that the union of the conjugates of H is strictly smaller than G, i.e.

$$\bigcup_{g \in G} gHg^{-1} \subsetneq G$$

3.5.3 c

Suppose G is a finite group acting transitively on a set S with at least 2 elements. Show that there is an element of G with no fixed points in S.

4 Groups: Classification

4.1 Spring 2020 #1 🚩

4.1.1 a

Show that any group of order 2020 is solvable.

4.1.2 a

Give (without proof) a classification of all abelian groups of order 2020.

4.1.3 c

Describe one nonabelian group of order 2020.

Work this problem.

4.2 Spring 2019 #3 ✨

How many isomorphism classes are there of groups of order 45?

Describe a representative from each class.

4.3 Spring 2012 #3 🚩

Let G be a group of order 70.

4.3.1 a

Show that G is not simple.

4.3.2 a

Exhibit 3 nonisomorphic groups of order 70 and prove that they are not isomorphic.

4.4 Fall 2016 #3 🚩

How many groups are there up to isomorphism of order pq where $p < q$ are prime integers?

4.5 Spring 2018 #1 ✨

4.5.1 a

Use the Class Equation (equivalently, the conjugation action of a group on itself) to prove that any p-group (a group whose order is a positive power of a prime integer p) has a nontrivial center.

4.5.2 b

Prove that any group of order p^2 (where p is prime) is abelian.

4.5.3 c

Prove that any group of order $5^2 \cdot 7^2$ is abelian.

4.5.4 d

Write down exactly one representative in each isomorphism class of groups of order $5^2 \cdot 7^2$.

5 | Groups: Simple and Solvable

5.1 ⋆ Fall 2016 #7 🚩

5.1.1 a

Define what it means for a group G to be *solvable*.

5.1.2 a

Show that every group G of order 36 is solvable.

Hint: you can use that S_4 is solvable.

5.2 Spring 2015 #4

Let N be a positive integer, and let G be a finite group of order N.

5.2.1 a

Let $\operatorname{Sym} G$ be the set of all bijections from $G \to G$ viewed as a group under composition. Note that $\operatorname{Sym} G \cong S_N$. Prove that the Cayley map

$$\begin{aligned} C : G &\to \operatorname{Sym} G \\ g &\mapsto (x \mapsto gx) \end{aligned}$$

is an injective homomorphism.

5.2.2 a

Let $\Phi : \operatorname{Sym} G \to S_N$ be an isomorphism. For $a \in G$ define $\varepsilon(a) \in \{\pm 1\}$ to be the sign of the permutation $\Phi(C(a))$. Suppose that a has order d. Prove that $\varepsilon(a) = -1 \iff d$ is even and N/d is odd.

5.2.3 c

Suppose $N > 2$ and $n \equiv 2 \pmod 4$. Prove that G is not simple.

Hint: use part (b).

5.3 Spring 2014 #1

Let p, n be integers such that p is prime and p does not divide n. Find a real number $k = k(p, n)$ such that for every integer $m \geq k$, every group of order $p^m n$ is not simple.

5.4 Fall 2013 #1

Let p, q be distinct primes.

5.4.1 a

Let $\bar{q} \in \mathbb{Z}_p$ be the class of $q \pmod{p}$ and let k denote the order of $\bar{q}$ as an element of $\mathbb{Z}_p^\times$.

Prove that no group of order pq^k is simple.

5.4.2 a

Let G be a group of order pq, and prove that G is not simple.

5.5 Spring 2013 #4

Define a *simple group*.

Prove that a group of order 56 can not be simple.

5.6 Fall 2019 Midterm #3

Show that there exist no simple groups of order 148.

6 Commutative Algebra

6.1 Spring 2020 #5

Let R be a ring and $f : M \to N$ and $g : N \to M$ be R-module homomorphisms such that $g \circ f = \mathrm{id}_M$. Show that $N \cong \operatorname{im} f \oplus \ker g$.

6.2 Fall 2019 #3

Let R be a ring with the property that for every $a \in R, a^2 = a$.

6.2.1 a

Prove that R has characteristic 2.

6.2.2 b

Prove that R is commutative.

6.3 Fall 2019 #6 ✨

Let R be a commutative ring with multiplicative identity. Assume Zorn's Lemma.

6.3.1 a

Show that

$$N = \{r \in R \mid r^n = 0 \text{ for some } n > 0\}$$

is an ideal which is contained in any prime ideal.

6.3.2 b

Let r be an element of R not in N. Let S be the collection of all proper ideals of R not containing any positive power of r. Use Zorn's Lemma to prove that there is a prime ideal in S.

6.3.3 c

Suppose that R has exactly one prime ideal P . Prove that every element r of R is either nilpotent or a unit.

6.4 Spring 2019 #6 ✨

Let R be a commutative ring with 1.

Recall that $x \in R$ is nilpotent iff $xn = 0$ for some positive integer n.

6.4.1 a

Show that every proper ideal of R is contained within a maximal ideal.

6.4.2 b

Let $J(R)$ denote the intersection of all maximal ideals of R.

Show that $x \in J(R) \iff 1 + rx$ is a unit for all $r \in R$.

6.4.3 c

Suppose now that R is finite. Show that in this case $J(R)$ consists precisely of the nilpotent elements in R.

6.5 Fall 2018 #7 ✨

Let R be a commutative ring.

6.5.1 a

Let $r \in R$. Show that the map

$$\begin{aligned} r\bullet : R &\to R \\ x &\mapsto rx. \end{aligned}$$

is an R-module endomorphism of R.

6.5.2 b

We say that r is a **zero-divisor** if r• is not injective. Show that if r is a zero-divisor and $r \neq 0$, then the kernel and image of R each consist of zero-divisors.

6.5.3 c

Let $n \geq 2$ be an integer. Show: if R has exactly n zero-divisors, then $\#R \leq n^2$.

6.5.4 d

Show that up to isomorphism there are exactly two commutative rings R with precisely 2 zero-divisors.

You may use without proof the following fact: every ring of order 4 is isomorphic to exactly one of the following:

$$\frac{\mathbb{Z}}{4\mathbb{Z}}, \quad \frac{\frac{\mathbb{Z}}{2\mathbb{Z}}[t]}{(t^2+t+1)}, \quad \frac{\frac{\mathbb{Z}}{2\mathbb{Z}}[t]}{(t^2-t)}, \quad \frac{\frac{\mathbb{Z}}{2\mathbb{Z}}[t]}{(t^2)}.$$

6.6 Spring 2018 #5

Let \$

$$M = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \quad \text{and} \quad N = \begin{pmatrix} x & u \\ -y & -v \end{pmatrix}$$

over a commutative ring R, where b and x are units of R. Prove that

$$MN = \begin{pmatrix} 0 & 0 \\ 0 & * \end{pmatrix} \implies MN = 0.$$

6.7 Spring 2018 #8

Let $R = C[0,1]$ be the ring of continuous real-valued functions on the interval $[0,1]$. Let I be an ideal of R.

6.7.1 a

Show that if $f \in I, a \in [0,1]$ are such that $f(a) \neq 0$, then there exists $g \in I$ such that $g(x) \geq 0$ for all $x \in [0,1]$, and $g(x) > 0$ for all x in some open neighborhood of a.

6.7.2 b

If $I \neq R$, show that the set $Z(I) = \{x \in [0,1] \mid f(x) = 0 \text{ for all } f \in I\}$ is nonempty.

6.7.3 c

Show that if I is maximal, then there exists $x_0 \in [0,1]$ such that $I = \{f \in R \mid f(x_0) = 0\}$.

6.8 Fall 2017 #5

A ring R is called *simple* if its only two-sided ideals are 0 and R.

6.8.1 a

Suppose R is a commutative ring with 1. Prove R is simple if and only if R is a field.

6.8.2 b

Let k be a field. Show the ring $M_n(k)$, $n \times n$ matrices with entries in k, is a simple ring.

6.9 Fall 2017 #6

For a ring R, let $U(R)$ denote the multiplicative group of units in R. Recall that in an integral domain R, $r \in R$ is called *irreducible* if r is not a unit in R, and the only divisors of r have the form ru with u a unit in R.

We call a non-zero, non-unit $r \in R$ *prime* in R if $r \mid ab \implies r \mid a$ or $r \mid b$. Consider the ring $R = \{a + b\sqrt{-5} \mid a, b \in Z\}$.

6.9.1 a

Prove R is an integral domain.

6.9.2 b

Show $U(R) = \{\pm 1\}$.

6.9.3 c

Show $3, 2 + \sqrt{-5}$, and $2 - \sqrt{-5}$ are irreducible in R.

6.9.4 d

Show 3 is not prime in R.

6.9.5 e

Conclude R is not a PID.

6.10 Spring 2017 #3

Let R be a commutative ring with 1. Suppose that M is a free R-module with a finite basis X.

6.10.1 a

Let $I \trianglelefteq R$ be a proper ideal. Prove that M/IM is a free R/I-module with basis X', where X' is the image of X under the canonical map $M \to M/IM$.

6.10.2 b

Prove that any two bases of M have the same number of elements. You may assume that the result is true when R is a field.

6.11 Spring 2017 #4

6.11.1 a

Let R be an integral domain with quotient field F. Suppose that $p(x), a(x), b(x)$ are monic polynomials in $F[x]$ with $p(x) = a(x)b(x)$ and with $p(x) \in R[x]$, $a(x)$ not in $R[x]$, and both $a(x), b(x)$ not constant.

Prove that R is not a UFD.

(You may assume Gauss' lemma)

6.11.2 b

Prove that $\mathbb{Z}[2\sqrt{2}]$ is not a UFD.

Hint: let $p(x) = x^2 - 2$.

6.12 Spring 2016 #8

Let R be a simple rng (a nonzero ring which is not assume to have a 1, whose only two-sided ideals are (0) and R) satisfying the following two conditions:

i. R has no zero divisors, and
ii. If $x \in R$ with $x \neq 0$ then $2x \neq 0$, where $2x \coloneqq x + x$.

Prove the following:

6.12.1 a

For each $x \in R$ there is one and only one element $y \in R$ such that $x = 2y$.

6.12.2 b

Suppose $x, y \in R$ such that $x \neq 0$ and $2(xy) = x$, then $yz = zy$ for all $z \in R$.

You can get partial credit for (b) by showing it in the case R has a 1.

6.13 Fall 2015 #3

Let R be a rng (a ring without 1) which contains an element u such that for all $y \in R$, there exists an $x \in R$ such that $xu = y$.

Prove that R contains a maximal left ideal.

Hint: imitate the proof (using Zorn's lemma) in the case where R does have a 1.

6.14 Fall 2015 #4

Let R be a PID and $(a_1) < (a_2) < \cdots$ be an ascending chain of ideals in R. Prove that for some n, we have $(a_j) = (a_n)$ for all $j \geq n$.

6.15 Spring 2015 #7

Let R be a commutative ring, and $S \subset R$ be a nonempty subset that does not contain 0 such that for all $x, y \in S$ we have $xy \in S$. Let $\mathcal{I}$ be the set of all ideals $I \trianglelefteq R$ such that $I \cap S = \emptyset$.

Show that for every ideal $I \in \mathcal{I}$, there is an ideal $J \in \mathcal{I}$ such that $I \subset J$ and J is not properly contained in any other ideal in $\mathcal{I}$.

Prove that every such ideal J is prime.

6.16 Fall 2014 #7

Give a careful proof that $\mathbb{C}[x, y]$ is not a PID.

6.17 Fall 2014 #8

Let R be a nonzero commutative ring without unit such that R does not contain a proper maximal ideal. Prove that for all $x \in R$, the ideal xR is proper.

You may assume the axiom of choice.

6.18 Spring 2014 #5

Let R be a commutative ring and $a \in R$. Prove that a is not nilpotent $\iff$ there exists a commutative ring S and a ring homomorphism $\varphi : R \to S$ such that $\varphi(a)$ is a unit.

Note: by definition, a is nilpotent $\iff$ there is a natural number n such that $a^n = 0$.

6.19 Spring 2014 #6

Let R be a commutative ring with identity and let n be a positive integer.

6.19.1 a

Prove that every surjective R-linear endomorphism $T : R^n \to R^n$ is injective.

6.19.2 b

Show that an injective R-linear endomorphism of R^n need not be surjective.

6.20 Fall 2013 #3

6.20.1 a

Define *prime ideal*, give an example of a nontrivial ideal in the ring $\mathbb{Z}$ that is not prime, and prove that it is not prime.

6.20.2 b

Define *maximal ideal*, give an example of a nontrivial maximal ideal in $\mathbb{Z}$ and prove that it is maximal.

6.21 Fall 2013 #4

Let R be a commutative ring with $1 \neq 0$. Recall that $x \in R$ is *nilpotent* iff $x^n = 0$ for some positive integer n.

6.21.1 a

Show that the collection of nilpotent elements in R forms an ideal.

6.21.2 b

Show that if x is nilpotent, then x is contained in every prime ideal of R.

6.21.3 c

Suppose $x \in R$ is not nilpotent and let $S = \left\{ x^n \;\middle|\; n \in \mathbb{N} \right\}$. There is at least on ideal of R disjoint from S, namely (0).

By Zorn's lemma the set of ideals disjoint from S has a maximal element with respect to inclusion, say I. In other words, I is disjoint from S and if J is any ideal disjoint from S with $I \subseteq J \subseteq R$ then $J = I$ or $J = R$.

Show that I is a prime ideal.

6.21.4 d

Deduce from (a) and (b) that the set of nilpotent elements of R is the intersection of all prime ideals of R.

6.22 Spring 2013 #1

Let R be a commutative ring.

6.22.1 a

Define a *maximal ideal* and prove that R has a maximal ideal.

6.22.2 b

Show than an element $r \in R$ is not invertible $\iff$ r is contained in a maximal ideal.

6.22.3 c

Let M be an R-module, and recall that for $0 \neq \mu \in M$, the *annihilator* of μ is the set

$$\operatorname{Ann}(\mu) = \left\{ r \in R \;\middle|\; r\mu = 0 \right\}.$$

Suppose that I is an ideal in R which is maximal with respect to the property that there exists an element $\mu \in M$ such that $I = \operatorname{Ann}(\mu)$ for some $\mu \in M$. In other words, $I = \operatorname{Ann}(\mu)$ but there does not exist $\nu \in M$ with $J = \operatorname{Ann}(\nu) \subsetneq R$ such that $I \subsetneq J$.

Prove that I is a prime ideal.

6.23 Spring 2013 #2

6.23.1 a

Define a *Euclidean domain*.

6.23.2 b

Define a *unique factorization domain*.

6.23.3 c

Is a Euclidean domain an UFD? Give either a proof or a counterexample with justification.

6.23.4 d

Is a UFD a Euclidean domain? Give either a proof or a counterexample with justification.

7 | Fields and Galois Theory

7.1 ⋆ Fall 2016 #5

How many monic irreducible polynomials over $\mathbb{F}_p$ of prime degree ℓ are there? Justify your answer.

7.2 ⋆ Fall 2013 #7

Let $F = \mathbb{F}_2$ and let $\overline{F}$ denote its algebraic closure.

7.2.1 a

Show that $\overline{F}$ is not a finite extension of F.

7.2.2 b

Suppose that $\alpha \in \overline{F}$ satisfies $\alpha^{17} = 1$ and $\alpha \neq 1$. Show that $F(\alpha)/F$ has degree 8.

7.3 Fall 2019 #4 ✨

Let F be a finite field with q elements.

Let n be a positive integer relatively prime to q and let ω be a primitive nth root of unity in an extension field of F.

Let $E = F[\omega]$ and let $k = [E : F]$.

7.3.1 a

Prove that n divides $q^k - 1$.

7.3.2 b

Let m be the order of q in $\mathbb{Z}/n\mathbb{Z}^\times$. Prove that m divides k.

7.3.3 c

Prove that $m = k$.

Revisit, tricky!

7.4 Fall 2019 #7 ✨

Let ζ_n denote a primitive nth root of $1 \in \mathbb{Q}$. You may assume the roots of the minimal polynomial $p_n(x)$ of ζ_n are exactly the primitive nth roots of 1.

Show that the field extension $\mathbb{Q}(\zeta_n)$ over $\mathbb{Q}$ is Galois and prove its Galois group is $(\mathbb{Z}/n\mathbb{Z})^\times$.

How many subfields are there of $\mathbb{Q}(\zeta_{20})$?

7.5 Spring 2019 #2 ✨

Let $F = \mathbb{F}_p$, where p is a prime number.

7.5.1 a

Show that if $\pi(x) \in F[x]$ is irreducible of degree d, then $\pi(x)$ divides $x^{p^d} - x$.

7.5.2 b

Show that if $\pi(x) \in F[x]$ is an irreducible polynomial that divides $x^{p^n} - x$, then $\deg \pi(x)$ divides n.

7.6 Spring 2019 #8 ✨

Let $\zeta = e^{2\pi i/8}$.

7.6.1 a

What is the degree of $\mathbb{Q}(\zeta)/\mathbb{Q}$?

7.6.2 b

How many quadratic subfields of $\mathbb{Q}(\zeta)$ are there?

7.6.3 c

What is the degree of $\mathbb{Q}(\zeta, \sqrt[4]{2})$ over $\mathbb{Q}$?

7.7 Fall 2018 #3 ✨

Let $F \subset K \subset L$ be finite degree field extensions. For each of the following assertions, give a proof or a counterexample.

7.7.1 a

If L/F is Galois, then so is K/F.

7.7.2 b

If L/F is Galois, then so is L/K.

7.7.3 c

If K/F and L/K are both Galois, then so is L/F.

7.8 Spring 2018 #2 ✨

Let $f(x) = x^4 - 4x^2 + 2 \in \mathbb{Q}[x]$.

7.8.1 a

Find the splitting field K of f, and compute $[K : \mathbb{Q}]$.

7.8.2 b

Find the Galois group G of f, both as an explicit group of automorphisms, and as a familiar abstract group to which it is isomorphic.

7.8.3 c

Exhibit explicitly the correspondence between subgroups of G and intermediate fields between $\mathbb{Q}$ and k.

Not the nicest proof! Would be better to replace the ad-hoc computations at the end.

7.9 Spring 2018 #3 ✨

Let K be a Galois extension of $\mathbb{Q}$ with Galois group G, and let E_1, E_2 be intermediate fields of K which are the splitting fields of irreducible $f_i(x) \in \mathbb{Q}[x]$.

Let $E = E_1E_2 \subset K$.

Let $H_i = \text{Gal}(K/E_i)$ and $H = \text{Gal}(K/E)$.

7.9.1 a

Show that $H = H_1 \cap H_2$.

7.9.2 b

Show that H_1H_2 is a subgroup of G.

7.9.3 c

Show that

$$\text{Gal}(K/(E_1 \cap E_2)) = H_1H_2.$$

7.10 Spring 2020 #4 🚩

Let $f(x) = x^4 - 2 \in \mathbb{Q}[x]$.

7.10.1 a

Define what it means for a finite extension field E of a field F to be a Galois extension.

7.10.2 b

Determine the Galois group $\text{Gal}(E/\mathbb{Q})$ for the polynomial $f(x)$, and justify your answer carefully.

7.10.3 c

Exhibit a subfield K in (b) such that $\mathbb{Q} \leq K \leq E$ with K not a Galois extension over $\mathbb{Q}$. Explain.

7.11 Spring 2020 #3

Let E be an extension field of F and $\alpha \in E$ be algebraic of odd degree over F.

7.11.1 a

Show that $F(\alpha) = F(\alpha^2)$.

7.11.2 b

Prove that α^{2020} is algebraic of odd degree over F.

7.12 Fall 2017 #4

7.12.1 a

Let $f(x)$ be an irreducible polynomial of degree 4 in $\mathbb{Q}[x]$ whose splitting field K over $\mathbb{Q}$ has Galois group $G = S_4$.

Let θ be a root of $f(x)$. Prove that $\mathbb{Q}[\theta]$ is an extension of $\mathbb{Q}$ of degree 4 and that there are no intermediate fields between $\mathbb{Q}$ and $\mathbb{Q}[\theta]$.

7.12.2 b

Prove that if K is a Galois extension of $\mathbb{Q}$ of degree 4, then there is an intermediate subfield between K and $\mathbb{Q}$.

7.13 Fall 2017 #3

Let F be a field. Let $f(x)$ be an irreducible polynomial in $F[x]$ of degree n and let $g(x)$ be any polynomial in $F[x]$. Let $p(x)$ be an irreducible factor (of degree m) of the polynomial $f(g(x))$.

Prove that n divides m. Use this to prove that if r is an integer which is not a perfect square, and n is a positive integer then every irreducible factor of $x^{2n} - r$ over $\mathbb{Q}[x]$ has even degree.

7.14 Spring 2017 #7

Let F be a field and let $f(x) \in F[x]$.

7.14.1 a

Define what a splitting field of $f(x)$ over F is.

7.14.2 b

Let F now be a finite field with q elements. Let E/F be a finite extension of degree $n > 0$. Exhibit an explicit polynomial $g(x) \in F[x]$ such that E/F is a splitting field of $g(x)$ over F. Fully justify your answer.

7.14.3 c

Show that the extension E/F in (b) is a Galois extension.

7.15 Spring 2017 #8

7.15.1 a

Let K denote the splitting field of $x^5 - 2$ over $\mathbb{Q}$. Show that the Galois group of $K/\mathbb{Q}$ is isomorphic to the group of invertible matrices

$$\begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix} \quad \text{where} \quad a \in \mathbb{F}_5^\times \text{ and } b \in \mathbb{F}_5.$$

7.15.2 b

Determine all intermediate fields between K and $\mathbb{Q}$ which are Galois over $\mathbb{Q}$.

7.16 Fall 2016 #4

Set $f(x) = x^3 - 5 \in \mathbb{Q}[x]$.

7.16.1 a

Find the splitting field K of $f(x)$ over $\mathbb{Q}$.

7.16.2 b

Find the Galois group G of K over $\mathbb{Q}$.

7.16.3 c

Exhibit explicitly the correspondence between subgroups of G and intermediate fields between $\mathbb{Q}$ and K.

7.17 Spring 2016 #2

Let $K = \mathbb{Q}[\sqrt{2} + \sqrt{5}]$.

7.17.1 a

Find $[K : \mathbb{Q}]$.

7.17.2 b

Show that $K/\mathbb{Q}$ is Galois, and find the Galois group G of $K/\mathbb{Q}$.

7.17.3 c

Exhibit explicitly the correspondence between subgroups of G and intermediate fields between $\mathbb{Q}$ and K.

7.18 Spring 2016 #6

Let K be a Galois extension of a field F with $[K : F] = 2015$. Prove that K is an extension by radicals of the field F.

7.19 Fall 2015 #5

Let $u = \sqrt{2 + \sqrt{2}}$, $v = \sqrt{2 - \sqrt{2}}$, and $E = \mathbb{Q}(u)$.

7.19.1 a

Find (with justification) the minimal polynomial $f(x)$ of u over $\mathbb{Q}$.

7.19.2 b

Show $v \in E$, and show that E is a splitting field of $f(x)$ over $\mathbb{Q}$.

7.19.3 c

Determine the Galois group of E over $\mathbb{Q}$ and determine all of the intermediate fields F such that $\mathbb{Q} \subset F \subset E$.

7.20 Fall 2015 #6

7.20.1 a

Let G be a finite group. Show that there exists a field extension K/F with $\mathrm{Gal}(K/F) = G$.

You may assume that for any natural number n there is a field extension with Galois group S_n.

7.20.2 b

Let K be a Galois extension of F with $|\mathrm{Gal}(K/F)| = 12$. Prove that there exists an intermediate field E of K/F with $[E : F] = 3$.

7.20.3 c

With K/F as in (b), does an intermediate field L necessarily exist satisfying $[L : F] = 2$? Give a proof or counterexample.

7.21 Spring 2015 #2

Let $\mathbb{F}$ be a finite field.

7.21.1 a

Give (with proof) the decomposition of the additive group $(\mathbb{F}, +)$ into a direct sum of cyclic groups.

7.21.2 b

The *exponent* of a finite group is the least common multiple of the orders of its elements. Prove that a finite abelian group has an element of order equal to its exponent.

7.21.3 c

Prove that the multiplicative group $(\mathbb{F}^\times, \cdot)$ is cyclic.

7.22 Spring 2015 #5

Let $f(x) = x^4 - 5 \in \mathbb{Q}[x]$.

7.22.1 a

Compute the Galois group of f over $\mathbb{Q}$.

7.22.2 b

Compute the Galois group of f over $\mathbb{Q}(\sqrt{5})$.

7.23 Fall 2014 #1

Let $f \in \mathbb{Q}[x]$ be an irreducible polynomial and L a finite Galois extension of $\mathbb{Q}$. Let $f(x) = g_1(x)g_2(x)\cdots g_r(x)$ be a factorization of f into irreducibles in $L[x]$.

7.23.1 a

Prove that each of the factors $g_i(x)$ has the same degree.

7.23.2 b

Give an example showing that if L is not Galois over $\mathbb{Q}$, the conclusion of part (a) need not hold.

7.24 Fall 2014 #3

Consider the polynomial $f(x) = x^4 - 7 \in \mathbb{Q}[x]$ and let $E/\mathbb{Q}$ be the splitting field of f.

7.24.1 a

What is the structure of the Galois group of $E/\mathbb{Q}$?

7.24.2 b

Give an explicit description of all of the intermediate subfields $\mathbb{Q} \subset K \subset E$ in the form $K = \mathbb{Q}(\alpha), \mathbb{Q}(\alpha, \beta), \cdots$ where α, β, etc are complex numbers. Describe the corresponding subgroups of

the Galois group.

7.25 Spring 2014 #3

Let $F \subset C$ be a field extension with C algebraically closed.

7.25.1 a

Prove that the intermediate field $C_{\text{alg}} \subset C$ consisting of elements algebraic over F is algebraically closed.

7.25.2 b

Prove that if $F \to E$ is an algebraic extension, there exists a homomorphism $E \to C$ that is the identity on F.

7.26 Spring 2014 #4

Let $E \subset \mathbb{C}$ denote the splitting field over $\mathbb{Q}$ of the polynomial $x^3 - 11$.

7.26.1 a

Prove that if n is a squarefree positive integer, then $\sqrt{n} \notin E$.

Hint: you can describe all quadratic extensions of $\mathbb{Q}$ contained in E.

7.26.2 b

Find the Galois group of $(x^3 - 11)(x^2 - 2)$ over $\mathbb{Q}$.

7.26.3 c

Prove that the minimal polynomial of $11^{1/3} + 2^{1/2}$ over $\mathbb{Q}$ has degree 6.

7.27 Fall 2013 #5

Let L/K be a finite extension of fields.

7.27.1 a

Define what it means for L/K to be *separable*.

7.27.2 b

Show that if K is a finite field, then L/K is always separable.

7.27.3 c

Give an example of a finite extension L/K that is not separable.

7.28 Fall 2013 #6

Let K be the splitting field of $x^4 - 2$ over $\mathbb{Q}$ and set $G = \mathrm{Gal}(K/\mathbb{Q})$.

7.28.1 a

Show that $K/\mathbb{Q}$ contains both $\mathbb{Q}(i)$ and $\mathbb{Q}(\sqrt[4]{2})$ and has degree 8 over $\mathbb{Q}/$

7.28.2 b

Let $N = \mathrm{Gal}(K/\mathbb{Q}(i))$ and $H = \mathrm{Gal}(K/\mathbb{Q}(\sqrt[4]{2}))$. Show that N is normal in G and $NH = G$.

Hint: what field is fixed by NH?

7.28.3 c

Show that $\mathrm{Gal}(K/\mathbb{Q})$ is generated by elements σ, τ, of orders 4 and 2 respectively, with $\tau\sigma\tau^{-1} = \sigma^{-1}$.

Equivalently, show it is the dihedral group of order 8.

7.28.4 d

How many distinct quartic subfields of K are there? Justify your answer.

7.29 Spring 2013 #7

Let $f(x) = g(x)h(x) \in \mathbb{Q}[x]$ and $E, B, C/\mathbb{Q}$ be the splitting fields of f, g, h respectively.

7.29.1 a

Prove that $\mathrm{Gal}(E/B)$ and $\mathrm{Gal}(E/C)$ are normal subgroups of $\mathrm{Gal}(E/\mathbb{Q})$.

7.29.2 b

Prove that $\mathrm{Gal}(E/B) \cap \mathrm{Gal}(E/C) = \{1\}$.

7.29.3 c

If $B \cap C = \mathbb{Q}$, show that $\mathrm{Gal}(E/B)\,\mathrm{Gal}(E/C) = \mathrm{Gal}(E/\mathbb{Q})$.

7.29.4 d

Under the hypothesis of (c), show that $\mathrm{Gal}(E/\mathbb{Q}) \cong \mathrm{Gal}(E/B) \times \mathrm{Gal}(E/C)$.

7.29.5 e

Use (d) to describe $\mathrm{Gal}(\mathbb{Q}[\alpha]/\mathbb{Q})$ where $\alpha = \sqrt{2} + \sqrt{3}$.

7.30 Spring 2013 #8

Let F be the field with 2 elements and K a splitting field of $f(x) = x^6 + x^3 + 1$ over F. You may assume that f is irreducible over F.

7.30.1 a

Show that if r is a root of f in K, then $r^9 = 1$ but $r^3 \neq 1$.

7.30.2 b

Find $\mathrm{Gal}(K/F)$ and express each intermediate field between F and K as $F(\beta)$ for an appropriate $\beta \in K$.

7.31 Fall 2012 #3

Let $f(x) \in \mathbb{Q}[x]$ be an irreducible polynomial of degree 5. Assume that f has all but two roots in $\mathbb{R}$. Compute the Galois group of $f(x)$ over $\mathbb{Q}$ and justify your answer.

7.32 Fall 2012 #4

Let $f(x) \in \mathbb{Q}[x]$ be a polynomial and K be a splitting field of f over $\mathbb{Q}$. Assume that $[K : \mathbb{Q}] = 1225$ and show that $f(x)$ is solvable by radicals.

7.33 Spring 2012 #1

Suppose that $F \subset E$ are fields such that E/F is Galois and $|\mathrm{Gal}(E/F)| = 14$.

7.33.1 a

Show that there exists a unique intermediate field K with $F \subset K \subset E$ such that $[K : F] = 2$.

7.33.2 b

Assume that there are at least two distinct intermediate subfields $F \subset L_1, L_2 \subset E$ with $[L_i : F] = 7$. Prove that $\mathrm{Gal}(E/F)$ is nonabelian.

7.34 Spring 2012 #4

Let $f(x) = x^7 - 3 \in \mathbb{Q}[x]$ and $E/\mathbb{Q}$ be a splitting field of f with $\alpha \in E$ a root of f.

7.34.1 a

Show that E contains a primitive 7th root of unity.

7.34.2 b

Show that $E \neq \mathbb{Q}(\alpha)$.

7.35 Fall 2019 Midterm #6

Compute the Galois group of $f(x) = x^3 - 3x - 3 \in \mathbb{Q}[x]/\mathbb{Q}$.

7.36 Fall 2019 Midterm #7

Show that a field k of characteristic $p \neq 0$ is perfect $\iff$ for every $x \in k$ there exists a $y \in k$ such that $y^p = x$.

7.37 Fall 2019 Midterm #8

Let k be a field of characteristic $p \neq 0$ and $f \in k[x]$ irreducible. Show that $f(x) = g(x^{p^d})$ where $g(x) \in k[x]$ is irreducible and separable.

Conclude that every root of f has the same multiplicity p^d in the splitting field of f over k.

7.38 Fall 2019 Midterm #9

Let $n \geq 3$ and ζ_n be a primitive nth root of unity. Show that $[\mathbb{Q}(\zeta_n + \zeta_n^{-1}) : \mathbb{Q}] = \varphi(n)/2$ for φ the totient function. 10.

Let L/K be a finite normal extension.

- Show that if L/K is cyclic and E/K is normal with $L/E/K$ then L/E and E/K are cyclic.
- Show that if L/K is cyclic then there exists exactly one extension E/K of degree n with $L/E/K$ for each divisor n of $[L : K]$.

7.39 Spring 2021 #4

Define

$$f(x) \coloneqq x^4 + 4x^2 + 64 \in \mathbb{Q}[x].$$

a. Find the splitting field K of f over $\mathbb{Q}$.

b. Find the Galois group G of f.

c. Exhibit explicitly the correspondence between subgroups of G and intermediate fields between $\mathbb{Q}$ and K.

7.40 Spring 2021 #7

Let p be a prime number and let F be a field of characteristic p. Show that if $a \in F$ is not a pth power in F, then $x^p - a \in F[x]$ is irreducible.

7.41 Fall 2020 #3

a. Define what it means for a finite extension of fields E over F to be a *Galois* extension.

b. Determine the Galois group of $f(x) = x^3 - 7$ over $\mathbb{Q}$, and justify your answer carefully.

c. Find all subfields of the splitting field of $f(x)$ over $\mathbb{Q}$.

7.42 Fall 2020 #4

Let K be a Galois extension of F, and let $F \subset E \subset K$ be inclusions of fields. Let $G \coloneqq \mathrm{Gal}(K/F)$ and $H \coloneqq \mathrm{Gal}(K/E)$, and suppose H contains $N_G(P)$, where P is a Sylow p-subgroup of G for p a prime. Prove that $[E : F] \equiv 1 \pmod{p}$.

8 Modules

8.1 General Questions

8.1.1 Fall 2018 #6 ✨

Let R be a commutative ring, and let M be an R-module. An R-submodule N of M is maximal if there is no R-module P with $N \subsetneq P \subsetneq M$.

a Show that an R-submodule N of M is maximal $\iff$ M/N is a simple R-module: i.e., M/N is nonzero and has no proper, nonzero R-submodules.

b Let M be a $\mathbb{Z}$-module. Show that a $\mathbb{Z}$-submodule N of M is maximal $\iff$ $\#M/N$ is a prime number.

c Let M be the $\mathbb{Z}$-module of all roots of unity in $\mathbb{C}$ under multiplication. Show that there is no maximal $\mathbb{Z}$-submodule of M.

8.1.2 Fall 2019 Final #2

Consider the $\mathbb{Z}$-submodule N of $\mathbb{Z}^3$ spanned by $f_1 = [-1, 0, 1], f_2 = [2, -3, 1], f_3 = [0, 3, 1], f_4 = [3, 1, 5]$. Find a basis for N and describe $\mathbb{Z}^3/N$.

8.1.3 Spring 2018 #6

Let

$$M = \{(w,x,y,z) \in \mathbb{Z}^4 \;\middle|\; w+x+y+z \in 2\mathbb{Z}\},$$

and

$$N = \{(w,x,y,z) \in \mathbb{Z}^4 \;\middle|\; 4 \,\middle|\, (w-x),\ 4 \,\middle|\, (x-y),\ 4 \,\middle|\, (y-z)\}.$$

a Show that N is a $\mathbb{Z}$-submodule of M .

b Find vectors $u_1, u_2, u_3, u_4 \in \mathbb{Z}^4$ and integers d_1, d_2, d_3, d_4 such that

$$\{u_1, u_2, u_3, u_4\}$$

is a free basis for M, and

$$\{d_1u_1,\ d_2u_2,\ d_3u_3,\ d_4u_4\}$$

is a free basis for N .

c Use the previous part to describe M/N as a direct sum of cyclic $\mathbb{Z}$-modules.

8.1.4 Spring 2018 #7

Let R be a PID and M be an R-module. Let p be a prime element of R. The module M is called $\langle p\rangle$ *-primary* if for every $m \in M$ there exists $k > 0$ such that $p^k m = 0$.

a Suppose M is $\langle p\rangle$ -primary. Show that if $m \in M$ and $t \in R,\ t \notin \langle p\rangle$, then there exists $a \in R$ such that $atm = m$.

b A submodule S of M is said to be *pure* if $S \cap rM = rS$ for all $r \in R$. Show that if M is $\langle p\rangle$ -primary, then S is pure if and only if $S \cap p^k M = p^k S$ for all $k \geq 0$.

8.1.5 Fall 2016 #6

Let R be a ring and $f : M \to N$ and $g : N \to M$ be R-module homomorphisms such that $g \circ f = \mathrm{id}_M$. Show that $N \cong \operatorname{im} f \oplus \ker g$.

8.1.6 Spring 2016 #4

Let R be a ring with the following commutative diagram of R-modules, where each row represents a short exact sequence of R-modules:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & A & \xrightarrow{f} & B & \xrightarrow{g} & C & \longrightarrow & 0 \\ & & \downarrow{\scriptstyle \alpha} & & \downarrow{\scriptstyle \beta} & & \downarrow{\scriptstyle \gamma} & & \\ 0 & \longrightarrow & A' & \xrightarrow{f'} & B' & \xrightarrow{g'} & C' & \longrightarrow & 0 \end{array}$$

Prove that if α and γ are isomorphisms then β is an isomorphism.

8.1.7 Spring 2015 #8

Let R be a PID and M a finitely generated R-module.

a Prove that there are R-submodules

$$0 = M_0 \subset M_1 \subset \cdots \subset M_n = M$$

such that for all $0 \leq i \leq n-1$, the module M_{i+1}/M_i is cyclic.

b Is the integer n in part (a) uniquely determined by M? Prove your answer.

8.1.8 Fall 2012 #6

Let R be a ring and M an R-module. Recall that M is *Noetherian* iff any strictly increasing chain of submodule $M_1 \subsetneq M_2 \subsetneq \cdots$ is finite. Call a proper submodule $M' \subsetneq M$ *intersection-decomposable* if it can not be written as the intersection of two proper submodules $M' = M_1 \cap M_2$ with $M_i \subsetneq M$.

Prove that for every Noetherian module M, any proper submodule $N \subsetneq M$ can be written as a finite intersection $N = N_1 \cap \cdots \cap N_k$ of intersection-indecomposable modules.

8.1.9 Fall 2019 Final #1

Let A be an abelian group, and show A is a $\mathbb{Z}$-module in a unique way.

8.1.10 Fall 2020 #6

Let R be a ring with 1 and let M be a left R-module. If I is a left ideal of R, define

$$IM \coloneqq \left\{ \sum_{i=1}^{N<\infty} a_i m_i \;\middle|\; a_i \in I, m_i \in M, n \in \mathbb{N} \right\},$$

i.e. the set of finite sums of of elements of the form am where $a \in I, m \in M$.

a. Prove that $IM \leq M$ is a submodule.

b. Let M, N be left R-modules, I a nilpotent left ideal of R, and $f : M \to N$ an R-module morphism. Prove that if the induced morphism $\overline{f} : M/IM \to N/IN$ is surjective, then f is surjective.

8.2 Torsion and the Structure Theorem

8.2.1 ⋆ Fall 2019 #5

Let R be a ring and M an R-module.

Recall that the set of torsion elements in M is defined by

$$\mathrm{Tor}(M) = \{m \in M \;|\; \exists r \in R,\ r \neq 0,\ rm = 0\}.$$

a Prove that if R is an integral domain, then $\mathrm{Tor}(M)$ is a submodule of M .

b Give an example where $\mathrm{Tor}(M)$ is not a submodule of M.

c If R has zero-divisors, prove that every non-zero R-module has non-zero torsion elements.

8.2.2 ⋆ Spring 2019 #5

Let R be an integral domain. Recall that if M is an R-module, the *rank* of M is defined to be the maximum number of R-linearly independent elements of M .

a Prove that for any R-module M, the rank of $\mathrm{Tor}(M)$ is 0.

b Prove that the rank of M is equal to the rank of of $M/\mathrm{Tor}(M)$.

c Suppose that M is a non-principal ideal of R.

Prove that M is torsion-free of rank 1 but not free.

8.2.3 ⋆ Spring 2020 #6 ✨

Let R be a ring with unity.

a Give a definition for a free module over R.

b Define what it means for an R-module to be torsion free.

c Prove that if F is a free module, then any short exact sequence of R-modules of the following form splits:

$$0 \to N \to M \to F \to 0.$$

d Let R be a PID. Show that any finitely generated R-module M can be expressed as a direct sum of a torsion module and a free module.

You may assume that a finitely generated torsionfree module over a PID is free.

8.2.4 Spring 2012 #5 🚩

Let M be a finitely generated module over a PID R.

a M_t be the set of torsion elements of M, and show that M_t is a submodule of M.

b Show that M/M_t is torsion free.

c Prove that $M \cong M_t \oplus F$ where F is a free module.

8.2.5 Spring 2017 #5

Let R be an integral domain and let M be a nonzero torsion R-module.

a Prove that if M is finitely generated then the annihilator in R of M is nonzero.

b Give an example of a non-finitely generated torsion R-module whose annihilator is (0), and justify your answer.

8.2.6 Fall 2019 Final #3

Let $R = k[x]$ for k a field and let M be the R-module given by

$$M = \frac{k[x]}{(x-1)^3} \oplus \frac{k[x]}{(x^2+1)^2} \oplus \frac{k[x]}{(x-1)(x^2+1)^4} \oplus \frac{k[x]}{(x+2)(x^2+1)^2}.$$

Describe the elementary divisors and invariant factors of M.

8.2.7 Fall 2019 Final #4

Let $I = (2, x)$ be an ideal in $R = \mathbb{Z}[x]$, and show that I is not a direct sum of nontrivial cyclic R-modules.

8.2.8 Fall 2019 Final #5

Let R be a PID.

a Classify irreducible R-modules up to isomorphism.

b Classify indecomposable R-modules up to isomorphism.

8.2.9 Fall 2019 Final #6

Let V be a finite-dimensional k-vector space and $T : V \to V$ a non-invertible k-linear map. Show that there exists a k-linear map $S : V \to V$ with $T \circ S = 0$ but $S \circ T \neq 0$.

8.2.10 Fall 2019 Final #7

Let $A \in M_n(\mathbb{C})$ with $A^2 = A$. Show that A is similar to a diagonal matrix, and exhibit an explicit diagonal matrix similar to A.

8.2.11 Fall 2019 Final #10

Show that the eigenvalues of a Hermitian matrix A are real and that $A = PDP^{-1}$ where P is an invertible matrix with orthogonal columns.

8.2.12 Fall 2020 #7

Let $A \in \mathrm{Mat}(n \times n, \mathbb{R})$ be arbitrary. Make $\mathbb{R}^n$ into an $\mathbb{R}[x]$-module by letting $f(x).\mathbf{v} \coloneqq f(A)(\mathbf{v})$ for $f(\mathbf{v}) \in \mathbb{R}[x]$ and $\mathbf{v} \in \mathbb{R}^n$. Suppose that this induces the following direct sum decomposition:

$$\mathbb{R}^n \cong \frac{\mathbb{R}[x]}{\langle (x-1)^3 \rangle} \oplus \frac{\mathbb{R}[x]}{\langle (x^2+1)^2 \rangle} \oplus \frac{\mathbb{R}[x]}{\langle (x-1)(x^2-1)(x^2+1)^4 \rangle} \oplus \frac{\mathbb{R}[x]}{\langle (x+2)(x^2+1)^2 \rangle}.$$

a. Determine the elementary divisors and invariant factors of A.

b. Determine the minimal polynomial of A.

c. Determine the characteristic polynomial of A.

9 | Ring Theory

9.1 Spring 2021 #5

Suppose that $f(x) \in (\mathbb{Z}/n\mathbb{Z})[x]$ is a zero divisor. Show that there is a nonzero $a \in \mathbb{Z}/n\mathbb{Z}$ with $af(x) = 0$.

9.2 Spring 2021 #6

a. Carefully state the definition of **Noetherian** for a commutative ring R.

b. Let R be a subset of $\mathbb{Z}[x]$ consisting of all polynomials

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_nx^n$$

such that a_k is even for $1 \leq k \leq n$. Show that R is a subring of $\mathbb{Z}[x]$.

c. Show that R is not Noetherian.

Hint: consider the ideal generated by $\left\{2x^k \;\middle|\; 1 \leq k \in \mathbb{Z}\right\}$.

10 Linear Algebra: Diagonalizability

10.1 Fall 2017 #7

Let F be a field and let V and W be vector spaces over F .

Make V and W into $F[x]$-modules via linear operators T on V and S on W by defining $X \cdot v = T(v)$ for all $v \in V$ and $X \cdot w = S(w)$ for all w ∈ W .

Denote the resulting $F[x]$-modules by V_T and W_S respectively.

10.1.1 a

Show that an $F[x]$-module homomorphism from V_T to W_S consists of an F-linear transformation $R : V \to W$ such that $RT = SR$.

10.1.2 b

Show that $VT \cong WS$ as $F[x]$-modules $\iff$ there is an F-linear isomorphism $P : V \to W$ such that $T = P^{-1}SP$.

10.1.3 c

Recall that a module M is *simple* if $M \neq 0$ and any proper submodule of M must be zero. Suppose that V has dimension 2. Give an example of F, T with V_T simple.

10.1.4 d

Assume F is algebraically closed. Prove that if V has dimension 2, then any V_T is not simple.

10.2 Spring 2015 #3

Let F be a field and V a finite dimensional F-vector space, and let $A, B : V \to V$ be commuting F-linear maps. Suppose there is a basis $\mathcal{B}_1$ with respect to which A is diagonalizable and a basis $\mathcal{B}_2$ with respect to which B is diagonalizable.

Prove that there is a basis $\mathcal{B}_3$ with respect to which A and B are both diagonalizable.

10.3 Fall 2016 #2

Let A, B be two $n \times n$ matrices with the property that $AB = BA$. Suppose that A and B are diagonalizable. Prove that A and B are *simultaneously* diagonalizable.

10.4 Spring 2019 #1

Let A be a square matrix over the complex numbers. Suppose that A is nonsingular and that A^{2019} is diagonalizable over $\mathbb{C}$.

Show that A is also diagonalizable over $\mathbb{C}$.

11 Linear Algebra: Misc

11.1 ⋆ Spring 2012 #6

Let k be a field and let the group $G = \mathrm{GL}(m,k) \times \mathrm{GL}(n,k)$ acts on the set of $m \times n$ matrices $M_{m,n}(k)$ as follows:

$$(A,B) \cdot X = AXB^{-1}$$

where $(A,B) \in G$ and $X \in M_{m,n}(k)$.

11.1.1 a

State what it means for a group to act on a set. Prove that the above definition yields a group action.

11.1.2 b

Exhibit with justification a subset S of $M_{m,n}(k)$ which contains precisely one element of each orbit under this action.

11.2 ⋆ Spring 2014 #7

Let $G = \mathrm{GL}(3, \mathbb{Q}[x])$ be the group of invertible 3×3 matrices over $\mathbb{Q}[x]$. For each $f \in \mathbb{Q}[x]$, let S_f be the set of 3×3 matrices A over $\mathbb{Q}[x]$ such that $\det(A) = cf(x)$ for some nonzero constant $c \in \mathbb{Q}$.

11.2.1 a

Show that for $(P,Q) \in G \times G$ and $A \in S_f$, the formula

$$(P,Q) \cdot A \coloneqq PAQ^{-1}$$

gives a well defined map $G \times G \times S_f \to S_f$ and show that this map gives a group action of $G \times G$ on S_f.

11.2.2 b

For $f(x) = x^3(x^2+1)^2$, give one representative from each orbit of the group action in (a), and justify your assertion.

11.3 Fall 2012 #7

Let k be a field of characteristic zero and $A, B \in M_n(k)$ be two square $n \times n$ matrices over k such that $AB - BA = A$. Prove that $\det A = 0$.

Moreover, when the characteristic of k is 2, find a counterexample to this statement.

11.4 Fall 2012 #8

Prove that any nondegenerate matrix $X \in M_n(\mathbb{R})$ can be written as $X = UT$ where U is orthogonal and T is upper triangular.

11.5 Fall 2012 #5

Let U be an infinite-dimensional vector space over a field k, $f : U \to U$ a linear map, and $\{u_1, \cdots, u_m\} \subset U$ vectors such that U is generated by $\left\{u_1, \cdots, u_m, f^d(u_1), \cdots, f^d(u_m)\right\}$ for some $d \in \mathbb{N}$.

Prove that U can be written as a direct sum $U \cong V \oplus W$ such that

1. V has a basis consisting of some vector $v_1, \cdots v_n, f^d(v_1), \cdots, f^d(v_n)$ for some $d \in \mathbb{N}$, and
2. W is finite-dimensional.

Moreover, prove that for any other decomposition $U \cong V' \oplus W'$, one has $W' \cong W$.

11.6 Fall 2015 #7

11.6.1 a

Show that two 3×3 matrices over $\mathbb{C}$ are similar $\iff$ their characteristic polynomials are equal and their minimal polynomials are equal.

11.6.2 b

Does the conclusion in (a) hold for 4×4 matrices? Justify your answer with a proof or counterexample.

11.7 Fall 2014 #4

Let F be a field and T an $n \times n$ matrix with entries in F. Let I be the ideal consisting of all polynomials $f \in F[x]$ such that $f(T) = 0$.

Show that the following statements are equivalent about a polynomial $g \in I$:

a. g is irreducible.

b. If $k \in F[x]$ is nonzero and of degree strictly less than g, then $k[T]$ is an invertible matrix.

11.8 Fall 2015 #8

Let V be a vector space over a field F and $V^\vee$ its dual. A *symmetric bilinear form* $(\,\cdot\,,\,\cdot\,)$ on V is a map $V \times V \to F$ satisfying

$$(av_1 + bv_2, w) = a(v_1, w) + b(v_2, w) \quad \text{and} \quad (v_1, v_2) = (v_2, v_1)$$

for all $a, b \in F$ and $v_1, v_2 \in V$. The form is *nondegenerate* if the only element $w \in V$ satisfying $(v, w) = 0$ for all $v \in V$ is $w = 0$.

Suppose $(\,\cdot\,,\,\cdot\,)$ is a nondegenerate symmetric bilinear form on V. If W is a subspace of V, define

$$W \perp := \left\{ v \in V \;\middle|\; (v, w) = 0 \text{ for all } w \in W \right\}.$$

11.8.1 a

Show that if X, Y are subspaces of V with $Y \subset X$, then $X \perp \subseteq Y \perp$.

11.8.2 b

Define an injective linear map

$$\psi : Y \perp / X \perp \hookrightarrow (X/Y)^\vee$$

which is an isomorphism if V is finite dimensional.

11.9 Fall 2018 #4 ✨

Let V be a finite dimensional vector space over a field (the field is not necessarily algebraically closed).

Let $\varphi : V \to V$ be a linear transformation. Prove that there exists a decomposition of V as $V = U \oplus W$, where U and W are φ-invariant subspaces of V , $\varphi|_U$ is nilpotent, and $\varphi|_W$ is nonsingular.

Revisit.

11.10 Fall 2018 #5 ✨

Let A be an $n \times n$ matrix.

11.10.1 a

Suppose that v is a column vector such that the set $\{v, Av, ..., A^{n-1}v\}$ is linearly independent. Show that any matrix B that commutes with A is a polynomial in A.

11.10.2 b

Show that there exists a column vector v such that the set $\{v, Av, ..., A^{n-1}v\}$ is linearly independent $\iff$ the characteristic polynomial of A equals the minimal polynomial of A.

11.11 Fall 2019 #8 🚩

Let $\{e_1, \cdots, e_n\}$ be a basis of a real vector space V and let

$$\Lambda := \left\{ \sum r_i e_i \;\middle|\; r_i \in \mathbb{Z} \right\}$$

Let $\cdot$ be a non-degenerate ($v \cdot w = 0$ for all $w \in V \iff v = 0$) symmetric bilinear form on V such that the Gram matrix $M = (e_i \cdot e_j)$ has integer entries.

Define the dual of Λ to be

$$\Lambda^\vee \coloneqq \{v \in V \;\middle|\; v \cdot x \in \mathbb{Z} \text{ for all } x \in \Lambda\}.$$

11.11.1 a

Show that $\Lambda \subset \Lambda^\vee$.

11.11.2 b

Prove that $\det M \neq 0$ and that the rows of M^{-1} span $\Lambda^\vee$.

11.11.3 c

Prove that $\det M = |\Lambda^\vee / \Lambda|$.

Todo, missing part (c).

11.12 Spring 2013 #6 ✨

Let V be a finite dimensional vector space over a field F and let $T : V \to V$ be a linear operator with characteristic polynomial $f(x) \in F[x]$.

11.12.1 a

Show that $f(x)$ is irreducible in $F[x] \iff$ there are no proper nonzero subspaces $W < V$ with $T(W) \subseteq W$.

11.12.2 b

If $f(x)$ is irreducible in $F[x]$ and the characteristic of F is 0, show that T is diagonalizable when we extend the field to its algebraic closure.

Is there a proof without matrices? What if V is infinite dimensional?

How to extend basis?

11.13 Fall 2020 #8

Let $A \in \mathrm{Mat}(n \times n, \mathbb{C})$ such that the group generated by A under multiplication is finite. Show that

$$\mathrm{Tr}(A^{-1}) = \overline{\mathrm{Tr}(A)},$$

where $\bar{\cdot}$ denotes taking the complex conjugate and $\mathrm{Tr}(\,\cdot\,)$ is the trace.

12 Linear Algebra: Canonical Forms

12.1 ⋆ Spring 2012 #8

Let V be a finite-dimensional vector space over a field k and $T : V \to V$ a linear transformation.

12.1.1 a

Provide a definition for the *minimal polynomial* in $k[x]$ for T.

12.1.2 b

Define the *characteristic polynomial* for T.

12.1.3 c

Prove the Cayley-Hamilton theorem: the linear transformation T satisfies its characteristic polynomial.

12.2 ⋆ Spring 2020 #8

Let $T : V \to V$ be a linear transformation where V is a finite-dimensional vector space over $\mathbb{C}$. Prove the Cayley-Hamilton theorem: if $p(x)$ is the characteristic polynomial of T, then $p(T) = 0$. You may use canonical forms.

12.3 ⋆ Spring 2012 #7

Consider the following matrix as a linear transformation from $V := \mathbb{C}^5$ to itself:

$$A = \begin{pmatrix} -1 & 1 & 0 & 0 & 0 \\ -4 & 3 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 2 \end{pmatrix}.$$

12.3.1 a

Find the invariant factors of A.

12.3.2 b

Express V in terms of a direct sum of indecomposable $\mathbb{C}[x]$-modules.

12.3.3 c

Find the Jordan canonical form of A.

12.4 Fall 2019 Final #8

Exhibit the rational canonical form for

- $A \in M_6(\mathbb{Q})$ with minimal polynomial $(x-1)(x^2+1)^2$.
- $A \in M_{10}(\mathbb{Q})$ with minimal polynomial $(x^2+1)^2(x^3+1)$.

12.5 Fall 2019 Final #9

Exhibit the rational and Jordan canonical forms for the following matrix $A \in M_4(\mathbb{C})$:

$$A = \begin{pmatrix} 2 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ -2 & -2 & 0 & 1 \\ -2 & 0 & -1 & -2 \end{pmatrix}.$$

12.6 Spring 2016 #7

Let $D = \mathbb{Q}[x]$ and let M be a $\mathbb{Q}[x]$-module such that

$$M \cong \frac{\mathbb{Q}[x]}{(x-1)^3} \oplus \frac{\mathbb{Q}[x]}{(x^2+1)^3} \oplus \frac{\mathbb{Q}[x]}{(x-1)(x^2+1)^5} \oplus \frac{\mathbb{Q}[x]}{(x+2)(x^2+1)^2}.$$

Determine the elementary divisors and invariant factors of M.

12.7 Spring 2020 #7

Let

$$A = \begin{bmatrix} 2 & 0 & 0 \\ 4 & 6 & 1 \\ -16 & -16 & -2 \end{bmatrix} \in M_3(\mathrm{C}).$$

12.7.1 a

Find the Jordan canonical form J of A.

12.7.2 b

Find an invertible matrix P such that $P^{-1}AP = J$. You should not need to compute P^{-1}.

12.7.3 c

Write down the minimal polynomial of A.

12.8 Spring 2019 #7 ✨

Let p be a prime number. Let A be a $p \times p$ matrix over a field F with 1 in all entries except 0 on the main diagonal.

Determine the Jordan canonical form (JCF) of A

12.8.1 a

When $F = \mathbb{Q}$,

12.8.2 b

When $F = \mathbb{F}_p$.

Hint: In both cases, all eigenvalues lie in the ground field. In each case find a matrix P such that $P^{-1}AP$ is in JCF.

12.9 Spring 2018 #4 🚩

Let

$$A = \begin{bmatrix} 0 & 1 & -2 \\ 1 & 1 & -3 \\ 1 & 2 & -4 \end{bmatrix} \in M_3(\mathbb{C})$$

12.9.1 a

Find the Jordan canonical form J of A.

12.9.2 b

Find an invertible matrix P such that $P^{-1}AP = J$.

You should not need to compute P^{-1}.

12.10 Spring 2017 #6

Let A be an $n \times n$ matrix with all entries equal to 0 except for the $n-1$ entries just above the diagonal being equal to 2.

12.10.1 a

What is the Jordan canonical form of A, viewed as a matrix in $M_n(\mathbb{C})$?

12.10.2 b

Find a nonzero matrix $P \in M_n(\mathbb{C})$ such that $P^{-1}AP$ is in Jordan canonical form.

12.11 Spring 2016 #1

Let

$$A = \begin{pmatrix} -3 & 3 & -2 \\ -7 & 6 & -3 \\ 1 & -1 & 2 \end{pmatrix} \in M_3(\mathrm{C}).$$

12.11.1 a

Find the Jordan canonical form J of A.

12.11.2 b

Find an invertible matrix P such that $P^{-1}AP = J$. You do not need to compute P^{-1}.

12.12 Spring 2015 #6

Let F be a field and n a positive integer, and consider

$$A = \begin{bmatrix} 1 & \dots & 1 \\ & \ddots & \\ 1 & \dots & 1 \end{bmatrix} \in M_n(F).$$

Show that A has a Jordan normal form over F and find it.

Hint: treat the cases $n \cdot 1 \neq 0$ in F and $n \cdot 1 = 0$ in F separately.

12.13 Fall 2014 #5

Let T be a 5×5 complex matrix with characteristic polynomial $\chi(x) = (x-3)^5$ and minimal polynomial $m(x) = (x-3)^2$. Determine all possible Jordan forms of T.

12.14 Spring 2013 #5

Let $T : V \to V$ be a linear map from a 5-dimensional $\mathbb{C}$-vector space to itself and suppose $f(T) = 0$ where $f(x) = x^2 + 2x + 1$.

12.14.1 a

Show that there does not exist any vector $v \in V$ such that $Tv = v$, but there *does* exist a vector $w \in V$ such that $T^2 w = w$.

12.14.2 b

Give all of the possible Jordan canonical forms of T.

12.15 Spring 2021 #1

Let m

$$A \coloneqq \begin{bmatrix} r & 1 & -1 \\ -6 & -1 & 2 \\ 2 & 1 & 1 \end{bmatrix} \in \mathrm{Mat}(3 \times 3, \mathbb{C}).$$

a. Find the Jordan canonical form J of A.

b. Find an invertible matrix P such that $J = P^{-1}AP$.

(You should not need to compute P^{-1})

c. Write down the minimal polynomial of A.

12.16 Fall 2020 #5

Consider the following matrix:

$$B \coloneqq \begin{bmatrix} 1 & 3 & 3 \\ 2 & 2 & 3 \\ -1 & -2 & -2 \end{bmatrix}.$$

a. Find the minimal polynomial of B.

b. Find a 3×3 matrix J in Jordan canonical form such that $B = JPJ^{-1}$ where P is an invertible matrix.