

Title

D. Zack Garza

Table of Contents

Contents

Table of Contents	2
1 The Cotangent Complex and Derived de Rham Cohomology	3
1.1 Motivation	3
1.1.1 Derived Schemes	4
1.2 Simplicial Rings	4
1.3 Examples of Simplicial Categories	5
1.3.1 Example: The Cotangent Complex	8
1.4 Derived de Rham Cohomology	10

1 | The Cotangent Complex and Derived de Rham Cohomology

Reference: MSRI Workshop on Derived AG, Birational Geometry, Moduli Spaces.

Video: <https://www.youtube.com/watch?v=zRPa-VAv16Q>

1.1 Motivation

Basic affine objects in AG: commutative rings, replace with simplicial commutative rings which we'll use as a base diagram. Later: derived stacks and geometric derived stacks.

An evolution of objects. and how we can think about them.

- Algebraic schemes/spaces, e.g. \mathbb{P}^n
 - Think of these as étale sheaves of sets (think functor of points), identified as discrete spaces $\mathcal{S}_{\leq 0}$, so every component is contractible and there are no higher homotopy groups
- The Picard stack $\underline{\mathrm{Pic}}_{X/k}$ for X a smooth and proper k -scheme, which is an Artin stack (a subclass Deligne-Mumford stacks). Note that this still has automorphisms given by global units on X .
 - Think of these as étale sheaves of groupoids $\mathcal{S}_{\leq 1}$.
 - Can take a fundamental groupoid Also note that this is the mapping stack $\underline{\mathrm{Map}}(X, K(\mathbb{G}_m, 1))$.
- $K(\mathbb{G}_m, n)$ a higher stack
 - A sheaf taking values in n -truncated spaces, i.e. a space where when basing at any point, there are no homotopy groups above degree n .
 - Stack with a single point, where the isotopy is $K(\mathbb{G}_m, n-1)$.

Note that these are all built from affine schemes with a few acceptable moves.

Example 1.1.1(?): We can realize

$$B\mathbb{G}_m = K(\mathbb{G}_m, 1) \cong [\{\mathrm{pt}\}/\mathbb{G}_m]$$

in stack notation. Similarly,

$$K(\mathbb{G}_m, 2) = [\{\mathrm{pt}\}/B\mathbb{G}_m],$$

which is a smooth Artin stack. Mapping into this gives the Picard groupoid of a scheme. It's a higher geometric stack that still has smoothness properties.

Question 1.1.2: What does it mean to give a map from a scheme X into a higher stack?

The world of étale schemes taking values in $\mathcal{S}_{\leq n}$ is enriched in topological spaces. There is a topological space

$$M := \mathrm{Map}(X, K(\mathbb{G}_m, n))$$

The homotopy groups are

$$\pi_i M = \begin{cases} H_{\mathrm{\acute{e}t}}^{n-i}(X, \mathbb{G}_m) & 0 \leq i \leq n \\ 0 & \text{else} \end{cases},$$

so this higher geometric stack that says something about higher cohomology groups. We thus have étale sheaves taking values in higher topological spaces, and has some geometric meaning. They're also built from geometric objects: iterating taking quotients by smooth actions. $K(\mathbb{G}_m, 1)$ is a quotient by a smooth algebraic group, $K(\mathbb{G}_m, 2)$ is now a smooth *Artin stack*, and we can keep going. This is the fundamental process for building geometric higher stacks.

Remark 1.1.3: Why derive things? Schemes are equipped with sheaves of commutative rings, so the basic idea is let the sheaves take values in groupoids, stacks, etc. So we can consider replacing the structure sheaf \mathcal{O}_X is itself a sheaf of spaces, and this is the fundamental idea of derived algebraic geometry.

1.1.1 Derived Schemes

Consider $\mathrm{Spec} k \otimes_{k[x]}^L k$, a derived tensor product. This is a simplicial commutative ring, and the basic version of an affine derived scheme. This is a complex C^\cdot with homology in degree 0 and 1, where $H_1 = \mathrm{Tor}^1(k \otimes_{k[x]} k)$. So analogously, we'll start with derived schemes and take quotients by smooth groups. In the end, we get derived stacks.

Example 1.1.4(Fundamental): An example is \mathcal{M}_φ , the moduli of objects in some DG category \mathcal{C} .

1.2 Simplicial Rings

We need to agree on what the local affine modules will look like. For our purposes, they'll be simplicial commutative rings. Consider the derived category $\mathcal{C} := D(\mathbb{Z})_{\geq 0}$ and its connective objects, which are chain complexes C^\cdot where $H_{<0}(C^\cdot) = 0$. There is a **derived tensor product** \otimes^L which makes \mathcal{C} into a symmetric monoidal category, . Basic idea: we want to look at commutative algebra objects in this symmetric monoidal category $(D(\mathbb{Z}), \otimes^L)$.¹ Note that of working in a symmetric monoidal abelian category, we will be looking at connective chain complexes, and simplicial rings are one way of studying commutative algebra objects here.

¹This is a familiar move: people in the 60s knew one could do AG in some ambient symmetric monoidal abelian category.

[^def:connective] **Connective** means $H_{<0} = 0$.

We have some choices for making sense of DAG:

- E_∞ -ring spectra
- Simplicial commutative rings, Note that this is what we will choose.
- Over \mathbb{Q} , \mathbb{Q} -commutative DGAs.

Definition 1.2.1 (Simplicial Commutative Ring)

Let Δ denote the *simplex category*, the category of non-empty finite ordered sets with order-preserving maps. We have the following situation:

$$\begin{array}{ccccc} [0] & \xrightarrow{\quad} & [1] & \xrightarrow{\quad} & [2] & \cdots \\ & \xleftarrow{\quad} & & \xleftarrow{\quad} & & \\ & \xrightarrow{\quad} & & \xrightarrow{\quad} & & \end{array}$$

$$\{0\} \quad \{0 \rightarrow 1\} \quad \{0 \rightarrow 1 \rightarrow 2\} \quad \cdots$$

The arrows going up are **face maps** (or **coface maps**), and the others are **degeneracy** maps. If \mathcal{C} is a category, then $s\mathcal{C} := \text{Fun}(\Delta^{\text{op}}, \mathcal{C})$ is the category of simplicial objects of \mathcal{C} .

1.3 Examples of Simplicial Categories

Example 1.3.1(a, simplicial sets and topological spaces.): $s\text{Sets} \simeq \text{Top}$: this is not an equivalence of categories, but rather they equivalent homotopy theories (theory up to weak equivalence). There are notions of weak equivalence (isomorphism on π_0 , and for each choice of basepoint, an isomorphism on all $\pi_{\geq 1}$ on each side).

Here there is an n -simplex on the LHS ($s\text{Sets}$),

$$\Delta^n = \text{hom}_\Delta(\cdot, [n])$$

and on the RHS we have

$$\Delta_{\text{Top}}^n := \left\{ [x_0, \dots, x_n] \in \mathbb{R}^n \mid x_i \geq 0, \sum x_i = 1 \right\}$$

If you make a functor $\Delta^n \rightleftharpoons \Delta_{\text{Top}}^n$, then by Yoneda the presheaf category $\text{Presh}(\Delta) := \text{Fun}(\Delta^{\text{op}}, \text{Set})$ is generated by representable objects. Everything on in $s\text{Sets}$ is generated by taking colimits of the Δ^n , so we can make some assignment and extend by colimits to get a functor $s\text{Sets} \rightarrow \text{Top}$. We have a notion of weak equivalence for Top , and so the notion of weak equivalence on $s\text{Sets}$ is just given by pullback along the functor $s\text{Sets} \rightarrow \text{Top}$, and this induces an equivalence of homotopy theories.

The functor back $\text{Top} \rightarrow s\text{Sets}$ is the **singular complex construction**. Considering Δ_{Top}^n , this is a cosimplicial object in Top .

Remark 1.3.2: \mathbf{Top} will denote that 1-category, while $\mathcal{T}\mathbf{op}$ will be its full ∞ -category.

So we have a natural cosimplicial object in \mathbf{Top} , so $\mathrm{Sing}(X) := \mathrm{Hom}_{\mathbf{Top}}(\Delta_{\mathbf{Top}}^\bullet, X)$ is a simplicial object in \mathbf{sSets} . As in singular homology, we can get a simplicial abelian group by taking the free abelian group $\mathbb{Z}[\mathrm{Sing}(X)]$. Note that this is just composing functors $\Delta^{\mathrm{op}} \rightarrow \mathbf{Set}$ and $\mathbf{Set} \rightarrow \mathbb{Z}\text{-mod}$. We can use this to create a chain complex $C.(\mathbb{Z}[\mathrm{Sing}(X)])$, and as expected, $H_i(C.) \cong H_i^{\mathrm{Sing}}(X, \mathbb{Z})$, the singular homology.

Example 1.3.3(b): We can take simplicial abelian groups \mathbf{sAb} and the connective objects $D(\mathbb{Z})_{\geq 0}$; this have equivalent homotopy theories. There is a notion of weak equivalence on the RHS, quasi-isomorphism, and asking if the literal underlying spaces on the LHS are weakly equivalence as spaces. A specific way of doing this is the **Dold-Kan** correspondence: suppose we have a simplicial abelian group $M.$, then we have face maps going to the left:

$$M_0 \begin{array}{c} \xleftarrow{\partial_1} \\ \xleftarrow{\partial_2} \end{array} M_1 \begin{array}{c} \xleftarrow{\quad} \\ \xleftarrow{\quad} \\ \xleftarrow{\quad} \end{array} M_2 \begin{array}{c} \xleftarrow{\quad} \\ \xleftarrow{\quad} \\ \xleftarrow{\quad} \end{array} \dots$$

We make this into a chain complex by setting the differential to a graded sum:

$$M_0 \xleftarrow{\Sigma(-)^i \partial_i} M_1 \xleftarrow{\Sigma(-)^i \partial_i} M_2 \xleftarrow{\quad} \dots$$

The homology of this complex turns out to be the same as the homotopy groups of the simplicial abelian group viewed as a topological space.

Example 1.3.4(c, *Simplicial Commutative k -Algebras*): Defined as

$$\mathbf{sCAlg}_k = \mathrm{Fun}(\Delta^{\mathrm{op}}, \mathbf{CAlg}_k),$$

where k is some commutative ring. This was studied by Quillen, and an impetus for model categories. Models give a notion of weak equivalence and a “right way” of computing: for the usual derived category of a ring, this is taking resolutions. So the LHS is sometimes called a **non-abelian derived category**.

Fact 1.3.5: For $R \in \mathbf{sCAlg}$, $\pi_* R$ has a graded commutative ring structure: $xy = (-1)^{|x||y|}yx$ and $x^2 = 0$ for elements x with $|x|$ odd. This is useful because it gives us some graded ring to associated to R .

Remark 1.3.6: The category of simplicial abelian groups is equivalent to $\mathrm{Ch}(\mathbb{Z})_{\geq 0}$; i.e. chain complexes of abelian groups concentrated in non-negative degree. This also yields an equivalence of homotopy theories.

A different perspective on simplicial commutative rings: there is an adjunction from sets to commutative k -algebras

$$\begin{aligned} \text{Set} &\rightleftarrows \text{CAlg}_k \\ S &\mapsto K[S] \\ &\cdot \end{aligned}$$

i.e. we send a set to the polynomial ring generated by S . Any time such an adjunction exists, given an $R \in \text{CAlg}_k$ we can construct a *simplicial resolution* S^\cdot and a map $S^\cdot \rightarrow R$. This resolution has the following structure:

$$S^0 = k[R], \text{ the free commutative algebra on } R.$$

Using the unit and counit maps of the adjunction, one obtains a canonical simplicial object,

$$k[R] \begin{array}{c} \xrightarrow{\quad} \\ \xleftarrow{\quad} \end{array} k[k[R]] \begin{array}{c} \xrightarrow{\quad} \\ \xleftarrow{\quad} \\ \xrightarrow{\quad} \\ \xleftarrow{\quad} \end{array} \dots$$

Moreover, $S^\cdot \xrightarrow{\sim} R$ is a homotopy equivalence. So we've taken an arbitrary k -algebra and replaced it with a *simplicial* k -algebra which is given by polynomial rings in each degree, typically in infinitely many variables, which has the same homology. This is the analog of a projective resolution.

Now define $\text{CAlg}_k^{\text{poly}}$ as the category of finitely generated polynomial rings, and suppose you have a functor

$$\text{CAlg}_k^{\text{poly}} \xrightarrow{F} \mathcal{C}$$

where \mathcal{C} is a “reasonable” category or possible in ∞ -category. We can consider the category $\text{Ind}(\text{CAlg}_k^{\text{poly}})$ given by formally adjoining filtered colimits. We have the following diagram,

$$\begin{array}{ccc} \text{CAlg}_k^{\text{poly}} & \xrightarrow{F} & \mathcal{C} \\ \downarrow & \nearrow \exists \tilde{F} & \uparrow \\ \text{Ind}(\text{CAlg}_k^{\text{poly}}) & & \\ \downarrow \text{Constant diagrams} & \nearrow \exists LF & \\ \text{sCAlg}_k & & \end{array}$$

where the bottom inclusion is given by viewing a commutative ring as the constant simplicial commutative ring, and the extension \tilde{F} exists by applying F to any colimit diagram. The functor LF is a derived functor that exists if \mathcal{C} has certain colimits.

So starting with a functor defined on finitely generated polynomial rings, i.e. affine spaces, we get a simplicial derived functor. For $R \in \text{CAlg}_k$, using $S^\cdot \xrightarrow{\sim} R$, we can apply F level-wise to get a new simplicial object $F(S^\cdot) \in \mathcal{C}$. Then $LF(R)$ is defined by taking the colimit over Δ yields the geometric realization, i.e.

$$LF(R) := \text{hocolim}_{\Delta} F(S^\cdot) = |F(S^\cdot)|$$

So we can promote functors on polynomial rings to functors on simplicial commutative rings. This ends up being a Kan extension.

There is a nice universal property here, namely that functors out of \mathbf{sAlg}_k are equivalent to functors out of $\mathbf{CAlg}_k^{\text{poly}}$ that satisfying some additional properties.

1.3.1 Example: The Cotangent Complex

There are 3 equivalent ways to view this complex.

Example 1.3.7(a): Use this derived functor approach: it will essentially be the derived functor of taking Ω^1 . Suppose $k \rightarrow R$ in commutative rings, denote by S^\cdot again the canonical resolution of R , yielding a diagram


$$\begin{array}{ccc} k & \xrightarrow{\quad} & R \\ & \searrow & \nearrow \sim \\ & S^\cdot & \end{array}$$

Take Kahler differentials degree-wise to get $\Omega_{S^\cdot/k}^1$. Now base-change along this map to get a simplicial R -module,

$$\Omega_{S^\cdot/k}^1 \otimes_{S^\cdot} R \in \mathbf{sMod}\text{-}R.$$

We can now use Dold-Kan to view this as a connective object in $D(R)_{\geq 0}$, the derived category of R , yielding $\mathbf{sMod}\text{-}R \simeq D(R)_{\geq 0}$, so we define the **cotangent complex of R/k** as

$$L_{R/k} := \Omega_{S^\cdot/k}^1 \otimes_{S^\cdot} R \in D(R)_{\geq 0}.$$


This turns out to work if you take a resolution other than the canonical one. A downside to this definition is that it's not clear how/why it might depend on the resolution. 

Example 1.3.8(b): We can define them by taking a map

$$\mathbf{CAlg}_k^{\text{poly}} \xrightarrow{\Omega^\cdot/k} D(k)_{\geq 0}$$

to the connective objects in the derived category of k . We can extend to get a diagram


$$\begin{array}{ccc} \mathbf{CAlg}_k^{\text{poly}} & \xrightarrow{\Omega^\cdot/k} & D(k)_{\geq 0} \\ \downarrow & \nearrow \Omega^\cdot/k & \\ \mathbf{sAlg}_k & & \end{array}$$

It then turns out that $L_{R/k} \simeq L\Omega_{R/k}^1$. Here it's not quite clear why this lands in k -mod instead of R -mod, or what the R -module structure works here. 

Example 1.3.9(c): As a universal property: Let $k \rightarrow R \rightarrow S$ where k is a commutative ring and R, S are ordinary (or simplicial) commutative rings. Let $M \in S\text{-mod}$ in $D(S)_{\geq 0}$. There is a natural enrichment in topological spaces, so we'll write Map for homs with their topological structure. In particular, these have homotopy groups. The universal property that the cotangent complex has comes from an equivalence

$$\text{Map}_R(L_{R/k}, M) \simeq \text{Map}_{\text{sCAlg}_{k//S}}(R, S \oplus M).$$

where the latter is in the category of simplicial commutative k -algebras with a *fixed* map to S , a bit like a comma category. This makes sense because $R \rightarrow S$ is a fixed map, and $S \oplus M$ has a projection to S , i.e. this is a square zero extension.

Remark 1.3.10: The point of simplicial commutative rings is that this extension still makes sense, even when M is a chain complex. It's still possible to make this into a simplicial commutative ring. 


This may look familiar from the exercises in Hartshorne, since it resembles the definition of Kahler differentials.

Exercise 1.3.11(?): Prove that

$$\pi_0 L_{R/k} \cong \Omega_{R/k}^1.$$


This is exactly lifting through the square zero extension. This is similar to having a lift

$$\begin{array}{ccc} & S \oplus M & \\ \tilde{f} \nearrow & \downarrow & \\ R & \xrightarrow{f} & S \end{array}$$

where giving \tilde{f} is like giving a derivation $R \rightarrow M$. So the RHS comma category mapping space in the previous equivalence is denoted the “space of derivations of R and M ” (whatever that means) 

Exercise 1.3.12(?): You can base-change the cotangent complex of R/K to S and obtain an exact triangle/cofiber sequence


$$S \otimes_R^L L_{R/k} \longrightarrow L_{S/k} \longrightarrow L_{S/R}$$

This follows from the universal property above, and is a form of “transitivity”. 

Exercise 1.3.13(?): Given a diagram

$$\begin{array}{ccc} K & \longrightarrow & R \\ \downarrow & & \downarrow \\ T & \longrightarrow & R \otimes_k^L T \end{array}$$

show that

$$T \otimes_k^L L_{R/k} \cong L_{R \otimes_k^L T/T}.$$


All of these follow from just the mapping space property, but require thinking about what it means to compute maps in a comma category.

Exercise 1.3.14(?): Let R be a perfect \mathbb{F}_p -algebra, so the Frobenius is an isomorphism:

$$F: R \xrightarrow{\sim} R \\ x \mapsto x^p.$$

Show that $L_{R/\mathbb{F}_p} \cong 0$.

1.4 Derived de Rham Cohomology

We can make a specific functor:

$$\mathrm{CAlg}_k^{\mathrm{poly}} \rightarrow D(k) \\ R \mapsto \mathrm{dR}_{R/k} := \left(R \rightarrow \Omega_{R/k}^1 \rightarrow \Omega_{R/k}^2 \rightarrow \cdots \right),$$

which takes a polynomial ring R to its algebraic de Rham complex. If R is finitely generated, then the complex is bounded. One can formally extend by deriving this functor to obtain

$$\begin{array}{ccc} \mathrm{CAlg}_k^{\mathrm{poly}} & \xrightarrow{\mathrm{dR}_{\cdot/k}} & D(k) \\ \downarrow & \nearrow L\mathrm{dR}_{\cdot/k} & \\ \mathrm{sCAlg}_k & & \end{array}$$

Remark 1.4.1: The point is that we know de Rham cohomology behaves well for smooth things, and we want to extend to non-smooth things.

⚠ Warning 1.4.2

Suppose $k \in \mathrm{Alg}_{\mathbb{Q}}$, noting that we're in characteristic zero, and let $R \in \mathrm{CAlg}_k^{\mathrm{poly}}$ be a polynomial ring. Then $\mathrm{dR}_{R/k} \cong k$ by the Poincaré lemma, just the cohomology of affine space. Then $L\mathrm{dR}_{S/k} \cong k$, which is disappointing.

How to fix this: we have the Hodge filtration $F_H \mathrm{dR}_{R/k}$, and the graded pieces are given by

$$\mathrm{gr}_H^i \mathrm{dR}_{R/k} \cong \Omega_{R/k}^i[-i]$$

Taking these gradings is compatible