

Exercises for Section 1.1: Norm and Inner Product

width=4in,totalheight=4in

1. Define the ℓ^1 -norm on n by

$$\|x\|_1 = \sum_{i=1}^n |x^i|,$$

and define the sup-norm on n by

$$\|x\|_\infty = \sup \{|x^i|\}.$$

Show that these satisfy Theorem 1.

Proof.

■

2. Prove that $\|x\| \leq \sum_{i=1}^n |x^i|$. In other words, the usual norm is no greater than the ℓ^1 -norm.

Proof.

■

3. Prove that $\|x - y\| \leq \|x\| + \|y\|$. (Compare this with part (2) of Theorem 1.) When does equality hold?

4. Prove that $\left| \|x\| - \|y\| \right| \leq \|x - y\|$.

5. The quantity $\|y - x\|$ is called the distance between x and y . Prove and interpret the “triangle inequality”:

$$\|z - x\| \leq \|z - y\| + \|y - x\|.$$

6. Let f and g be integrable on $[a, b]$.

- a) Prove the integral version of the Cauchy-Schwarz inequality:

$$\left| \int_a^b fg \right| \leq \left(\int_a^b f^2 \right)^{1/2} \left(\int_a^b g^2 \right)^{1/2}.$$

Hint: Consider separately the cases $0 = \int_a^b (f - tg)^2$ for some $t \in \mathbb{R}$, and $0 < \int_a^b (f - tg)^2$ for all $t \in \mathbb{R}$.

- b) If equality holds, must $f = tg$ for some $t \in \mathbb{R}$? What if f and g are continuous?

c) Show that the Cauchy-Schwarz inequality is a special case of (a).

7. A linear transformation $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is norm preserving if

$$\|T(x)\| = \|x\|,$$

for all $x \in \mathbb{R}^n$, and inner product preserving if

$$\langle Tx, Ty \rangle = \langle x, y \rangle,$$

for all $x, y \in \mathbb{R}^n$.

a) Prove that T is norm preserving if and only if it is inner product preserving.

b) Prove that such a linear transformation is 1-1, and T^{-1} is norm preserving (and inner product preserving).

8. If $T : \mathbb{R}^m \rightarrow \mathbb{R}^n$ is a linear transformation, show that there is a number M such that $\|T(h)\| \leq M\|h\|$ for all $h \in \mathbb{R}^m$. Hint: Estimate $\|T(h)\|$ in terms of $\|h\|$ and the entries in the matrix for T .

9. If $x, y \in \mathbb{R}^n$, and $z, w \in \mathbb{R}^m$, show that $ds\langle (x, z), (y, w) \rangle = \langle x, y \rangle + \langle z, w \rangle$, and $ds\|(x, z)\| = \sqrt{\|x\|^2 + \|z\|^2}$. Note that (x, z) and (y, w) denote points in $\mathbb{R}^{n+m}$.

10. If $x, y \in \mathbb{R}^n$, then x and y are called perpendicular (or orthogonal), and we write $x \perp y$, if $\langle x, y \rangle = 0$. If $x \perp y$, prove that $\|x + y\|^2 = \|x\|^2 + \|y\|^2$.

Exercises for Section 1.2: More Topology: Open and Closed Sets in $\mathbb{R}^n$

width=4in,totalheight=4in

1. Prove that the union of any (even infinite) number of open sets is open. Prove that the intersection of two (and hence of finitely many) open sets is open. Give a counterexample for the intersection of infinitely many open sets.

2. If $A \subset B \subset \mathbb{R}^n$, prove that

$$\text{cl}A \subset \text{cl}B, \quad \text{and} \quad A \subset B.$$

3. Prove that if B is an open subset of A , then $B \subset \text{int}(A)$. Note that this says that $\text{int}(A)$ is the largest open subset of A .

4. Prove that the n -dimensional ball centered at a of radius r ,

$$dsB^n(a; r) = \{x \in \mathbb{R}^n : \|x - a\| < r\}$$

is open.

-
5. Find the interior, exterior, and boundary of the sets:

$$B^n = \{x \in^n: \|x\| \leq 1\},$$

$$S^{n-1} = \{x \in^n: \|x\| = 1\},$$

$$^n = \{x \in^n: \text{ each } x^i \text{ is rational}\}.$$

6. If $A \subset [0, 1]$ is the union of open intervals (a_i, b_i) such that each rational number in $(0, 1)$ is contained in some (a_i, b_i) , show that $\partial A = [0, 1] - A$.
7. If A is a closed set that contains every rational number $r \in [0, 1]$, show that $[0, 1] \subset A$.
8. Graph generic open balls in 2 with respect to each of the “non-Euclidean” norms, $\|\cdot\|_1$ and $\|\cdot\|_\infty$. What shapes are they?