

Notes on Mathematics

David K. Zhang

Last modified January 15, 2022

List of Definitions

| | |
|---|---|
| Cartesian Product, $A \times B$ | 1 |
| Relation, Binary Relation, $a R b$ | 1 |
| Function, Domain, Codomain, $f : A \rightarrow B, f(x)$ | 1 |
| Identity Function, id_A | 1 |
| Cartesian Product, $\prod_{i \in I} A_i$ | 1 |
| Metric, Distance Function | 3 |
| Metric Space | 3 |
| Continuity | 3 |
| Group | 4 |
| Abelian Group | 4 |
| Rng | 5 |
| Zero Rng, Trivial Rng | 6 |
| Nonzero Rng, Nontrivial Rng | 6 |
| Ring | 6 |
| Commutative Ring | 6 |

List of Theorems

| | |
|---|---|
| Multiplying by Zero Yields Zero | 5 |
|---|---|

Chapter 1

Set Theory

Definition: Cartesian Product, $A \times B$

Let A and B be sets. The **Cartesian product** of A and B , denoted by $A \times B$, is the set of all ordered pairs (a, b) consisting of an element $a \in A$ followed by an element $b \in B$.

$$A \times B := \{(a, b) : a \in A \text{ and } b \in B\}$$

Definition: Relation, Binary Relation, $a R b$

Let A and B be sets. A **relation** between A and B , also known as a **binary relation** between A and B , is a subset $R \subseteq A \times B$ of their Cartesian product.

Given $a \in A$ and $b \in B$, we write $a R b$ to denote that $(a, b) \in R$. If this is the case, then we say that R *relates* a to b .

If $A = B$, then instead of saying that R is a relation *between* A and A , we simply say that R is a binary relation *on* A .

For example, the less-than relation $<$ is a binary relation on the set \mathbb{N} of natural numbers. We write $3 < 4$ to indicate that the ordered pair $(3, 4) \in \mathbb{N} \times \mathbb{N}$ is an element of the relation $<$.

Definition: Function, Domain, Codomain, $f : A \rightarrow B, f(x)$

Let A and B be sets. A **function** from A to B is a binary relation $f \subseteq A \times B$ between A and B that has the following property: for each $x \in A$, there exists a unique $y \in B$ such that $x f y$. We call the set A the **domain** of the function f , and we call B the **codomain** of f .

We write $f : A \rightarrow B$ to denote that f is a function from A to B , and given $x \in A$, we write $f(x)$ to denote the unique element of B that is related to x by f .

Definition: Identity Function, id_A

Let A be a set. The **identity function** on A is the function $\text{id}_A : A \rightarrow A$ defined by $\text{id}_A(a) := a$ for all $a \in A$.

Definition: Cartesian Product, $\prod_{i \in I} A_i$

Let I be an index set, and let $\{A_i\}_{i \in I}$ be an indexed family of sets. The ***Cartesian product*** of $\{A_i\}_{i \in I}$, denoted by $\prod_{i \in I} A_i$, is the set of all functions $f : I \rightarrow \bigcup_{i \in I} A_i$ such that $f(i) \in A_i$ for all $i \in I$.

$$\prod_{i \in I} A_i := \left\{ f : I \rightarrow \bigcup_{i \in I} A_i \mid \forall i \in I : f(i) \in A_i \right\}$$

Chapter 2

Topology

Definition: Metric, Distance Function

Let X be a set. A **metric** on X , also known as a **distance function** on X , is a function $d : X \times X \rightarrow \mathbb{R}$ that satisfies the following requirements:

- **Positive-definiteness:** $d(x, y) \geq 0$ for all $x, y \in X$, and $d(x, y) = 0$ if and only if $x = y$.
- **Symmetry:** $d(x, y) = d(y, x)$ for all $x, y \in X$.
- **Triangle inequality:** $d(x, z) \leq d(x, y) + d(y, z)$ for all $x, y, z \in X$.

Definition: Metric Space

A **metric space** is an ordered pair (X, d) consisting of a set X and a metric d on X .

Definition: Continuity

Let (X, d_X) and (Y, d_Y) be metric spaces. A function $f : X \rightarrow Y$ is **continuous** at a point $x_0 \in X$ if, for every $\epsilon > 0$, there exists $\delta > 0$, such that for all $x \in X$, if $d_X(x, x_0) < \delta$, then $d_Y(f(x), f(x_0)) < \epsilon$.

Chapter 3

Group Theory

Definition: Group

A **group** is an algebraic structure $\langle G; 1, ^{-1}, \cdot \rangle$ consisting of:

- a set G , called the **underlying set**;
- a distinguished element $1 \in G$, called the **identity element**;
- a unary operation $^{-1} : G \rightarrow G$, written as $x \mapsto x^{-1}$, called **inversion**;
- a binary operation $\cdot : G \times G \rightarrow G$, written as $(x, y) \mapsto x \cdot y$, called the **group operation** or **group product**;

satisfying the following requirements:

- **Associative property**: $(x \cdot y) \cdot z = x \cdot (y \cdot z)$ for all $x, y, z \in G$.
- **Identity property**: $1 \cdot x = x \cdot 1 = x$ for all $x \in G$.
- **Inverse property**: $x \cdot x^{-1} = x^{-1} \cdot x = 1$ for all $x \in G$.

Definition: Abelian Group

An **abelian group** is a group $\langle G; 1, ^{-1}, \cdot \rangle$ that satisfies the following additional requirement:

- **Commutative property**: $x \cdot y = y \cdot x$ for all $x, y \in G$.

Chapter 4

Ring Theory

In this chapter, we introduce a new class of algebraic structures, called rngs and rings, whose study is collectively called **ring theory**. Rngs and rings are more complicated than groups because their definition involves not one, but two binary operations.

Definition: Rng

A **rng** (pronounced as “*rung*”) is an algebraic structure $\langle R; 0, -, +, \cdot \rangle$ consisting of:

- a set R , called the **underlying set**;
- a distinguished element $0 \in R$, called the **zero element**;
- a unary operation $- : R \rightarrow R$, written as $x \mapsto -x$, called **negation**;
- a binary operation $+ : R \times R \rightarrow R$, written as $(x, y) \mapsto x + y$, called **addition**;
- a binary operation $\cdot : R \times R \rightarrow R$, written as $(x, y) \mapsto x \cdot y$, called **multiplication**;

satisfying the following requirements:

- **Additive structure**: $\langle R; 0, -, + \rangle$ is an abelian group.
- **Associativity**: $(x \cdot y) \cdot z = x \cdot (y \cdot z)$ for all $x, y, z \in R$.
- **Left distributivity**: $x \cdot (y + z) = (x \cdot y) + (x \cdot z)$ for all $x, y, z \in R$.
- **Right distributivity**: $(x + y) \cdot z = (x \cdot z) + (y \cdot z)$ for all $x, y, z \in R$.

The key ingredient in the definition of a rng is **distributivity**, which establishes a link between two different binary operations. We begin our study of rngs by proving a simple (but important) result to demonstrate the utility of the distributive property.

Multiplying by Zero Yields Zero

Theorem: Let $\langle R; 0, -, +, \cdot \rangle$ be a rng. For any $x \in R$, we have $0 \cdot x = x \cdot 0 = 0$.

Proof: Let $x \in R$ be given. Because 0 is the identity element of the abelian group $\langle R; 0, -, + \rangle$, we have $0 = 0 + 0$. Using left distributivity, it follows that $0 \cdot x = (0 + 0) \cdot x = (0 \cdot x) + (0 \cdot x)$, and by canceling one copy of $0 \cdot x$ on both sides, we conclude that $0 \cdot x = 0$. We similarly apply

right distributivity to the expression $x \cdot 0 = x \cdot (0 + 0) = (x \cdot 0) + (x \cdot 0)$ to conclude that $x \cdot 0 = 0$.

Definition: Zero Rng, Trivial Rng

The *zero rng* or *trivial rng* is the rng $\langle \{0\}; 0, -, +, \cdot \rangle$ whose underlying set is a singleton containing the distinguished element 0, and whose operations are defined by $-0 = 0 + 0 = 0 \cdot 0 = 0$.

Definition: Nonzero Rng, Nontrivial Rng

A rng is *nonzero* or *nontrivial* if its underlying set contains more than one element.

Definition: Ring

A *ring* is an algebraic structure $\langle R; 0, 1, -, +, \cdot \rangle$ consisting of:

- a set R , called the *underlying set*;
- a distinguished element $0 \in R$, called the *zero element*;
- a distinguished element $1 \in R$, called the *identity element*;
- a unary operation $- : R \rightarrow R$, written as $x \mapsto -x$, called *negation*;
- a binary operation $+ : R \times R \rightarrow R$, written as $(x, y) \mapsto x + y$, called *addition*;
- a binary operation $\cdot : R \times R \rightarrow R$, written as $(x, y) \mapsto x \cdot y$, called *multiplication*;

satisfying the following requirements:

- *Rng structure*: $\langle R; 0, -, +, \cdot \rangle$ is a rng.
- *Identity*: $1 \cdot x = x \cdot 1 = x$ for all $x \in R$.

Definition: Commutative Ring

A *commutative ring* is a ring R that satisfies the following additional requirement:

- *Commutativity*: $x \cdot y = y \cdot x$ for all $x, y \in R$.