

### Problem 1

E) The equation for the line through  $(x_0, y_0)$  with slope  $a$  is:  $y(x) = y_0 + a(x - x_0)$ . The intersection  $x_s$  of the line with the  $x$ -axis is  $x_s = x_0 - y_0/a$ , because  $y(x_s) = 0$ .

For the Newton-method we have  $y_0 = f(x_0)$  and  $a = f'(x_0)$ , so  $x_1 = x_s = x_0 - f(x_0)/f'(x_0)$ . Repeating this gives the iteration method

$$x_{k+1} = x_k - \frac{f(x_k)}{f'(x_k)}.$$


The secant method needs two initial values,  $x_0$  and  $x_1$ . The next value is the intersection  $x_2$  between the line through  $(x_0, f(x_0))$  and  $(x_1, f(x_1))$  (secant) with the  $x$ -axis. The secant is the line through  $(x_1, f(x_1))$  with the slope  $\frac{f(x_1)-f(x_0)}{x_1-x_0}$  so we have

$$x_2 = x_1 - \frac{f(x_1)}{\frac{f(x_1)-f(x_0)}{x_1-x_0}} = \frac{x_0 f(x_1) - x_1 f(x_0)}{f(x_1) - f(x_0)}$$

The rest is explained in the lecture notes.

### Problem 2

E) The equation  $x = -2\ln(x)$  has exactly one solution  $x_*$ , which is in the interval  $]0, 1[$ , see the Figure.



The derivative of  $\phi(x) = -2\ln(x)$  is  $\phi'(x) = -2/x$ . It is  $|\phi'(x)| > 1$  for  $x \in ]0, 1[$ . In particular we have  $|\phi'(x_*)| > 1$ . So  $x_*$  is an repelling fix point and cannot be found by the iteration method  $x_{k+1} = \phi(x_k)$ . We have

$$x = -2\ln(x) \quad \Leftrightarrow \quad -x/2 = \ln(x) \quad \Leftrightarrow \quad \underbrace{e^{-x/2}}_{=\psi(x)} = x$$

We have  $|\psi'(x)| < 1$  for  $x \in ]0, 1[$ . So  $x_*$  is a attracting fix point of  $\psi$ . The iteration method  $x_{k+1} = \psi(x_k)$  converges to  $x_* \approx 0.70347$  if we start in the interval  $(0, 1)$ .

**Problem 3**

E) The Newton-method for  $f(x) = 0$  with  $f(x) = x^2 - a$  is

$$x_{k+1} = x_k - f(x_k)/f'(x_k) = x_k - (x_k^2 - a)/(2x_k) = \underbrace{\frac{1}{2}(x_k + a/x_k)}_{=\phi(x_k)}.$$

The derivative of the function  $\phi$  at the fix point  $x_* = \sqrt{a}$  is

$$\phi'(\sqrt{a}) = \frac{1}{2}(1 - a/x^2)|_{x=\sqrt{a}} = 0$$

We conclude that the iteration has order of convergence 2, see lecture slides.

**Problem 4**

E)

$$\begin{aligned} \begin{bmatrix} x_{k+1} \\ y_{k+1} \end{bmatrix} &= \begin{bmatrix} x_k \\ y_k \end{bmatrix} - \begin{bmatrix} y_k + 1 & x_k - 1 \\ y_k^2 & 2x_k y_k \end{bmatrix}^{-1} \begin{bmatrix} x_k y_k + x_k - y_k + 1 \\ x_k y_k^2 + 5 \end{bmatrix} \\ &= \begin{bmatrix} x_k \\ y_k \end{bmatrix} - \frac{\begin{bmatrix} 2x_k y_k & -x_k + 1 \\ -y_k^2 & y_k + 1 \end{bmatrix}}{2(y_k + 1)x_k y_k - (x_k - 1)y_k^2} \begin{bmatrix} x_k y_k + x_k - y_k - 1 \\ x_k y_k^2 + 5 \end{bmatrix}. \\ \begin{bmatrix} x_0 \\ y_0 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix} &\Rightarrow \begin{bmatrix} x_1 \\ y_1 \end{bmatrix} = \begin{bmatrix} 1 \\ -2 \end{bmatrix}. \end{aligned}$$