

# Math Camp

Justin Grimmer

Professor  
Department of Political Science  
Stanford University

September 6th, 2018

# Optimization

Social scientists are often concerned with finding **extrema**: **maxima** or **minima**

# Optimization

Social scientists are often concerned with finding **extrema**: **maxima** or **minima**

- Given data, **most** likely value of a parameter

# Optimization

Social scientists are often concerned with finding **extrema**: **maxima** or **minima**

- Given data, **most** likely value of a parameter
- Game theory: given other player's strategy, action that **maximizes** utility

# Optimization

Social scientists are often concerned with finding **extrema**: **maxima** or **minima**

- Given data, **most** likely value of a parameter
- Game theory: given other player's strategy, action that **maximizes** utility
- Across substantive areas: what is the **optimal** action, strategy, prediction?

# Optimization

Social scientists are often concerned with finding **extrema**: **maxima** or **minima**

- Given data, **most** likely value of a parameter
- Game theory: given other player's strategy, action that **maximizes** utility
- Across substantive areas: what is the **optimal** action, strategy, prediction?

## How to Optimize

# Optimization

Social scientists are often concerned with finding **extrema**: **maxima** or **minima**

- Given data, **most** likely value of a parameter
- Game theory: given other player's strategy, action that **maximizes** utility
- Across substantive areas: what is the **optimal** action, strategy, prediction?

## How to Optimize

- When functions are **well behaved** and **known**  $\leadsto$  analytic solutions

# Optimization

Social scientists are often concerned with finding **extrema**: **maxima** or **minima**

- Given data, **most** likely value of a parameter
- Game theory: given other player's strategy, action that **maximizes** utility
- Across substantive areas: what is the **optimal** action, strategy, prediction?

## How to Optimize

- When functions are **well behaved** and **known**  $\leadsto$  analytic solutions
  - Differentiate, set equal to zero, solve



# Optimization

Social scientists are often concerned with finding **extrema**: **maxima** or **minima**

- Given data, **most** likely value of a parameter
- Game theory: given other player's strategy, action that **maximizes** utility
- Across substantive areas: what is the **optimal** action, strategy, prediction?

## How to Optimize

- When functions are **well behaved** and **known**  $\leadsto$  analytic solutions
  - Differentiate, set equal to zero, solve
  - Check end points and use second derivative test

# Optimization

Social scientists are often concerned with finding **extrema**: **maxima** or **minima**

- Given data, **most** likely value of a parameter
- Game theory: given other player's strategy, action that **maximizes** utility
- Across substantive areas: what is the **optimal** action, strategy, prediction?

## How to Optimize

- When functions are **well behaved** and **known**  $\rightsquigarrow$  analytic solutions
  - Differentiate, set equal to zero, solve
  - Check end points and use second derivative test
- More difficult problems  $\rightsquigarrow$  computational solutions

# Optimization


Social scientists are often concerned with finding **extrema**: **maxima** or **minima**

- Given data, **most** likely value of a parameter
- Game theory: given other player's strategy, action that **maximizes** utility
- Across substantive areas: what is the **optimal** action, strategy, prediction?


## How to Optimize

- When functions are **well behaved** and **known**  $\rightsquigarrow$  analytic solutions
  - Differentiate, set equal to zero, solve
  - Check end points and use second derivative test
- More difficult problems  $\rightsquigarrow$  computational solutions

# Intuition: Optimization with Derivatives, **Known** well behaved functions




# Intuition: Optimization with Derivatives, **Known** well behaved functions




- Rolle's theorem guarantee's that, at some point,  $f'(x) = 0$

# Intuition: Optimization with Derivatives, **Known** well behaved functions




- Rolle's theorem guarantee's that, at some point,  $f'(x) = 0$
- **Intuition from proof**—what happens as we approach from the left?

# Intuition: Optimization with Derivatives, **Known** well behaved functions



- Rolle's theorem guarantee's that, at some point,  $f'(x) = 0$
- **Intuition from proof**—what happens as we approach from the left?
- **Intuition from proof**—what happens as we approach from the right?

# Intuition: Optimization with Derivatives, **Known** well behaved functions



- Rolle's theorem guarantee's that, at some point,  $f'(x) = 0$
- **Intuition from proof**—what happens as we approach from the left?
- **Intuition from proof**—what happens as we approach from the right?
- **critical intuition** first, second derivatives



# Second Derivatives

## Definition

*Suppose  $f : \mathbb{R} \rightarrow \mathbb{R}$  is differentiable. Recall we write this as  $f'$  and suppose that  $f' : \mathbb{R} \rightarrow \mathbb{R}$ . Then if the limit,*

$$\lim_{x \rightarrow x_0} R(x) = \frac{f'(x) - f'(x_0)}{x - x_0}$$

*exists, we call this the **second derivative** at  $x_0$ ,  $f''(x_0)$ .*

# Example of Second Derivatives

$$f(x) = x$$

$$f'(x) = 1$$

$$f''(x) = 0$$

# Example of Second Derivatives

$$\begin{aligned}f(x) &= e^x \\f'(x) &= e^x \\f''(x) &= e^x\end{aligned}$$

# Example of Second Derivatives

$$f(x) = \log(x)$$

$$f'(x) = \frac{1}{x}$$

$$f''(x) = \frac{-1}{x^2}$$

# Example of Second Derivatives

$$f(x) = \frac{1}{x}$$

$$f'(x) = \frac{-1}{x^2}$$

$$f''(x) = \frac{2}{x^3}$$

# Example of Second Derivatives

$$\begin{aligned}f(x) &= -x^2 + 20 \\f'(x) &= -2x \\f''(x) &= -2\end{aligned}$$

# Approximating functions and second order conditions

## Theorem

**Taylor's Theorem** Suppose  $f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $f(x)$  is infinitely differentiable function. Then, the taylor expansion of  $f(x)$  around  $a$  is given by

$$f(x) = f(a) + \frac{f'(a)}{1!}(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f'''(a)}{3!}(x-a)^3 + \dots$$



$$f(x) = \sum_{n=0}^{\infty} \frac{f^n(a)}{n!}(x-a)^n$$

R Code!





# Concavity, Convexity, Inflections

Second derivatives provide further information about functions



# Concavity, Convexity, Inflections

Second derivatives provide further information about functions



# Concave Up/ Convex

## Definition


Suppose  $f : [a, b] \rightarrow \mathbb{R}$  is a **twice** differentiable function. If, for all  $x \in [a, b]$  and  $y \in [a, b]$  and  $t \in (0, 1)$

$$f((1 - t)x + ty) < (1 - t)f(x) + tf(y)$$

We say that  $f$  is strictly **concave up** or **convex**. Equivalently if  $f''(x) > 0$  for all  $x \in [a, b]$ , we say that  $f$  is strictly **concave up**.


# Concave Up, Graphical Test

$$f(x) = e^x, [1, 4]$$




# Concave Up, Graphical Test

$$f(x) = e^x, [1, 4]$$




# Concave Up, Graphical Test

$$f(x) = e^x, [1, 4]$$




# Concave Up, Graphical Test

$$f(x) = e^x, [1, 4]$$



# Concave Up, Graphical Test


$$f(x) = e^x, [1, 4]$$






# Concave Up, Graphical Test

$$f(x) = e^x, [1, 4]$$




# Concave Up, Graphical Test


$$f(x) = e^x, [1, 4]$$



# Concave Up, Second Derivative




# Concave Up, Second Derivative



$$f(x) = e^x$$


# Concave Up, Second Derivative



$$f(x) = e^x$$

$$f'(x) = e^x$$

# Concave Up, Second Derivative




$$f(x) = e^x$$

$$f'(x) = e^x$$

$$f''(x) = e^x$$

# Concave Up, Second Derivative



$e^x > 0$  for all  $x \in [1, 4]$

# Concave Down

## Definition


Suppose  $f : [a, b] \rightarrow \mathbb{R}$  is a **twice** differentiable function. If, for all  $x \in [a, b]$  and  $y \in [a, b]$  and  $t \in (0, 1)$

$$f((1 - t)x + ty) > (1 - t)f(x) + tf(y)$$

We say that  $f$  is strictly **concave down**. Equivalently if  $f''(x) < 0$  for all  $x \in [a, b]$ , we say that  $f$  is strictly **concave down**.



## Concave Down



- Show Concave down with graph test for  $x \in [1, 4]$
- Show concave down with second derivative test for  $x \in [1, 4]$

# Optimization


## Theorem

**Extreme Value Theorem** Suppose  $f : [a, b] \rightarrow \mathbb{R}$  and that  $f$  is continuous. Then  $f$  obtains its extreme value on  $[a, b]$ .

## Corollary

Suppose  $f : [a, b] \rightarrow \mathbb{R}$ , that  $f$  is continuous and differentiable, and that  $f(a)$  nor  $f(b)$  is the extreme value. Then  $f$  obtains its maximum on  $(a, b)$  and if  $f(x_0)$  is the extreme value of  $f$   $x_0 \in (a, b)$  then,  $f'(x_0) = 0$ .


# Extrema on End Points



# Maximum in Middle, Concave Down


$$f(x) = -x^2 + 5.$$

**Rolle's Theorem**




# Minimum in Interior, Concave Up

$$f(x) = x^2 + 9x + 9$$




# Local Optima

$$f(x) = \sin(x)$$



# Inflection points

$$f(x) = x^3$$



# Framework for Optimization

Recipe for optimization



# Framework for Optimization

Recipe for optimization

- Find  $f'(x)$ .

# Framework for Optimization

Recipe for optimization

- Find  $f'(x)$ .
- Set  $f'(x) = 0$  and solve for  $x$ . Call all  $x_0$  such that  $f'(x_0) = 0$  **critical values**.

# Framework for Optimization

## Recipe for optimization

- Find  $f'(x)$ .
- Set  $f'(x) = 0$  and solve for  $x$ . Call all  $x_0$  such that  $f'(x_0) = 0$  **critical values**.
- Find  $f''(x)$ . Evaluate at each  $x_0$ .

# Framework for Optimization

## Recipe for optimization

- Find  $f'(x)$ .
- Set  $f'(x) = 0$  and solve for  $x$ . Call all  $x_0$  such that  $f'(x_0) = 0$  **critical values**.
- Find  $f''(x)$ . Evaluate at each  $x_0$ .
  - If  $f''(x) > 0$ , Concave up, **local** minimum

# Framework for Optimization

## Recipe for optimization

- Find  $f'(x)$ .
- Set  $f'(x) = 0$  and solve for  $x$ . Call all  $x_0$  such that  $f'(x_0) = 0$  **critical values**.
- Find  $f''(x)$ . Evaluate at each  $x_0$ .
  - If  $f''(x) > 0$ , Concave up, **local** minimum
  - If  $f''(x) < 0$ , Concave down, **local** maximum

# Framework for Optimization

## Recipe for optimization


- Find  $f'(x)$ .
- Set  $f'(x) = 0$  and solve for  $x$ . Call all  $x_0$  such that  $f'(x_0) = 0$  **critical values**.
- Find  $f''(x)$ . Evaluate at each  $x_0$ .
  - If  $f''(x) > 0$ , Concave up, **local** minimum
  - If  $f''(x) < 0$ , Concave down, **local** maximum
  - If  $f''(x) = 0$ , No knowledge—local minimum, maximum, or inflection point

# Framework for Optimization

## Recipe for optimization

- Find  $f'(x)$ .
- Set  $f'(x) = 0$  and solve for  $x$ . Call all  $x_0$  such that  $f'(x_0) = 0$  **critical values**.
- Find  $f''(x)$ . Evaluate at each  $x_0$ .
  - If  $f''(x) > 0$ , Concave up, **local** minimum
  - If  $f''(x) < 0$ , Concave down, **local** maximum
  - If  $f''(x) = 0$ , No knowledge—local minimum, maximum, or inflection point
- **Check End Points!**

Example 1:  $f(x) = -x^2$ ,  $x \in [-3, 3]$





Example 1:  $f(x) = -x^2$ ,  $x \in [-3, 3]$

Example 1:  $f(x) = -x^2$ ,  $x \in [-3, 3]$

1) Critical Value:

$$\begin{aligned}f'(x) &= -2x \\0 &= -2x^* \\x^* &= 0\end{aligned}$$

Example 1:  $f(x) = -x^2$ ,  $x \in [-3, 3]$

1) Critical Value:

$$\begin{aligned}f'(x) &= -2x \\ 0 &= -2x^* \\ x^* &= 0\end{aligned}$$

2) Second Derivative:

$$\begin{aligned}f'(x) &= -2x \\ f''(x) &= -2\end{aligned}$$

$f''(x) < 0$ , local maximum

Example 1:  $f(x) = -x^2$ ,  $x \in [-3, 3]$

1) Critical Value:

$$\begin{aligned}f'(x) &= -2x \\ 0 &= -2x^* \\ x^* &= 0\end{aligned}$$

2) Second Derivative:

$$\begin{aligned}f'(x) &= -2x \\ f''(x) &= -2\end{aligned}$$

$f''(x) < 0$ , local maximum

Example 1:  $f(x) = -x^2$ ,  $x \in [-3, 3]$


3) Check end points

$$f(0) = -0^2 = 0$$

$$f(-3) = -(-3)^2 = -9$$

$$f(3) = -(3)^2 = -9$$

Example 2:  $f(x) = x^3$ ,  $x \in [-3, 3]$



Example 2:  $f(x) = x^3$ ,  $x \in [-3, 3]$

1) Critical Values:

$$f'(x) = 3x^2$$

$$0 = 3(x^*)^2$$

$$x^* = 0$$

Example 2:  $f(x) = x^3$ ,  $x \in [-3, 3]$

1) Critical Values:

$$\begin{aligned}f'(x) &= 3x^2 \\ 0 &= 3(x^*)^2 \\ x^* &= 0\end{aligned}$$

2) Second Derivative:

$$\begin{aligned}f''(x) &= 6x \\ f''(0) &= 0\end{aligned}$$

No information



Example 2:  $f(x) = x^3$ ,  $x \in [-3, 3]$

3) Check End Points:

$$f(0) = 0^3 = 0$$

$$f(-3) = -3^3 = -27$$

$$f(3) = 3^3 = 27$$

Neither maximum nor minimum, **saddle point**

## Example 3: Spatial Model

A large literature in Congress supposes legislators and policies can be situated in **policy space**

## Example 3: Spatial Model

A large literature in Congress supposes legislators and policies can be situated in **policy space**

Suppose legislator  $i$  and policies  $x, i \in \mathcal{R}$ .

### Example 3: Spatial Model

A large literature in Congress supposes legislators and policies can be situated in **policy space**

Suppose legislator  $i$  and policies  $x, i \in \mathcal{R}$ .

Define legislator  $i$ 's utility as,  $U : \mathcal{R} \rightarrow \mathcal{R}$ ,

### Example 3: Spatial Model

A large literature in Congress supposes legislators and policies can be situated in **policy space**

Suppose legislator  $i$  and policies  $x, i \in \mathfrak{R}$ .

Define legislator  $i$ 's utility as,  $U : \mathfrak{R} \rightarrow \mathfrak{R}$ ,

$$U_i(x) = -(x - \mu)^2$$

### Example 3: Spatial Model

A large literature in Congress supposes legislators and policies can be situated in **policy space**

Suppose legislator  $i$  and policies  $x, i \in \mathfrak{R}$ .

Define legislator  $i$ 's utility as,  $U : \mathfrak{R} \rightarrow \mathfrak{R}$ ,

$$U_i(x) = -(x - \mu)^2$$

$$U_i(x) = -x^2 + 2x\mu - \mu^2$$

### Example 3: Spatial Model

A large literature in Congress supposes legislators and policies can be situated in **policy space**

Suppose legislator  $i$  and policies  $x, i \in \mathfrak{R}$ .

Define legislator  $i$ 's utility as,  $U : \mathfrak{R} \rightarrow \mathfrak{R}$ ,

$$U_i(x) = -(x - \mu)^2$$

$$U_i(x) = -x^2 + 2x\mu - \mu^2$$

What is  $i$ 's optimal policy over the range  $x \in [\mu - 2, \mu + 2]$ ?

### Example 3: Spatial Model

A large literature in Congress supposes legislators and policies can be situated in **policy space**

Suppose legislator  $i$  and policies  $x, i \in \mathfrak{R}$ .

Define legislator  $i$ 's utility as,  $U : \mathfrak{R} \rightarrow \mathfrak{R}$ ,

$$U_i(x) = -(x - \mu)^2$$

$$U_i(x) = -x^2 + 2x\mu - \mu^2$$

What is  $i$ 's optimal policy over the range  $x \in [\mu - 2, \mu + 2]$ ?

$$U'_i(x) = -2(x - \mu)$$



### Example 3: Spatial Model

A large literature in Congress supposes legislators and policies can be situated in **policy space**

Suppose legislator  $i$  and policies  $x, i \in \mathfrak{R}$ .

Define legislator  $i$ 's utility as,  $U : \mathfrak{R} \rightarrow \mathfrak{R}$ ,

$$U_i(x) = -(x - \mu)^2$$

$$U_i(x) = -x^2 + 2x\mu - \mu^2$$

What is  $i$ 's optimal policy over the range  $x \in [\mu - 2, \mu + 2]$ ?

$$U'_i(x) = -2(x - \mu)$$

$$0 = -2x^* + 2\mu$$

### Example 3: Spatial Model

A large literature in Congress supposes legislators and policies can be situated in **policy space**

Suppose legislator  $i$  and policies  $x, i \in \mathfrak{R}$ .

Define legislator  $i$ 's utility as,  $U : \mathfrak{R} \rightarrow \mathfrak{R}$ ,

$$U_i(x) = -(x - \mu)^2$$

$$U_i(x) = -x^2 + 2x\mu - \mu^2$$

What is  $i$ 's optimal policy over the range  $x \in [\mu - 2, \mu + 2]$ ?

$$U'_i(x) = -2(x - \mu)$$

$$0 = -2x^* + 2\mu$$

$$x^* = \mu$$

### Example 3: Spatial Model

A large literature in Congress supposes legislators and policies can be situated in **policy space**

Suppose legislator  $i$  and policies  $x, i \in \mathfrak{R}$ .

Define legislator  $i$ 's utility as,  $U : \mathfrak{R} \rightarrow \mathfrak{R}$ ,

$$U_i(x) = -(x - \mu)^2$$

$$U_i(x) = -x^2 + 2x\mu - \mu^2$$

What is  $i$ 's optimal policy over the range  $x \in [\mu - 2, \mu + 2]$ ?

$$U'_i(x) = -2(x - \mu)$$

$$0 = -2x^* + 2\mu$$

$$x^* = \mu$$

### Second Derivative Test

### Example 3: Spatial Model

A large literature in Congress supposes legislators and policies can be situated in **policy space**

Suppose legislator  $i$  and policies  $x, i \in \mathfrak{R}$ .

Define legislator  $i$ 's utility as,  $U : \mathfrak{R} \rightarrow \mathfrak{R}$ ,

$$U_i(x) = -(x - \mu)^2$$

$$U_i(x) = -x^2 + 2x\mu - \mu^2$$

What is  $i$ 's optimal policy over the range  $x \in [\mu - 2, \mu + 2]$ ?

$$U'_i(x) = -2(x - \mu)$$

$$0 = -2x^* + 2\mu$$

$$x^* = \mu$$

### Second Derivative Test

$$U''_i(x) = -2 < 0 \rightarrow \text{Concave Down}$$

### Example 3: Spatial Model

A large literature in Congress supposes legislators and policies can be situated in **policy space**

Suppose legislator  $i$  and policies  $x, i \in \mathfrak{R}$ .

Define legislator  $i$ 's utility as,  $U : \mathfrak{R} \rightarrow \mathfrak{R}$ ,

$$U_i(x) = -(x - \mu)^2$$

$$U_i(x) = -x^2 + 2x\mu - \mu^2$$

What is  $i$ 's optimal policy over the range  $x \in [\mu - 2, \mu + 2]$ ?

$$U'_i(x) = -2(x - \mu)$$

$$0 = -2x^* + 2\mu$$

$$x^* = \mu$$

### Second Derivative Test

$$U''_i(x) = -2 < 0 \rightarrow \text{Concave Down}$$

We call  $\mu$  legislator  $i$ 's **ideal point**

## Example 3: Spatial Model

$$\begin{aligned}U_i(\mu) &= -(\mu - \mu)^2 = 0 \\U_i(\mu - 2) &= -(\mu - 2 - \mu)^2 = -4 \\U_i(\mu + 2) &= -(\mu + 2 - \mu)^2 = -4\end{aligned}$$

Maximize utility at  $\mu$

## Example 4: Maximum Likelihood Estimation

In statistics classes we'll learn about **parameters** from **data**.

## Example 4: Maximum Likelihood Estimation

In statistics classes we'll learn about **parameters** from **data**.

Here is an example **likelihood** function: We want to find the **Maximum likelihood estimate**



## Example 4: Maximum Likelihood Estimation

In statistics classes we'll learn about **parameters** from **data**.

Here is an example **likelihood** function: We want to find the **Maximum likelihood estimate**

$$f(\mu) = \prod_{i=1}^N \exp\left(\frac{-(Y_i - \mu)^2}{2}\right)$$

## Example 4: Maximum Likelihood Estimation

In statistics classes we'll learn about **parameters** from **data**.

Here is an example **likelihood** function: We want to find the **Maximum likelihood estimate**

$$\begin{aligned} f(\mu) &= \prod_{i=1}^N \exp\left(-\frac{(Y_i - \mu)^2}{2}\right) \\ &= \exp\left(-\frac{(Y_1 - \mu)^2}{2}\right) \times \dots \times \exp\left(-\frac{(Y_N - \mu)^2}{2}\right) \end{aligned}$$

## Example 4: Maximum Likelihood Estimation

In statistics classes we'll learn about **parameters** from **data**.

Here is an example **likelihood** function: We want to find the **Maximum likelihood estimate**

$$\begin{aligned} f(\mu) &= \prod_{i=1}^N \exp\left(-\frac{(Y_i - \mu)^2}{2}\right) \\ &= \exp\left(-\frac{(Y_1 - \mu)^2}{2}\right) \times \dots \times \exp\left(-\frac{(Y_N - \mu)^2}{2}\right) \\ &= \exp\left(-\frac{\sum_{i=1}^N (Y_i - \mu)^2}{2}\right) \end{aligned}$$

## Example 4: Maximum Likelihood Estimation

In statistics classes we'll learn about **parameters** from **data**.

Here is an example **likelihood** function: We want to find the **Maximum likelihood estimate**

$$\begin{aligned}f(\mu) &= \prod_{i=1}^N \exp\left(-\frac{(Y_i - \mu)^2}{2}\right) \\&= \exp\left(-\frac{(Y_1 - \mu)^2}{2}\right) \times \dots \times \exp\left(-\frac{(Y_N - \mu)^2}{2}\right) \\&= \exp\left(-\frac{\sum_{i=1}^N (Y_i - \mu)^2}{2}\right)\end{aligned}$$

### Theorem

*Suppose  $f : \mathbb{R} \rightarrow (0, \infty)$ . If  $x_0$  maximizes  $f$ , then  $x_0$  maximizes  $\log(f(x))$ .*

## Example 4: Maximum Likelihood Estimation

## Example 4: Maximum Likelihood Estimation

$$\log f(\mu) = \log \left( \exp \left( -\frac{\sum_{i=1}^N (Y_i - \mu)^2}{2} \right) \right)$$

## Example 4: Maximum Likelihood Estimation

$$\begin{aligned}\log f(\mu) &= \log \left( \exp \left( -\frac{\sum_{i=1}^N (Y_i - \mu)^2}{2} \right) \right) \\ &= -\frac{\sum_{i=1}^N (Y_i - \mu)^2}{2}\end{aligned}$$

## Example 4: Maximum Likelihood Estimation

$$\begin{aligned}\log f(\mu) &= \log \left( \exp \left( -\frac{\sum_{i=1}^N (Y_i - \mu)^2}{2} \right) \right) \\ &= -\frac{\sum_{i=1}^N (Y_i - \mu)^2}{2} \\ &= -\frac{1}{2} \left( \sum_{i=1}^N Y_i^2 - 2\mu \sum_{i=1}^N Y_i + N \times \mu^2 \right)\end{aligned}$$



## Example 4: Maximum Likelihood Estimation

$$\begin{aligned}\log f(\mu) &= \log \left( \exp \left( -\frac{\sum_{i=1}^N (Y_i - \mu)^2}{2} \right) \right) \\ &= -\frac{\sum_{i=1}^N (Y_i - \mu)^2}{2} \\ &= -\frac{1}{2} \left( \sum_{i=1}^N Y_i^2 - 2\mu \sum_{i=1}^N Y_i + N \times \mu^2 \right) \\ \frac{\partial \log f(\mu)}{\partial \mu} &= -\frac{1}{2} \left( -2 \sum_{i=1}^N Y_i + 2N\mu \right)\end{aligned}$$

## Example 4: Maximum Likelihood Estimation

## Example 4: Maximum Likelihood Estimation

$$0 = -\frac{1}{2} \left( -\sum_{i=1}^N Y_i + 2N\mu^* \right)$$

## Example 4: Maximum Likelihood Estimation

$$0 = -\frac{1}{2} \left( -\sum_{i=1}^N Y_i + 2N\mu^* \right)$$

$$2 \sum_{i=1}^N Y_i = 2N\mu^*$$

## Example 4: Maximum Likelihood Estimation

$$0 = -\frac{1}{2} \left( -\sum_{i=1}^N Y_i + 2N\mu^* \right)$$

$$2 \sum_{i=1}^N Y_i = 2N\mu^*$$

$$\frac{\sum_{i=1}^N Y_i}{N} = \mu^*$$

## Example 4: Maximum Likelihood Estimation

$$0 = -\frac{1}{2} \left( -\sum_{i=1}^N Y_i + 2N\mu^* \right)$$

$$2 \sum_{i=1}^N Y_i = 2N\mu^*$$

$$\frac{\sum_{i=1}^N Y_i}{N} = \mu^*$$

$$\bar{Y} = \mu^*$$

## Example 4: Maximum Likelihood Estimation

$$0 = -\frac{1}{2} \left( -\sum_{i=1}^N Y_i + 2N\mu^* \right)$$

$$2 \sum_{i=1}^N Y_i = 2N\mu^*$$

$$\frac{\sum_{i=1}^N Y_i}{N} = \mu^*$$

$$\bar{Y} = \mu^*$$

## Second Derivative Test

## Example 4: Maximum Likelihood Estimation

$$0 = -\frac{1}{2} \left( -\sum_{i=1}^N Y_i + 2N\mu^* \right)$$

$$2 \sum_{i=1}^N Y_i = 2N\mu^*$$

$$\frac{\sum_{i=1}^N Y_i}{N} = \mu^*$$

$$\bar{Y} = \mu^*$$

### Second Derivative Test

$$f'(\mu) = -\frac{1}{2} \left( -2 \sum_{i=1}^N Y_i + 2N\mu \right)$$



## Example 4: Maximum Likelihood Estimation

$$0 = -\frac{1}{2} \left( -\sum_{i=1}^N Y_i + 2N\mu^* \right)$$

$$2 \sum_{i=1}^N Y_i = 2N\mu^*$$

$$\frac{\sum_{i=1}^N Y_i}{N} = \mu^*$$

$$\bar{Y} = \mu^*$$

### Second Derivative Test

$$f'(\mu) = -\frac{1}{2} \left( -2 \sum_{i=1}^N Y_i + 2N\mu \right)$$

$$f''(\mu) = -N$$

## Example 5: IR Bargaining (from Jim Fearon, Part 1)

Countries fight wars, usually to get stuff.

## Example 5: IR Bargaining (from Jim Fearon, Part 1)

Countries fight wars, usually to get stuff.

- Suppose two countries 1, 2 are fighting for something they value at  $v$ .

## Example 5: IR Bargaining (from Jim Fearon, Part 1)

Countries fight wars, usually to get stuff.

- Suppose two countries 1, 2 are fighting for something they value at  $v$ .
- Each country decides to invest  $a_1 \in [0, 1]$  and  $a_2 \in [0, 1]$ .

## Example 5: IR Bargaining (from Jim Fearon, Part 1)

Countries fight wars, usually to get stuff.

- Suppose two countries 1, 2 are fighting for something they value at  $v$ .
- Each country decides to invest  $a_1 \in [0, 1]$  and  $a_2 \in [0, 1]$ .
- The probability of country 1 winning the war is

## Example 5: IR Bargaining (from Jim Fearon, Part 1)

Countries fight wars, usually to get stuff.

- Suppose two countries 1, 2 are fighting for something they value at  $v$ .
- Each country decides to invest  $a_1 \in [0, 1]$  and  $a_2 \in [0, 1]$ .
- The probability of country 1 winning the war is

$$p(a_1, a_2) = \frac{a_1^n}{a_1^n + a_2^n}$$

## Example 5: IR Bargaining (from Jim Fearon, Part 1)

Countries fight wars, usually to get stuff.

- Suppose two countries 1, 2 are fighting for something they value at  $v$ .
- Each country decides to invest  $a_1 \in [0, 1]$  and  $a_2 \in [0, 1]$ .
- The probability of country 1 winning the war is

$$p(a_1, a_2) = \frac{a_1^n}{a_1^n + a_2^n}$$

- Country 1's utility is given by

## Example 5: IR Bargaining (from Jim Fearon, Part 1)

Countries fight wars, usually to get stuff.

- Suppose two countries 1, 2 are fighting for something they value at  $v$ .
- Each country decides to invest  $a_1 \in [0, 1]$  and  $a_2 \in [0, 1]$ .
- The probability of country 1 winning the war is

$$p(a_1, a_2) = \frac{a_1^n}{a_1^n + a_2^n}$$

- Country 1's utility is given by

$$\begin{aligned} U_1(a_1) &= \underbrace{1 - a_1}_{\text{cost}} + \underbrace{p(a_1, a_2)v}_{\text{Expected Benefit}} \\ &= 1 - a_1 + \frac{a_1^n}{a_1^n + a_2^n} v \end{aligned}$$



## Example 5: IR Bargaining (from Jim Fearon, Part 1)

Countries fight wars, usually to get stuff.

- Suppose two countries 1, 2 are fighting for something they value at  $v$ .
- Each country decides to invest  $a_1 \in [0, 1]$  and  $a_2 \in [0, 1]$ .
- The probability of country 1 winning the war is

$$p(a_1, a_2) = \frac{a_1^n}{a_1^n + a_2^n}$$


- Country 1's utility is given by

$$\begin{aligned} U_1(a_1) &= \underbrace{1 - a_1}_{\text{cost}} + \underbrace{p(a_1, a_2)v}_{\text{Expected Benefit}} \\ &= 1 - a_1 + \frac{a_1^n}{a_1^n + a_2^n} v \end{aligned}$$

- Suppose country 2 selected value  $x$ . What should country 1 invest to maximize utility?

## Example 5: IR Bargaining (from Jim Fearon, Part 1)

$n = 1, v = 0.5$



## Example 5: IR War (from Jim Fearon, Part 1)

$$\begin{aligned}\frac{\partial U_1(a_1)}{\partial a_1} &= -1 + \frac{na_1^{n-1}(a_1^n + x^n) - (na_1^{n-1}a_1^n)}{(a_1^n + x^n)^2}v \\ &= -1 + \frac{na_1^{n-1}x^n}{(a_1^n + x^n)^2}v\end{aligned}$$

Set  $n = 1$  (for simplicity)

$$\begin{aligned}0 &= -1 + \frac{x}{(a_1 + x)^2}v \\ a_1^* &= \sqrt{v}\sqrt{x} - x\end{aligned}\tag{0.1}$$

Second derivative!

$$U_1''(a_1) = \frac{-2vx}{(a_1 + x)^3}$$

## Example 5: IR Bargaining (from Jim Fearon, Part 1)

One more—check endpoints

$$a_1^* = 0, \text{ if } \sqrt{v}\sqrt{x} - x < 0$$

$$a_1^* = 0, \text{ if } \sqrt{v} < \sqrt{x}$$


$$a_1^* = \sqrt{v}\sqrt{x} - x \text{ otherwise}$$

# Optimization Challenge Problem

- Suppose a candidate is attempting to mobilize voters. Suppose that for each investment of  $x \in [0, \infty)$  the candidate receives return of  $x^{1/2}$ , but incurs cost of  $ax$ . So, candidate utility is,

$$U_i = x^{1/2} - ax$$

What is the optimal investment  $x^*$ ?



# Computational Optimization Approaches

**Analytic** (Closed form)  $\rightsquigarrow$  Often difficult, impractical, or unavailable

# Computational Optimization Approaches

**Analytic** (Closed form)  $\rightsquigarrow$  Often difficult, impractical, or unavailable  
**Computational**

# Computational Optimization Approaches

**Analytic** (Closed form)  $\rightsquigarrow$  Often difficult, impractical, or unavailable  
**Computational**  $\rightsquigarrow$  **iterative** algorithm that converges to a solution  
(hopefully the right one!)



# Computational Optimization Approaches

- Analytic** (Closed form)  $\rightsquigarrow$  Often difficult, impractical, or unavailable
- Computational**  $\rightsquigarrow$  **iterative** algorithm that converges to a solution (hopefully the right one!)
- Methods for optimization:

# Computational Optimization Approaches

**Analytic** (Closed form)  $\rightsquigarrow$  Often difficult, impractical, or unavailable  
**Computational**  $\rightsquigarrow$  **iterative** algorithm that converges to a solution  
(hopefully the right one!)

- Methods for optimization:
  - **Newton's method** and related methods

# Computational Optimization Approaches

**Analytic** (Closed form)  $\rightsquigarrow$  Often difficult, impractical, or unavailable  
**Computational**  $\rightsquigarrow$  **iterative** algorithm that converges to a solution  
(hopefully the right one!)

- Methods for optimization:
  - **Newton's method** and related methods
  - Gradient descent (ascent)

# Computational Optimization Approaches

**Analytic** (Closed form)  $\rightsquigarrow$  Often difficult, impractical, or unavailable  
**Computational**  $\rightsquigarrow$  **iterative** algorithm that converges to a solution  
(hopefully the right one!)

- Methods for optimization:

- **Newton's method** and related methods
- Gradient descent (ascent)
- Expectation Maximization

# Computational Optimization Approaches

**Analytic** (Closed form)  $\rightsquigarrow$  Often difficult, impractical, or unavailable  
**Computational**  $\rightsquigarrow$  **iterative** algorithm that converges to a solution  
(hopefully the right one!)

- Methods for optimization:

- **Newton's method** and related methods
- Gradient descent (ascent)
- Expectation Maximization
- Genetic Optimization

# Computational Optimization Approaches

**Analytic** (Closed form)  $\rightsquigarrow$  Often difficult, impractical, or unavailable  
**Computational**  $\rightsquigarrow$  **iterative** algorithm that converges to a solution  
(hopefully the right one!)

- Methods for optimization:

- **Newton's method** and related methods
- Gradient descent (ascent)
- Expectation Maximization
- Genetic Optimization
- Branch and Bound ...

# Newton-Raphson Method

Iterative procedure to find a **root**

# Newton-Raphson Method

Iterative procedure to find a **root**

Often solving for  $x$  when  $f(x) = 0$  is hard  $\rightsquigarrow$  complicated function



# Newton-Raphson Method

Iterative procedure to find a **root**

Often solving for  $x$  when  $f(x) = 0$  is hard  $\rightsquigarrow$  complicated function

Solving for  $x$  when  $f(x)$  is linear  $\rightsquigarrow$  easy

# Newton-Raphson Method


Iterative procedure to find a **root**

Often solving for  $x$  when  $f(x) = 0$  is hard  $\rightsquigarrow$  complicated function


Solving for  $x$  when  $f(x)$  is linear  $\rightsquigarrow$  easy

Approximate with **tangent line**, iteratively update


# Tangent Line




# Tangent Line




# Tangent Line




# Tangent Line



# Tangent Line




# Tangent Line







# Tangent Line




# Tangent Line




# Tangent Line




# Tangent Line




# Tangent Line




# Tangent Line



# Tangent Line




# Tangent Line







# Tangent Line




# Tangent Line




# Tangent Line




# Tangent Line




# Tangent Line




# Tangent Line



# Tangent Line




# Tangent Line







# Tangent Line




# Tangent Line



# Tangent Line



# Tangent Line



# Tangent Line

Formula for Tangent line at  $x_0$ :

# Tangent Line

Formula for Tangent line at  $x_0$ :

$$g(x) = f'(x_0)(x - x_0) + f(x_0)$$

# Tangent Line

Formula for Tangent line at  $x_0$ :

$$g(x) = f'(x_0)(x - x_0) + f(x_0)$$

# Tangent Line

Formula for Tangent line at  $x_0$ :

$$g(x) = f'(x_0)(x - x_0) + f(x_0)$$



# Tangent Line

Formula for Tangent line at  $x_0$ :

$$g(x) = f'(x_0)(x - x_0) + f(x_0)$$

# Newton-Raphson Method

Suppose we have some initial guess  $x_0$ . We're going to approximate  $f'(x)$  with the tangent line to generate a new guess

# Newton-Raphson Method

Suppose we have some initial guess  $x_0$ . We're going to approximate  $f'(x)$  with the tangent line to generate a new guess

$$g(x) = f''(x_0)(x - x_0) + f'(x_0)$$

# Newton-Raphson Method

Suppose we have some initial guess  $x_0$ . We're going to approximate  $f'(x)$  with the tangent line to generate a new guess

$$\begin{aligned}g(x) &= f''(x_0)(x - x_0) + f'(x_0) \\ 0 &= f''(x_0)(x_1 - x_0) + f'(x_0)\end{aligned}$$

# Newton-Raphson Method


Suppose we have some initial guess  $x_0$ . We're going to approximate  $f'(x)$  with the tangent line to generate a new guess

$$\begin{aligned}g(x) &= f''(x_0)(x - x_0) + f'(x_0) \\0 &= f''(x_0)(x_1 - x_0) + f'(x_0) \\x_1 &= x_0 - \frac{f'(x_0)}{f''(x_0)}\end{aligned}$$

# Example Function

$f(x) = x^3 + 2x^2 - 1$  find  $x$  that maximizes  $f(x)$  with  $x \in [-3, 0]$

$$x^3 + 2x^2 - 1$$




$$f'(x) = 3x^2 + 4x$$

$$f''(x) = 6x + 4$$

Suppose we have guess  $x_t$  then the next step is:


$$x_{t+1} = x_t - \frac{3x_t^2 + 4x_t}{6x_t + 4}$$

$$x^3 + 2x^2 - 1$$







$$x^3 + 2x^2 - 1$$




$$x^3 + 2x^2 - 1$$




$$x^3 + 2x^2 - 1$$




$$x^3 + 2x^2 - 1$$




$$x^3 + 2x^2 - 1$$




$$x^* = -1.3333$$



# What is Happening with the Roots




# What is Happening with the Roots







# What is Happening with the Roots




# What is Happening with the Roots




# What is Happening with the Roots




# What is Happening with the Roots




# What is Happening with the Roots




# What is Happening with the Roots



# What is Happening with the Roots




# What is Happening with the Roots







# What is Happening with the Roots




# What is Happening with the Roots




# What is Happening with the Roots




$$x^3 + 2x^2 - 1$$




$$x^3 + 2x^2 - 1$$




$$x^3 + 2x^2 - 1$$



$$x^3 + 2x^2 - 1$$




$$x^3 + 2x^2 - 1$$







$$x^3 + 2x^2 - 1$$




$$x^3 + 2x^2 - 1$$




$$x^3 + 2x^2 - 1$$




$$x^3 + 2x^2 - 1$$




$$x^3 + 2x^2 - 1$$



$$x^3 + 2x^2 - 1$$



$$x^3 + 2x^2 - 1$$



To the R Code!



# Today/Tomorrow

- A Framework for optimization
  - Analytic: pencil and paper math
  - Computational: iterative algorithm that aids in solution
- Integration: antidifferentiation/area finding