

Chapter 5: Estimation

(Ott & Longnecker Sections: 4.12, 4.14 and 5.2)

Duzhe Wang

<https://dzwang91.github.io/stat371/>

Part 4



WISCONSIN
UNIVERSITY OF WISCONSIN-MADISON

What do we study?



Key Concepts: t-distribution, confidence intervals

¹Some of the slides in this lecture have been adapted/borrowed from materials developed by Cecile Ane.




1 Review of point estimation

2 The t-distribution

3 Confidence interval

Big bend lizards tail length





- Our goal is to estimate μ , the population mean tail length in the entire Big bend lizards.

- Our goal is to estimate μ , the population mean tail length in the entire Big bend lizards.

```
> bigbend
[1] 8.8 9.7 10.8 7.1 6.6 9.9 10.2 8.6 10.4 11.9 7.6 8.0 8.5
[16] 7.4 8.3 9.1 9.2 7.9 8.4 11.3 6.2 8.8
> mean(bigbend)
[1] 8.895833
> sd(bigbend)
[1] 1.429953
> length(bigbend)
[1] 24
```

- Our goal is to estimate μ , the population mean tail length in the entire Big bend lizards.

```
> bigbend
[1] 8.8 9.7 10.8 7.1 6.6 9.9 10.2 8.6 10.4 11.9 7.6 8.0 8.5
[16] 7.4 8.3 9.1 9.2 7.9 8.4 11.3 6.2 8.8
> mean(bigbend)
[1] 8.895833
> sd(bigbend)
[1] 1.429953
> length(bigbend)
[1] 24
```

- $\bar{X} = 8.896$ cm is one estimate for μ .

- Our goal is to estimate μ , the population mean tail length in the entire Big bend lizards.

```
> bigbend
[1] 8.8 9.7 10.8 7.1 6.6 9.9 10.2 8.6 10.4 11.9 7.6 8.0 8.5
[16] 7.4 8.3 9.1 9.2 7.9 8.4 11.3 6.2 8.8
> mean(bigbend)
[1] 8.895833
> sd(bigbend)
[1] 1.429953
> length(bigbend)
[1] 24
```

- $\bar{X} = 8.896$ cm is one estimate for μ .
- How good is this estimate? How far is μ from 8.896 cm?



- The standard error of \bar{X} is $\frac{\sigma}{\sqrt{n}}$, but we don't know σ .



- The standard error of \bar{X} is $\frac{\sigma}{\sqrt{n}}$, but we don't know σ .
- estimated $SE = \frac{s}{\sqrt{n}}$ is the estimated standard error of the mean.
- $s=1.43$ and $n=24$, so estimated $SE = \frac{1.43}{\sqrt{24}} = 0.292$.
- The estimated SE gives us an idea of how far \bar{X} is from μ typically.



1 Review of point estimation

2 The t-distribution

3 Confidence interval

- If X_1, \dots, X_n have a normal distribution, then

$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$$

- If X_1, \dots, X_n have a normal distribution, then


$$\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$$

- When we replace σ/\sqrt{n} by estimated $SE=s/\sqrt{n}$,

$$\frac{\bar{X} - \mu}{s/\sqrt{n}} \sim T_v$$

where $v = n - 1$ is called **degrees of freedom** and T_v is called t-distribution with degrees of freedom v .

The t-distribution




It looks very similar to a standard normal: it's symmetric and bell-shaped, but it is a little more spread out. The amount of additional spread decreases as the degrees of freedom (the sample size) increases.



How do we find a value t s.t. $P(T_{10} \geq t) = 0.17$.

How do we find a value t s.t. $P(T_{10} \geq t) = 0.17$.

Table A.8 t Curve Tail Areas



The diagram shows a bell-shaped curve representing a t-distribution. The horizontal axis is labeled with 0 at the center and t to the right. The area under the curve to the right of t is shaded and labeled 'Area to the right of t'. The curve itself is labeled 't curve'.

t	ν	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
0.0		.500	.500	.500	.500	.500	.500	.500	.500	.500	.500	.500	.500	.500	.500	.500	.500	.500	.500
0.1		.468	.465	.463	.463	.462	.462	.462	.461	.461	.461	.461	.461	.461	.461	.461	.461	.461	.461
0.2		.437	.430	.427	.426	.425	.424	.424	.423	.423	.423	.423	.422	.422	.422	.422	.422	.422	.422
0.3		.407	.396	.392	.390	.388	.387	.386	.386	.385	.385	.385	.385	.384	.384	.384	.384	.384	.384
0.4		.379	.364	.358	.355	.353	.352	.351	.350	.349	.349	.348	.348	.348	.347	.347	.347	.347	.347
0.5		.352	.333	.326	.322	.319	.317	.316	.315	.315	.314	.313	.313	.313	.312	.312	.312	.312	.312
0.6		.328	.305	.295	.290	.287	.285	.284	.283	.282	.281	.280	.280	.279	.279	.279	.278	.278	.278
0.7		.306	.278	.267	.261	.258	.255	.253	.252	.251	.250	.249	.249	.248	.247	.247	.247	.247	.246
0.8		.285	.254	.241	.234	.230	.227	.225	.223	.222	.221	.220	.220	.219	.218	.218	.218	.217	.217
0.9		.267	.232	.217	.210	.205	.201	.199	.197	.196	.195	.194	.193	.192	.191	.191	.191	.190	.190
1.0		.250	.211	.196	.187	.182	.178	.175	.173	.172	.170	.169	.169	.168	.167	.167	.166	.166	.165
1.1		.235	.193	.176	.167	.162	.157	.154	.152	.150	.149	.147	.146	.146	.144	.144	.144	.143	.143
1.2		.221	.177	.158	.148	.142	.138	.135	.132	.130	.129	.128	.127	.126	.124	.124	.124	.123	.123
1.3		.209	.162	.142	.132	.125	.121	.117	.115	.113	.111	.110	.109	.108	.107	.107	.106	.105	.105
1.4		.197	.148	.128	.117	.110	.106	.102	.100	.098	.096	.095	.093	.092	.091	.091	.090	.090	.089
1.5		.187	.136	.115	.104	.097	.092	.089	.086	.084	.082	.081	.080	.079	.077	.077	.077	.076	.075
1.6		.178	.125	.104	.092	.085	.080	.077	.074	.072	.070	.069	.068	.067	.065	.065	.065	.064	.064
1.7		.169	.116	.094	.082	.075	.070	.065	.064	.062	.060	.059	.057	.056	.055	.055	.054	.054	.053
1.8		.161	.107	.085	.073	.066	.061	.057	.055	.053	.051	.050	.049	.048	.046	.046	.045	.045	.044
1.9		.154	.099	.077	.065	.058	.053	.050	.047	.045	.043	.042	.041	.040	.038	.038	.038	.037	.037
2.0		.148	.092	.070	.058	.051	.046	.043	.040	.038	.037	.035	.034	.033	.032	.032	.031	.031	.030


Download T table from <https://dzwang91.github.io/stat371/resource/>.



1 Review of point estimation


2 The t-distribution

3 Confidence interval



- Point estimates are almost always wrong.
- Why not collect a lot of good guesses which form an interval, and let the interval cover the population mean with high probability?

Interpretation of a confidence interval



A 95% confidence interval indicates that 19 out of 20 samples (95%) from the same population will produce confidence intervals that contain the population parameter.


In confidence interval, the population mean μ is a **fixed** unknown constant, the interval is **random**.

Mechanics of a confidence interval: case 1



If we know the population standard deviation σ ,

- 1 Choose a confidence level $1 - \alpha$. Typically, if we require 95% confidence level, then $\alpha = 0.05$.
- 2 Use z table to find the $z_{\frac{\alpha}{2}}$ critical value such that $P(-z_{\frac{\alpha}{2}} \leq Z \leq z_{\frac{\alpha}{2}}) = 1 - \alpha$.



- 3 Construct the interval: (L, U) , where $L = \bar{X} - z_{\frac{\alpha}{2}} * \frac{\sigma}{\sqrt{n}}$, $U = \bar{X} + z_{\frac{\alpha}{2}} * \frac{\sigma}{\sqrt{n}}$. (Why do we construct this way?)
- 4 Conclude: $P(L \leq \mu \leq U) = 1 - \alpha$. We are $(1 - \alpha) \times 100\%$ confident that the population mean is between (L, U) .

If we don't know the population standard deviation σ ,

- 1 Choose a confidence level $1 - \alpha$. Typically, if we require 95% confidence level, then $\alpha = 0.05$.
- 2 Find the value t such that $P(-t \leq T_{n-1} \leq t) = 1 - \alpha$. It also means $P(T_{n-1} \geq t) = \frac{\alpha}{2}$. Use t table with degrees of freedom $n-1$. We denote the value t as $t_{n-1, \alpha/2}$.
- 3 Construct the interval: (L, U) , where $L = \bar{X} - t_{n-1, \alpha/2} \frac{S}{\sqrt{n}}$, $U = \bar{X} + t_{n-1, \alpha/2} \frac{S}{\sqrt{n}}$. (Why do we construct this way?)
- 4 Conclude: $P(L \leq \mu \leq U) = 1 - \alpha$. We are $(1 - \alpha) \times 100\%$ confident that the population mean is between (L, U) .

Example: big bend lizards tail length



- 1 We want to construct a 90% confidence interval for the population mean tail length, so $\alpha = 0.1$.

Example: big bend lizards tail length



- 1 We want to construct a 90% confidence interval for the population mean tail length, so $\alpha = 0.1$.
- 2 Find the value t such that $P(T_{23} \geq t) = \frac{\alpha}{2} = 0.05$. Use t table with degrees of freedom $n-1=24-1=23$.

Example: big bend lizards tail length



- 1 We want to construct a 90% confidence interval for the population mean tail length, so $\alpha = 0.1$.
- 2 Find the value t such that $P(T_{23} \geq t) = \frac{\alpha}{2} = 0.05$. Use t table with degrees of freedom $n-1=24-1=23$.
t-table gives: $t=1.71$

Example: big bend lizards tail length



- 1 We want to construct a 90% confidence interval for the population mean tail length, so $\alpha = 0.1$.
- 2 Find the value t such that $P(T_{23} \geq t) = \frac{\alpha}{2} = 0.05$. Use t table with degrees of freedom $n-1=24-1=23$.
t-table gives: $t=1.71$
or use R: `qt(0.95, df=23)`

Example: big bend lizards tail length



- 1 We want to construct a 90% confidence interval for the population mean tail length, so $\alpha = 0.1$.
- 2 Find the value t such that $P(T_{23} \geq t) = \frac{\alpha}{2} = 0.05$. Use t table with degrees of freedom $n-1=24-1=23$.
t-table gives: $t=1.71$
or use R: `qt(0.95, df=23)`
- 3 Construct the interval: (L, U) , where

$$L = \bar{X} - t_{n-1, \alpha/2} \frac{S}{\sqrt{n}} = 8.896 - 1.71 * \frac{1.43}{\sqrt{24}} = 8.396,$$

$$U = \bar{X} + t_{n-1, \alpha/2} \frac{S}{\sqrt{n}} = 8.896 + 1.71 * \frac{1.43}{\sqrt{24}} = 9.396$$

Example: big bend lizards tail length



- 1 We want to construct a 90% confidence interval for the population mean tail length, so $\alpha = 0.1$.
- 2 Find the value t such that $P(T_{23} \geq t) = \frac{\alpha}{2} = 0.05$. Use t table with degrees of freedom $n-1=24-1=23$.
t-table gives: $t=1.71$
or use R: `qt(0.95, df=23)`
- 3 Construct the interval: (L, U) , where

$$L = \bar{X} - t_{n-1, \alpha/2} \frac{S}{\sqrt{n}} = 8.896 - 1.71 * \frac{1.43}{\sqrt{24}} = 8.396,$$

$$U = \bar{X} + t_{n-1, \alpha/2} \frac{S}{\sqrt{n}} = 8.896 + 1.71 * \frac{1.43}{\sqrt{24}} = 9.396$$

- 4 Conclude.



See R codes from the course webpage.

What's the next?



In the next lecture, we'll discuss sample size and population proportions.