

Poster Oral presentation82 Sixth European Hemiptera Congress – 25-29 June 2012,
Blagoevgrad, Bulgaria

**Contribution to the knowledge of Psylloidea (Hemiptera) of
Southeastern Anatolia, Turkey**

İnanç Özgen¹, Celalettin Gözüağık², Daniel Burckhardt

¹ Fırat University, Turkey

² Plant Protection Research Station, Diyarbakır, Turkey

³ Naturhistorisches Museum Augustinergasse 2, CH-4052 Basel, Switzerland

Key words: Psylloidea, fauna, Turkey

A study on the psyllid fauna of ten plant species was carried out in 2007 and 2008 in Southeastern Anatolia, Turkey. In total, 12 species were identified. These species were *Cyamophila glycyrrhizae* (Becker, 1864), *Spanioneura pechai* (Klimaszewski & Lodos, 1977), *Homotoma ficus* (Linnaeus, 1758), *Euphyllura straminea* Loginova, 1973, *Trioza magnisetosa* Loginova, 1964, *Aphalara grandicula* (Gegechkori, 1981), *Trioza neglecta* Loginova, 1978, *Trioza remota* Foerster, 1848, *Cacopsylla permixta* Burckhardt & Hodkinson, 1986, *Cacopsylla bidens* (Šulc, 1907), *Cacopsylla marianne* (Baeva, 1966), and *Agonoscena pistaciae* Burckhardt & Lauterer, 1989