




# ARTmie: Fast and Flexible Computation of Aerosol Optical Properties for Atmospheric Modeling and Remote Sensing

Enrico P. Metzner  <sup>†¶</sup>

<sup>†</sup> Karlsruhe Institute of Technology (KIT), Germany <sup>¶</sup> Corresponding author

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

## Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Open Journals](#) 

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

## Summary

In atmospheric science, astronomy, and related fields, accurate computation of particle optical properties — such as mass extinction coefficient, single-scattering albedo, asymmetry factor, and phase function — is essential for modeling interactions with radiation. For spherical particles, these properties can be calculated using Mie theory, a well-established analytical solution. The foundational formulas and early computational algorithms for Mie scattering were first formalized and improved by Bohren & Huffman (1983).

ARTmie implements these Mie formulas to compute extinction, scattering, absorption, and backscattering efficiencies, as well as the corresponding cross sections. It ports key routines from (Mätzler, 2002) (originally in MATLAB) and Bessel function implementations from (Amos, 1986) (Fortran 77) into optimized C++.

ARTmie was developed with a strong focus on computational speed, particularly for applications involving particle size distributions and angle-weighted backscattering efficiencies — important for lidar-based remote sensing. Its performance gains stem not only from the C++ backend, but also from algorithmic optimizations that reduce redundant calculations, especially when computing ensemble-averaged properties (e.g., for log-normal distributions).

Compared to widely used Mie codes, ARTmie offers significantly faster execution, making it ideal for large-scale aerosol optical property calculations in atmospheric models, remote sensing applications, and the creation of high-resolution datasets for machine learning (ML). It supports standard optical property outputs and is designed for seamless integration into modeling pipelines or data-driven workflows.

## Statement of need

Accurate calculation of aerosol optical properties is fundamental to research in physics, atmospheric science, and astronomy. As the complexity of aerosol mixtures increases and the number of particles considered grows, the computational demand for Mie scattering calculations becomes significant. Traditional Mie codes written in FORTRAN or Python often struggle with performance in such scenarios.

ARTmie addresses this challenge by offering a fast and efficient solution for computing Mie scattering properties. It significantly outperforms pure Python implementations, making it suitable for large-scale applications, including high-resolution look-up table generation and ensemble simulations.

Beyond the standard Mie efficiencies for extinction, scattering, and absorption, ARTmie also computes the backscatter efficiency through a weighted average over the backward scattering angles — an important metric for interpreting lidar-based remote sensing measurements.

40 Despite its performance-oriented design, ARTmie remains user-friendly and requires minimal  
41 setup, enabling easy integration into research workflows.

42 ARTmie has already been successfully applied in the ICON-ART module of the ICON numerical  
43 weather and climate prediction model ((MPI-M et al., 2025; Rieger et al., 2015)). It was used  
44 to recalculate the optical properties of aerosols routinely used in ICON-ART and to extend the  
45 model with additional aerosol types and coating configurations (Hoshyaripour et al., 2025).

## 46 State of the field

47 There are many Mie scattering libraries available in a variety of programming languages. Python  
48 is probably the most accessible programming language for quickly and easily analyzing and  
49 testing data within the scientific community. ARTmie is not the first Python module for Mie  
50 scattering; therefore, it will be compared to three other commonly used Python libraries:

### 51 1. miepython (Prah, 2025) :

52 miepython is a minimal, pure Python implementation of classical Mie theory that provides  
53 efficient scattering, absorption and extinction calculations for homogeneous spheres only.  
54 While it is ideal for simple, educational or small-scale use cases, it lacks support for  
55 coated particles or size distributions. Its performance is limited due to the absence of a  
56 compiled backend accelerator.

### 57 2. PyMieScatt (Sumlin et al., 2018) :

58 PyMieScatt is a feature-rich Python library that supports homogeneous and single-coated  
59 spheres. It also offers additional tools for data analysis and angular scattering, as well  
60 as inversion methods. However, like miepython, it is implemented entirely in Python,  
61 making it significantly slower than ARTmie for coated particles and polydisperse systems,  
62 where computational demands are higher.

### 63 3. PyMieSim (Sivry, 2022; Sivry-Houle et al., 2023) :

64 Designed for simulating optical systems, PyMieSim provides a modular interface for  
65 defining sources, detectors and particle configurations. It supports both homogeneous  
66 and single-coated spheres. Although it uses a C++ backend to optimize performance,  
67 the modular design introduces additional setup overhead. PyMieSim is best suited to  
68 integrating Mie scattering into broader optical modelling workflows but does not yet  
69 support polydisperse particles.

## 70 Acknowledgement

71 This work was done during the funding by the Deutsche Forschungsgemeinschaft (DFG) under  
72 the project HO 5275/6-1.

## 73 References

- 74 Amos, D. E. (1986). Algorithm 644: A portable package for bessel functions of a complex  
75 argument and nonnegative order. *ACM Trans. Math. Softw.*, 12(3), 265–273. <https://doi.org/10.1145/7921.214331>  
76
- 77 Bohren, C. F., & Huffman, D. R. (1983). Absorption and scattering of light by small particles.  
78 *Wiley-VCH*.
- 79 Hoshyaripour, G. A., Baer, A., Bierbauer, S., Bruckert, J., Brunner, D., Diekmann, C., Förstner,  
80 J., Hamzehloo, A., Hanft, V., Keller, C., Klose, M., Ludwig, P., Metzner, E. P., Muth,  
81 L., Pauling, A., Ramezani, M., Ziarani, Reddmann, T., Reißig, L., ... Hoose, C. (2025).

- 82 The atmospheric composition component of the ICON modeling framework: ICON-ART  
83 version 2025.04. *EGUsphere*, 2025, 1–54. <https://doi.org/10.5194/egusphere-2025-3400>
- 84 Mätzler, C. (2002). MATLAB functions for mie scattering and absorption, version 2. *IAP*  
85 *Research Report*, 2002-11.
- 86 (MPI-M, I. partnership, DWD, DKRZ, KIT, & C2SM). (2025). *ICON release 2025.04*.  
87 World Data Center for Climate (WDCC) at DKRZ. [https://doi.org/10.35089/wdcc/](https://doi.org/10.35089/wdcc/iconrelease2025.04)  
88 [iconrelease2025.04](https://doi.org/10.35089/wdcc/iconrelease2025.04)
- 89 Prah, S. (2025). *Miepython: Pure python calculation of mie scattering* (Version 3.0.0).  
90 Zenodo. <https://doi.org/10.5281/zenodo.15035350>
- 91 Rieger, D., Bangert, M., Bischoff-Gauss, I., Förstner, J., Lundgren, K., Reinert, D., Schröter,  
92 J., Vogel, H., Zängl, G., Ruhnke, R., & Vogel, B. (2015). ICON-ART 1.0 – a new online-  
93 coupled model system from the global to regional scale. *Geoscientific Model Development*,  
94 8(6), 1659–1676. <https://doi.org/10.5194/gmd-8-1659-2015>
- 95 Sivry, M. P. de. (2022). *MartinPdeS/PyMieSim: PyMieSim beta (2.0.0)* (Version v2.0.0).  
96 Zenodo. <https://doi.org/10.5281/zenodo.7407114>
- 97 Sivry-Houle, M. P. de, Godbout, N., & Boudoux, C. (2023). PyMieSim: An open-source  
98 library for fast and flexible far-field mie scattering simulations. *Opt. Continuum*, 2(3),  
99 520–534. <https://doi.org/10.1364/OPTCON.473102>
- 100 Sumlin, B. J., Heinson, W. R., & Chakrabarty, R. K. (2018). Retrieving the aerosol complex  
101 refractive index using PyMieScatt: A mie computational package with visualization  
102 capabilities. *Journal of Quantitative Spectroscopy and Radiative Transfer*, 205, 127–134.  
103 <https://doi.org/10.1016/j.jqsrt.2017.10.012>