

# The Conjugate Gradient Method

Michael Overton

Convex and Nonsmooth Optimization, Spring 2020

The Conjugate Gradient (CG) method (or the linear conjugate gradient method, or the method of conjugate gradients) is the standard iterative method for minimizing the quadratic function

$$\frac{1}{2}x^T Ax - b^T x$$

where  $A$  is an  $n \times n$  symmetric positive definite matrix, or equivalently, solving the linear system of equations  $Ax = b$ . The method is defined in many books. Here we use the notation in [Chapter 7 of \*A First Course in Numerical Methods\* by Ascher and Greif \(login with NYU netid\)](#) (p. 184). The main theorem about the method is on p. 187 of the same chapter. The proof is not given there, as it is a little long, but it can be found in many books, including Trefethen and Bau's *Numerical Linear Algebra* (however there the notation is different, and the assumption that  $x_0 = 0$  is made there for convenience). Here, we prove something quite simple, which is still an important basic property of the method.

Let  $\text{span}(v_1, \dots, v_m)$  denote the linear span of vectors  $v_1, \dots, v_m$ , that is:

$$\text{span}(v_1, \dots, v_m) = \{w : w = \gamma_1 v_1 + \dots + \gamma_m v_m \text{ for some } \gamma_1, \dots, \gamma_m \in \mathbb{R}\}.$$

Define the  $k$ th Krylov space of  $A$  with respect to  $b$  as

$$\mathcal{K}_k = \text{span}(b, Ab, A^2b, \dots, A^{k-1}b).$$

Exercise: show that if  $w \in \mathcal{K}_k$ , then  $Aw \in \mathcal{K}_{k+1}$ .

**Theorem.** Assume  $x_0 = 0$ . Then, for  $k = 1, 2, \dots$ , the following statements hold:

1. The  $(k-1)$ th residual vector  $r_{k-1}$  is in  $\mathcal{K}_k$
2. The  $(k-1)$ th direction vector  $p_{k-1}$  is in  $\mathcal{K}_k$
3. The  $k$ th solution approximation vector  $x_k$  is in  $\mathcal{K}_k$ .

The proof is by induction. For  $k = 0$ , we have by definition that  $p_0 = r_0 = b$  and  $x_1 = 0 + \alpha_0 p_0$ , so, since  $\alpha_0$  is a real scalar, the result holds.

Now assume the inductive hypothesis, namely, these three properties hold for given  $k$ ; we must show they also hold when  $k$  is replaced by  $k + 1$ . We have

$$r_k = r_{k-1} - \alpha_{k-1} A p_{k-1}$$

so, since  $r_{k-1}$  and  $p_{k-1}$  are both in  $\mathcal{K}_k$  by the inductive hypothesis, and  $A$  times any vector in  $\mathcal{K}_k$  is in  $\mathcal{K}_{k+1}$ , we have that  $r_k \in \mathcal{K}_{k+1}$ . Furthermore, we have

$$p_k = r_k + \frac{\delta_k}{\delta_{k-1}} p_{k-1}$$

so, since  $p_{k-1} \in \mathcal{K}_k$ , and  $r_k \in \mathcal{K}_{k+1}$ , then  $p_k \in \mathcal{K}_{k+1}$ . Finally, we have

$$x_{k+1} = x_k + \alpha_k p_k$$

so, since  $x_k \in \mathcal{K}_k$ , and  $p_k \in \mathcal{K}_{k+1}$ , then  $x_{k+1} \in \mathcal{K}_{k+1}$ . This proves the theorem.