

I don't know, are you sure you want to do this?

T. Thorarinsdottir¹, P. Guttorp¹, M. Drews²

Corresponding author: Name, Address. (email)

¹Norwegian Computing Centere

²Danish Technological University

³Affiliation three

² **Abstract.** ...

1. Introduction

2. Sea level projections (PG)

2.1. Global sea level

Most climate models do not explicitly provide sea level as an output of the calculations.

We will use the empirical approach of Rahmstorf and collaborators [Rahmstorf2007, @; Rahmstorf, Perrette, and VermeerRahmstorf et al.2011, @], employing the statistical modeling of Bolin et al. [2014] to relate global annual mean temperature anomalies [Hansen et al. , 2001] to global mean sea level anomalies [Church and White , 2011].

We then apply the estimated historical relationship to projected temperatures from the CMIP5 experiment [Taylor et al., 2012] to obtain projected global annual mean sea level.

2.2. Local sea level

In order to get from global sea level projections to local ones, it is important to note that sea level rise not is uniform over the globe. Glacial and land ice melting affect the local sea level differently depending on where the melted ice is located. Another major effect in Fennoskandia is the land rise due to isostatic effects of the glaciers from the last ice age. Again, we will use historical data to relate global sea level to isostatically corrected local sea level using a time series regression model.

The local sea level projections are then obtained by first relating projected temperature to global sea level, and then relating the global sea level to the local one. Each climate model temperature projection yields a different local sea level projection.

2.3. Uncertainty assessment

Following the approach of Guttorp et al. [2014] we assess the uncertainty in the local sea level projections taking into account the variability between the climate projections used, the uncertainties in the regressions of global mean temperature on global mean sealevel and of global on local sea level. We express the sea level projection uncertainty in terms of a confidence band that is simultaneously of the intended level for all projection years. This allows us, for example, to get a confidence band for the years when a given sea level rise is obtained.

2.4. Limitations of the sea level projections

The main assumption is using historical relationships in statistical projections of the type used in this paper is that there is no major change in how temperature relates to sea level, globally and locally. Among the factors that may invalidate this approach are changes in water storage on land, changes in the rates of glacial and land ice melt, and changes in Earth's gravitational field due to transfer of mass from land ice to ocean water. For example, the rate of ice melt on Greenland may suddenly increase substantially due to intense warming of both air and sea water. Our current climate models are not able to resolve the ice processes sufficiently to include such so called tipping points into the projections.

3. Decision tools (KdB, MD, TT)

3.1. Timing of adaptation measures

We consider adaptation decision making related to the timing of proactive adaptation measures. That is, the goal is to adapt to sea level rise before major damages occur. In a

cost-benefit framework, an investment should be delayed as long as the benefits of delay (avoided investment costs) are greater than the associated costs (higher climate change damages) [Fankhauser et al., 1999].

Fankhauser et al. [1999] describe a deterministic framework where an adaptation investment of C^0 now (at time $n = 0$) leads to unmitigated damage of d_0^0 in period 0, and a stream of partially mitigated damages d_t^0 in periods $t = 1, 2, \dots$. If r is the discount rate, the net present value damage, D^0 , associated with this investment is

$$D^0 = C^0 + d_0^0 + \frac{d_1^0}{1+r} + \frac{d_2^0}{(1+r)^2} + \dots \quad (1)$$

In comparison, postponing the adaptation investment to time period $n = 1$ would lead to unmitigated damages in periods 0 and 1, and partially mitigated damages, d_t^1 , thereafter.

The delay would be preferable if

$$C^0 - \frac{C^1}{(1+r)} > (d_0^1 - d_0^0) + \frac{d_1^1 - d_1^0}{1+r} + \frac{d_2^1 - d_2^0}{(1+r)^2} + \dots$$

Here, the expression on the left describes the benefits of the delay while the expression on the right describes the cost of the delay. In the simplest case, there is no change in investment costs ($C^0 = C^1 = C$) and the delay has no lasting effects beyond period 1 ($d_t^1 = d_t^0$ for $t > 1$). In this case, the comparison is between the expected return r earned on the capital while implementation is delayed and one additional time period of unmitigated damage,

$$rC > d_1^1 - d_1^0.$$

3.2. Limitations of the decision framework

4. Case studies

4.1. Data (PG)

4.2. Timing of adaptation measures (KdB, TT)

56 A case study focusing on and comparing different cities in Norway.

4.3. Selection of adaptation measures(?) (MD)

57 A case study focusing on Denmark.

5. Conclusions

58 **Acknowledgments.** This work was funded by NordForsk through project number
 59 74456 “Statistical Analysis of Climate Projections” (eSACP) and The Research Council
 60 of Norway through project number 243953 “Physical and Statistical Analysis of Climate
 61 Extremes in Large Datasets” (ClimateXL). The source code for the analysis is imple-
 62 mented in the statistical programming language R (<http://www.R-project.org>) and is
 63 available on GitHub at <http://github.com/eSACP/...>

References

- 64 Bolin, D., P. Guttorp, A. Januzzi, M. Novak, H. Podschwit, L. Richardson, C. Sowder, A.
 65 Särkkä and A. Zimmerman (2014), Statistical prediction of global sea level from global
 66 temperature. *Statistica Sinica*, *25*, 351–367.
- 67 Church, J. and N. White (2011), Sea-level rise from the late 19th to the early 21st century.
 68 *Surveys in Geophysics* *32*, 585–602.
- 69 Fankhauser, S., J. B. Smith, and R. S. J. Tol (1999), Weathering climate change: some
 70 simple rules to guide adaptation decisions, *Ecological Economics*, *30*, 67–78.

- 71 Guttorp, P., D. Bolin, A. Januzzi, D. Jones, M. Novak, H. Podschwit, L. Richardson,
72 A. Särkkä, C. Sowder and A Zimmerman (2014), Assessing the uncertainty in project-
73 ing local mean sea level from global temperature. *Journal of Applied Meteorology and*
74 *Climatology*. 53, 2163–2170.
- 75 Hansen, J., R. Rued, M. Sato, M. Imhoff, W. Lawrence, D. Easterling, T. Peterson, and
76 T. Karl (2001), A closer look at United States and global surface temperature change.
77 *J. Geophys. Res.* 106, 23947–23963.
- 78 Rahmstorf, S. (2007). A semi-empirical approach to projecting future sea-level rise. *Sci-*
79 *ence* 315, 368–370.
- 80 Rahmstorf, S., M. Perrette, and M. Vermeer (2011), Testing the robustness of semi-
81 empirical sea level projections. climate dynamics. *Climate Dynamics* 39, 861–875.
- 82 Robert, C. P., and G. Casella (2004), *Monte Carlo Statistical Methods*, 2nd ed., Springer,
83 New York.
- 84 Rue, H., S. Martino, and N. Chopin (2009), Approximate Bayesian Inference for Latent
85 Gaussian Models Using Integrated Nested Laplace Approximations (with Discussion),
86 *Journal of the Royal Statistical Society, Series B*, 71, 319–392.
- 87 Taylor, K., R. Stouffer, and G. Meehl, 2012: An overview of CMIP5 and the experiment
88 design. *Bull. Amer. Meteor. Soc.*, **93**, 485–498.