

THÈSE PRÉSENTÉE  
POUR OBTENIR LE GRADE DE  
**DOCTEUR**  
**DE L'UNIVERSITÉ DE BORDEAUX**  
ECOLE DOCTORALE SCIENCES ET ENVIRONNEMENTS  
ÉCOLOGIE ÉVOLUTIVE, FONCTIONNELLE ET DES COMMUNAUTÉS

Par **Maxime Lavaud**

Confined Brownian Motion

Sous la direction de : **Thomas Salez**  
Co-direction : **Yacine Amarouchene**

Soutenue le 25 décembre 2019

Membres du jury :

Mme. Aude ALPHA	Directrice de Recherche	Université	Rapporteur
M. Bernard BETA	Directeur de Recherche	Université	Rapporteur
M. Georges GAMMA	Directeur de Recherche	Université	Président
Mme. Dominique DELTA	Chargée de Recherche	Université	Examinatrice
M. Eric EPSILON	Ingénieur de Recherche	Université	Examineur
Mme. Jane DOE	Directrice de Recherche	Université	Directrice
Mme. Simone UNTEL	Ingénieure de Recherche	Université	Invitée

## Abstract

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

# Table of Contents

Abstract	i
Table of Contents	ii
List of Figures	iii
List of Tables	iv
1 Introduction	v

## List of Figures

## List of Tables

# 1 Introduction

Since the observations of Gordon Moore in the 60's we know that the technological progress is bound to our ability to miniaturize. It's indeed due to the miniaturization that we are able to have more computational power leading to the rise of new technologies like the Deep Learning that showed the need of large computational capabilities by having the computer program *AlphaGo* beating *Lee Sedol* one of the greatest player of *Go* in 2016. Since this powerful demonstration AIs using the same technologies are showing up in every field, from the language translator to autonomous cars and is now starting to be extensively used in physics with in 2020 the first focus session on machine learning at the *March Meeting* that continued this year with presentations at every sessions. The success of Deep learning is not due to the fact that it's new and fancy algorithm since it known for several decade but only the fact that the miniaturization permitted to do the stunning amount of computation needed to have a smart AI. Our ability to use this technologies is finally bound to our ability to understand the surface physics at the nanometer scale.

On another side we have microfluidic since the 80s which is an incredible multidisciplinary field involving chemistry, engineering, soft matter physics and also biotechnology. Microfluidic permitted the development of daily life technologies like the ink-jet printers or more advanced tools such as DNA chips or lab-on-a-chip technology. The ability to compose with a lot of different system to build microfluidic systems is a wonderful playground for physicists which gave a lot of complex systems in confinement to study and understand how different boundaries can change the dynamic properties of a system. At a time of miniaturization and nanotechnologies, the need of tools permitting the systematic study of complex confined system is a key.

In order to address these challenges my work in the past three years focused on using the confined Brownian motion. Brownian Motion is a central paradigm in modern science. It has implications in fundamental physics, biology, and even finance, to name a few. By understanding that the apparent erratic motion of colloids is a direct consequence of the thermal motion of surrounding fluid molecules, pioneers like Einstein and Perrin provided decisive evidence for the existence of atoms, Specifically, free Brownian motion in the bulk is characterized by a typical spatial extent evolving as the square root of time, as well as Gaussian displacements. At a time of miniaturization and interfacial science, and moving beyond the idealized bulk picture, it is relevant to consider the added roles of boundaries to the above context. Indeed, Brownian motion at interfaces and in confinement is a widespread practical situation in microbiology and nanofluidics. In such case, surface effects become dominant and alter drastically the Brownian statistics, with key

implications towards: i) the understanding and smart control of the interfacial dynamics of microscale entities; and ii) high-resolution measurements of surface forces at equilibrium. Interestingly, a confined colloid will exhibit non-Gaussian statistics in displacements, due to the presence of multiplicative noises induced by the hindered mobility near the wall. Besides, the particle can be subjected to electrostatic or Van der Waals forces exerted by the interface, and might experience slippage too. Considering the two-body problem, the nearby boundary can also induce some effective interaction. Previous studies have designed novel methods to measure the diffusion coefficient of confined colloids, or to infer surface forces.