

Supplementary material for: When do we need multipel infectious disease models? Agreement between projection rank and magnitude in a multi-model setting

La Keisha Wade-Malone, The Pennsylvania State University

Emily Howerton, The Pennsylvania State University

William J.M. Probert, University of Oxford

Michael C. Runge, U.S. Geological Survey

Cecile Viboud, National Institutes of Health

Katriona Shea, The Pennsylvania State University




Fig. S1: SIR dynamics of infected individuals for varying transmission rate (β) and recovery rate (γ) parameters. Dynamics are shown for three scenarios: no intervention (solid line), non-pharmaceutical interventions (NPIs) that reduce transmission by 30% (short dashed line), and vaccination of approximately 1% of the susceptible population per day (long dashed line).




Fig. S2: Results from Fig. 1B repeated for varying intervention efficacies (vaccination rate ranging from 0.5% to 2% of the population per day, and NPI reduction of the transmission rate ranging from 10% to 40%). Each panel shows a large part of the space of possible models across biological uncertainties (transmission rate, β , and recovery rate, γ). Each position on the graph represents an individual model, the color of the tile represents the recommended intervention (orange: non-pharmaceutical interventions, NPIs; purple: vaccination), and the intensity of the color is the magnitude of the projected difference in cumulative infections. Colors have been rescaled from Fig. 1. The gray line shows where the difference in cumulative infections is 0, or where neither intervention is recommended over the other. Black contours show sample values of individual model $R_0 = 2, 3.5, 6$, and 10.




Fig. S3: Model results for minimizing peak infections. Results are shown for varying intervention efficacies (vaccination rate ranging from 0.5% to 2% of the population per day, and NPI reduction of the transmission rate ranging from 10% to 40%). Each panel shows a large part of the space of possible models across biological uncertainties (transmission rate, β , and recovery rate, γ). Each position on the graph represents an individual model, the color of the tile represents the recommended intervention (orange: non-pharmaceutical interventions, NPIs; purple: vaccination), and the intensity of the color is the magnitude of the projected difference in peak infections. Colors have been rescaled from Figure 1. The gray line shows where the difference in peak infections is 0, or where neither intervention is recommended over the other. Black contours show sample values of individual model $R_0 = 2, 3.5, 6$, and 10 .




Fig. S4: Probability of models agreeing about intervention scenario ranking or estimates of cumulative infections without intervention for different sampling schemes (columns) and number of models agreeing (rows). Probabilities are calculated for a set of models of size two to size twelve, either when choosing randomly among all models or among models with “similar” assumptions, defined as those with transmission and recovery rate assumptions within ± 0.1 , ± 0.05 or ± 0.15 of a given model. Agreement probability of at least 2, 4, 6, or 8 models is shown for estimates of cumulative cases within 20 infections (2% of population, red), within 50 infections (5% of population, yellow), within 100 infections (10% of population, blue), or intervention scenario ranking (black).




Fig. S5: Probability of models agreeing about intervention scenario ranking or estimates of cumulative infections without intervention (columns) for different degrees of model similarity (rows). Probabilities are calculated for a set of models of size two to size twelve, either when choosing randomly among all models or among models with “similar” assumptions, defined as those with transmission and recovery rate assumptions within ± 0.1 , ± 0.05 or ± 0.15 of a given model. Agreement probability of at least 2, 4, 6, or 8 models is shown for estimates of cumulative cases within 20 infections (2% of population), within 50 infections (5% of population), within 100 infections (10% of population), or intervention scenario ranking. Breakpoints where agreement probability is at least 0.5 is shown with a solid black line and agreement probability is at least 0.9 is shown with a dashed black line.




Fig. S6: Results from Fig. 3 replicated with agreement assessed for only two intervention scenarios. For a varying number of models in a set, agreement is measured for projection magnitude across intervention scenarios (using intraclass correlation coefficient, ICC) and ranking of intervention scenarios (using Kendall's W). Results are shown when models in the set come from all possible models considered (red), and when models are similar (blue). Each point represents one set of randomly selected models from either all possible models considered (red) or a neighborhood of “similar” models (blue, those with transmission and recovery rate assumptions within ± 0.1 of a given model). Ellipses show the area within which 95% of points fall. Dotted horizontal lines show agreement that is significant with a p-value of 0.05.

Round 1 (horizon: 26 weeks)



Fig. S7: COVID-19 Scenario Modeling Hub projections of cumulative hospitalizations for Round 1. Projections are collected from 4 models for four scenarios. Across locations (panels), projections are shown for each scenario (separated by horizontal black lines) from each model (individual points), points are colored based on the rank of that scenario for each model (blue is lowest projection, and red is highest projection).

Round 2 (horizon: 26 weeks)




Fig. S8: COVID-19 Scenario Modeling Hub projections of cumulative hospitalizations for Round 2. Projections are collected from 5 models for four scenarios. Across locations (panels), projections are shown for each scenario (separated by horizontal black lines) from each model (individual points), points are colored based on the rank of that scenario for each model (blue is lowest projection, and red is highest projection).

Round 3 (horizon: 26 weeks)




Fig. S9: COVID-19 Scenario Modeling Hub projections of cumulative hospitalizations for Round 3. Projections are collected from 4 models for four scenarios. Across locations (panels), projections are shown for each scenario (separated by horizontal black lines) from each model (individual points), points are colored based on the rank of that scenario for each model (blue is lowest projection, and red is highest projection).

Round 4 (horizon: 26 weeks)




Fig. S10: COVID-19 Scenario Modeling Hub projections of cumulative hospitalizations for Round 4. Projections are collected from 6 models for four scenarios. Across locations (panels), projections are shown for each scenario (separated by horizontal black lines) from each model (individual points), points are colored based on the rank of that scenario for each model (blue is lowest projection, and red is highest projection).

Round 5 (horizon: 26 weeks)




Fig. S11: COVID-19 Scenario Modeling Hub projections of cumulative hospitalizations for Round 5. Projections are collected from 7 models for four scenarios. Across locations (panels), projections are shown for each scenario (separated by horizontal black lines) from each model (individual points), points are colored based on the rank of that scenario for each model (blue is lowest projection, and red is highest projection).

Round 6 (horizon: 26 weeks)



Fig. S12: COVID-19 Scenario Modeling Hub projections of cumulative hospitalizations for Round 6. Projections are collected from 8 models for four scenarios. Across locations (panels), projections are shown for each scenario (separated by horizontal black lines) from each model (individual points), points are colored based on the rank of that scenario for each model (blue is lowest projection, and red is highest projection).

Round 7 (horizon: 26 weeks)



Fig. S13: COVID-19 Scenario Modeling Hub projections of cumulative hospitalizations for Round 7. Projections are collected from 8 models for four scenarios. Across locations (panels), projections are shown for each scenario (separated by horizontal black lines) from each model (individual points), points are colored based on the rank of that scenario for each model (blue is lowest projection, and red is highest projection).

Round 9 (horizon: 26 weeks)




Fig. S14: NULL

Round 11 (horizon: 12 weeks)



Fig. S15: COVID-19 Scenario Modeling Hub projections of cumulative hospitalizations for Round 9. Projections are collected from 8 models for four scenarios. Across locations (panels), projections are shown for each scenario (separated by horizontal black lines) from each model (individual points), points are colored based on the rank of that scenario for each model (blue is lowest projection, and red is highest projection).

Round 12 (horizon: 12 weeks)



Fig. S16: NULL

Round 13 (horizon: 52 weeks)




Fig. S17: COVID-19 Scenario Modeling Hub projections of cumulative hospitalizations for Round 11. Projections are collected from 6 models for four scenarios. Across locations (panels), projections are shown for each scenario (separated by horizontal black lines) from each model (individual points), points are colored based on the rank of that scenario for each model (blue is lowest projection, and red is highest projection).

Round 14 (horizon: 50 weeks)




Fig. S18: COVID-19 Scenario Modeling Hub projections of cumulative hospitalizations for Round 12. Projections are collected from 6 models for four scenarios. Across locations (panels), projections are shown for each scenario (separated by horizontal black lines) from each model (individual points), points are colored based on the rank of that scenario for each model (blue is lowest projection, and red is highest projection).

Round 15 (horizon: 40 weeks)




Fig. S19: COVID-19 Scenario Modeling Hub projections of cumulative hospitalizations for Round 13. Projections are collected from 8 models for four scenarios. Across locations (panels), projections are shown for each scenario (separated by horizontal black lines) from each model (individual points), points are colored based on the rank of that scenario for each model (blue is lowest projection, and red is highest projection).

Round 16 (horizon: 26 weeks)




Fig. S20: COVID-19 Scenario Modeling Hub projections of cumulative hospitalizations for Round 14. Projections are collected from 8 models for four scenarios. Across locations (panels), projections are shown for each scenario (separated by horizontal black lines) from each model (individual points), points are colored based on the rank of that scenario for each model (blue is lowest projection, and red is highest projection).

Round 1 (horizon: 26 weeks)




Fig. S21: Agreement for 1,000 null model simulations and the corresponding SMH projections of cumulative hospitalizations for Round 1. Agreement is calculated for projection magnitude (as measured by intraclass correlation coefficient) and scenario ranking (as measured by Kendall's W). Across locations (panels), null model results are shown with black points and SMH results are shown with a red point.

Round 2 (horizon: 26 weeks)




Fig. S22: Agreement for 1,000 null model simulations and the corresponding SMH projections of cumulative hospitalizations for Round 2. Agreement is calculated for projection magnitude (as measured by intraclass correlation coefficient) and scenario ranking (as measured by Kendall's W). Across locations (panels), null model results are shown with black points and SMH results are shown with a red point.

Round 3 (horizon: 26 weeks)




Fig. S23: Agreement for 1,000 null model simulations and the corresponding SMH projections of cumulative hospitalizations for Round 3. Agreement is calculated for projection magnitude (as measured by intraclass correlation coefficient) and scenario ranking (as measured by Kendall's W). Across locations (panels), null model results are shown with black points and SMH results are shown with a red point.

Round 4 (horizon: 26 weeks)




Fig. S24: Agreement for 1,000 null model simulations and the corresponding SMH projections of cumulative hospitalizations for Round 4. Agreement is calculated for projection magnitude (as measured by intraclass correlation coefficient) and scenario ranking (as measured by Kendall's W). Across locations (panels), null model results are shown with black points and SMH results are shown with a red point.

Round 5 (horizon: 26 weeks)




Fig. S25: Agreement for 1,000 null model simulations and the corresponding SMH projections of cumulative hospitalizations for Round 5. Agreement is calculated for projection magnitude (as measured by intraclass correlation coefficient) and scenario ranking (as measured by Kendall's W). Across locations (panels), null model results are shown with black points and SMH results are shown with a red point.

Round 6 (horizon: 26 weeks)




Fig. S26: Agreement for 1,000 null model simulations and the corresponding SMH projections of cumulative hospitalizations for Round 6. Agreement is calculated for projection magnitude (as measured by intraclass correlation coefficient) and scenario ranking (as measured by Kendall's W). Across locations (panels), null model results are shown with black points and SMH results are shown with a red point.

Round 7 (horizon: 26 weeks)




Fig. S27: Agreement for 1,000 null model simulations and the corresponding SMH projections of cumulative hospitalizations for Round 7. Agreement is calculated for projection magnitude (as measured by intraclass correlation coefficient) and scenario ranking (as measured by Kendall's W). Across locations (panels), null model results are shown with black points and SMH results are shown with a red point.

Round 9 (horizon: 26 weeks)




Fig. S28: Agreement for 1,000 null model simulations and the corresponding SMH projections of cumulative hospitalizations for Round 9. Agreement is calculated for projection magnitude (as measured by intraclass correlation coefficient) and scenario ranking (as measured by Kendall's W). Across locations (panels), null model results are shown with black points and SMH results are shown with a red point.

Round 11 (horizon: 12 weeks)




Fig. S29: Agreement for 1,000 null model simulations and the corresponding SMH projections of cumulative hospitalizations for Round 11. Agreement is calculated for projection magnitude (as measured by intraclass correlation coefficient) and scenario ranking (as measured by Kendall's W). Across locations (panels), null model results are shown with black points and SMH results are shown with a red point.

Round 12 (horizon: 12 weeks)




Fig. S30: Agreement for 1,000 null model simulations and the corresponding SMH projections of cumulative hospitalizations for Round 12. Agreement is calculated for projection magnitude (as measured by intraclass correlation coefficient) and scenario ranking (as measured by Kendall's W). Across locations (panels), null model results are shown with black points and SMH results are shown with a red point.

Round 13 (horizon: 52 weeks)




Fig. S31: Agreement for 1,000 null model simulations and the corresponding SMH projections of cumulative hospitalizations for Round 13. Agreement is calculated for projection magnitude (as measured by intraclass correlation coefficient) and scenario ranking (as measured by Kendall's W). Across locations (panels), null model results are shown with black points and SMH results are shown with a red point.

Round 14 (horizon: 50 weeks)




Fig. S32: Agreement for 1,000 null model simulations and the corresponding SMH projections of cumulative hospitalizations for Round 14. Agreement is calculated for projection magnitude (as measured by intraclass correlation coefficient) and scenario ranking (as measured by Kendall's W). Across locations (panels), null model results are shown with black points and SMH results are shown with a red point.

Round 15 (horizon: 40 weeks)




Fig. S33: Agreement for 1,000 null model simulations and the corresponding SMH projections of cumulative hospitalizations for Round 15. Agreement is calculated for projection magnitude (as measured by intraclass correlation coefficient) and scenario ranking (as measured by Kendall's W). Across locations (panels), null model results are shown with black points and SMH results are shown with a red point.

Round 16 (horizon: 26 weeks)




Fig. S34: Agreement for 1,000 null model simulations and the corresponding SMH projections of cumulative hospitalizations for Round 16. Agreement is calculated for projection magnitude (as measured by intraclass correlation coefficient) and scenario ranking (as measured by Kendall's W). Across locations (panels), null model results are shown with black points and SMH results are shown with a red point.




Fig. S35: Agreement between projection magnitude and ranks across evaluated SMH rounds (columns) and SMH projection horizons (rows) for SMH Q25. Agreement between projection magnitude is measured by intraclass correlation coefficient, and agreement between scenario ranks is measured by Kendall's W. Each point represents results for a set of projections from a single round, target, location, week. Maximum projection horizon is shown for each round, but horizon varies by round so not all horizons are available for all rounds.




Fig. S36: Agreement between projection magnitude and ranks across evaluated SMH rounds (columns) and SMH projection horizons (rows) for SMH Q50. Agreement between projection magnitude is measured by intraclass correlation coefficient, and agreement between scenario ranks is measured by Kendall's W. Each point represents results for a set of projections from a single round, target, location, week. Maximum projection horizon is shown for each round, but horizon varies by round so not all horizons are available for all rounds.




Fig. S37: Agreement between projection magnitude and ranks across evaluated SMH rounds (columns) and SMH projection horizons (rows) for SMH Q75. Agreement between projection magnitude is measured by intraclass correlation coefficient, and agreement between scenario ranks is measured by Kendall's W. Each point represents results for a set of projections from a single round, target, location, week. Maximum projection horizon is shown for each round, but horizon varies by round so not all horizons are available for all rounds.




Fig. S38: Comparison of intraclass correlation coefficient (ICC) and Kendall's W between quantiles. Each metric was assessed on the projected median, as well as Q25 and Q50. The left panel shows comparisons between Q25 and Q50, and the right panel shows comparisons between Q75 and Q50. Each point represents one projection (for a single round, location).