

**Instructions:** (10 points total - 5 pts each) Show all work for credit. You may use your book, but no other resource. **GS:** Scan ~~THREE~~ FOUR pages for your solutions.

1. Consider the two dimensional vector field

$$\mathbf{F}(x, y) = \left\langle e^{xy}(y \sin(x) + \cos(x)), xe^{xy} \sin(x) + \frac{1}{y} \right\rangle$$

defined on all of  $\mathbb{R}^2$ . *the upper half-plane in  $\mathbb{R}^2$ .*

(a) Prove that  $\mathbf{F}$  is conservative, then find its potential function  $f(x, y)$ .

$$\begin{aligned} \text{With } P &= e^{xy}(y \sin x + \cos x), \quad \frac{\partial P}{\partial y} = e^{xy}[\sin x] + xe^{xy}y \sin x + xe^{xy} \cos x \\ &= e^{xy}(\sin x + xy \sin x + x \cos x) \end{aligned}$$

$$\begin{aligned} Q &= xe^{xy} \sin x + \frac{1}{y}, \quad \frac{\partial Q}{\partial x} = [xe^{xy}] \cos x + [xe^{xy}y + (1)e^{xy}] \sin x \\ &= e^{xy}(x \cos x + xy \sin x + \sin x) \end{aligned}$$

Then  $\frac{\partial P}{\partial y} = \frac{\partial Q}{\partial x}$  on The upper-half plane (simply-connected, open, etc.)

therefore,  $\mathbf{F}$  is conservative.

Moreover, if  $Q = \frac{\partial f}{\partial y} = xe^{xy} \sin x + \frac{1}{y}$ , then  $\int \frac{\partial f}{\partial y} dy = \int xe^{xy} \sin x + \frac{1}{y} dy$

$= e^{xy} \sin x + \ln(y) + c(x) = f(x, y)$  where  $c(x)$  is a function of  $x$  alone.

Then  $\frac{\partial f}{\partial x} = e^{xy}(y \sin x + \cos x) + \ln(y) + c'(x) = P \Rightarrow c(x) = C$ . Thus

(b) Letting  $C$  be the line segment joining  $(0, 1)$  to the point  $(0, \frac{\pi}{2})$ , compute the line integral  $\int_C \mathbf{F} \cdot d\mathbf{r}$ .

$$f(x, y) = e^{xy}(y \sin x + \cos x) + \ln(y) + C$$

For (b) use  $f(x, y)$  the potential function.

$$\begin{aligned} \int_C \mathbf{F} \cdot d\mathbf{r} &= f(0, \frac{\pi}{2}) - f(0, 1) = e^{0(\pi/2)} \left( \frac{\pi}{2} \sin(0) + \cos(0) + \ln\left(\frac{\pi}{2}\right) \right) - \left[ e^{0(1)} (1 \sin(0) + \cos(0)) + \ln(1) \right] \\ &= (1 + \ln(\frac{\pi}{2})) - (1 + 0) = \boxed{\ln(\frac{\pi}{2})} \end{aligned}$$

2. In this problem you will show that the line integral

$$\oint_C \mathbf{F} \cdot d\mathbf{r} = 2\pi$$

for the vector field  $\mathbf{F}(x, y) = \left\langle \frac{-y}{x^2+y^2}, \frac{x}{x^2+y^2} \right\rangle$  and  $C$  any positively oriented simple closed circle enclosing the origin. **Note** that the vector field  $\mathbf{F}$  is not defined at the origin, so the domain is the punctured plane.


(a) Letting  $C_r$  and  $C_R$  denote the circles of radius  $r < R$ , first compute by parameterizing the circle that  $\oint_{C_R} \mathbf{F} \cdot d\mathbf{r} = 2\pi$ .

$$\vec{r}(t) = \langle R \cos t, R \sin t \rangle \quad 0 \leq t \leq 2\pi \quad \vec{r}'(t) = \langle -R \sin t, R \cos t \rangle \quad \text{Note: } x^2 + y^2 = R^2$$

$$\vec{F}(\vec{r}(t)) = \left\langle \frac{-R \sin t}{R^2}, \frac{R \cos t}{R^2} \right\rangle = \frac{1}{R} \langle -\sin t, \cos t \rangle$$

$$\oint_C \vec{F} \cdot d\vec{r} = \int_0^{2\pi} \left\langle \frac{-1}{R} \sin t, \frac{1}{R} \cos t \right\rangle \cdot \langle -R \sin t, R \cos t \rangle dt = \int_0^{2\pi} \sin^2 t + \cos^2 t dt = \boxed{2\pi}$$

(b) Now use the extended Green's theorem to compute that  $\oint_{C_r} \mathbf{F} \cdot d\mathbf{r} = \oint_{C_R} \mathbf{F} \cdot d\mathbf{r}$ . See picture.



$C_r$  = inner circle

$C_R$  = outer circle

Red line segments help with extension

Caution: Watch orientation of  $C_r, C_R$  in your computations.

But the annulus into two semi-annular regions so that Green's Theorem can be applied.  $D_2$  = bottom annulus  $D_1$  = top. I will use  $A$  for the annular region

$$\oint_{\partial D_1} \vec{F} \cdot d\vec{r} + \oint_{\partial D_2} \vec{F} \cdot d\vec{r} = \iint_{D_1} \left( \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA + \iint_{D_2} \left( \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dA \quad \text{by Green's Theorem}$$

$$= \oint_{C_R} \vec{F} \cdot d\vec{r} - \oint_{C_r} \vec{F} \cdot d\vec{r} = I$$

(continued on next page ...)

Notice: In applying Green's Theorem,  $C_r$  is traversed in the

negative orientation.

Computing:

•  $P = -y(x^2 + y^2)^{-1}$ , then

$$\frac{\partial P}{\partial y} = -y(-1)(x^2 + y^2)^{-2}(2y) - (x^2 + y^2)^{-1}$$

$$= \frac{2y^2}{(x^2 + y^2)^2} - \frac{1}{(x^2 + y^2)}$$

$$= \frac{2y^2 - (x^2 + y^2)}{(x^2 + y^2)^2} = \frac{y^2 - x^2}{(x^2 + y^2)^2}$$

•  $Q = x(x^2 + y^2)^{-1}$ , then  $\frac{\partial Q}{\partial x} = x(-1)(x^2 + y^2)^{-2}(2x) + (1)(x^2 + y^2)^{-1}$

$$= (x^2 + y^2)^{-2}(-2x^2 + (x^2 + y^2))$$
$$= \frac{y^2 - x^2}{(x^2 + y^2)^2}$$

and (miraculously)  $\frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} = 0$ .

Thus,  $I = 0$ ! and since  $I = \oint_{C_R} \vec{F} \cdot d\vec{r} - \oint_{C_r} \vec{F} \cdot d\vec{r} = 0$

$$\oint_{C_R} \vec{F} \cdot d\vec{r} = 2\pi = \oint_{C_r} \vec{F} \cdot d\vec{r}$$

(c) Green's Theorem requires an open simply-connected domain and here there is a "hole" or "puncture" at  $(0,0)$  since  $\vec{F}$  is not defined there.