

Top-down estimates of ammonia (NH_3) emissions to characterise the influence on local air quality





Eloise Marais

University College London (UCL), 11 June 2025



with Martin Van Damme, Lieven Clarisse, Christine Wiedinmyer, Killian Murphy, Guido van der Werf, Alok Pandey, Pierre-François Coheur, Mark Shephard, Karen Cady-Pereira, Tom Misselbrook, Lei Zhu, Gan Luo, Fangqun Yu


Agricultural and fire emissions of ammonia affect air quality, human health, and ecosystem vitality




Ammonia is a precursor of fine particles and deposit nitrogen to sensitive habitats

Ammonia Emissions from Agriculture

UK NH₃ Emissions [thousands tonnes]




Contribution of management practices



[UK Clean Air Strategy, 2019]

>80% of total NH₃ emissions in the UK

Ammonia Emissions from Burning Biomass



Observationally-derived (Top-Down) NH_3 Emissions

Convert atmospheric **column concentrations** to surface **emissions** using a model

IASI COLUMNS




Conversion Factor



EMISSIONS

Satellite columns
(IASI)



Column-to-Emission ratio
(GEOS-Chem)



Satellite-derived
Surface Emissions

Emission




Simple mass-balance inversion method made possible by relatively short NH_3 lifetime

Preprocess IASI to Finer Resolution Grid than Instrument

Use so-called oversampling to enhance spatial resolution relative to native resolution of instrument


Oversampling Technique




Weights pixel by area of overlap

Oversampling technique over London

Satellite pixel resolution



Fixed (~10 km) grid




Lose time (temporal) resolution; gain spatial resolution

Improve resolution from 12-40 km to 10 km for an instrument observing ammonia (NH_3)

Multiyear means from the IASI version 3 ammonia

Multiyear (2008-2018) monthly means for warmer months of the year



Oversampling used to
finer scales (~ 10 km) than
instrument native
resolution (~ 12 km)




[Marais et al., 2021]

Climatological mean to be consistent with bottom-up ammonia emissions

IASI-derived multiyear (2008-2018) monthly mean NH₃ emissions






Focus on **Mar-Sep** when warm temperatures and clearer conditions increase sensitivity to surface NH₃

IASI: morning
overpass



Total monthly
emissions

1 Gg = 1 kilotonne




[Marais et al., 2021]

Monthly emissions for March-September from **IASI**-derived estimates sum to **271.5 Gg**
Uncertainty range is 11-36% and bottom-up is ~27% less than IASI-derived emissions

Satellite vs inventory NH₃ emissions: spatial distribution


Comparison of months with peak emissions according to IASI (April and July)




Large July difference over locations dominated by dairy cattle. Inventory is 27-49% less than the satellite values.

Biomass Burning Ammonia Emissions in Southern Africa

Monthly bottom-up June-October 2019 emissions



MODIS Landcover



Tropical forest
↓
Woody savanna
↓
Savanna

Inventory differences:


- Emission factors
- Fire data
- Time resolution
- Spatial resolution
- Injection height
- Small fires
- Landcover classes

Mostly savanna fires. Some tropical forest fires.

Apply all 3 inventories **GEOS-Chem** to compare to IASI for NH₃

Reprocess IASI version 4 with local GEOS-Chem priors

IASI columns for July-October 2019. Prior from GEOS-Chem with FINN emissions



Overall decline in columns with local a priori, as more NH₃ placed higher up


Less noisy over Atlantic Ocean east of Angola and northern Namibia

More retrievals pass quality checks




Evaluation of Inventories with Satellite Observations

NH₃ vertical column densities for Jul-Oct 2019 [10^{15} molecules cm⁻²]

IASI with
GEOS-Chem
prior:



GEOS-Chem:






June excluded, as no inventories consistent with IASI observations ($R < 0.5$)


Evaluation of Inventories with Satellite Observations

NH₃ vertical column densities for Jul-Oct 2019 [10^{15} molecules cm⁻²]

IASI with
GEOS-Chem
prior:



GEOS-Chem:




June excluded, as no inventories consistent with IASI observations ($R < 0.5$)


Top-down Emissions with Best Performing Inventories

Calculate emissions for GEOS-Chem gridsquares with > 50% biomass burning emissions


(a) Top-down [kt]



(b) Bottom-up [kt]



(b) minus (a)



Uses GEOS-Chem driven with FINN inventory, as yields best agreement with IASI

Emissions long-tailed distribution and degrades correlation with NO_x emissions from $R > 0.8$ to $R < 0.4$

Emissions peak in August, just like inventory. Error estimate on emissions is 1.2 ± 0.4 Tg.

In Summary

- Spatial resolutions of ~12-14 km enhanced by oversampling for UK NH₃, but lots of data needed
- Estimate UK agricultural emissions that are consistent with spring fertilizer location and timing
- Identify large bottom-up and top-down UK agricultural emissions discrepancies in summer over cattle farming intensive regions to steer further research
- FINN inventory better than GFED and GFAS at representing smouldering emissions of NH₃
- Suggests would outperform other inventories for other smouldering pollutants (VOCs, CO, organic aerosols, methane)
- Inventories collocate NH₃ and NO_x emissions (smouldering and flaming fires), but these are mostly separate in top-down estimates
- Need independent observations for biomass burning in Africa. Ideally in National Parks
- Geostationary infrared instrument soon to launch over Europe to observe NH₃ every 30 min (including over Africa) offers tremendous opportunity to advance top-down estimates further
- UK NH₃ paper: <https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2021JD035237>
- Southern Africa NH₃ work in review.