Long-term trends in air quality in tropical megacities using Earth observations

Karn Vohra (kxv745@bham.ac.uk), E. A. Marais, S. Suckra, L. Kramer, W. J. Bloss, R. Sahu, A. Gaur, S. N. Tripathi, M. Van Damme, L. Clarisse, P. F. Coheur






Long-term trends in air quality in tropical megacities using Earth observations


Karn Vohra (kxv745@bham.ac.uk), E. A. Marais, S. Suckra, L. Kramer, W. J. Bloss, R. Sahu, A. Gaur, S. N. Tripathi, M. Van Damme, L. Clarisse, P. F. Coheur

Tropics are the next frontier in air pollution



51 cities within the tropics will be megacities by 2100 [Hoornweg & Pope, 2016]


Assessing Earth observations & trends




- Surface monitoring of air pollutants have limitations
- We assess Earth observations against available surface data
- We then analyse trends from the long-record of Earth observations

Satellite versus surface NO₂

London (2005-2018)



Trend in London NMVOCs




[Vohra et al., ACPD, in review]

Long-term trends in air quality in tropical megacities

Trends in NO_2 (2005-2018)

NASA OMI Level-2 Tropospheric column NO₂ version 3.0




NO₂ has increased in 46 out of 51 cities

Long-term trends in air quality in tropical megacities

Trends in reactive NMVOCs (2005-2018)

QA4ECV OMI Level-2 Total column HCHO version 1.2




HCHO (reactive NMVOCs) has increased in 37 out of 51 cities

Long-term trends in air quality in tropical megacities

Trends in $PM_{2.5}$ (2005-2018)

NASA MODIS Level-2 Dark Target AOD Collection 6.1



AOD has increased in 25 out of 33 cities

Next steps

- Interpret the drivers of these trends
- Compare trends to widely used global emission inventories