
Satellites up above monitor the air down below!!!



The Problem

Exposure to fine particles caused 4.2 millions deaths worldwide in 2015 representing 7.6% of global deaths

Cohen et al., 2017

Estimated excess mortality attributed to air pollution in Europe European Heart Journal, 2019

Pollution map reveals unsafe air quality at almost 2,000 UK sites The Guardian, 2019

Study links air pollution exposure to miscarriages

Zhang et al., 2019

Air pollution may be damaging 'every organ in the body'

Schraufnagel et al, 2019

Air pollution takes decade off memory, study suggests The Times, 2019


Transport: 47% Industries: 38%

What's common???


Nitrogen dioxide (NO₂)

But where is it coming from???

NOx Emissions (2017)




Data from National Atmospheric Emissions Inventory




Surface monitoring networks have their limitations


☐ Expensive to set up and maintain




- ☐ Expensive to set up and maintain
- ☐ Limited spatial and temporal coverage





- ☐ Expensive to set up and maintain
- ☐ Limited spatial and temporal coverage
- □ Limited pollutants monitored





- ☐ Expensive to set up and maintain
- ☐ Limited spatial and temporal coverage
- □ Limited pollutants monitored
- ☐ Issues with data quality










































The Solution


Satellites are the only solution to address this global challenge

Sensors in space have been providing us with petabytes of data for more than 2 decades

Gases

Particles






Future

Completed


On-going


Space-based instruments provide extensive data coverage




Space-based instruments provide extensive data coverage

We develop our tool focusing on 4 dynamic cities





Satellite columns reflect changes in surface pollution




Air pollution is at the surface (~1-2 km)


Column is sensitive to changes in surface concentration

And conduct careful assessment with surface monitors (where available)

Satellite versus surface NO₂ in London




Satellite versus surface NH₃ in Harwell



Points are monthly averages


Pearson's correlation coefficient (R-value) indicates consistency

We use robust statistics to derive trends




NO₂ (NO_x) over London decreased by **35** % from 2005 to 2018

Air pollution control measures are working in the UK While air quality has been deteriorating over time in India




Next steps

High spatial resolution satellite data




TROPOMI NO₂ for April 2018 [Geffen et al., 2019]

Global 3D Chemical Transport Model




Interpret data using model to understand drivers of trends and inter-annual variability in air pollutants



A global problem

No routine monitoring of air pollutants in many parts of the world


Publicly available NO₂ data for 14th October, 2019



A scalable solution

The algorithms that we have developed can be applied to any city in the world

Global annual average NO₂ concentrations observed with OMI for 2005-2006



Acknowledgements























Thank you and everyone for your attention!!!