

Health Burdens and Racial-Ethnic Disparities of Air Pollution from US Oil and Gas Lifecycle Stages



Karn Vohra, Eloise A. Marais, Ploy Achakulwisut, Susan Anenberg, Colin Harkins

Sierra Club

13 November 2025

Acknowledgements

Funding:



- Schmidt Family Foundation's 11th Hour Project program
- European Research Council (**ERC**) under the European Union's Horizon 2020 research and innovation programme

Multi-institution Collaboration:

- University College London (UK)
- Stockholm Environment Institute (Seattle)
- George Washington University (DC)
- University of Colorado, Boulder
- US National Oceanographic and Atmospheric Administration (NOAA)

National Contributions to Global Natural Gas Production

Top natural gas production countries and regions in 2023
billion cubic feet per day




[Energy Information Administration (EIA), <https://www.eia.gov/todayinenergy/index.php>]

US largest global producer of natural gas (and oil)

Trends in US Crude Oil Production


U.S. crude oil production by state (2010-2019)

million barrels per day (b/d)



Annual change (2018-2019)


thousand barrels per day



[EIA, <https://www.eia.gov/todayinenergy/detail.php?id=43015>]

Steep growth since 2011 in oil (and natural gas) production spurred by unconventional oil and gas (O&G) and phase-out of coal


Trends in US Energy Consumption



[EIA, <https://www.eia.gov/todayinenergy/detail.php?id=52959>]


Growth in demand for O&G continues to grow, despite pledges to Paris Climate Accord

Major Stages of the Oil and Gas Lifecycle



End use includes combusted crude and refined fuels and non-energy products (waxes, plastics)

Activities in All Stages Produce Pollutants



BTEX: benzene, toluene, ethylbenzene, xylene

HAPs: Hazardous air pollutants

VOCs: volatile organic compounds

NO_x: nitrogen oxides (NO + NO₂)


PM_{2.5}: Small particles

[Adapted from
Donaghy et al. 2023]

Directly emitted: NO_x, primary PM_{2.5}, most VOCs/HAPs


Formed via chemical reactions: secondary PM_{2.5}, VOCs/HAPs such as formaldehyde, ozone

Air pollution impacts every organ of the human body




Source: <https://airqoon.com/>

Air pollution affects people throughout their lifetime



Fine Particles (PM_{2.5})

Particles that are mixtures of multiple chemicals that are air small enough to penetrate deep into the lungs



Exposure to PM_{2.5} causes **respiratory** and **cardiovascular** diseases that may lead to early death

Exposure to PM_{2.5} also associated with preterm birth

Ground-level ozone (O_3)

- Secondary pollutant (not emitted)
- Ingredients for its formation include sunlight, VOCs (or CO) and NO_x
- Long-term exposure associated with chronic obstructive pulmonary disease (COPD) leading to premature death across all population age groups



Source: <https://www.ourair.org/gooduphigh-badnearby-ozone-infographic/>

Nitrogen dioxide (NO₂)

- NO₂ is one of the two NO_x compounds
- Well established that long-term NO₂ exposure linked to onset of asthma amongst children
- Increasingly accepted that long-term NO₂ exposure also linked to premature mortality amongst elderly

First UK case with air pollution on death certificate




Source:

https://www.london.gov.uk/sites/default/files/hia_asthma_air_pollution_in_london.pdf

Foundation created in her name
campaigning for clean air for all:
<https://www.ellaroberta.org/>

Volatile Organic Compounds (VOCs)

VOCs that are known carcinogens



X-axis:
Unit Risk Estimate or
one in a million risk of
developing cancer


For example,
There is a 2 in a million
chance to develop cancer
from sustained lifetime
exposure to $1 \mu\text{g m}^{-3}$
acetaldehyde

Benzene is emitted, others
are emitted and produced
from chemical reactions

[Loh et al., 2007]

Unjust Exposure to O&G Air Pollution

Example past study assessing unjust health burden from natural gas flaring



OR: odds ratio

Hispanic population closer to flaring wells, so greater likelihood than non-Hispanic white population of preterm birth (by 0.9-3.4 days)

Unjust exposure from:

- placement of infrastructure (extraction wells, pipelines, compressor stations, downstream chemical manufacturing plants)
- proximity to more intense flaring
- proximity to busy roadways

[Cushing et al., 2020]

Motivation for Our Study

So much already known about O&G harm to health, so why another study on this topic?


Past studies not sufficiently holistic:

- Focused on a single O&G lifecycle stage (mostly production)
- Focused on a single pollutant (typically PM_{2.5})
- Focused on a specific state or natural gas basin
- Combined O&G with coal for fossil fuel focused analysis
- Used reduced complexity chemistry transport models that may not be suitably representing harmful secondary pollutants
- Environmental injustice typically focus on inequitable exposure, quantification of inequitable health outcomes is rare

Vital for directing policies that prioritize the most adversely affected communities

We construct a unified inventory of air pollutant emissions for 2017

US EPA National Emission Inventory (NEI) 2017




Fuel-based Inventory for Vehicular Emissions (FIVE)



Aviation Emissions Inventory Code (AEIC)



Global Fuel Exploitation Inventory (GFEI) for methane




Community Emissions Data System (CEDS) for shipping and outside US





Our inventory is for 2017 as most contemporary datasets were available for 2017

Air pollutant emissions from the US oil and gas lifecycle




Oil and gas activities in the US accounts for more than half of CO, NO_x, formaldehyde, acetaldehyde, benzene, and BC anthropogenic emissions

Computer models are used to determine impact of individual sources on air quality and health




→ Estimate baseline of
air pollution

Computer models are used to determine impact of individual sources on air quality and health






Model runs






→ Estimate contribution of upstream and midstream activities

Health-harming pollutant concentrations from oil and gas activities in 2017




Upstream + Midstream



Downstream




End-use



Largest enhancements from end-use activities, mainly in California and eastern US

Combined upstream/midstream in Texas and neighboring states, downstream in Texas and Louisiana


Annual mean HAPs concentrations linked to oil and gas activities




**Contribution of oil and gas activities to benzene is less than formaldehyde and acetaldehyde.
Lower benzene emissions and formaldehyde/acetaldehyde also formed from other VOCs**

Estimating health burden of O&G air pollution exposure


Health-risk assessment models relate long-term air pollution exposure to adverse health outcomes




Air pollutant concentrations with and without the O&G lifecycle stage




Proportion of health outcome linked to the O&G lifecycle stage



Health burden estimates



Baseline health data from US CDC



Population data from WorldPop

Annual premature mortality linked to oil and gas air pollution


Non-accidental mortality in adults from PM_{2.5} and elderly from NO₂, COPD mortality from MDA8 O₃

Upstream + Midstream

PM_{2.5}-attributable deaths in adults (25+ years)



NO₂-attributable deaths in elderly (65+ years)



MDA8 O₃-attributable deaths across all ages


Downstream

End-use




Oil and gas lifecycle


Oil and gas lifecycle



US totals:
47,200



US totals:
39,000



US totals:
4,750

91,000 premature deaths in CONUS linked to oil and gas activities in the US.
1,170 premature deaths in southern Canada and 440 in northern Mexico




2-3% of all early deaths in the US

Preterm births, asthma and cancer incidences from oil and gas air pollution

Preterm births from PM_{2.5}, childhood-onset asthma from NO₂, lifetime cancers from HAPs

Upstream + Midstream

PM_{2.5}-attributable preterm birth incidences






Downstream



End-use



Oil and gas lifecycle




US oil and gas air pollution is linked to 3% of preterm births, 10% of pediatric asthma incidences, and 4% of all respiratory and hematologic cancers in the US


Absolute disparities in oil and gas air pollution health burden

We estimate population-standardized health burdens for each racial and ethnic group as well as the national average for the total population. Subgroup population demographic data from ACS estimates.


 **Upstream +
Midstream**



 **Downstream**





 **End-use**



Absolute racial and ethnic disparities for the end-use stage are much higher than the other stages

The inequitable health burden of oil and gas air pollution

Population-standardized health burdens for each racial and ethnic group normalized to the national value



Air pollution health burden in Native Americans is 20-50% greater than the national average mortality rates. Burden in the Black population is 20-70% higher than the national estimates. Air pollution asthma and cancer incidence rates in Asian population are 35-50% higher than the national value. Greatest relative disparities almost all occur in the downstream stage

Explore the health burden results in our interactive dashboard

https://bit.ly/US_oilgas_healthburden_dashboard

Severe health burden of air pollution from the US oil and gas sector


Step 1: Select an adverse health outcome.

Early deaths in adults (25+ years) from annual exposure to fine particles (PM2.5)

Step 2 (Optional): Choose one, more or all states.


(All)

Upstream and midstream activities
(Exploration and production) **Totals: 1,294**



Estimate 0 217

Up to 15 worst affected counties




Downstream activities
(Processing or refining) **Totals: 1,616**



State: California
Lifecycle stage: Downstream
Health outcome: Early deaths in adults (25+ years) from annual exposure to fine particles (PM2.5)
Estimate: 375


Estimate 0 375

End-use activities
(Consumption or combustion) **Totals: 44,301**



Estimate 0 375

Up to 15 worst affected counties




For more information, feel free to contact [Dr Karn Vohra](#) or [Prof Eloise Marais](#). Suggested citation: K. Vohra, E. A. Marais, P. Achakulwisut, S. Anenberg, C. Harkins, **The health burden and racial-ethnic disparities of air pollution from the major oil and gas lifecycle stages in the United States**, [doi:10.1126/sciadv.adu2241](https://doi.org/10.1126/sciadv.adu2241), *Science Advances*, 2025.

OR



SCAN ME

The severe and unjust health burden of oil and gas air pollution!



UK ▾

The Guardian

Air pollution from oil and gas causes 90,000 premature US deaths each year, says new study

Study analyzed health impacts of fossil fuels from exploration to end use, and found communities of color bear brunt of harm

<https://www.theguardian.com/us-news/2025/aug/22/air-pollution-oil-gas-health-study>

<https://richardconniff.substack.com/p/oil-and-gas-air-pollution-is-a-killer>

[https://www.thelancet.com/journals/lanplh/article/PIIS2542-5196\(25\)00264-5/fulltext](https://www.thelancet.com/journals/lanplh/article/PIIS2542-5196(25)00264-5/fulltext)

Glass half full, our health burden estimates support substantial health benefits and reductions in unjust exposure that can be achieved with a transition away from oil and gas to cleaner alternatives.