

Spectral asymptotics and scattering theory in the nilpotent Lie group setting

Edward McDonald

(based on joint work with Z. Fan, J. Li, F. Sukochev and D. Zanin)

Penn State University

March 6, 2023

This talk is based on a series of preprints by myself with Zhijie Fan (Wuhan), Ji Li (Macquarie), Fedor Sukochev (UNSW) and Dmitriy Zanin (UNSW).

The first two papers are available:

- a Spectral estimates and asymptotics for stratified Lie groups
arXiv:2201.12349 (with Sukochev and Zanin)
- b Endpoint weak Schatten class estimates and trace formula for commutators of Riesz transforms with multipliers on Heisenberg groups
arXiv:2201.12350 (with Fan, Li, Sukochev and Zanin)

There will also be other papers (currently in preparation).

Plan for this talk

- ① Some elementary background on scattering theory (why do we care?)
- ② Stratified lie groups and recent developments
- ③ The future(?)

We have some technical results on the spectra of operators of the form

$$M_f D^{-1} : L_2(G) \rightarrow L_2(G)$$

where G is a stratified Lie group, M_f is the operator of pointwise multiplication by a function f on G and D is a positive maximally hypoelliptic differential operator on G .

Specifically, we now have a much better understanding of the singular values of these operators.

We have some technical results on the spectra of operators of the form

$$M_f D^{-1} : L_2(G) \rightarrow L_2(G)$$

where G is a stratified Lie group, M_f is the operator of pointwise multiplication by a function f on G and D is a positive maximally hypoelliptic differential operator on G .

Specifically, we now have a much better understanding of the singular values of these operators.

These results are interesting on their own, but I will discuss a program (mostly unfinished) to do scattering theory (in the style of Birman-Kato) for maximally hypoelliptic operators (in the style of Helffer-Nourigat, Androulidakis-Mohsen-Yuncken).

Why do we want to know?

Singular value estimates for operators like $M_f D^{-1}$ have several applications. For example:

- Bound state problems: estimate the number of eigenvalues of $D + M_f$.
- Scattering theory: compare the effect of M_f on the evolution of $\exp(it(D + M_f))$,
- Spectral theory: determine the Weyl asymptotics of general maximally hypoelliptic differential operators.

Very elementary scattering theory

If Q is an elliptic and symmetric differential operator

$$Q : C^\infty(X, E) \rightarrow C^\infty(X, E)$$

where X is compact and Riemannian, and E is some Hermitian vector bundle, then Q is self-adjoint and has a discrete spectral decomposition

$$Q = \sum_{n=0}^{\infty} \lambda(n, Q) P_n$$

where P_n is a finite rank $L_2(X, E)$ -orthogonal projection, and $\{\lambda(n, Q)\}_{n=0}^{\infty}$ enumerates the spectrum in increasing order of absolute value.

Very elementary scattering theory

If Q is an elliptic and symmetric differential operator

$$Q : C^\infty(X, E) \rightarrow C^\infty(X, E)$$

where X is compact and Riemannian, and E is some Hermitian vector bundle, then Q is self-adjoint and has a discrete spectral decomposition

$$Q = \sum_{n=0}^{\infty} \lambda(n, Q) P_n$$

where P_n is a finite rank $L_2(X, E)$ -orthogonal projection, and $\{\lambda(n, Q)\}_{n=0}^{\infty}$ enumerates the spectrum in increasing order of absolute value.

If X is not compact, this is of course not true.

Very elementary scattering theory

Suppose that X is not compact (later, we will simply take $X = \mathbb{R}^d$). If we assume that the geometry of X and E are not so bad and that the coefficients of Q are uniformly bounded in the correct sense, then Q is still self-adjoint but its spectrum is complicated.

Very elementary scattering theory

Suppose that X is not compact (later, we will simply take $X = \mathbb{R}^d$). If we assume that the geometry of X and E are not so bad and that the coefficients of Q are uniformly bounded in the correct sense, then Q is still self-adjoint but its spectrum is complicated.

In this case scattering theory can provide a more useful description.

Very elementary scattering theory

A conventional situation is that we have a symmetric differential operator (on \mathbb{R}^d),

$$D_1 = \sum_{|\alpha| \leq m} a_\alpha(x) \partial^\alpha$$

with smooth coefficients $\{a_\alpha\}$ that are constant outside of a compact set, say $a_\alpha(x) = c_\alpha$. If we define

$$D_0 = \sum_{|\alpha| \leq m} c_\alpha \partial^\alpha$$

The spectral theory of D_0 is easy to understand using the Fourier transform. We expect that D_1 “looks similar” to D_0 in some sense. Ideally, D_1 is relatively compact with respect to D_0 .

Another question we might want to ask about D_1 : does it have a discrete spectrum? If so, where is it?

Uses of scattering theory

The Birman-Kato theory is well-known in geometry and topology. It has several applications.

Uses of scattering theory

The Birman-Kato theory is well-known in geometry and topology. It has several applications.

- Relative index theorems: Suppose that D_1 and D_0 are odd self-adjoint operators on a $\mathbb{Z}/2\mathbb{Z}$ -graded Hilbert space H . The relative index of D_1 with respect to D_0 is

$$\mathrm{ind}(D_1, D_0) = \mathrm{Str}(e^{-tD_1^2} - e^{-tD_0^2})$$

(provided it exists). The relative index is the differences of the indices of D_1 and D_0 , plus an extra term coming from the continuous spectrum. See Eichhorn *Relative Index Theory* (2008), and also Borisov-Müller-Schrader "Relative Index Theorems and Supersymmetric Scattering Theory" (1988).

Uses of scattering theory

The Birman-Kato theory is well-known in geometry and topology. It has several applications.

- Relative index theorems: Suppose that D_1 and D_0 are odd self-adjoint operators on a $\mathbb{Z}/2\mathbb{Z}$ -graded Hilbert space H . The relative index of D_1 with respect to D_0 is

$$\mathrm{ind}(D_1, D_0) = \mathrm{Str}(e^{-tD_1^2} - e^{-tD_0^2})$$

(provided it exists). The relative index is the differences of the indices of D_1 and D_0 , plus an extra term coming from the continuous spectrum. See Eichhorn *Relative Index Theory* (2008), and also Borisov-Müller-Schrader "Relative Index Theorems and Supersymmetric Scattering Theory" (1988).

- Witten index: It is conceivable that one could have a non-Fredholm operator D such that

$$\mathrm{wind}(D) := \lim_{t \rightarrow \infty} \mathrm{Tr}(e^{-tD^*D} - e^{-tDD^*})$$

exists. This is called the Witten index, and can be expressed in terms

Uses of scattering theory

The Birman-Kato theory is well-known in geometry and topology. It has several applications.

- Relative index theorems: Suppose that D_1 and D_0 are odd self-adjoint operators on a $\mathbb{Z}/2\mathbb{Z}$ -graded Hilbert space H . The relative index of D_1 with respect to D_0 is

$$\mathrm{ind}(D_1, D_0) = \mathrm{Str}(e^{-tD_1^2} - e^{-tD_0^2})$$

(provided it exists). The relative index is the differences of the indices of D_1 and D_0 , plus an extra term coming from the continuous spectrum. See Eichhorn *Relative Index Theory* (2008), and also Borisov-Müller-Schrader "Relative Index Theorems and Supersymmetric Scattering Theory" (1988).

- Witten index: It is conceivable that one could have a non-Fredholm operator D such that

$$\mathrm{wind}(D) := \lim_{t \rightarrow \infty} \mathrm{Tr}(e^{-tD^*D} - e^{-tDD^*})$$

exists. This is called the Witten index, and can be expressed in terms

Very elementary scattering theory

Scattering theory is about the solutions to the equation

$$\frac{\partial u}{\partial t} = iD_1 u.$$

Or $u(t) = \exp(itD_1)u(0)$. We want to know when there exists u_+ such that

$$\lim_{t \rightarrow \infty} \|\exp(itD_1)u(0) - \exp(itD_0)u_+\| = 0.$$

Or, alternatively, when there exists a strong limit

$$W_+(D_1, D_0) := s\text{-}\lim_{t \rightarrow \infty} e^{-itD_0} e^{itD_1}.$$

(actually, we are interested in a slight modification of this).

Very elementary scattering theory

With a little more effort, we can compare the solutions of the wave equations

$$\frac{\partial^2 u}{\partial t^2} = D_1 u, \quad \frac{\partial^2 u}{\partial t^2} = D_0 u.$$

(this is called acoustical scattering; see Reed-Simon Volume III.)

Very elementary scattering theory

Let D_0, D_1 be self-adjoint operators on some Hilbert space H , and let $P_{ac}(D_1)$ be the projection onto the absolutely continuous subspace of D_1 . Define two operators $W_{\pm}(D_0, D_1)$ by

$$W_{\pm}(D_0, D_1) := s\text{-}\lim_{t \rightarrow \pm\infty} e^{-itD_0} e^{itD_1} P_{ac}(D_1).$$

These are called the wave operators. We say that the wave operators (if they exist) are *complete* if

$$\text{ran}(W_{\pm}(D_0, D_1)) = P_{ac}(D_0).$$

Birman's theorem

Suppose that A_1, A_0 are self-adjoint operators on a Hilbert space H . If for any bounded interval $I \subset \mathbb{R}$ we have

$$\chi_I(A_1)(A_1 - A_0)\chi_I(A_0) \in \mathcal{L}_1(H)$$

then the wave operators $W_{\pm}(A_1, A_0)$ exist and are complete.

Birman's theorem

Suppose that A_1, A_0 are self-adjoint operators on a Hilbert space H . If for any bounded interval $I \subset \mathbb{R}$ we have

$$\chi_I(A_1)(A_1 - A_0)\chi_I(A_0) \in \mathcal{L}_1(H)$$

then the wave operators $W_{\pm}(A_1, A_0)$ exist and are complete.

There are some additional technical assumptions on A_1, A_0 which I will not address here.

Birman's theorem

Suppose that A_1, A_0 are self-adjoint operators on a Hilbert space H . If for any bounded interval $I \subset \mathbb{R}$ we have

$$\chi_I(A_1)(A_1 - A_0)\chi_I(A_0) \in \mathcal{L}_1(H)$$

then the wave operators $W_{\pm}(A_1, A_0)$ exist and are complete.

There are some additional technical assumptions on A_1, A_0 which I will not address here.

It suffices, for example, to have

$$(A_1 - A_0)(1 + A_0^2)^{-N} \in \mathcal{L}_1(H)$$

for sufficiently large N .

Using Birman's theorem

Consider the pair

$$A_1 = c(x)\Delta, A_0 = \Delta = \sum_{j=1}^d \partial_{x_j}^2$$

on \mathbb{R}^d , where c is a smooth positive function equal to 1 outside a compact set. Then

$$(A_1 - A_0)(1 - \Delta)^{-N} = (c(x) - 1)\Delta(1 - \Delta)^{-N}$$

This belongs to \mathcal{L}_1 for sufficiently large N , thanks to some old results of Birman-Solomyak.

Stratified Lie groups

Let \mathfrak{g} be a Lie algebra which admits a direct sum decomposition

$$\mathfrak{g} = \bigoplus_{n=1}^{\infty} \mathfrak{g}_n$$

where $[\mathfrak{g}_k, \mathfrak{g}_n] \subseteq \mathfrak{g}_{k+n}$ and \mathfrak{g}_1 generates \mathfrak{g} . This is called a stratified Lie algebra.

The number

$$Q := \sum_{n=1}^{\infty} n \dim(\mathfrak{g}_n)$$

is called the homogeneous dimension of \mathfrak{g} .

Exponentiating \mathfrak{g} , we get a simply connected nilpotent Lie group

$$G = \exp(\mathfrak{g}).$$

This is a homeomorphism, and the Lebesgue measure of \mathfrak{g} pushes forward to the Haar measure of G . Suppose that \mathfrak{g}_1 has a basis $\{X_1, \dots, X_m\}$, and G is essentially a Euclidean space \mathbb{R}^d equipped with a family of vector fields

$$X_1, \dots, X_m$$

with polynomial coefficients satisfying the Hörmander condition at every point.

Ellipticity on stratified Lie groups

The stratification of \mathfrak{g} defines a grading on the algebra of invariant differential operators, $\mathcal{U}(\mathfrak{g})$, on G . Say that an operator $P \in \mathcal{U}(\mathfrak{g})$ has order k if the highest degree term in P is homogeneous of degree k .

Theorem (Helffer-Nourigat, Rockland)

Let $P \in \mathcal{U}(\mathfrak{g})$ have degree k . If for every $\pi \in \widehat{G}_u$ (the unitary dual of G), $\pi(P)$ is injective on H_π^∞ (the smooth vectors), then for every Q of degree less than or equal to k we have

$$\|Qu\|_{L_2(G)} \lesssim \|Pu\|_{L_2(G)} + \|u\|_{L_2(G)}, \quad u \in L_2(G).$$

In particular, P is hypoelliptic.

Recently some substantial advances have been made in the study of ellipticity on Heisenberg manifolds and more general foliated manifolds.

Recently some substantial advances have been made in the study of ellipticity on Heisenberg manifolds and more general foliated manifolds. This opens up the possibility to study the scattering theory for maximally hypoelliptic operators.

Some results

Recall that $\{X_1, \dots, X_m\}$ denotes a basis for \mathfrak{g}_1 , the first layer of our stratified Lie algebra. By assumption X_1, X_2, \dots, X_m generate \mathfrak{g} . Let

$$\Delta := \sum_{j=1}^m X_j^2.$$

This is hypoelliptic.

Some results

Recall that $\{X_1, \dots, X_m\}$ denotes a basis for \mathfrak{g}_1 , the first layer of our stratified Lie algebra. By assumption X_1, X_2, \dots, X_m generate \mathfrak{g} . Let

$$\Delta := \sum_{j=1}^m X_j^2.$$

This is hypoelliptic.

Given a function f on G , denote by M_f the (possibly unbounded) operator of pointwise multiplication by f . We want to understand the operators

$$M_f(1 - \Delta)^{-N}, \quad (1 - \Delta)^{-N} M_f(1 - \Delta)^{-N}$$

and their trace ideal properties.

Some results

Recall that $\{X_1, \dots, X_m\}$ denotes a basis for \mathfrak{g}_1 , the first layer of our stratified Lie algebra. By assumption X_1, X_2, \dots, X_m generate \mathfrak{g} . Let

$$\Delta := \sum_{j=1}^m X_j^2.$$

This is hypoelliptic.

Given a function f on G , denote by M_f the (possibly unbounded) operator of pointwise multiplication by f . We want to understand the operators

$$M_f(1 - \Delta)^{-N}, \quad (1 - \Delta)^{-N} M_f(1 - \Delta)^{-N}$$

and their trace ideal properties.

Why is this?

Some results

Recall that $\{X_1, \dots, X_m\}$ denotes a basis for \mathfrak{g}_1 , the first layer of our stratified Lie algebra. By assumption X_1, X_2, \dots, X_m generate \mathfrak{g} . Let

$$\Delta := \sum_{j=1}^m X_j^2.$$

This is hypoelliptic.

Given a function f on G , denote by M_f the (possibly unbounded) operator of pointwise multiplication by f . We want to understand the operators

$$M_f(1 - \Delta)^{-N}, \quad (1 - \Delta)^{-N} M_f(1 - \Delta)^{-N}$$

and their trace ideal properties.

Why is this?

Among other things, so we can use Birman's theorem to study the scattering of differential operators on G .

A first result

One not-entirely-trivial results we obtained is the following.

Theorem (M.-Sukochev-Zanin)

Let $r > Q$ (recall that Q is the homogeneous dimension) and let $q > 2$. Given $f \in \ell_1(L_q)(G)$ (*a function space on G*), the operator

$$M_f(1 - \Delta)^{-\frac{r}{2}} : L_2(G) \rightarrow L_2(G)$$

is trace class.

This is a stratified version of an old result of Birman and Solomyak.

Theorem

Let G be a stratified Lie group with stratification $\mathfrak{g} = \bigoplus_{n=1}^{\infty} \mathfrak{g}_n$, homogeneous dimension $d_{\text{hom}} = \sum_{n=1}^{\infty} n \cdot \dim(\mathfrak{g}_n)$ and a fixed sub-Laplacian $\Delta = \sum_{j=1}^m X_j^2$, where $\{X_j\}_{j=1}^m$ is a basis for \mathfrak{g}_1 .

i) if $p > 2$, then

$$\|M_f(-\Delta)^{-\frac{d_{\text{hom}}}{2p}}\|_{p,\infty} \leq c_p \|f\|_{L_p(G)}$$

ii) if $p < 2$ and $q > 2$, then

$$\|M_f(1 - \Delta)^{-\frac{d_{\text{hom}}}{2p}}\|_{p,\infty} \leq c_{p,q} \|f\|_{\ell_p(L_q)(G)}.$$

iii) if $p = 2$ and $q > 2$, then

$$\|M_f(1 - \Delta)^{-\frac{d_{\text{hom}}}{2p}}\|_{p,\infty} \leq c_q \|f\|_{\ell_2 \log(L_q)(G)}.$$

Birman's theorem for stratified Lie groups

Suppose that

$$D_1 = \sum_{|\alpha|_h \leq m} a_\alpha(x) X^\alpha$$

where each a_α is a smooth function on G equal to a constant (say, c_α) outside a compact set. Then we expect that

$$D_0 = \sum_{|\alpha|_h \leq m} c_\alpha X^\alpha$$

is a good model for D_1 asymptotically, since $D_1 - D_0$ is a differential operator with compactly supported coefficients.

Spectral asymptotics

Related to these estimates we have spectral asymptotics. In the following theorem, μ denotes the singular value function. In particular, the sequence $\{\mu(n, T)\}_{n=0}^{\infty}$ is the sequence of singular values of a compact operator T . We give a precise definition of μ in the next section.

Theorem

Let G be a non-abelian stratified Lie group with stratification $\mathfrak{g} = \bigoplus_{n=1}^{\infty} \mathfrak{g}_n$, homogeneous dimension $d_{\text{hom}} = \sum_{n=1}^{\infty} n \cdot \dim(\mathfrak{g}_n)$ and a fixed sub-Laplacian $\Delta = \sum_{j=1}^m X_j^2$, where $\{X_j\}_{j=1}^m$ is a basis for \mathfrak{g}_1 . Let $k \in \mathbb{N}$ and let $p = \frac{d_{\text{hom}}}{k}$. If one of the following conditions holds

- ❶ $p > 1$ and $0 \leq f \in L_p(G)$;
- ❷ $p < 1$ and $0 \leq f \in \ell_p(L_q)(G)$ for some $q > 1$;
- ❸ $p = 1$ and $0 \leq f \in \ell_{1, \log}(L_q)(G)$ for some $q > 1$;

then there exists the limit

$$\lim_{t \rightarrow 0} t \mu(t, (1 - \Delta)^{-\frac{k}{4}} M_f (1 - \Delta)^{-\frac{k}{4}})^p = c_G \int f^p.$$

Corollary

Let $G \neq \mathbb{H}^1$ be a stratified Lie group with stratification $\mathfrak{g} = \bigoplus_{n=1}^{\infty} \mathfrak{g}_n$, homogeneous dimension $d_{\text{hom}} = \sum_{n=1}^{\infty} n \cdot \dim(\mathfrak{g}_n)$ and a fixed sub-Laplacian $\Delta = \sum_{j=1}^m X_j^2$, where $\{X_j\}_{j=1}^m$ is a basis for \mathfrak{g}_1 . Assume that $V \in L_{\frac{d_{\text{hom}}}{2}}(G)$ is real-valued. For $h > 0$, the operator $-h^2\Delta + M_V$ can be defined in the sense of quadratic forms. There exists a constant $c_G > 0$ such that

$$\lim_{h \rightarrow 0} h^{d_{\text{hom}}} \text{Tr}(\chi_{(-\infty, 0)}(-h^2\Delta + M_V)) = c_G \int_G V_-^{\frac{d_{\text{hom}}}{2}}.$$

Here, $V_- = \frac{1}{2}(|V| - V)$ is the negative part of V .

Theorem (Cwikel–Lieb–Rozenblum estimate)

Assume that $d_{\text{hom}} > 2$. Let $V \in L_{\frac{d_{\text{hom}}}{2}}(G)$ be real-valued. The quadratic form sum

$$-\Delta \dot{+} M_V$$

is well-defined on the form domain $W_2^1(G)$, and defines an unbounded self-adjoint operator on $L_2(G)$ with essential spectrum $[0, \infty)$. The operator $-\Delta \dot{+} M_V$ has finitely many negative eigenvalues, and the total number of eigenvalues less than $-t$ for $t \geq 0$ is bounded by

$$\text{Tr}(\chi_{(-\infty, -t)}(-\Delta \dot{+} M_V)) \leq C_G \int_G (V + t)_-^{\frac{d_{\text{hom}}}{2}}.$$

Theorem

Let G be a stratified Lie group with $d_{\text{hom}} > 4$, and let $f \in (L_{\frac{d_{\text{hom}}}{2}} \cap L_1)(G)$ be real valued. The wave operators exist and are complete for the couple $(-\Delta, -\Delta \dot{+} M_f)$, where $-\Delta \dot{+} M_f$ denotes the quadratic form sum.

We can make similar assertions about pairs of the form (D_1, D_0) , where D_1 and D_0 are maximally hypoelliptic differential operators whose coefficients are asymptotically equal.

These estimates are suboptimal for a number of reasons, one of them being that we state the results for functions on G rather than a general Heisenberg manifold (or an even more general filtered manifold). This is probably not a significant restriction.

Thank you for listening!