

# Lipschitz estimates in quasi-Banach Schatten ideals

Edward McDonald (Penn State University)

George Mason University, January 2025

January 28, 2025

This talk is mostly about the paper

M., Sukochev, Lipschitz estimates in quasi-Banach Schatten ideals. *Math. Ann.* 383 (2022), no.1–2, 571–619.

# Plan for this talk

- ① Operator Lipschitz functions: some basic concepts and history.
- ② Some very light background on Schatten ideals: the problem with  $0 < p < 1$ .
- ③ Schur multipliers in  $0 < p < 1$ .
- ④ Besov spaces and wavelets
- ⑤ Future directions

# Operator Lipschitz functions

Let  $H$  be a (complex and separable) Hilbert space, and denote the operator norm by  $\|\cdot\|_\infty$ . A function  $f : \mathbb{R} \rightarrow \mathbb{C}$  is said to be *operator Lipschitz* if there exists a constant  $C_f$  such that

$$\|f(A) - f(B)\|_\infty \leq C_f \|A - B\|_\infty, \quad A, B \in \mathcal{B}_{\text{sa}}(H)$$

## Question (from Krein)

Is every Lipschitz function operator Lipschitz?

# Operator Lipschitz functions

Let  $H$  be a (complex and separable) Hilbert space, and denote the operator norm by  $\|\cdot\|_\infty$ . A function  $f : \mathbb{R} \rightarrow \mathbb{C}$  is said to be *operator Lipschitz* if there exists a constant  $C_f$  such that

$$\|f(A) - f(B)\|_\infty \leq C_f \|A - B\|_\infty, \quad A, B \in \mathcal{B}_{\text{sa}}(H)$$

## Question (from Krein)

Is every Lipschitz function operator Lipschitz? That is, does  $|f(t) - f(s)| \lesssim |t - s|$  imply that  $\|f(A) - f(B)\|_\infty \lesssim \|A - B\|_\infty$ ?

# Operator Lipschitz functions

Answer

No.

# Operator Lipschitz functions

## Answer

No.

Farforovskaya (1968): There exist Lipschitz functions that are not operator Lipschitz

# Operator Lipschitz functions

## Answer

No.

Farforovskaya (1968): There exist Lipschitz functions that are not operator Lipschitz

Kato (1973): The absolute value function  $f(t) = |t|$  is not operator Lipschitz



# Operator Lipschitz functions

## Answer

No.

Farforovskaya (1968): There exist Lipschitz functions that are not operator Lipschitz

Kato (1973): The absolute value function  $f(t) = |t|$  is not operator Lipschitz

Johnson & Williams (1975): An operator Lipschitz function is differentiable.

If  $H$  is  $N$ -dimensional, then

$$\|f(A) - f(B)\|_{\infty} \leq C_{\text{abs}} \log(1 + N) \|f\|_{\text{Lip}} \|A - B\|_{\infty}$$

where  $C_{\text{abs}}$  is an absolute constant. This is sharp in the order of growth as  $N \rightarrow \infty$ . I do not know if a sharp estimate for  $C_{\text{abs}}$  is known.

# Operator Lipschitz functions

It is easy to check that sufficiently good functions are operator Lipschitz. Let's check the function  $f(x) = e^{i\xi x}$  for  $\xi \in \mathbb{R}$ . We have

$$e^{i\xi A} - e^{i\xi B} = i\xi \int_0^1 e^{i\xi(1-\theta)A} (A - B) e^{i\xi\theta B} d\theta.$$

The integral converges in the Bochner sense. The triangle inequality implies

$$\|e^{i\xi A} - e^{i\xi B}\|_\infty \leq |\xi| \|A - B\|_\infty.$$

# Operator Lipschitz functions

It is easy to check that sufficiently good functions are operator Lipschitz. Let's check the function  $f(x) = e^{i\xi x}$  for  $\xi \in \mathbb{R}$ . We have

$$e^{i\xi A} - e^{i\xi B} = i\xi \int_0^1 e^{i\xi(1-\theta)A} (A - B) e^{i\xi\theta B} d\theta.$$

The integral converges in the Bochner sense. The triangle inequality implies

$$\|e^{i\xi A} - e^{i\xi B}\|_\infty \leq |\xi| \|A - B\|_\infty.$$

By Fourier inversion,

$$\|f(A) - f(B)\|_\infty \leq \|A - B\|_\infty \cdot 2\pi \|\widehat{\partial f}\|_1.$$

# Operator Lipschitz functions

It is easy to check that sufficiently good functions are operator Lipschitz. Let's check the function  $f(x) = e^{i\xi x}$  for  $\xi \in \mathbb{R}$ . We have

$$e^{i\xi A} - e^{i\xi B} = i\xi \int_0^1 e^{i\xi(1-\theta)A} (A - B) e^{i\xi\theta B} d\theta.$$

The integral converges in the Bochner sense. The triangle inequality implies

$$\|e^{i\xi A} - e^{i\xi B}\|_\infty \leq |\xi| \|A - B\|_\infty.$$

By Fourier inversion,

$$\|f(A) - f(B)\|_\infty \leq \|A - B\|_\infty \cdot 2\pi \|\widehat{\partial f}\|_1.$$

By Cauchy-Schwarz,  $\|\widehat{\partial f}\|_1 \leq \|f'\|_2 + \|f''\|_2$ . This is a “good enough” sufficient condition for most purposes.

# Peller's theorem

The previous computation was based on Fourier inversion of  $f$  and a description of  $e^{i\xi A} - e^{i\xi B}$  as an integral (Duhamel's integral).

# Peller's theorem

The previous computation was based on Fourier inversion of  $f$  and a description of  $e^{i\xi A} - e^{i\xi B}$  as an integral (Duhamel's integral). Using a more subtle description of  $e^{i\xi A} - e^{i\xi B}$ , and handling the Littlewood-Paley components of  $f$  individually, V. V. Peller has proved the following:

## Theorem (Peller (1990))

*If  $f$  is Lipschitz and belongs to the homogeneous Besov class  $\dot{B}_{\infty,1}^1(\mathbb{R})$  then  $f$  is operator Lipschitz.*

# Peller's theorem

The previous computation was based on Fourier inversion of  $f$  and a description of  $e^{i\xi A} - e^{i\xi B}$  as an integral (Duhamel's integral). Using a more subtle description of  $e^{i\xi A} - e^{i\xi B}$ , and handling the Littlewood-Paley components of  $f$  individually, V. V. Peller has proved the following:

## Theorem (Peller (1990))

*If  $f$  is Lipschitz and belongs to the homogeneous Besov class  $\dot{B}_{\infty,1}^1(\mathbb{R})$  then  $f$  is operator Lipschitz.*

In other words, if  $f$  is Lipschitz and

$$\int_0^\infty \sup_{t \in \mathbb{R}} \frac{|f(t-h) - 2f(t) + f(t+h)|}{h^2} dh < \infty$$

then  $f$  is operator Lipschitz.



# Peller's theorem

The previous computation was based on Fourier inversion of  $f$  and a description of  $e^{i\xi A} - e^{i\xi B}$  as an integral (Duhamel's integral). Using a more subtle description of  $e^{i\xi A} - e^{i\xi B}$ , and handling the Littlewood-Paley components of  $f$  individually, V. V. Peller has proved the following:

## Theorem (Peller (1990))

*If  $f$  is Lipschitz and belongs to the homogeneous Besov class  $\dot{B}_{\infty,1}^1(\mathbb{R})$  then  $f$  is operator Lipschitz.*

In other words, if  $f$  is Lipschitz and

$$\int_0^\infty \sup_{t \in \mathbb{R}} \frac{|f(t-h) - 2f(t) + f(t+h)|}{h^2} dh < \infty$$

then  $f$  is operator Lipschitz. For example, if  $f', f'' \in L_\infty(\mathbb{R})$  then  $f$  is operator Lipschitz.

# Peller's operator Bernstein inequality

The classical Bernstein inequality states that if  $f \in L_\infty(\mathbb{R})$  has Fourier transform supported in the interval  $[-\sigma, \sigma]$ , then

$$\|f\|_{\text{Lip}} \leq C\sigma \|f\|_\infty.$$

# Peller's operator Bernstein inequality

The classical Bernstein inequality states that if  $f \in L_\infty(\mathbb{R})$  has Fourier transform supported in the interval  $[-\sigma, \sigma]$ , then

$$\|f\|_{\text{Lip}} \leq C\sigma \|f\|_\infty.$$

Peller's theorem is a consequence of his *operator Bernstein inequality*.

## Theorem (Peller (1990))

If  $f \in L_\infty(\mathbb{R})$  has Fourier transform supported in the interval  $[-\sigma, \sigma]$ , then

$$\|f\|_{\text{O-Lip}} \leq C\sigma \|f\|_\infty.$$

Here  $\|f\|_{\text{O-Lip}}$  is the operator Lipschitz seminorm, i.e.

$$\|f\|_{\text{O-Lip}} := \sup_{A=A^*, B=B^* \in \mathcal{B}(H)} \frac{\|f(A) - f(B)\|_\infty}{\|A - B\|_\infty}.$$

If  $T$  is a compact operator on  $H$ , the singular value sequence of  $T$  is defined as

$$\mu(k, T) := \inf\{\|T - R\|_\infty : \text{rank}(R) \leq k\}, \quad k \geq 0.$$

(Equivalently,  $\mu(T) = \{\mu(k, T)\}_{k=0}^\infty$  is the sequence of eigenvalues of the absolute value  $|T|$  arranged in non-increasing order with multiplicities.)

If  $T$  is a compact operator on  $H$ , the singular value sequence of  $T$  is defined as

$$\mu(k, T) := \inf\{\|T - R\|_\infty : \text{rank}(R) \leq k\}, \quad k \geq 0.$$

(Equivalently,  $\mu(T) = \{\mu(k, T)\}_{k=0}^\infty$  is the sequence of eigenvalues of the absolute value  $|T|$  arranged in non-increasing order with multiplicities.)

Note that  $\|T\|_\infty = \mu(0, T) = \|\mu(T)\|_{\ell_\infty}$ .

If  $T$  is a compact operator on  $H$ , the singular value sequence of  $T$  is defined as

$$\mu(k, T) := \inf\{\|T - R\|_\infty : \text{rank}(R) \leq k\}, \quad k \geq 0.$$

(Equivalently,  $\mu(T) = \{\mu(k, T)\}_{k=0}^\infty$  is the sequence of eigenvalues of the absolute value  $|T|$  arranged in non-increasing order with multiplicities.)

Note that  $\|T\|_\infty = \mu(0, T) = \|\mu(T)\|_{\ell_\infty}$ . For  $1 \leq p < \infty$ , the Schatten  $\mathcal{L}_p$ -norm of a compact operator  $T$  is

$$\|T\|_p := \|\mu(T)\|_{\ell_p} = \left( \sum_{k=0}^{\infty} \mu(k, T)^p \right)^{\frac{1}{p}}.$$

Equivalently,  $\|T\|_p = \text{Tr}(|T|^p)^{1/p}$ . It is not obvious, but this is a norm (i.e.  $\|T + S\|_p \leq \|T\|_p + \|S\|_p$ .)

## $\mathcal{L}_p$ -operator Lipschitz functions

A function  $f$  on  $\mathbb{R}$  is said to be  $\mathcal{L}_p$ -operator Lipschitz if there exists a constant  $C_f > 0$  such that

$$\|f(A) - f(B)\|_p \leq C_f \|A - B\|_p, \quad A, B \in \mathcal{B}_{\text{sa}}(H).$$

By a duality argument,  $\mathcal{L}_1$ -operator Lipschitz is the same thing as operator Lipschitz.

## $\mathcal{L}_p$ -operator Lipschitz functions

A function  $f$  on  $\mathbb{R}$  is said to be  $\mathcal{L}_p$ -operator Lipschitz if there exists a constant  $C_f > 0$  such that

$$\|f(A) - f(B)\|_p \leq C_f \|A - B\|_p, \quad A, B \in \mathcal{B}_{\text{sa}}(H).$$

By a duality argument,  $\mathcal{L}_1$ -operator Lipschitz is the same thing as operator Lipschitz.

What about  $1 < p < \infty$ ?



# $\mathcal{L}_p$ -operator Lipschitz functions

A function  $f$  on  $\mathbb{R}$  is said to be  $\mathcal{L}_p$ -operator Lipschitz if there exists a constant  $C_f > 0$  such that

$$\|f(A) - f(B)\|_p \leq C_f \|A - B\|_p, \quad A, B \in \mathcal{B}_{\text{sa}}(H).$$

By a duality argument,  $\mathcal{L}_1$ -operator Lipschitz is the same thing as operator Lipschitz.

What about  $1 < p < \infty$ ?

**Theorem (Potapov and Sukochev (2010))**

*For  $1 < p < \infty$ , all Lipschitz functions are  $\mathcal{L}_p$ -operator Lipschitz.*

# $\mathcal{L}_p$ -operator Lipschitz functions

A function  $f$  on  $\mathbb{R}$  is said to be  $\mathcal{L}_p$ -operator Lipschitz if there exists a constant  $C_f > 0$  such that

$$\|f(A) - f(B)\|_p \leq C_f \|A - B\|_p, \quad A, B \in \mathcal{B}_{\text{sa}}(H).$$

By a duality argument,  $\mathcal{L}_1$ -operator Lipschitz is the same thing as operator Lipschitz.

What about  $1 < p < \infty$ ?

**Theorem (Potapov and Sukochev (2010))**

*For  $1 < p < \infty$ , all Lipschitz functions are  $\mathcal{L}_p$ -operator Lipschitz.*

For  $p = 2$  this is almost trivial and has been known for approx. 110 years.  
For  $p \neq 2$ , this requires some very deep harmonic analysis.

# $\mathcal{L}_p$ -operator Lipschitz functions

A function  $f$  on  $\mathbb{R}$  is said to be  $\mathcal{L}_p$ -operator Lipschitz if there exists a constant  $C_f > 0$  such that

$$\|f(A) - f(B)\|_p \leq C_f \|A - B\|_p, \quad A, B \in \mathcal{B}_{\text{sa}}(H).$$

By a duality argument,  $\mathcal{L}_1$ -operator Lipschitz is the same thing as operator Lipschitz.

What about  $1 < p < \infty$ ?

**Theorem (Potapov and Sukochev (2010))**

*For  $1 < p < \infty$ , all Lipschitz functions are  $\mathcal{L}_p$ -operator Lipschitz.*

For  $p = 2$  this is almost trivial and has been known for approx. 110 years. For  $p \neq 2$ , this requires some very deep harmonic analysis. There are now essentially four proofs of Potapov-Sukochev. The most recent is due to Conde-Alonso, González-Pérez, Parcet and Tablate and uses operator-valued harmonic analysis.

# What about $0 < p < 1$ ?

For  $0 < p < 1$ , we can still define

$$\|T\|_p := \|\mu(T)\|_{\ell_p} = \text{Tr}(|T|^p)^{\frac{1}{p}}.$$

This is not a norm, merely a quasi-norm. There is no triangle inequality, merely a quasi-triangle inequality

$$\|T + S\|_p \leq 2^{\frac{1}{p}-1}(\|T\|_p + \|S\|_p).$$

# What about $0 < p < 1$ ?

For  $0 < p < 1$ , we can still define

$$\|T\|_p := \|\mu(T)\|_{\ell_p} = \text{Tr}(|T|^p)^{\frac{1}{p}}.$$

This is not a norm, merely a quasi-norm. There is no triangle inequality, merely a quasi-triangle inequality

$$\|T + S\|_p \leq 2^{\frac{1}{p}-1}(\|T\|_p + \|S\|_p).$$

Nonetheless, we have

$$\|T + S\|_p^p \leq \|T\|_p^p + \|S\|_p^p.$$

# Geometry in $\mathcal{L}_p$ .

The unit ball  $B = \{T : \|T\|_p \leq 1\}$  in  $\mathcal{L}_p$  is not convex.  
i.e., if  $\xi_1, \dots, \xi_n \in B$  then it might happen that

$$\theta_1 \xi_1 + \dots + \theta_n \xi_n \notin B, \quad |\theta_1| + \dots + |\theta_n| \leq 1.$$

For this reason the theory of integration  $\mathcal{L}_p$ -valued functions is not straightforward. We could have continuous functions  $f \in C([0, 1], \mathcal{L}_p)$  whose integral is not in  $\mathcal{L}_p$ .

# Geometry in $\mathcal{L}_p$ .

The unit ball  $B = \{T : \|T\|_p \leq 1\}$  in  $\mathcal{L}_p$  is not convex.  
i.e., if  $\xi_1, \dots, \xi_n \in B$  then it might happen that

$$\theta_1 \xi_1 + \dots + \theta_n \xi_n \notin B, \quad |\theta_1| + \dots + |\theta_n| \leq 1.$$

For this reason the theory of integration  $\mathcal{L}_p$ -valued functions is not straightforward. We could have continuous functions  $f \in C([0, 1], \mathcal{L}_p)$  whose integral is not in  $\mathcal{L}_p$ .

Instead,  $B$  is only closed under  $p$ -convex combinations, i.e.

$$\theta_1 \xi_1 + \dots + \theta_n \xi_n \in B, \quad |\theta_1|^p + \dots + |\theta_n|^p \leq 1.$$

## $\mathcal{L}_p$ -Lipschitz functions for $0 < p < 1$ .

Which functions are Lipschitz in  $\mathcal{L}_p$  when  $0 < p < 1$ ?



## $\mathcal{L}_p$ -Lipschitz functions for $0 < p < 1$ .

Which functions are Lipschitz in  $\mathcal{L}_p$  when  $0 < p < 1$ ?

At least some functions are, for example  $f(t) = (t + \lambda)^{-1}$ ,  $\lambda \in \mathbb{C} \setminus \mathbb{R}$ .

# $\mathcal{L}_p$ -Lipschitz functions for $0 < p < 1$ .

Which functions are Lipschitz in  $\mathcal{L}_p$  when  $0 < p < 1$ ?

At least some functions are, for example  $f(t) = (t + \lambda)^{-1}$ ,  $\lambda \in \mathbb{C} \setminus \mathbb{R}$ .

What about  $f(t) = \exp(it\xi)$  for  $\xi \in \mathbb{R}$ ?

# Schur multipliers

The usual method to prove  $\mathcal{L}_p$ -estimates in  $0 < p < 1$  is the same as for  $p \geq 1$ , to use Schur multipliers.

If  $A = \{A_{j,k}\}_{j,k}$  and  $B = \{B_{j,k}\}_{j,k}$  are matrices of the same size, then  $A \circ B := \{A_{j,k} B_{j,k}\}_{j,k}$ .

## Definition

Let  $m = \{m_{j,k}\}_{j,k=1}^n$  be an  $n \times n$  matrix. Define

$$\|m\|_{\mathfrak{m}_p} := \sup_{\|B\|_p \leq 1} \|m \circ B\|_p.$$

In general, let  $m : \mathbb{R}^2 \rightarrow \mathbb{C}$  be bounded and define

$$\|m\|_{\mathfrak{m}_p} := \sup_{x_1, \dots, x_n, y_1, \dots, y_n \in \mathbb{R}} \|\{m(x_j, y_k)\}_{j,k=1}^n\|_{\mathfrak{m}_p}.$$

# Schur multipliers and operator-Lipschitz functions

A folk result:

Theorem (Hadamard(?), Schur(?), Löwner(?), Daletskii-Krein(?))

Let  $1 \leq p \leq \infty$ , and let  $f$  be a measurable function on  $\mathbb{R}$ . Define

$$f^{[1]}(\lambda, \mu) = \begin{cases} \frac{f(\lambda) - f(\mu)}{\lambda - \mu}, & \lambda \neq \mu \\ f'(\lambda), & \lambda = \mu. \end{cases}$$

Then  $f$  is  $\mathcal{L}_p$ -Lipschitz if and only if  $f^{[1]}$  is a bounded Schur multiplier.

# Schur multipliers and operator-Lipschitz functions

A folk result:

Theorem (Hadamard(?), Schur(?), Löwner(?), Daletskii-Krein(?))

Let  $1 \leq p \leq \infty$ , and let  $f$  be a measurable function on  $\mathbb{R}$ . Define

$$f^{[1]}(\lambda, \mu) = \begin{cases} \frac{f(\lambda) - f(\mu)}{\lambda - \mu}, & \lambda \neq \mu \\ f'(\lambda), & \lambda = \mu. \end{cases}$$

Then  $f$  is  $\mathcal{L}_p$ -Lipschitz if and only if  $f^{[1]}$  is a bounded Schur multiplier.

If  $f$  is not differentiable then  $f'(\lambda)$  does not make sense everywhere, but this is no big deal. Bounded diagonal matrices are Schur multipliers in  $\mathcal{L}_p$  for any  $1 \leq p \leq \infty$  so we can make  $f^{[1]}(\lambda, \lambda)$  any bounded function of  $\lambda$  without changing the result.

# Schur multipliers in $\mathcal{L}_p$

One noteworthy difference between  $p = 1$  and  $p < 1$  is the following example:

## Example

Let  $I_n = \{\delta_{j,k}\}_{j,k=0}^{n-1}$  be the  $n \times n$  identity matrix. Then

$$\|I_n\|_{\mathfrak{M}_p} = n^{\frac{1}{p}-1}.$$

To see this, compute  $I_n \circ (\xi \otimes \xi)$  where  $\xi = (1, \dots, 1)$ .

# Schur multipliers in $\mathcal{L}_p$

One noteworthy difference between  $p = 1$  and  $p < 1$  is the following example:

## Example

Let  $I_n = \{\delta_{j,k}\}_{j,k=0}^{n-1}$  be the  $n \times n$  identity matrix. Then

$$\|I_n\|_{\mathfrak{M}_p} = n^{\frac{1}{p}-1}.$$

To see this, compute  $I_n \circ (\xi \otimes \xi)$  where  $\xi = (1, \dots, 1)$ .

What this means is that restriction to the diagonal

$$\{A_{j,k}\}_{j,k \geq 0} \mapsto \{A_{j,j}\delta_{j,k}\}_{j,k=0}^{\infty}$$

is not bounded in  $\mathcal{L}_p$  for any  $0 < p < 1$ !

# Schur multipliers and operator-Lipschitz functions in $\mathcal{L}_p$ for $0 < p < 1$ .

## Theorem

Let  $0 < p \leq \infty$ . and let  $f$  be a measurable function on  $\mathbb{R}$ . Then  $f$  is  $\mathcal{L}_p$ -Lipschitz if and only if

$$\sup_{\{\lambda_j\}, \{\mu_j\}} \|\{f^{[1]}(\lambda_j, \mu_k)\}_{j,k}\|_{\mathfrak{m}_p} < \infty$$

where the supremum is over all disjoint sequences  $\{\lambda_j\}_j$  and  $\{\mu_k\}_k$ .

Since we only consider disjoint sequences, the diagonal does not enter the picture.



# An example

Consider the function

$$f(x) = \sin(x)$$

and  $\mu_j = \lambda_j = 2\pi j$ . Then

$$f^{[1]}(\lambda_j, \mu_k) = \delta_{j,k}.$$

But this is not a Schur multiplier of  $\mathcal{L}_p$ ! The same reasoning applies to any periodic function  $f$  with  $f'(\lambda) \neq 0$  for some  $\lambda$ .

# An example

Consider the function

$$f(x) = \sin(x)$$

and  $\mu_j = \lambda_j = 2\pi j$ . Then

$$f^{[1]}(\lambda_j, \mu_k) = \delta_{j,k}.$$

But this is not a Schur multiplier of  $\mathcal{L}_p$ ! The same reasoning applies to any periodic function  $f$  with  $f'(\lambda) \neq 0$  for some  $\lambda$ . This argument is obviously flawed, we are supposed to consider *disjoint* sequences.

## A corrected example

We can fix the previous (wrong) argument by shifting one of the sequences by  $\varepsilon > 0$ . Let  $f$  be a 1-periodic function. Consider the sequences

$$\lambda_j = j + \varepsilon, \mu_k = k, \quad j, k \geq 0.$$

Then

$$f^{[1]}(\lambda_j, \mu_k) = \frac{f(j + \varepsilon) - f(k)}{j - k + \varepsilon} = (f(\varepsilon) - f(0)) \frac{1}{j - k + \varepsilon}, \quad j, k \geq 0.$$

If  $f(\varepsilon) \neq f(0)$ , then we need to consider the matrix

$$\{(j - k + \varepsilon)^{-1}\}_{j, k \geq 0}.$$

This matrix is not diagonal, but a straightforward modification of the argument for  $\{\delta_{j,k}\}_{j,k \geq 0}$  shows that it is not a Schur multiplier of  $\mathcal{L}_p$  for any  $0 < p < 1$ .

## A corrected example

We can fix the previous (wrong) argument by shifting one of the sequences by  $\varepsilon > 0$ . Let  $f$  be a 1-periodic function. Consider the sequences

$$\lambda_j = j + \varepsilon, \mu_k = k, \quad j, k \geq 0.$$

Then

$$f^{[1]}(\lambda_j, \mu_k) = \frac{f(j + \varepsilon) - f(k)}{j - k + \varepsilon} = (f(\varepsilon) - f(0)) \frac{1}{j - k + \varepsilon}, \quad j, k \geq 0.$$

If  $f(\varepsilon) \neq f(0)$ , then we need to consider the matrix

$$\{(j - k + \varepsilon)^{-1}\}_{j, k \geq 0}.$$

This matrix is not diagonal, but a straightforward modification of the argument for  $\{\delta_{j, k}\}_{j, k \geq 0}$  shows that it is not a Schur multiplier of  $\mathcal{L}_p$  for any  $0 < p < 1$ . Aleksandrov and Peller have characterised Schur multipliers of  $\mathcal{L}_p$  of the Herz-Toeplitz form  $m(j - k)$ , so we could also use their result.

Periodic functions are not  $\mathcal{L}_p$ -Lipschitz for  $0 < p < 1$ .

Summarising the preceding reasoning:

**Lemma (M. and Sukochev (2022))**

*Let  $0 < p < 1$ , and let  $f$  be a periodic function on  $\mathbb{R}$ . Then  $f$  is  $\mathcal{L}_p$ -Lipschitz if and only if it is constant.*

What does this imply?

Periodic functions are not  $\mathcal{L}_p$ -Lipschitz for  $0 < p < 1$ .

Summarising the preceding reasoning:

**Lemma (M. and Sukochev (2022))**

*Let  $0 < p < 1$ , and let  $f$  be a periodic function on  $\mathbb{R}$ . Then  $f$  is  $\mathcal{L}_p$ -Lipschitz if and only if it is constant.*

What does this imply?

- Even  $C^\infty$  functions with all derivatives bounded may not be  $\mathcal{L}_p$ -Lipschitz;
- In particular  $f(t) = \exp(it\xi)$ ,  $\xi \neq 0$  is not  $\mathcal{L}_p$ -Lipschitz for any  $0 < p < 1$ . This means that methods based on a Fourier decomposition are unlikely to work.

# An idea

Consider the matrix

$$\left\{ \frac{c_j - c_k}{j - k + \varepsilon} \right\}_{j,k \geq 0} \quad (0.1)$$

where  $c_j$  is a scalar sequence. This is approximately a model for  $f^{[1]}(j + \varepsilon, k)$  where  $f$  is a function of the form

$$f(x) = \sum_{k \in \mathbb{Z}} c_k \psi(x - k)$$

where  $\psi$  is some bump function.

## Lemma

*If  $\sum_j |c_j|^{\frac{p}{1-p}} < \infty$  then (0.1) is a Schur multiplier of  $\mathcal{L}_p$ .*

# An idea

Consider the matrix

$$\left\{ \frac{c_j - c_k}{j - k + \varepsilon} \right\}_{j,k \geq 0} \quad (0.1)$$

where  $c_j$  is a scalar sequence. This is approximately a model for  $f^{[1]}(j + \varepsilon, k)$  where  $f$  is a function of the form

$$f(x) = \sum_{k \in \mathbb{Z}} c_k \psi(x - k)$$

where  $\psi$  is some bump function.

## Lemma

*If  $\sum_j |c_j|^{\frac{p}{1-p}} < \infty$  then (0.1) is a Schur multiplier of  $\mathcal{L}_p$ .*

Why is it  $\frac{p}{1-p}$ ? It comes down to the inequality

$$\left( \sum_j |a_j b_j|^p \right)^{\frac{1}{p}} \leq \left( \sum_j |a_j|^{\frac{p}{1-p}} \right)^{\frac{1-p}{p}} \left( \sum_j |b_j| \right)$$



# Sums of shifted bump functions

With considerably more effort, it is possible to prove the following:

## Theorem

Let  $\psi \in C_c^k(\mathbb{R})$ , where  $k > \frac{2}{p} - 1$ . If  $\{c_k\}_{k \in \mathbb{Z}}$  is some scalar sequence, and

$$f(x) = \sum_{k \in \mathbb{Z}} c_k \psi(x - k)$$

then

$$\|f^{[1]}\|_{m_p} \lesssim \|\{c_k\}_{k \in \mathbb{Z}}\|_{\ell_{\frac{p}{1-p}}}.$$

What kind of functions can we build out of functions like this?

What is a good way of approximating a general function from compactly supported  $C^k$ -functions?

What is a good way of approximating a general function from compactly supported  $C^k$ -functions?

## Theorem (Daubechies (1988))

*For all  $k > 0$ , there exists a compactly supported  $C^k$  function  $\psi$  such that the system of translations and dilations*

$$\psi_{j,k}(t) := 2^{\frac{j}{2}} \psi(2^j t - k), \quad j, k \in \mathbb{Z}$$

*forms an orthonormal basis of  $L_2(\mathbb{R})$ .*

Wavelets are analogous to Fourier series, in the sense that if

$$f(t) = \sum_{j \in \mathbb{Z}} \sum_{k \in \mathbb{Z}} c_{j,k} \psi_{j,k}(t)$$

then the coefficients  $c_{j,k}$  for  $j > N$  represent oscillations of  $f$  on the scale  $\sim 2^{-N}$ . A function of the form

$$f(t) = \sum_{j < N} \sum_{k \in \mathbb{Z}} c_{j,k} \psi_{j,k}(t)$$

does not oscillate greatly on scales smaller than  $2^{-N}$ . This is similar to functions with Fourier transform supported in  $[-2^N, 2^N]$ .

# An $\mathcal{L}_p$ -Lipschitz Bernstein inequality

## Theorem (M.-Sukochev (2022))

Let  $f \in L_{\frac{p}{1-p}}(\mathbb{R})$  have Wavelet expansion

$$f(t) = \sum_{j \in \mathbb{Z}} \sum_{k \in \mathbb{Z}} c_{j,k} \psi_{j,k}(t)$$

where  $c_{j,k} = 0$  for  $k > N$ . Then

$$\|f^{[1]}\|_{\mathfrak{m}_p} \leq C 2^{\frac{N}{p}} \|f\|_{\frac{p}{1-p}}.$$

With  $p = 1$ , this is the wavelet analogy of Peller's operator Bernstein inequality. For  $p < 1$  it is new.

It follows from the Wavelet Bernstein inequality that Besov spaces have a very simple characterisation in terms of wavelet coefficients.

## Theorem (Meyer (1986))

*Let  $s \in \mathbb{R}$  and  $p, q \in (0, \infty]$ . Let  $\psi$  be a compactly supported  $C^k$  wavelet where  $k > -s$ . Then a distribution  $f \in \mathcal{D}'(\mathbb{R})$  belongs to the homogeneous Besov space  $\dot{B}_{p,q}^s(\mathbb{R})$  if and only if*

$$\|f\|_{\dot{B}_{p,q}^s} \approx \sum_{j \in \mathbb{Z}} 2^{jq(s + \frac{1}{2} - \frac{1}{p})} \left( \sum_{k \in \mathbb{Z}} |\langle f, \psi_{j,k} \rangle|^p \right)^{\frac{q}{p}} < \infty.$$

# A new result

Using the  $p$ -triangle inequality and the  $\mathcal{L}_p$ -Lipschitz Bernstein inequality, we get the following:

## Theorem (M. and Sukochev (2022))

Let  $0 < p < 1$ . Let  $f \in \dot{B}_{\frac{p}{1-p}, p}^{\frac{1}{p}}(\mathbb{R})$  be Lipschitz continuous. Then  $f$  is  $\mathcal{L}_p$ -Lipschitz and

$$\|f(A) - f(B)\|_p \leq C_p(\|f'\|_\infty + \|f\|_{\dot{B}_{\frac{p}{1-p}, p}^{\frac{1}{p}}(\mathbb{R})})\|A - B\|_p, \quad A, B \in \mathcal{B}_{\text{sa}}(H).$$

# A new result

Using the  $p$ -triangle inequality and the  $\mathcal{L}_p$ -Lipschitz Bernstein inequality, we get the following:

## Theorem (M. and Sukochev (2022))

Let  $0 < p < 1$ . Let  $f \in \dot{B}_{\frac{p}{1-p}, p}^{\frac{1}{p}}(\mathbb{R})$  be Lipschitz continuous. Then  $f$  is  $\mathcal{L}_p$ -Lipschitz and

$$\|f(A) - f(B)\|_p \leq C_p(\|f'\|_\infty + \|f\|_{\dot{B}_{\frac{p}{1-p}, p}^{\frac{1}{p}}(\mathbb{R})})\|A - B\|_p, \quad A, B \in \mathcal{B}_{\text{sa}}(H).$$

In other words, we require that  $f$  be Lipschitz and for some  $n > \frac{1}{p}$  that

$$\int_0^\infty \left( \int_{-\infty}^\infty \left| \sum_{k=0}^n \binom{n}{k} (-1)^{n-k} f(t + kh) \right|^{\frac{p}{1-p}} dt \right)^{1-p} \frac{dh}{h^2} < \infty.$$



# A new result

Using the  $p$ -triangle inequality and the  $\mathcal{L}_p$ -Lipschitz Bernstein inequality, we get the following:

## Theorem (M. and Sukochev (2022))

Let  $0 < p < 1$ . Let  $f \in \dot{B}_{\frac{p}{1-p}, p}^{\frac{1}{p}}(\mathbb{R})$  be Lipschitz continuous. Then  $f$  is  $\mathcal{L}_p$ -Lipschitz and

$$\|f(A) - f(B)\|_p \leq C_p(\|f'\|_\infty + \|f\|_{\dot{B}_{\frac{p}{1-p}, p}^{\frac{1}{p}}(\mathbb{R})})\|A - B\|_p, \quad A, B \in \mathcal{B}_{\text{sa}}(H).$$

In other words, we require that  $f$  be Lipschitz and for some  $n > \frac{1}{p}$  that

$$\int_0^\infty \left( \int_{-\infty}^\infty \left| \sum_{k=0}^n \binom{n}{k} (-1)^{n-k} f(t + kh) \right|^{\frac{p}{1-p}} dt \right)^{1-p} \frac{dh}{h^2} < \infty.$$

For example,  $f', \dots, f^{(k)} \in L_{\frac{p}{1-p}}(\mathbb{R})$  where  $k > \frac{1}{p} - 1$  is sufficient.

Thank you for listening!