

# Progress and Questions in QHA

Edward McDonald (Penn State University)

Ghent University

January 12, 2025

# Plan for this talk

In this talk I will go over some of the recent work

# Prologue: Meyer's decomposition

Let  $F \in C^\infty(\mathbb{R})$ . The mapping of functional composition

$$L_\infty(\mathbb{R}^d, \mathbb{R}) \rightarrow L_\infty(\mathbb{R}^d), \quad u \mapsto F(u)$$

is sometimes called the *Nemytskij operator*.

Let  $F \in C^\infty(\mathbb{R})$ . The mapping of functional composition

$$L_\infty(\mathbb{R}^d, \mathbb{R}) \rightarrow L_\infty(\mathbb{R}^d), \quad u \mapsto F(u)$$

is sometimes called the *Nemytskij operator*. **Question:** In terms of function spaces what are the mapping properties of  $u \mapsto F(u)$ ? (This is needed for nonlinear PDE).

How can we study Nemytskij operators from the point of view of harmonic analysis?

# Meyer's decomposition

How can we study Nemytskij operators from the point of view of harmonic analysis?

One method relates Nemytskij operators to pseudodifferential operators via the *Meyer decomposition*.

# Meyer's decomposition

How can we study Nemytskij operators from the point of view of harmonic analysis?

One method relates Nemytskij operators to pseudodifferential operators via the *Meyer decomposition*.

Let  $\{\Delta_j\}_{j=0}^\infty$  be an inhomogeneous Littlewood-Paley decomposition for  $\mathbb{R}^d$ . That is,  $\Delta_j = \Psi_j(-i\nabla)$  where  $\Psi_j$  is supported in the ball of radius  $2^j$  and

$$\sum_{j=0}^{\infty} \Psi_j = 1.$$

Let  $S_n = \sum_{j=0}^n \Delta_j$ .



# Meyer's decomposition (cont.)

Assume initially that  $u = S_n u$  for some  $n \geq 1$ . We have

$$F(u) = F(0) + F(S_n u) - F(0) = F(0) + \sum_{j=0}^{\infty} F(S_j u) - F(S_{j-1} u)$$

where  $S_{-1} u = 0$ . Therefore

$$F(u) = F(0) + \sum_{j=0}^{\infty} \int_0^1 F'(S_{j-1} u + \theta \Delta_j u) \Delta_j u d\theta =: F(0) + m(F, u)u$$

where  $m(F, u)$  is the *paradifferential operator*

$$m(F, u) = \sum_{j=0}^{\infty} \int_0^1 F'(S_{j-1} u + \theta \Delta_j u) \Delta_j d\theta.$$

# Bourdaud's theorem

Some elementary estimates show for any  $u \in L_\infty(\mathbb{R}^d, \mathbb{R})$  and  $F \in C^\infty(\mathbb{R})$ ,  $m(F, u)$  is a pseudodifferential operator in the Hörmander class  $\Psi_{1,1}^0(\mathbb{R}^d)$  (Stein's "forbidden symbols").

## Theorem (Bourdaud (1988))

*If  $T \in \Psi_{1,1}^0(\mathbb{R}^d)$ , then  $T$  is bounded on the Sobolev space  $W_p^s(\mathbb{R}^d)$  for all  $s > 0$  and  $1 < p < \infty$ .*

# Bourdaud's theorem

Some elementary estimates show for any  $u \in L_\infty(\mathbb{R}^d, \mathbb{R})$  and  $F \in C^\infty(\mathbb{R})$ ,  $m(F, u)$  is a pseudodifferential operator in the Hörmander class  $\Psi_{1,1}^0(\mathbb{R}^d)$  (Stein's "forbidden symbols").

## Theorem (Bourdaud (1988))

*If  $T \in \Psi_{1,1}^0(\mathbb{R}^d)$ , then  $T$  is bounded on the Sobolev space  $W_p^s(\mathbb{R}^d)$  for all  $s > 0$  and  $1 < p < \infty$ .*

Conclusion: If  $F(0) = 0$  and  $u \in W_p^s(\mathbb{R}^d, \mathbb{R})$  for some  $s > 0$  and  $1 < p < \infty$  then  $F(u) \in W_p^s(\mathbb{R}^d)$ .

# Section 2: Quantum Harmonic Analysis

# Quantum Euclidean spaces

Let  $d \geq 1$  and let  $\theta$  be a  $d \times d$  antisymmetric real matrix. The quantum Euclidean space is an attempt to make rigorous mathematical sense of a space  $\mathbb{R}_\theta^d$  with coordinates functions

$$\{x_1, \dots, x_d\}$$

which do not commute, but instead

$$x_j x_k - x_k x_j = i\theta_{j,k}.$$

The construction goes by many names (Noncommutative Euclidean space, Quantum Euclidean space, Moyal  $d$ -space, Groenwald space, and others).

# Quantum Euclidean spaces

Let  $d \geq 1$  and let  $\theta$  be a  $d \times d$  antisymmetric real matrix. The quantum Euclidean space is an attempt to make rigorous mathematical sense of a space  $\mathbb{R}_\theta^d$  with coordinates functions

$$\{x_1, \dots, x_d\}$$

which do not commute, but instead

$$x_j x_k - x_k x_j = i\theta_{j,k}.$$

The construction goes by many names (Noncommutative Euclidean space, Quantum Euclidean space, Moyal  $d$ -space, Groenwald space, and others). Much of the literature motivates this from the relation  $[q, p] = i\hbar$  from quantum mechanics.

# Uncertainty principle




Fig. 1.5 Precise  $q$  implying large uncertainty in  $p$ .

(Similar pictures can be found in many physics books. This one is from Müller-Kirsten's "Introduction to Quantum mechanics")

# Definition of $\mathbb{R}_\theta^d$

$\mathbb{R}_\theta^d$  is a space in the sense of noncommutative geometry: we do not define it directly, but instead we define it in terms of its function spaces  $L_\infty(\mathbb{R}_\theta^d)$ ,  $L_2(\mathbb{R}_\theta^d)$ ,  $\mathcal{S}(\mathbb{R}_\theta^d)$ , etc.



## Definition of $\mathbb{R}_\theta^d$

$\mathbb{R}_\theta^d$  is a space in the sense of noncommutative geometry: we do not define it directly, but instead we define it in terms of its function spaces  $L_\infty(\mathbb{R}_\theta^d)$ ,  $L_2(\mathbb{R}_\theta^d)$ ,  $\mathcal{S}(\mathbb{R}_\theta^d)$ , etc. The following definition is expedient:

### Definition

For  $t \in \mathbb{R}^d$ , let  $\lambda_\theta(t)$  be the unitary operator on  $L_2(\mathbb{R}^d)$  given by

$$\lambda_\theta(t)u(s) := e^{i(t,s)}u(s - \frac{1}{2}\theta t), \quad u \in L_2(\mathbb{R}^d).$$

The von Neumann subalgebra of  $\mathcal{B}(L_2(\mathbb{R}^d))$  generated by  $\{\lambda_\theta(t)\}_{t \in \mathbb{R}^d}$  is denoted  $L_\infty(\mathbb{R}_\theta^d)$ .

# Remarks on the definition

- ① Note that if  $\theta = 0$ , this reduces to the description of  $L_\infty(\mathbb{R}^d)$  represented as bounded multipliers on  $L_2(\mathbb{R}^d)$ , and  $\lambda_0$  is the operator of pointwise multiplication by the exponential function  $s \mapsto \exp(i(t, s))$ .
- ② A simple computation shows that

$$\lambda_\theta(t + s) = \exp\left(\frac{1}{2}i(t, \theta s)\right)\lambda_\theta(t)\lambda_\theta(s), \quad t, s \in \mathbb{R}^d.$$

This is the Weyl form of the canonical commutation relations. It is also a twisted unitary representation of the group  $\mathbb{R}^d$  by a 2-cocycle.

- ③ Heuristically, we have

$$\lambda_\theta(t) = \exp(it_1x_1 + \cdots + it_dx_d)$$

where

$$[x_j, x_k] = i\theta_{j,k}.$$

# Identification of $L_\infty(\mathbb{R}_\theta^d)$

As a von Neumann algebra,  $L_\infty(\mathbb{R}_\theta^d)$  is not very complicated: it has type I.

**Theorem (Essentially due to Stone and von Neumann)**

*There is a \*-algebra isomorphism*

$$L_\infty(\mathbb{R}_\theta^d) \approx L_\infty(\mathbb{R}^{\ker(\theta)}) \otimes \mathcal{B}(L_2(\mathbb{R}^{\frac{\text{rank}(\theta)}{2}})).$$

When  $\theta = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ , this is recognisable as the Schrödinger representation;

$$x_1 = M_x, \quad x_2 = -i\partial_x, \quad \text{on } L_2(\mathbb{R}_x).$$

There is a group action of  $\mathbb{R}^d$  on  $L_\infty(\mathbb{R}_\theta^d)$  given on the generators by

$$T_t \lambda_\theta(s) = e^{i(t,s)} \lambda_\theta(s), \quad t, s \in \mathbb{R}^d.$$

This is the action of *translation*. Say that  $x \in L_\infty(\mathbb{R}_\theta^d)$  is uniformly smooth if

$$t \mapsto T_t x, \quad \mathbb{R}^d \rightarrow L_\infty(\mathbb{R}_\theta^d)$$

is smooth. We can define the derivatives

$$\partial_j x = \frac{d}{dt} T_{te_j}(x)|_{t=0}.$$

Given  $f \in \mathcal{S}(\mathbb{R}^d)$ , let

$$\lambda_\theta(f) = \int_{\mathbb{R}^d} f(t) \lambda_\theta(t) dt.$$

The space  $\mathcal{S}(\mathbb{R}_\theta^d)$  is defined as the image  $\lambda_\theta(\mathcal{S}(\mathbb{R}^d))$ . This is a  $*$ -algebra, multiplication is given by twisted convolution

$$\lambda_\theta(f) \lambda_\theta(g) = \lambda_\theta(f *_\theta g).$$

# Noncommutative integral

The noncommutative integral  $\tau_\theta$  is defined on  $\mathcal{S}(\mathbb{R}_\theta^d)$  by

$$\tau_\theta(\lambda_\theta(f)) := (2\pi)^d f(0).$$

## Theorem

*$\tau_\theta$  extends to a semifinite normal trace on  $L_\infty(\mathbb{R}_\theta^d)$ , giving the pair  $(L_\infty(\mathbb{R}_\theta^d), \tau_\theta)$  the structure of a semifinite von Neumann algebra.*

We can therefore define  $L_p$  spaces,  $L_p(L_\infty(\mathbb{R}_\theta^d), \tau_\theta)$ , which we abbreviate as  $L_p(\mathbb{R}_\theta^d)$ .

# Harmonic analysis questions

Many authors (notably G. Hong) and most recently M. Ruzhansky and S. Shaimardan and K. Tulenov have addressed the problem of Fourier multipliers on  $\mathbb{R}_\theta^d$ . I.e., the boundedness of functions

$$m(-i\nabla) : L_p(\mathbb{R}_\theta^d) \rightarrow L_q(\mathbb{R}_\theta^d)$$

and associated questions such as Sobolev embedding and elliptic regularity.

# Harmonic analysis questions

Many authors (notably G. Hong) and most recently M. Ruzhansky and S. Shaimardan and K. Tulenov have addressed the problem of Fourier multipliers on  $\mathbb{R}_\theta^d$ . I.e., the boundedness of functions

$$m(-i\nabla) : L_p(\mathbb{R}_\theta^d) \rightarrow L_q(\mathbb{R}_\theta^d)$$

and associated questions such as Sobolev embedding and elliptic regularity. We can also define Sobolev spaces  $W_p^s(\mathbb{R}_\theta^d)$  (Gonzales-Perez, Junge, Parcet) and Besov spaces  $B_{p,q}^s(\mathbb{R}_\theta^d)$  (Lafleche). Triebel spaces  $F_{p,q}^s(\mathbb{R}_\theta^d)$  are more challenging.



# Section 3: PDE on quantum Euclidean spaces

Linear PDE with constant coefficients

$$P(-i\nabla)u = f$$

are best studied with Fourier multipliers. Here pretty much everything works as in the commutative case.

# Linear PDE, variable coefficients

There is a theory (Gonzales-Perez, Junge, Parcet [GJP]) of elliptic PDE defined in terms of *left* multipliers:

$$\sum_{|\alpha| \leq m} a_\alpha \cdot \partial^\alpha u = f$$

where  $a_\alpha \in C^\infty(\mathbb{R}_\theta^d)$ .

# Linear PDE, variable coefficients

There is a theory (Gonzales-Perez, Junge, Parcet [GJP]) of elliptic PDE defined in terms of *left* multipliers:

$$\sum_{|\alpha| \leq m} a_\alpha \cdot \partial^\alpha u = f$$

where  $a_\alpha \in C^\infty(\mathbb{R}_\theta^d)$ .

To analyse these, they defined left pseudodifferential operators on  $u \in \mathcal{S}(\mathbb{R}_\theta^d)$  by

$$T_\sigma u = (2\pi)^{-d} \int_{\mathbb{R}^d} \sigma(\xi) \lambda_\theta(\xi) \tau_\theta(\lambda_\theta(\xi)^* u) d\xi.$$

Here,  $\sigma : \mathbb{R}^d \rightarrow L_\infty(\mathbb{R}_\theta^d)$  is a smooth function satisfying

$$\|\partial_\xi^\alpha \partial^\beta \sigma(x, \xi)\|_\infty \lesssim (1 + |\xi|)^{m - \rho|\alpha| + \delta|\beta|}.$$

# Linear PDE, variable coefficients

There is a theory (Gonzales-Perez, Junge, Parcet [GJP]) of elliptic PDE defined in terms of *left* multipliers:

$$\sum_{|\alpha| \leq m} a_\alpha \cdot \partial^\alpha u = f$$

where  $a_\alpha \in C^\infty(\mathbb{R}_\theta^d)$ .

To analyse these, they defined left pseudodifferential operators on  $u \in \mathcal{S}(\mathbb{R}_\theta^d)$  by

$$T_\sigma u = (2\pi)^{-d} \int_{\mathbb{R}^d} \sigma(\xi) \lambda_\theta(\xi) \tau_\theta(\lambda_\theta(\xi)^* u) d\xi.$$

Here,  $\sigma : \mathbb{R}^d \rightarrow L_\infty(\mathbb{R}_\theta^d)$  is a smooth function satisfying

$$\|\partial_\xi^\alpha \partial^\beta \sigma(x, \xi)\|_\infty \lesssim (1 + |\xi|)^{m - \rho|\alpha| + \delta|\beta|}.$$

Using these operators, [GJP] analysed the  $L_p$ -theory of linear elliptic equations.

# Unaddressed question: two-sided linear PDE

A trivial modification of [GJP] allows to study *right* PDE

$$\sum_{|\alpha| \leq m} (\partial^\alpha u) a_\alpha = f$$

where  $a_\alpha \in C^\infty(\mathbb{R}_\theta^d)$ .

Something that no-one has addressed (to my knowledge) is the question of two-sided linear PDE, something like

$$\sum_{|\alpha| \leq m} a_\alpha (\partial^\alpha u) b_\alpha = f.$$

This is a new phenomenon in the noncommutative case.

# Unaddressed question: two-sided linear PDE

A trivial modification of [GJP] allows to study *right* PDE

$$\sum_{|\alpha| \leq m} (\partial^\alpha u) a_\alpha = f$$

where  $a_\alpha \in C^\infty(\mathbb{R}_\theta^d)$ .

Something that no-one has addressed (to my knowledge) is the question of two-sided linear PDE, something like

$$\sum_{|\alpha| \leq m} a_\alpha (\partial^\alpha u) b_\alpha = f.$$

This is a new phenomenon in the noncommutative case. To analyse it, we would need some kind of theory of *bilateral pseudodifferential operators*.

# Section 4: Towards a theory of bilateral pseudodifferential operators



# Noncommutative Meyer decomposition?

If we want to study nonlinear PDE, we need to consider Nemytskij operators

$$u \mapsto F(u)$$

where now  $F(u)$  is defined for an operator  $u$  using functional calculus. Can we do something similar to Meyer's decomposition?

# Noncommutative Meyer decomposition?

If we want to study nonlinear PDE, we need to consider Nemytskij operators

$$u \mapsto F(u)$$

where now  $F(u)$  is defined for an operator  $u$  using functional calculus. Can we do something similar to Meyer's decomposition? Answer: Yes, but we need pseudodifferential operators formed out of both left and right multipliers.

# Simple case of a noncommutative Meyer decomposition

Consider the case  $F(u) = u^2$ . Then

$$F(u) = \sum_{j=0}^{\infty} F(S_j u) - F(S_{j-1} u) = \sum_{j=0}^{\infty} S_{j-1} u \Delta_j u + \Delta_j u S_j u.$$

So the Meyer operator should be the following:

$$m(F, u)v = \sum_{j=0}^{\infty} S_{j-1} u \Delta_j v + \Delta_j v S_j u.$$

Note that here we need both *left* and *right* multiplication.

# Bilateral multiplication

The algebra  $L_\infty(\mathbb{R}_\theta^d)^{\text{op}}$  is the opposite algebra. Given  $A \in L_\infty(\mathbb{R}_\theta^d) \otimes L_\infty(\mathbb{R}_\theta^d)^{\text{op}}$  and  $u \in L_1(\mathbb{R}_\theta^d)$ , denote  $A\sharp u$  for the linear extension of the mapping  $(a \otimes b)\sharp u = aub$ . The Haagerup tensor product  $L_\infty(\mathbb{R}_\theta^d) \otimes_h L_\infty(\mathbb{R}_\theta^d)$  has the property that  $\sharp$  has a continuous extension

$$\sharp : (L_\infty(\mathbb{R}_\theta^d) \otimes_h L_\infty(\mathbb{R}_\theta^d)^{\text{op}}) \times L_\infty(\mathbb{R}_\theta^d) \rightarrow L_\infty(\mathbb{R}_\theta^d).$$

# Proposed double symbol definition

## Definition

Let  $\rho, \delta_1, \delta_2 \in [0, 1]$  and  $m \in \mathbb{R}$ . A *bisymbol* in the class  $S_{\rho, \delta_1, \delta_2}^m(\mathbb{R}^d \times \mathbb{R}_\theta^d)$  is a function:

$$\sigma : \mathbb{R}^d \rightarrow L_\infty(\mathbb{R}_\theta^d) \otimes_h L_\infty(\mathbb{R}_\theta^d)^{\text{op}}.$$

Such that for all multi-indices  $\alpha, \beta_1, \beta_2 \in \mathbb{N}^d$  we have:

$$\|\partial_\xi^\alpha (\partial_x^{\beta_1} \otimes \partial_x^{\beta_2}) \sigma(\xi)\|_{L_\infty(\mathbb{R}_\theta^d) \otimes_h L_\infty(\mathbb{R}_\theta^d)} \leq C_{\alpha, \beta_1, \beta_2} (1 + |\xi|)^{m - \rho|\alpha| + \delta_1|\beta_1| + \delta_2|\beta_2|}$$

# Bilateral pseudodifferential operators

Let  $\sigma \in S_{\rho, \delta_1, \delta_2}^m(\mathbb{R}^d \times \mathbb{R}_\theta^d)$ . Define an operator  $T_\sigma$  on  $u \in \mathcal{S}(\mathbb{R}_\theta^d)$  by

$$T_\sigma u := (2\pi)^{-d} \int_{\mathbb{R}^d} (\sigma(\xi) \sharp \lambda_\theta(\xi)) \tau_\theta(\lambda_\theta(\xi)^* u) d\xi$$

Denote space of such operators by  $\Psi_{\rho, \delta_1, \delta_2}^m(\mathbb{R}_\theta^d)$ .

# Bilateral pseudodifferential operators

Let  $\sigma \in S_{\rho, \delta_1, \delta_2}^m(\mathbb{R}^d \times \mathbb{R}_\theta^d)$ . Define an operator  $T_\sigma$  on  $u \in \mathcal{S}(\mathbb{R}_\theta^d)$  by

$$T_\sigma u := (2\pi)^{-d} \int_{\mathbb{R}^d} (\sigma(\xi) \sharp \lambda_\theta(\xi)) \tau_\theta(\lambda_\theta(\xi)^* u) d\xi$$

Denote space of such operators by  $\Psi_{\rho, \delta_1, \delta_2}^m(\mathbb{R}_\theta^d)$ .

## Conjecture

- For  $\delta_1 + \delta_2 < \rho$ ,  $\Psi_{\rho, \delta_1, \delta_2}^m(\mathbb{R}_\theta^d)$  has similar properties of adjointability, composition and asymptotic convergence to traditional pseudodifferential operators.

# Bilateral pseudodifferential operators

Let  $\sigma \in S_{\rho, \delta_1, \delta_2}^m(\mathbb{R}^d \times \mathbb{R}_\theta^d)$ . Define an operator  $T_\sigma$  on  $u \in \mathcal{S}(\mathbb{R}_\theta^d)$  by

$$T_\sigma u := (2\pi)^{-d} \int_{\mathbb{R}^d} (\sigma(\xi) \sharp \lambda_\theta(\xi)) \tau_\theta(\lambda_\theta(\xi)^* u) d\xi$$

Denote space of such operators by  $\Psi_{\rho, \delta_1, \delta_2}^m(\mathbb{R}_\theta^d)$ .

## Conjecture

- For  $\delta_1 + \delta_2 < \rho$ ,  $\Psi_{\rho, \delta_1, \delta_2}^m(\mathbb{R}_\theta^d)$  has similar properties of adjointability, composition and asymptotic convergence to traditional pseudodifferential operators.
- For  $\rho = 1 > \delta_1 + \delta_2$ , and  $m = 0$ , we have boundedness on  $L_p(\mathbb{R}_\theta^d)$  for  $1 < p < \infty$ .
- In general for  $m = 0$ , we have boundedness on  $W_p^s(\mathbb{R}_\theta^d)$  for  $s > 0$ .



# Why is it interesting?

- If the conjecture is correct, we would have a *genuinely noncommutative* theory of pseudodifferential operators which would allow us to study bilateral PDE and also nonlinear PDE over  $\mathbb{R}_\theta^d$ .
- Following the standard commutative proofs does not seem to work.

Thank you for listening!