School of Physics
Trinity College Dublin
Dublin 2
Ireland
+353 (0)85 720 3415
eogorma@tcd.ie

November 21, 2013

NAASC Postdoctoral Fellow Charlottesville, VA, USA

To whom it may concern:

I am currently in the final stages of completing a Ph.D. in astrophysics at Trinity College Dublin, Ireland, under the supervision of Prof. Graham Harper. On September 30, 2013, I submitted my thesis (Title: Radio Interferometric Studies of Cool Evolved Stellar Winds), and I successfully defended my thesis on November 15, 2013. I am due to graduate from Trinity College Dublin in April 2014 and envisage to continue with my research in stellar astrophysics thereafter. I can assure you that my technical experience in radio interferometry, scientific knowledge, and communication skills, make me a strong candidate for the NAASC postdoctoral fellowship.

My Ph.D. research has focused on understanding the atmospheres of red giants and red supergiants through multi-wavelength radio interferometric studies (both continuum and line emission). These stars have substantial mass-loss rates, but the mechanisms which heat and launch their outflows remain an enduring mystery. Progress in this field continues to be driven by observations, and this NAASC postdoctoral fellowship at Charlottesville would enable me to pursue such observations.

I have extensive experience in reducing interferometric data at both millimeter and centimeter wavelengths, all of which have been carried out within CASA. During my Ph.D., I have been a successful P.I. on proposals to use the Combined Array for Research in Millimeter-wave Astronomy (CARMA) array, the Karl G. Jansky Very Large Array (VLA), and the Giant Metrewave Radio Telescope (GMRT). I have designed and planned all such observations and therefore have intensive experience in creating observing scripts and using online submission tools. I also have a keen interest in public outreach and would love to continue this interest into my postdoctoral career.

Thank you for taking the time to consider my credentials and I look forward to hearing from you.

Eamon O'Gorman

Eamon O'Gorman - CV

CONTACT Information School of Physics Trinity College Dublin

Dublin 2 Ireland office: +353 (0)1 896 2157 mobile: +353 (0)85 720 3415 e-mail: eogorma@tcd.ie website: maths.tcd.ie/ \sim eogorma

EDUCATION

Trinity College Dublin, Dublin, Ireland

Doctor of Philosophy

October 2009 – present

- Advisor: Professor Graham Harper
- Thesis title: Radio Interferometric Studies of Cool Evolved Stars
- Thesis successfully defended on 15th November 2013
 - External Examiner: Professor Tom Millar (Queen's University Belfast, Northern Ireland)
 - Internal Examiner: Professor Peter Gallagher

International Space University (ISU), Strasbourg, France

M.Sc., Space Studies

September 2008 – August 2009

University College Dublin (UCD), Dublin, Ireland

B.Sc., Theoretical Physics, (First class honours)

September 2003 – June 2007

Honours and Awards

- Enterprise Ireland/European Space Agency scholarship to study at the ISU, 2008
- UCD Entrance Scholar, 2003

Conference Presentations

Oral:

O'Gorman, E. Radio Interferometric Studies of Cool Evolved Stellar Mass Outflows. DIAS Seminar, Dublin, Dublin Institute for Advanced Studies, Ireland, February 2013.

O'Gorman, E., et al. Probing the Thermodynamics of Red Giant Mass Outflows with the JVLA. Astronomical Science Group of Ireland, Galway, Ireland, November 2012.

O'Gorman, E., et al. Probing the Thermodynamics of Red Giant Mass Outflows with the JVLA. Radio Stars and Their Lives in the Galaxy, MIT Haystack Observatory, MA, USA, October 2012.

O'Gorman, E. CO in the Circumstellar Envelope of Betelgeuse with CARMA. The 41st Young European Radio Astronomers Conference, University of Manchester/Jodrell Bank Observatory, UK, July 2011.

Poster:

O'Gorman, E., Harper, G. M. What is Heating Arcturus' Wind?

16th Cambridge Workshop on Cool Stars, Stellar Systems, and the Sun, University of Washington, Seattle, USA, August 2010.

ACCEPTED PROPOSALS AS PI O'Gorman, E., et al. Beta Gem b: An alternative candidate in the search for extrasolar planetary radio emission (II), GMRT, 2013, ID: 25_039

O'Gorman, E., et al. Beta Gem b: An alternative candidate in the search for extrasolar planetary radio emission, GMRT, 2013, ID: 24_013

O'Gorman, E., et al. Thermal Continuum Mapping of Red Giant Chromospheres, CARMA, 2012, ID: c1038

O'Gorman, E., et al. L and S band Continuum Observations of Arcturus: Completing a Clean Sweep, VLA, 2012, ID: VLA-12A-472

SHORT-TERM RESEARCH STAYS

National Centre for Radio Astrophysics, India, 2013: Collaboration with Dr. Sandeep Sirothia to prepare our GMRT 150 MHz observations and carry out initial data analysis.

Harvard-Smithsonian Center for Astrophysics, USA, 2011: Collaboration with Dr. Joanna Brown to carry out initial analysis on our CARMA millimeter data using CASA.

NASA Goddard Space Flight Center, USA, 2009: Three month student internship with Dr. Drake Deming in the area of transiting exoplanet characterization. Analyzed data from the FLAMINGOS infrared camera on the 2.1 m Kitt Peak National Observatory Telescope.

TEACHING EXPERIENCE

- September 2010 April 2012: Bi-weekly physics tutorials for undergraduate engineering students.
- September 2009 April 2010: Teaching assistant for undergraduate physics students.

Outreach

- Throughout the year we carry out a "Build a Comet in the Lab" workshop for both secondary school students interested in pursuing physics in college, and for primary school students from disadvantaged backgrounds.
- Regularly visit secondary schools to discuss career opportunities in physics and astrophysics to prospective students.
- Active member in "SunSpotter", a new project which aims to enlist the help of the public to readily identify and characterize sunspots in NASA satellite image.

Programming

IDL, Python, CASA, \LaTeX 2ε , PHP, BASH, and UNIX-like operating systems.

Professional Organisations

- Fellow of the Royal Astronomical Society
- Associate Member of the Institute of Physics

OTHER Interests

I love the outdoors and playing sport. I have played hurling (a traditional Gaelic sport) since I was a child. In 2011 I was named Trinity College senior hurling player of the year.

Referees

Prof. Graham Harper School of Physics Trinity College Dublin Dublin 2, Ireland phone: +353 (0)1 896 3257 e-mail: graham.harper@tcd.ie

Trinity College Dublin
Dublin 2, Ireland
phone: +353 (0)1 896 1300
e-mail: peter.gallagher@tcd.ie

Prof. Peter Gallagher

School of Physics

Prof. Hugh Hill International Space University 1 rue Jean–Dominique Cassini Strasbourg, France phone: +33 (0)3 88 65 54 39

phone: +33 (0)3 88 65 54 39e-mail: hill@isu.isunet.edu

Refereed Publications

O'Gorman, E., Harper, G. M., Brown, A., Drake, S., Richards, A. M. S. Multi-wavelength Radio Continuum Emission Studies of Dust-free Red Giants, 2013, AJ, 146, 98.

Richards, A. M. S., Davis, R. J., Decin, L., Etoka, S., Harper, G. M., Lim, J. J., Garrington, S. T., Gray, M. D., McDonald, I., **O'Gorman, E.**, Wittkowski, M. e-MERLIN resolves Betelgeuse at λ 5 cm: hotspots at 5 R_{\star} , 2013, MNRAS, 432, L61.

O'Gorman, E., Harper, G. M., Brown, J. M., Brown, A., Redfield, S., Richter, M. J., Requena-Torres, M. A. $CARMA\ CO(J=2-1)$ Observations of the Circumstellar Envelope of Betelgeuse, 2012, AJ, 144, 36.

Sada, P. V., Deming, D., Jennings, D. E., Jackson, B. K., Hamilton, C. M., Fraine, J., Peterson, S. W., Hasse, F., Bays, K., Lunsford, A., **O'Gorman, E.** Extrasolar Planet Transits Observed at Kitt Peak National Observatory, 2012, PASP, 124, 212.

Sada, P. V., Deming, D., Jackson, B., Jennings, D. E., Peterson, S. W., Haase, F., Bays, K., O'Gorman, E., Lundsford, A. Recent Transits of the Super-Earth Exoplanet GJ 1214b, 2010, ApJ, 720, L215.

Non-refereed Publications

O'Gorman, E., & Harper, G. M., What is Heating Arcturus' Wind?, 2011, Proceedings of the 16th Cambridge Workshop on Cool Stars, Stellar Systems and the Sun. Astronomical Society of the Pacific Conference Series, 448, 691.

Research Accomplishments

My main research achievements to date have been associated with gaining a broader understanding into the dynamics and thermodynamics of the atmospheres of red giants (non-AGB red giants) and red supergiants (RSGs), through radio interferometric techniques. The mass-loss from these stars plays a crucial role in galactic evolution and ultimately provides much of the material required for the next generation of stars and planets. Also, a substantial fraction of the star's initial mass can be dispersed to the interstellar medium during these post main sequence evolutionary stages and this mass-loss is therefore a crucial factor governing stellar evolution, and also in explaining the frequency of supernovae in the galaxy. Despite the importance of this phenomenon, and decades of study, the specific mechanisms that drive winds from evolved spectral-type K through mid-M stars remain unknown. There is insufficient atomic, molecular, or dust opacity to drive a radiationdriven outflow and acoustic/pulsation models cannot drive the observed mass-loss rates. UV and optical observations reveal an absence of significant hot wind plasma, and the winds are thus too cool to be Parker-type thermally-driven flows. Magnetic fields are most likely involved in the massloss process, although current magnetic models are also unable to explain spectral diagnostics. Traditionally, observations have provided only limited disk-averaged information about the outflow environments of these stars, making it difficult to infer the wind properties. My primary research to date has focused on using the latest suite of millimeter and centimeter interferometers to provide essential spatial information on these outflow environments, to gain a better understanding of the entire mass-loss process.

Betelgeuse is one of the few nearby RSGs that can be studied in great detail across most of the electromagnetic spectrum. The extended atmosphere of this oxygen-rich RSG provides an ideal testbed for studying the poorly understood processes that drive mass-loss from K and M supergiants. Part of my research to date has focused on explaining and reconciling the puzzling results from previous single dish millimeter observations of this star's circumstellar environment. Ro-vibrational absorption lines of ¹²C¹⁶O and ¹³C¹⁶O by Bernat et al. (1979) revealed that Betelgeuse has undergone at least two distinct phases of mass-loss in its recent past. However, previous millimeter observations were only able to detect one component of the outflow, while the spatial extent of both components were unknown. Using multiple array configurations of the CARMA interferometer, we successfully imaged the CO(J=2-1) emission line at sub-arcsecond resolution, and for the first time were able to find the spatial extent of both outflow components, allowing their ages to be calculated (O'Gorman et al., 2012, AJ). We also found both components to be inhomogeneous, far from the existing spherically symmetric models of its circumstellar environment, indicating a chaotic mass-loss loss process. These millimeter line emission data were obtained with the CARMA heterogeneous interferometer at Ceder Flat, California, in three different array configurations. The raw data were flagged, calibrated, and imaged using CASA. The analysis required knowledge of interferometric spatial scales and imaging techniques such as CLEAN and multi-scale CLEAN.

The wind acceleration region is where most of the momentum and heat which drive the mass-loss in RSGs is deposited. Thermal millimeter and centimeter continuum emission directly sample this region which extends out to only a few R_{\star} . Spatially resolved VLA cm-continuum radio maps (Lim et al. 1998) revealed that Betelgeuse's extended atmosphere $(2-7R_{\star})$ was significantly cooler and much less ionized than predicted by magnetic wave models. I have been heavily involved in a collaboration which has used e-MERLIN to image the thermal continuum emission from this inner region of Betelgeuse's atmosphere at 6 cm (Richards et al., 2013). We have discovered two chromospheric "hotspots" (Figure 1, left) separated by $4R_{\star}$, hidden just beyond the spatial resolution of the VLA at 6 cm. Using the astrometric solution of Harper et al. (2008), the brighter hotspot ($T_e > 5400 \,\mathrm{K}$) is $\sim 3.5 \,R_{\star}$ from the photosphere. Inspired by this new discovery, I have analyzed multi-wavelength,

multi-epoch, centimeter old VLA data which contains the Pie Town VLBA antenna, to look for signatures of these hotspots. At the highest available frequencies, these data have even superior spatial resolution than e-MERLIN (at 6 cm). In Figure 1 (right) a preliminary map at 0.7 cm is presented, and evidence of at least two features separated by only $2 R_{\star}$ is shown. These features may be generated by giant photospheric convective cells, although it is unlikely that such features could be accountable for the "hotspots" seen with e-MERLIN (O'Gorman, 2013, *Thesis*).

We have recently used the Jansky Very Large Array to observe two non-dusty, non-pulsating, K spectral-type red giants at multiple radio wavelengths $(0.7 - 20 \,\mathrm{cm})$ (O'Gorman et al., 2013, AJ). Such stars are feeble emitters at these wavelengths however, and previous observations have provided only a small number of modest S/N measurements slowly accumulated over three decades. Our observations of each star were carried out over just a few days, so that we obtained an essentially consistent snapshot of the different stellar atmospheric layers sampled at different wavelengths. We found that our observations were in disagreement with the existing atmospheric models for these stars, which were based only on UV diagnostics. We also found evidence for a rapidly cooling stellar wind for one of the targets which allowed us to develop a new semi-empirical wind model for the star. This model was then used as the basis to compute a thermal energy balance of the star's outflow by investigating the various heating and cooling processes that control its thermal structure (O'Gorman, 2013, Thesis). As part of this project, I planned and prepared two sets of observations of multi-frequency centimeter continuum observations with the Jansky Very Large Array. Different frequencies required different observing strategies, such as choice of calibrator (flux and phase) and length of observing scans. All data analysis was again carried out in CASA. The large bandwidth now provided by the Jansky VLA required additional steps to mitigate RFI, and to create wideband images.




Figure 1. Left: e-MERLIN 6 cm image of Betelgeuse showing the two chromospheric "hotspots". The red filled circle marks the expected photospheric position. Right: Old VLA + Pie Town 0.7 cm image of Betelgeuse's inner asymmetric atmosphere. The presence of giant photospheric convective cells can account for the asymmetries in the 0.7 cm image, but cannot account for the hotspot at $\sim 3.5 R_{\star}$ in the 6 cm image.

Future Research Plans

The latest suite of radio interferometers such as the Jansky VLA, e-MERLIN, and ALMA now have (or will have in the near future) the capability of providing spatially resolved sensitive millimeter and centimeter observations of the atmospheres of the closest RSGs and red giants. A NAASC postdoctoral fellowship would provide me with both the time and resources to carry out these observations. Such observations would contribute exciting new insights into the currently unknown mass-loss process in RSGs and red giants. I am part of a collaboration that has been awarded ALMA

cycle 1 time (PI: P. Kervella) which will trace CO emission around Betelgeuse at a resolution 10 times better than we achieved with CARMA. This project will also provide insight into the role dust plays in the mass-loss of Betelgeuse, a relatively non-dusty M supergiant. These observations may be carried over to Cycle 2 if our project is not fully observed during Cycle 1 (i.e., by the end of May 2014). My past experience with analyzing millimeter interferometric data will be of great importance to this project and working with the first ALMA observations of a RSG may be my first project as a NAASC postdoctoral fellow.

Our recent findings with e-MERLIN (described above) raise more questions about Betelgeuse than answers. Is the current astrometric solution accurate? What mass-loss process could cause such features? What are the *hotspots*? On what time scales do they evolve? Multi-wavelength high spatial resolution monitoring of Betelgeuse with e-MERLIN is required to solve this puzzling evidence and this project would be one of my main priorities in my first year as a NAASC postdoctoral fellow. The next call for e-MERLIN proposals is in spring 2014 and I plan to submit a strong proposal which will observe Betelgeuse at both 6 cm (C band) and 1.3 cm (K band) for at least two epochs. I also plan to apply for Jansky VLA A-configuration time in August 2014 to observe Betelgeuse again at 6 cm and 1.3 cm with the intention of combining these data with data from e-MERLIN. Such data would produce the highest spatial resolution radio image ever of the thermal emission from any star other than our Sun.

The Jansky Very Large Array now provides over an order of magnitude increase in continuum sensitivity. Another goal in the first year as a NAASC postdoctoral fellow would be to utilize this new capability and carry out a survey ($\sim 10-15$ targets) of nearby coronal red giants to detect their total radio flux density at 3.5 cm and 6 cm. Such measurements will provide estimates of their mass-loss rates, which are notoriously difficult to estimate at other wavelengths due to their optically thin winds. Plotting mass-loss rates as a function of various stellar parameters will for the first time allow empirical mass-loss formulas to be constructed for these late-type stars. To avoid confusion from near-by extragalactic objects, the second most extended B-configuration of the Jansky VLA will be required and so I again plan on proposing for time in August 2014.

The atmospheres of red giants cannot currently be spatially resolved at millimeter or centimeter wavelengths, but this will change in the next few years when ALMA and e-MERLIN achieve sufficient spatial resolution. The atmospheric properties of these stars are still poorly understood and this ultimately leads to a lack of understanding into the mechanism by which they lose mass. As well as revealing any large scale structure that may be present in their wind acceleration zones, spatially resolved thermal free-free millimeter and centimeter emission can directly provide a measurement of the gas temperature. It is the gas temperature which controls the heating rates in the wind, which in turn provides insights into the unknown mass-loss mechanism. I envisage to carry out such studies in my second and third year as a NAASC postdoctoral fellow.

My interest in astrophysics is wide and varied however, and I would also be very much interested in starting new collaborations at NAASC and indeed at NRAO where I could utilize my experience in millimeter and centimeter interferometry in other areas of astrophysics. An example of this is my ongoing project to detect radio emission from β Gem b, an exoplanet orbiting the closet red giant, Pollux (PI: E. OGorman, ID: 24 013). Previous searches for exoplanet radio emission have focused on close orbiting planets, the so-called *hot Jupiters*. Our target is further away from its host star and free from tidal locking, which may reduce the internal magnetic field of the hot Jupiters, thus reducing the strength of the radio emission. We expect the relatively large mass-loss of the host red giant to be a key driver in detecting this emission. We are currently using the GMRT to search for this emission at 150 MHz. In June 2013, I stayed at the GMRT while our long wavelength radio observations were being carried out. While there I prepared observing scripts and carried out initial data analysis. At NAASC, I would envisage to carry out further observations of this interesting planetary system at lower frequencies and higher sensitivity with LOFAR.