

Radio Interferometric Studies of Cool Evolved Stellar Winds

A dissertation submitted to the University of Dublin
for the degree of Doctor of Philosophy

Eamon O’Gorman

Supervisor: Prof. Graham M. Harper
Trinity College Dublin, September 2013

SCHOOL OF PHYSICS
UNIVERSITY OF DUBLIN
TRINITY COLLEGE



Declaration

I declare that this thesis has not been submitted as an exercise for a degree at this or any other university and it is entirely my own work.

I agree to deposit this thesis in the University's open access institutional repository or allow the library to do so on my behalf, subject to Irish Copyright Legislation and Trinity College Library conditions of use and acknowledgement.

Name: Eamon O'Gorman

Signature: **Date:**

Summary

Mass-loss becomes significant for most stars as they approach the end of their lives and become either red giants or red supergiants. This continuous mass-loss, which occurs in the form of a relatively dense and slow-moving wind, can have a significant impact on the evolution of gas and dust in galaxies, on surrounding planets, and indeed on the very evolution of the star itself. Despite the importance of this phenomenon and decades of study, the fundamental mechanisms responsible for producing these winds remain unknown. The main reason for this is due to our lack of understanding of the dynamics and thermodynamics of the stellar outflow environment. Isolated giants and supergiants do not contain the expected additional complexities encountered by binaries, making them ideal targets for understanding the nature of these outflows. Traditionally, observations have provided only limited disk-averaged information about the outflow environments of these stars, making it difficult to infer the outflow properties. However, the latest suite of radio interferometers now have the capability to provide essential spatial information on these outflow environments.

This thesis first presents the results of a radio interferometric study into the dynamics of the two unique flows in the circumstellar environment of the M2 red supergiant, Betelgeuse. The Combined Array for Research in Millimeter-wave Astronomy (CARMA) was used in multiple configurations to observe the $\text{CO}(J = 2 - 1)$ emission line allowing spatial scales as small as $0''.9$ ($\sim 40 R_\star$) to be traced over a $32''$ ($\sim 1500 R_\star$) field of view. The outer flow known as S2, was found to have outflow velocities of -15.4 and $+13.2 \text{ km s}^{-1}$ with respect to the stellar rest frame and extend out to $17''$, while the inner

flow known as S1, was found to have outflow velocities of -9.0 and $+10.6 \text{ km s}^{-1}$ and extend out to between $4 - 6''$. Both flows were found to be inhomogeneous down to the resolution limit, but when azimuthally averaged, their intensity falloff was found to be consistent with an optically thin, spherically symmetric, constant velocity outflow. High resolution multi-epoch centimeter continuum observations of Betelgeuse which probe its inner atmosphere ($< 10 R_{\star}$) are also presented. The radio flux density is found to vary on time scales of $\lesssim 14$ months at all wavelengths, and again evidence for inhomogeneities in the outflow is found.

This thesis also presents Karl G. Jansky Very Large Array (VLA) multi-wavelength ($0.7 - 20 \text{ cm}$) observations of two non-dusty, non-pulsating K spectral-type red giants, Arcturus and Aldebaran. Detections at 10 cm (3.0 GHz : S-band) and 20 cm (1.5 GHz : L-band) represent the first isolated luminosity class III red giants to be detected at these long wavelengths. These thermal continuum observations provide a snapshot of the different stellar atmospheric layers and are independent of any long-term variability. The long wavelength data sample Arcturus' outer atmosphere where the wind velocity is approaching its terminal value and the ionization balance is becoming *frozen-in*. For Aldebaran, the data samples its inner atmosphere where the wind is still accelerating. Our data is in conflict with published semi-empirical models based on ultraviolet data. Spectral indices are used to discuss the possible properties of the stellar atmospheres. Evidence for a rapidly cooling wind in the case of Arcturus is found and a new analytical wind model is developed for this star. This model is used as the basis to compute a thermal energy balance of Arcturus' outflow by investigating the various heating and cooling processes that control its thermal structure. The analysis focuses on distances between $1.2 \rightarrow 10 R_{\star}$ and includes the wind acceleration zone. We find that an additional substantial heating mechanism is required to maintain the inner thermal structure of the outflow.

*For Mum and Dad,
a constant source of inspiration and guidance.*

Acknowledgements

I am immensely grateful to my supervisor Graham Harper. His enthusiasm, dedication, patience, and encouragement, have been a huge help to me over the past four years. His knowledge and passion for the subject is beyond belief.

I would like to thank Joanna Brown, Alex Brown, and Anita Richards, for their help and advice with the data analysis aspects of this thesis.

I would also like to thank all past and present members of the Astrophysics Research Group. It has been great getting to know everybody and you have made the last four years an unforgettable experience.

A huge thanks to Sarah for proof reading this thesis, to Alex for providing me with a roof over my head during the write-up, and to Claire for keeping me motivated all the way from Brussels. Finally and most importantly, I would like to thank Mum and Dad for everything.

List of Publications

Refereed

1. **O’Gorman, E.**, Harper, G. M., Brown, A., Brown, A., Drake, S., and Richards, A. M. S.
“Multi-wavelength Radio Continuum Emission Studies of Dust-free Red Giants”
The Astronomical Journal, 146, 98 (2013)
2. Richards, A. M. S., Davis, R. J., Decin, L., Etoke, S., Harper, G. M., Lim, J. J., Garrington, S. T., Gray, M. D., McDonald, I., **O’Gorman, E.**, Wittkowski, M.
“e-MERLIN resolves Betelgeuse at wavelength 5 cm”
Monthly Notices of the Royal Astronomical Society Letters, 432, L61 (2013)
3. **O’Gorman, E.**, Harper, G. M., Brown, J. M., Brown, A., Redfield, S., Richter, M. J., and Requena-Torres, M. A.
“CARMA CO(J = 2 - 1) Observations of the Circumstellar Envelope of Betelgeuse”
The Astronomical Journal, 144, 36 (2012)
4. Sada, P. V., Deming, D., Jennings, D. E., Jackson, B. K., Hamilton, C. M., Fraine, J., Peterson, S. W., Haase, F., Bays, K., Lunsford, A., and **O’Gorman, E.**
“Extrasolar Planet Transits Observed at Kitt Peak National Observatory”
Publications of the Astronomical Society of the Pacific, 124, 212 (2012)

-
5. Sada, P. V., Deming, D., Jackson, B. K., Jennings, D. E., Peterson, S. W., Haase, F., Bays, K., **O’Gorman, E.**, and Lundsford, A.
“Recent Transits of the Super-Earth Exoplanet GJ 1214b”
The Astrophysical Journal Letters, 720, L215 (2010)

Non-Refereed

1. **O’Gorman, E.**, & Harper, G. M.
“What is Heating Arcturus’ Wind?”,
Proceedings of the 16th Cambridge Workshop on Cool Stars, Stellar Systems and the Sun. Astronomical Society of the Pacific Conference Series, 448, 691 (2011)

Contents

List of Publications	vii
List of Figures	xiv
List of Tables	xvii
A List of Abbreviations Used in this Thesis.	1
B Discrete Absorption Feature	3
C Ambipolar Diffusion Heating	5
D Turbulent Heating	8
References	10

List of Figures

List of Tables

A.1 List of Abbreviations 1



List of Abbreviations Used in this Thesis.

Table A.1: List of Abbreviations

Acronym	Meaning
ALMA	The Atacama Large Millimeter/submillimeter Array
AGB	Asymptotic Giant Branch
ALC	Automatic Level Control
BIMA	Berkeley Illinois Maryland Association
CARMA	Combined Array for Research in Millimeter-wave Astronomy
CASA	Common Astronomy Software Application
CSE	Circumstellar Envelope
DDT	Director's Discretionary Time
e-MERLIN	e-Multi-Element Radio Linked Interferometer Network
FFT	Fast Fourier Transform
FITS	Flexible Image Transport System
FOV	Field of View
GBT	Robert C. Byrd Green Bank Telescope
GHRS	Goddard High-Resolution Spectrograph
GREAT	German Receiver for Astronomy at Terahertz Frequencies
HIFI	Heterodyne Instrument for the Far Infrared
HPBW	Half Power Beamwidth
H-R	Hertzsprung-Russell

Continued on next page

Table A.1 – *Continued from previous page*

Acronym	Meaning
HST	Hubble Space Telescope
IF	Intermediate Frequency
IOTA	Infrared Optical Telescope Array
IR	Infrared
IRAM	Institut de Radioastronomie Millimétrique
ISM	Interstellar Medium
IUE	International Ultraviolet Explorer
LNA	Low Noise Amplifier
LO	Local Oscillator
LSR	Local Standard of Rest
LTE	Local Thermodynamic Equilibrium
MEM	Maximum Entropy Method
MERLIN	Multi-Element Radio Linked Interferometer Network
MHD	Magnetohydrodynamic
OVRO	Owens Valley Radio Observatory
OSRO	Open Shared Risk Observing
RF	Radio Frequency
RFI	Radio Frequency Interference
RGC	Red Giant Clump
RGB	Red Giant Branch
RSG	Red Supergiant
S/N	signal-to-noise ratio
SB	Scheduling Block
SGB	Subgiant Branch
SOFIA	Stratospheric Observatory for Infrared Astronomy
SMA	Submillimeter Array
SZA	Sunyaev-Zel'dovich Array
SIS	Superconductor Insulator Superconductor
UV	Ultraviolet
VLA	Karl G. Jansky Very Large Array
VLBA	Very Long Baseline Array
VLT	Very Large Telescope
W-R	Wolf-Rayet

B

Discrete Absorption Feature

During this study we analyzed *HST* STIS spectra of Arcturus from the online StarCAT catalog (Ayres, 2010b). The Mg II *h* and *k* lines from data obtained in 2001 show a wind velocity $\sim 30\text{--}40\text{ km s}^{-1}$, which is similar to that adopted in the Drake models for this star Drake (1985). A narrow discrete absorption feature was found at -49 km s^{-1} in the broad blue-shifted wind absorption component of both lines as shown in Figure B.1. For this discrete feature we find a most probable turbulent velocity of 3.4 km s^{-1} and a Mg II column density¹ of $1.4 \times 10^{12}\text{ cm}^{-2}$. A Mg II column density of 10^{15} cm^{-2} is required to produce the blueward absorption components in the *h* and *k* lines (?). Therefore, this discrete absorption feature accounts for only $\sim 0.1\%$ of the total wind column density and is probably a result of a discrete ejection event.

¹Assuming all Mg to be Mg II





Figure B.1: Analysis on the absorption feature found in the Mg II *h* and *k* lines. A function composed of a linear combination of a Gaussian and a quadratic fitted the discrete absorption feature the best. The red data in the upper row shows the data that is used in this analysis.



Ambipolar Diffusion Heating

Considering a steady flow accelerating in an inertial frame, the equation of motion can be written as

$$\rho_n \mathbf{a}_n = \rho_n \mathbf{g} + \mathbf{f}_d \quad (\text{C.1})$$

for the neutral species n , and

$$\rho_i \mathbf{a}_i = \rho_i \mathbf{g} - \mathbf{f}_d + \mathbf{f}_L \quad (\text{C.2})$$

for the ion species, i . The flow acceleration is defined as $\mathbf{a} \equiv (\mathbf{v} \cdot \nabla) \mathbf{v}$ and the gravitational acceleration is $\mathbf{g} \equiv \nabla(GM_\star/r)$. The volumetric drag force of the ions on the neutrals is defined as

$$\mathbf{f}_d = \gamma \rho_n \rho_i (\mathbf{v}_i - \mathbf{v}_n) \quad (\text{C.3})$$

and \mathbf{f}_L is the volumetric Lorentz force. The equation of motion for the combined ion-neutral fluid is found by the addition of Equations C.1 and C.2

$$\rho \mathbf{a} = \rho \mathbf{g} + \mathbf{f}_L \quad (\text{C.4})$$

where $\rho \equiv \rho_n + \rho_i$ is the mass density without the electrons and $\mathbf{a} \equiv (\rho_n \mathbf{a}_n + \rho_i \mathbf{a}_i) / \rho$ is the total acceleration.

The gravitational acceleration term can then be eliminated from Equations C.1 and C.2 to give

$$\mathbf{a}_n - \mathbf{a}_i = \left(\frac{1}{\rho_n} + \frac{1}{\rho_i} \right) \mathbf{f}_d - \frac{1}{\rho_i} \mathbf{f}_L. \quad (\text{C.5})$$

Assuming then that the acceleration of the neutrals and ions are the same we get

$$\mathbf{f}_d = \left(\frac{\rho_n}{\rho_n + \rho_i} \right) \mathbf{f}_L. \quad (\text{C.6})$$

This equation tells us that for a lightly ionized outflow the drag force is almost equal to the Lorentz force. We can now obtain an expression for the slip velocity, \mathbf{w} , by subbing this equation into Equation C.3

$$\mathbf{w} = \mathbf{v}_i - \mathbf{v}_n = \frac{\mathbf{f}_L}{\gamma \rho_i (\rho_n + \rho_i)}. \quad (\text{C.7})$$

The slip velocity becomes large when the ion density becomes small, but does not become large when the neutral density becomes small because the large density of ions drag the few neutrals that are present along with the rest of the mostly ionized plasma. The heating rate per unit volume due to ambipolar diffusion heating is

$$\Gamma = \mathbf{f}_d \cdot \mathbf{w} \quad (\text{C.8})$$

and substitution of Equations C.6 and C.7 gives

$$\Gamma = \frac{\rho_n |\mathbf{f}_L|^2}{\gamma \rho_i (\rho_n + \rho_i)^2}, \quad (\text{C.9})$$

and so for a completely ionized plasma, $\Gamma = 0$.

In order to calculate the ambipolar diffusion heating, we need to find a value for the ion-neutral momentum transfer coefficient, γ (in units $\text{cm}^3 \text{s}^{-1} \text{g}^{-1}$) which depends on the collisional coefficient rates, cross sections, slip speed, and gas

composition. Shang *et al.* (2002) give the following expression

$$\gamma = \frac{2.13 \times 10^{14}}{1 - 0.714x_e} \left(\left[3.23 + 41.0T_4^{0.5} \times \left(1 + 1.338 \times 10^{-3} \frac{w_5^2}{T_4} \right)^{0.5} \right] x_{HI} + 0.243 \right) \quad (\text{C.10})$$

where T_4 is the temperature in units of 10^4 K, and w_5 is the slip velocity in units of km s^{-1} . We have assumed no molecular hydrogen to be present and the fractional abundance of He, $x_{He} = 0.1$. Subbing Equation C.7 into Equation C.10 gives a quartic equation for γ , i.e.,

$$\gamma^4 - (2AE + 2ABx_{HI})\gamma^3 + (A^2E^2 + 2A^2BE x_{HI} + A^2B^2x_{HI}^2 - A^2C^2x_{HI}^2)\gamma^2 - GA^2C^2x_{HI}^2 = 0$$

where

$A = \frac{2.13 \times 10^{14}}{1 - 0.714x_e}$, $B = 3.23$, $C = 41.0T_4^{0.5}$, $D = \frac{1.338 \times 10^{-3}}{T_4}$, $E = 0.243$, $F = \frac{\mathbf{f}_L}{\rho_i(\rho_n + \rho_i)}$, and $G = \frac{DF^2}{1 \times 10^{10}}$. Finally the components of the Lorentz force and thus the corresponding volumetric ambipolar heating rates can be calculated by using the following expressions for the flow and gravitational accelerations:

$$\mathbf{a} = v \frac{dv}{dr} \mathbf{r} + \frac{v}{r} \frac{dv}{d\theta} \boldsymbol{\theta} + \frac{v}{r \sin \theta} \frac{dv}{d\phi} \boldsymbol{\phi} \quad (\text{C.11})$$

and

$$\mathbf{g} = -\frac{GM_\star}{r^2} \mathbf{r} + \frac{1}{r} \frac{d}{d\theta} \left(\frac{GM_\star}{r} \right) \boldsymbol{\theta} + \frac{1}{r \sin \theta} \frac{d}{d\phi} \left(\frac{GM_\star}{r} \right) \boldsymbol{\phi}. \quad (\text{C.12})$$

D

Turbulent Heating

By including time dependence and shear viscosity, μ , into the stellar wind momentum equation we get the Navier-Stokes equation

$$\rho \left(\frac{\partial v}{\partial t} + v \cdot \nabla v \right) = -\nabla P + \mu \nabla^2 v - \frac{GM_\star}{r} + f_{\text{other}} \quad (\text{D.1})$$

where f_{other} represents all other forces, and ∇^2 is the vector Laplace operator. In most astronomical settings we expect the viscous term to be unimportant. This can be seen by considering the ratio of the inertial term to the viscous term in the Navier-Stokes equation ?:

$$\frac{\rho v \cdot \nabla v}{\mu \nabla^2 v} \sim \frac{\rho U^2 / L}{\mu U / L^2} = \frac{UL}{\nu} \equiv Re, \quad (\text{D.2})$$

where U is a typical flow speed, L is the macroscopic length of the problem, Re is the Reynolds number, and $\nu \equiv \mu/\rho$ is called the kinematic viscosity. The shear viscosity is defined as

$$\mu \sim mv_T/\sigma \quad (\text{D.3})$$

where v_T is the thermal speed and σ is the typical collision cross section. Therefore the kinematic viscosity has the unit of velocity times length where,

$$\nu \sim v_T l \quad (\text{D.4})$$

and l is the collision mean free path. The Reynolds number can then be written as

$$Re \sim \frac{UL}{v_T l} \gg 1 \quad \text{when} \quad U \sim v_T. \quad (\text{D.5})$$

In other words, in stellar winds where the flow speeds are sonic or supersonic, the Reynolds number must be large and viscous forces are much less important than inertial effects. These high Reynolds number values (i.e., $Re \gg 10^3$ to 10^4) are associated with the onset of turbulence (?) and produce turbulent flows resulting in the formation of eddies and other flow instabilities.

The rate at which energy is fed into the largest eddies per unit mass equals

$$\epsilon \sim \frac{U^2}{L/U} = \frac{U^3}{L}. \quad (\text{D.6})$$

This energy can neither accumulate nor dissipate viscously, so the only other route for it is to be progressively transferred to eddies of smaller and smaller scales. If these smaller eddies have a scale λ and velocity v_λ then the energy that cascades from the large to the small per unit mass is

$$\epsilon \sim v_\lambda^3 / \lambda. \quad (\text{D.7})$$

Combining equation D.6 and D.7 gives Kolmogorov's law:

$$v_\lambda \sim U \left(\frac{\lambda}{L} \right)^{1/3} \quad (\text{D.8})$$

The process of transferring energy from large eddies down to smaller eddies continues until the scale of the turbulence is small enough for viscous action to become important and dissipation as heat to occur.

References

- ABIA, C., PALMERINI, S., BUSO, M. & CRISTALLO, S. (2012). Carbon and oxygen isotopic ratios in Arcturus and Aldebaran. Constraining the parameters for non-convective mixing on the red giant branch. *Astronomy & Astrophysics*, **548**, A55. (Not cited.)
- ADAMS, W.S. & MACCORMACK, E. (1935). Systematic Displacements of Lines in the Spectra of Certain Bright Stars. *Astrophysical Journal*, **81**, 119. (Not cited.)
- AIRAPETIAN, V., CARPENTER, K.G. & OFMAN, L. (2010). Winds from Luminous Late-type Stars. II. Broadband Frequency Distribution of Alfvén Waves. *Astrophysical Journal*, **723**, 1210–1218. (Not cited.)
- ALTENHOFF, W.J., MEZGER, P.G., WENDKER, H.J. & WESTERHOUT, G. (1960). Veröff. Sternwarte, Bonn. **59**, 48. (Not cited.)
- ALTENHOFF, W.J., HUCHTMEIER, W.K., SCHMIDT, J., SCHRAML, J.B. & STUMPF, P. (1986). Radio continuum observations of comet Halley. *Astronomy & Astrophysics*, **164**, 227–230. (Not cited.)
- ALTENHOFF, W.J., THUM, C. & WENDKER, H.J. (1994). Radio emission from stars: A survey at 250 GHz. *Astronomy & Astrophysics*, **281**, 161–183. (Not cited.)
- ASPLUND, M., GREVESSE, N., SAUVAL, A.J. & SCOTT, P. (2009). The Chemical Composition of the Sun. *Annual Review of Astronomy & Astrophysics*, **47**, 481–522. (Not cited.)
- AYRES, T. (2010a). A stellar perspective on chromospheres. *Memorie della Societa Astronomia Italiana*, **81**, 553. (Not cited.)
- AYRES, T.R. (1996). Thermal bifurcation as a driver of stellar surface inhomogeneities (review). In K.G. Strassmeier & J.L. Linsky, eds., *Stellar Surface Structure*, vol. 176 of *IAU Symposium*, 371. (Not cited.)
- AYRES, T.R. (2010b). StarCAT: A Catalog of Space Telescope Imaging Spectrograph Ultraviolet Echelle Spectra of Stars. *Astrophysical Journal Supplemental Series*, **187**, 149–171. (Cited on page 3.)
- AYRES, T.R. & LINSKY, J.L. (1975). Stellar model chromospheres. III - Arcturus /K2 III/. *Astrophysical Journal*, **200**, 660–674. (Not cited.)
- AYRES, T.R., LINSKY, J.L., VAIANA, G.S., GOLUB, L. & ROSNER, R. (1981). The cool half of the H-R diagram in soft X-rays. *Astrophysical Journal*, **250**, 293–299. (Not cited.)

REFERENCES

- AYRES, T.R., BROWN, A., HARPER, G.M., BENNETT, P.D., LINSKY, J.L., CARPENTER, K.G. & ROBINSON, R.D. (1997). Digging Deeper in the Coronal Graveyard. *Astrophysical Journal*, **491**, 876. (Not cited.)
- AYRES, T.R., BROWN, A. & HARPER, G.M. (2003). Buried Alive in the Coronal Graveyard. *Astrophysical Journal*, **598**, 610–625. (Not cited.)
- BAADE, R., KIRSCH, T., REIMERS, D., TOUSSAINT, F., BENNETT, P.D., BROWN, A. & HARPER, G.M. (1996). The Wind Outflow of zeta Aurigae: A Model Revision Using Hubble Space Telescope Spectra. *Astrophysical Journal*, **466**, 979. (Not cited.)
- BEASLEY, A.J., STEWART, R.T. & CARTER, B.D. (1992). Non-thermal radio emission from post-main-sequence stars. *Monthly Notices of the Royal Astronomical Society*, **254**, 1–6. (Not cited.)
- BECK, P.G., BEDDING, T.R., MOSSER, B., STELLO, D., GARCIA, R.A., KALLINGER, T., HEKKER, S., ELSWORTH, Y., FRANDSEN, S., CARRIER, F., DE RIDDER, J., AERTS, C., WHITE, T.R., HUBER, D., DUPRET, M.A., MONTALBÁN, J., MIGLIO, A., NOELS, A., CHAPLIN, W.J., KJELDSSEN, H., CHRISTENSEN-DALSGAARD, J., GILLILAND, R.L., BROWN, T.M., KAWALER, S.D., MATHUR, S. & JENKINS, J.M. (2011). Kepler Detected Gravity-Mode Period Spacings in a Red Giant Star. *Science*, **332**, 205–. (Not cited.)
- BECKLIN, E.E. & GEHRZ, R.D. (2009). SOFIA: Stratospheric Observatory for Infrared Astronomy. In D. C. Lis, J. E. Vaillancourt, P. F. Goldsmith, T. A. Bell, N. Z. Scoville, & J. Zmuidzinas, ed., *Submillimeter Astrophysics and Technology: a Symposium Honoring Thomas G. Phillips*, vol. 417 of *Astronomical Society of the Pacific Conference Series*, 101. (Not cited.)
- BEDDING, T.R., MOSSER, B., HUBER, D., MONTALBÁN, J., BECK, P., CHRISTENSEN-DALSGAARD, J., ELSWORTH, Y.P., GARCÍA, R.A., MIGLIO, A., STELLO, D., WHITE, T.R., DE RIDDER, J., HEKKER, S., AERTS, C., BARBAN, C., BELKACEM, K., BROOMHALL, A.M., BROWN, T.M., BUZASI, D.L., CARRIER, F., CHAPLIN, W.J., DI MAURO, M.P., DUPRET, M.A., FRANDSEN, S., GILLILAND, R.L., GOUPIL, M.J., JENKINS, J.M., KALLINGER, T., KAWALER, S., KJELDSSEN, H., MATHUR, S., NOELS, A., AGUIRRE, S.V. & VENTURA, P. (2011). Gravity modes as a way to distinguish between hydrogen- and helium-burning red giant stars. *Nature*, **471**, 608–611. (Not cited.)
- BEDECARRAX, I., PETIT, P., AURIÈRE, M., GRUNHUT, J., WADE, G., CHIAVASSA, A., DONATI, J.F., KONSTANTINOVA-ANTOVA, R. & PERRIN, G. (2013). Long-term spectropolarimetric monitoring of the cool supergiant betelgeuse. In *EAS Publications Series*, vol. 60 of *EAS Publications Series*, 161–165. (Not cited.)
- BERGER, T.E. & TITLE, A.M. (2001). On the Relation of G-Band Bright Points to the Photospheric Magnetic Field. *Astrophysical Journal*, **553**, 449–469. (Not cited.)
- BERIO, P., MERLE, T., THÉVENIN, F., BONNEAU, D., MOURARD, D., CHESNEAU, O., DELAAS, O., LIGI, R., NARDETTO, N., PERRAUT, K., PICHON, B., STEE, P., TALLON-BOSC, I., CLAUSSE, J.M., SPANG, A., MCALISTER, H., TEN BRUMMELAAR, T., STURMANN, J., STURMANN, L., TURNER, N., FARRINGTON, C. & GOLDFINGER, P.J. (2011). Chromosphere of K giant stars. Geometrical extent and spatial structure detection. *Astronomy & Astrophysics*, **535**, A59. (Not cited.)

REFERENCES

- BERNAT, A.P. & LAMBERT, D.L. (1976). K I 7699 A emission from the Betelgeuse shell. *Astrophysical Journal*, **210**, 395–401. (Not cited.)
- BERNAT, A.P., HALL, D.N.B., HINKLE, K.H. & RIDGWAY, S.T. (1979). Observations of CO circumstellar absorption in the 4.6 micron spectrum of Alpha Orionis. *Astrophysical Journal Letters*, **233**, L135–L139. (Not cited.)
- BLACKWELL, D.E., BOOTH, A.J., PETFORD, A.D., LEGGETT, S.K., MOUNTAIN, C.M. & SELBY, M.J. (1986). The infrared flux method and its use for study of Alpha Boo, MU HER and Beta Dra; relation to the VEGA 1.2-5 micron infrared excess. *Monthly Notices of the Royal Astronomical Society*, **221**, 427–443. (Not cited.)
- BLUNDELL, R. (2007). The Submillimeter Array. In *IEEE MTT-S International Microwave Symposium Digest*, 1857–1860. (Not cited.)
- BOCK, D.C.J. (2006). CARMA: Combined Array for Research in Millimeter-Wave Astronomy. In D.C. Backer, J.M. Moran & J.L. Turner, eds., *Revealing the Molecular Universe: One Antenna is Never Enough*, vol. 356 of *Astronomical Society of the Pacific Conference Series*, 17. (Not cited.)
- BOICE, D.C., KUHN, J.R., ROBINSON, R.D. & WORDEN, S.P. (1981). Possible radio flaring activity on a late-type giant star, Alpha Ceti. *Astrophysical Journal Letters*, **245**, L71–L74. (Not cited.)
- BOOKBINDER, J.A., STENCEL, R.E., DRAKE, S.A., SIMON, T., LINSKY, J.L. & FLORKOWSKI, D. (1987). VLA Observations of Rapid 6 cm Flux Variations in α Ori. In J. L. Linsky & R. E. Stencel, ed., *Cool Stars, Stellar Systems and the Sun*, vol. 291 of *Lecture Notes in Physics*, Berlin Springer Verlag, 337. (Not cited.)
- BOOTHROYD, A.I. & SACKMANN, I.J. (1999). The CNO Isotopes: Deep Circulation in Red Giants and First and Second Dredge-up. *Astrophysical Journal*, **510**, 232–250. (Not cited.)
- BOTTLINGER, C.F. (1911). Die Bahn des spektroskopischen Doppelsternes α Orionis. *Astronomische Nachrichten*, **187**, 33. (Not cited.)
- BOWERS, P.F. & KNAPP, G.R. (1987). Detection of H I emission from the circumstellar envelope of Alpha Orionis. *Astrophysical Journal*, **315**, 305–314. (Not cited.)
- BRACEWELL, R.N. (2000). *The Fourier Transform and Its Applications*. Columbus, OH: McGraw Hill, 3rd edn. (Not cited.)
- BROWN, K.I.T. (2007). Long-Term Spectroscopic and Precise Radial Velocity Monitoring of Arcturus. *Publications of the Astronomical Society of the Pacific*, **119**, 237–237. (Not cited.)
- BROWN, K.I.T., GRAY, D.F. & BALIUNAS, S.L. (2008). Long-Term Spectroscopic Monitoring of Arcturus. *Astrophysical Journal*, **679**, 1531–1540. (Not cited.)
- BUSCHER, D.F., BALDWIN, J.E., WARNER, P.J. & HANIFF, C.A. (1990). Detection of a bright feature on the surface of Betelgeuse. *Monthly Notices of the Royal Astronomical Society*, **245**, 7P–11P. (Not cited.)

REFERENCES

- CARILLI, C.L., HOLDAWAY, M.A. & SOWINSKI, K.P. (1996). Faster Switching at the VLA. In *VLA Scientific Memo. No. 169*, vol. 169 of *VLA memos*. (Not cited.)
- CARLSON, B. (2001). Refined EVLA WIDAR Correlator Architecture. In *NRC-EVLA Memo 014*, vol. 14 of *NRC-EVLA Memos*. (Not cited.)
- CARLSSON, M. (1986). A computer program for solving multi-level non-LTE radiative transfer-problems in moving or static atmospheres. *Uppsala Astronomical Observatory Reports*, **33**. (Not cited.)
- CARPENTER, K.G. & ROBINSON, R.D. (1996). HST Studies of Carbon and K-M Giant/Supergiant Stars. In P. Benvenuti, F.D. Macchetto & E.J. Schreier, eds., *Science with the Hubble Space Telescope - II*, 418. (Not cited.)
- CARPENTER, K.G. & ROBINSON, R.D. (1997). GHRS Observations of Cool, Low-Gravity Stars. III. Plasma Flows and Turbulence in the Outer Atmosphere of alpha Orionis (M2 Iab). *Astrophysical Journal*, **479**, 970–+. (Not cited.)
- CARPENTER, K.G., ROBINSON, R.D., HARPER, G.M., BENNETT, P.D., BROWN, A. & MULLAN, D.J. (1999). GHRS Observations of Cool, Low-Gravity Stars. V. The Outer Atmosphere and Wind of the Nearby K Supergiant lambda Velorum. *Astrophysical Journal*, **521**, 382–406. (Not cited.)
- CASSINELLI, J.P. & HARTMANN, L. (1977). The effect of winds and coronae of hot stars on the infrared and radio continua. *Astrophysical Journal*, **212**, 488–493. (Not cited.)
- CASTOR, J.I. (2004). *Radiation Hydrodynamics*. (Not cited.)
- CASTOR, J.I., ABBOTT, D.C. & KLEIN, R.I. (1975). Radiation-driven winds in Of stars. *Astrophysical Journal*, **195**, 157–174. (Not cited.)
- CHAPMAN, R.D. (1981). The 1979-1980 eclipse of Zeta Aurigae. I - The circumstellar envelope. *Astrophysical Journal*, **248**, 1043–1052. (Not cited.)
- CHAPMAN, S. & ZIRIN, H. (1957). Notes on the Solar Corona and the Terrestrial Ionosphere. *Smithsonian Contributions to Astrophysics*, **2**, 1. (Not cited.)
- CHARBONNEAU, P. & MACGREGOR, K.B. (1995). Stellar Winds with Non-WKB Alfven Waves. II. Wind Models for Cool, Evolved Stars. *Astrophysical Journal*, **454**, 901. (Not cited.)
- CHARBONNEL, C. (1995). A Consistent Explanation for $^{12}\text{C}/^{13}\text{C}$, ^7Li and ^3He Anomalies in Red Giant Stars. *Astrophysical Journal Letters*, **453**, L41. (Not cited.)
- CHIAVASSA, A., HAUBOIS, X., YOUNG, J.S., PLEZ, B., JOSSELIN, E., PERRIN, G. & FREYTAG, B. (2010). Radiative hydrodynamics simulations of red supergiant stars. II. Simulations of convection on Betelgeuse match interferometric observations. *Astronomy & Astrophysics*, **515**, A12. (Not cited.)
- CHIOSI, C. & MAEDER, A. (1986). The evolution of massive stars with mass loss. *Annual Review of Astronomy & Astrophysics*, **24**, 329–375. (Not cited.)

REFERENCES

- COHEN, M., CARBON, D.F., WELCH, W.J., LIM, T., SCHULZ, B., MCMURRY, A.D., FORSTER, J.R. & GOORVITCH, D. (2005). Far-Infrared and Millimeter Continuum Studies of K Giants: α Bootis and α Tauri. *Astronomical Journal*, **129**, 2836–2848. (Not cited.)
- CORNWELL, T., BRAUN, R. & BRIGGS, D.S. (1999). Deconvolution. In G.B. Taylor, C.L. Carilli & R.A. Perley, eds., *Synthesis Imaging in Radio Astronomy II*, vol. 180 of *Astronomical Society of the Pacific Conference Series*, 151. (Not cited.)
- CORNWELL, T.J. (2008). Multiscale CLEAN Deconvolution of Radio Synthesis Images. *IEEE Journal of Selected Topics in Signal Processing*, **2**, 793–801. (Not cited.)
- CRANMER, S.R. (2012). Self-Consistent Models of the Solar Wind. *Space Science Reviews*, **172**, 145–156. (Not cited.)
- CRANMER, S.R. & SAAR, S.H. (2011). Testing a Predictive Theoretical Model for the Mass Loss Rates of Cool Stars. *Astrophysical Journal*, **741**, 54. (Not cited.)
- CRANMER, S.R. & VAN BALLEGOOIJEN, A.A. (2005). On the Generation, Propagation, and Reflection of Alfvén Waves from the Solar Photosphere to the Distant Heliosphere. *Astrophysical Journal Supplemental Series*, **156**, 265–293. (Not cited.)
- CRANMER, S.R., VAN BALLEGOOIJEN, A.A. & EDGAR, R.J. (2007). Self-consistent Coronal Heating and Solar Wind Acceleration from Anisotropic Magnetohydrodynamic Turbulence. *Astrophysical Journal Supplemental Series*, **171**, 520–551. (Not cited.)
- CROWLEY, C., ESPEY, B.R. & MCCANDLISS, S.R. (2008). EG And: Far Ultraviolet Spectroscopic Explorer and Hubble Space Telescope STIS Monitoring of an Eclipsing Symbiotic Binary. *Astrophysical Journal*, **675**, 711–722. (Not cited.)
- CROWLEY, C., ESPEY, B.R., HARPER, G.M. & ROCHE, J. (2009). Winds and Chromospheres of Cool (Super-) Giants. In E. Stempels, ed., *15th Cambridge Workshop on Cool Stars, Stellar Systems, and the Sun*, vol. 1094 of *American Institute of Physics Conference Series*, 267–274. (Not cited.)
- DALGARNO, A. & LAYZER, D. (1987). *Spectroscopy of Astrophysical Plasmas*. (Not cited.)
- DALGARNO, A. & MCCRAY, R.A. (1972). Heating and Ionization of HI Regions. *Annual Review of Astronomy & Astrophysics*, **10**, 375. (Not cited.)
- DANCHI, W.C., BESTER, M., DEGIACOMI, C.G., GREENHILL, L.J. & TOWNES, C.H. (1994). Characteristics of dust shells around 13 late-type stars. *Astronomical Journal*, **107**, 1469–1513. (Not cited.)
- DE GRAAUW, T., HELMICH, F.P., PHILLIPS, T.G., STUTZKI, J., CAUX, E., WHYBORN, N.D., DIELEMAN, P., ROELFSEMA, P.R., AARTS, H., ASSENDORP, R., BACHILLER, R., BAECHTOLD, W., BARCIA, A., BEINTEMA, D.A., BELITSKY, V., BENZ, A.O., BIEBER, R., BOOGERT, A., BORYS, C., BUMBLE, B., CAÏS, P., CARIS, M., CERULLI-IRELLI, P., CHATTOPADHYAY, G., CHEREDNICHENKO, S., CIECHANOWICZ, M., COEUR-JOLY, O., COMITO, C., CROS, A., DE JONGE, A., DE LANGE, G., DELFORGES, B., DELORME, Y., DEN BOGGENDE, T., DESBAT, J.M., DIEZ-GONZÁLEZ, C., DI GIORGIO, A.M., DUBBELDAM, L., EDWARDS, K., EGGENS, M., ERICKSON, N., EVERS, J., FICH, M., FINN, T.,

REFERENCES

- FRANKE, B., GAIER, T., GAL, C., GAO, J.R., GALLEGO, J.D., GAUFFRE, S., GILL, J.J., GLENZ, S., GOLSTEIN, H., GOULOOZE, H., GUNSING, T., GÜSTEN, R., HARTOGH, P., HATCH, W.A., HIGGINS, R., HONINGH, E.C., HUISMAN, R., JACKSON, B.D., JACOBS, H., JACOBS, K., JARCHOW, C., JAVADI, H., JELLEMA, W., JUSTEN, M., KARPOV, A., KASEMANN, C., KAWAMURA, J., KEIZER, G., KESTER, D., KLAPWIJK, T.M., KLEIN, T., KOLLBERG, E., KOOI, J., KOOIMAN, P.P., KOPF, B., KRAUSE, M., KRIEG, J.M., KRAMER, C., KRUIZENGA, B., KUHN, T., LAAUWEN, W., LAI, R., LARSSON, B., LEDUC, H.G., LEINZ, C., LIN, R.H., LISEAU, R., LIU, G.S., LOOSE, A., LÓPEZ-FERNANDEZ, I., LORD, S., LUINGE, W., MARSTON, A., MARTÍN-PINTADO, J., MAESTRINI, A., MAIWALD, F.W., MCCOEY, C., MEHDI, I., MEGEJ, A., MELCHIOR, M., MEINSMAN, L., MERKEL, H., MICHALSKA, M., MONSTEIN, C., MORATSCHKE, D., MORRIS, P., MULLER, H., MURPHY, J.A., NABER, A., NATALE, E., NOWOSIELSKI, W., NUZZOLO, F., OLBERG, M., OLBRICH, M., ORFEI, R., ORLEANSKI, P., OSSENKOPF, V., PEACOCK, T., PEARSON, J.C., PERON, I., PHILLIP-MAY, S., PIAZZO, L., PLANESAS, P., RATAJ, M., RAVERA, L., RISACHER, C., SALEZ, M., SAMOSKA, L.A., SARACENO, P., SCHIEDER, R., SCHLECHT, E., SCHLÖDER, F., SCHMÜLLING, F., SCHULTZ, M., SCHUSTER, K., SIEBERTZ, O., SMIT, H., SZCZERBA, R., SHIPMAN, R., STEINMETZ, E., STERN, J.A., STOKROOS, M., TEIPEN, R., TEYSSIER, D., TILS, T., TRAPPE, N., VAN BAAREN, C., VAN LEEUWEN, B.J., VAN DE STADT, H., VISSER, H., WILDEMAN, K.J., WAFELBAKKER, C.K., WARD, J.S., WESSELIUS, P., WILD, W., WULFF, S., WUNSCH, H.J., TIELENS, X., ZAAL, P., ZIRATH, H., ZMUIDZINAS, J. & ZWART, F. (2010). The Herschel-Heterodyne Instrument for the Far-Infrared (HIFI). *Astronomy & Astrophysics*, **518**, L6. (Not cited.)
- DECIN, L., VANDENBUSSCHE, B., WAELEKENS, C., DECIN, G., ERIKSSON, K., GUSTAFSSON, B., PLEZ, B. & SAUVAL, A.J. (2003). ISO-SWS calibration and the accurate modelling of cool-star atmospheres. IV. G9 to M2 stars. *Astronomy & Astrophysics*, **400**, 709–727. (Not cited.)
- DECIN, L., COX, N.L.J., ROYER, P., VAN MARLE, A.J., VANDENBUSSCHE, B., LADJAL, D., KERSCHBAUM, F., OTTENSAMER, R., BARLOW, M.J., BLOMMAERT, J.A.D.L., GOMEZ, H.L., GROENEWEGEN, M.A.T., LIM, T., SWINYARD, B.M., WAELEKENS, C. & TIELENS, A.G.G.M. (2012). The enigmatic nature of the circumstellar envelope and bow shock surrounding Betelgeuse as revealed by Herschel. I. Evidence of clumps, multiple arcs, and a linear bar-like structure. *Astronomy & Astrophysics*, **548**, A113. (Not cited.)
- DEHAES, S., BAUWENS, E., DECIN, L., ERIKSSON, K., RASKIN, G., BUTLER, B., DOWELL, C.D., ALI, B. & BLOMMAERT, J.A.D.L. (2011). Structure of the outer layers of cool standard stars. *Astronomy & Astrophysics*, **533**, A107. (Not cited.)
- DEUTSCH, A.J. (1956). The Circumstellar Envelope of Alpha Herculis. *Astrophysical Journal*, **123**, 210. (Not cited.)
- DI BENEDETTO, G.P. (1993). Empirical effective temperatures and angular diameters of stars cooler than the sun. *Astronomy & Astrophysics*, **270**, 315–334. (Not cited.)
- DRAINE, B.T. (2011). *Physics of the Interstellar and Intergalactic Medium*. (Not cited.)
- DRAKE, S. & LINSKY, J. (1986). Radio continuum emission from winds, chromospheres, and coronae of cool giants and supergiants. *Astronomical Journal*, **91**, 602–620. (Not cited.)

REFERENCES

- DRAKE, S.A. (1985). Modeling lines formed in the expanding chromospheres of red giants. In J.E. Beckman & L. Crivellari, eds., *Progress in stellar spectral line formation theory; Proceedings of the Advanced Research Workshop, Trieste, Italy, September 4-7, 1984 (A86-37976 17-90)*. Dordrecht, D. Reidel Publishing Co., 1985, p. 351-357., 351–357. (Cited on page 3.)
- DRAKE, S.A. & LINSKY, J.L. (1983a). First detection of winds in red giants by microwave continuum techniques. *Astrophysical Journal Letters*, **274**, L77–L81. (Not cited.)
- DRAKE, S.A. & LINSKY, J.L. (1983b). The formation of emission lines in the expanding chromospheres of luminous cool stars. I - The importance of atmospheric extension and partial redistribution effects. *Astrophysical Journal*, **273**, 299–308. (Not cited.)
- DRAKE, S.A., BOOKBINDER, J.A., FLORKOWSKI, D.R., LINSKY, J.L., SIMON, T. & STENCEL, R.E. (1992). Four Years of Monitoring a Orionis with the VLA: Where have all the Flares Gone? In M. S. Giampapa & J. A. Bookbinder, ed., *Cool Stars, Stellar Systems, and the Sun*, vol. 26 of *Astronomical Society of the Pacific Conference Series*, 455. (Not cited.)
- DULK, G.A. (1985). Radio emission from the sun and stars. *Annual Review of Astronomy & Astrophysics*, **23**, 169–224. (Not cited.)
- DUPREE, A.K., BALIUNAS, S.L., HARTMANN, L., GUINAN, E.F. & SONNEBORN, G. (1990). Alpha ORI - Evidence for pulsation. In C. Cacciari & G. Clementini, eds., *Confrontation Between Stellar Pulsation and Evolution*, vol. 11 of *Astronomical Society of the Pacific Conference Series*, 468–471. (Not cited.)
- DUPREE, A.K., LOBEL, A., YOUNG, P.R., AKE, T.B., LINSKY, J.L. & REDFIELD, S. (2005). A Far-Ultraviolet Spectroscopic Survey of Luminous Cool Stars. *Astrophysical Journal*, **622**, 629–652. (Not cited.)
- DZIEMBOWSKI, W.A., GOUGH, D.O., HOUEK, G. & SIENKIEWICZ, R. (2001). Oscillations of α UMa and other red giants. *Monthly Notices of the Royal Astronomical Society*, **328**, 601–610. (Not cited.)
- EATON, J.A. (2008). A Model for the Chromosphere/Wind of 31 Cygni and Its Implications for Single Stars. *Astronomical Journal*, **136**, 1964–1979. (Not cited.)
- EGGEN, O.J. (1971). The Arcturus Group. *Publications of the Astronomical Society of the Pacific*, **83**, 271. (Not cited.)
- EGGEN, O.J. (1996). Star Streams and Galactic Structure. *Astronomical Journal*, **112**, 1595. (Not cited.)
- EGGLETON, P.P., DEARBORN, D.S.P. & LATTANZIO, J.C. (2007). Compulsory Deep Mixing of ^3He and CNO Isotopes on the First Giant Branch. In R.J. Stancliffe, G. Houdek, R.G. Martin & C.A. Tout, eds., *Unsolved Problems in Stellar Physics: A Conference in Honor of Douglas Gough*, vol. 948 of *American Institute of Physics Conference Series*, 27–34. (Not cited.)
- EKERS, R. (1982). Common Map Defects. In A.R. Thompson & L.R. D’Addario, eds., *Synthesis Mapping*, 12. (Not cited.)

REFERENCES

- ENGARGIOLA, G.A., NAVARRINI, A., PLAMBECK, R.L. & WADEFALK, N. (2004). Simple 1-mm receivers with fixed tuned double sideband SIS mixer and wideband InP MMIC amplifier. In C.M. Bradford, P.A.R. Ade, J.E. Aguirre, J.J. Bock, M. Dragovan, L. Duband, L. Earle, J. Glenn, H. Matsuhara, B.J. Naylor, H.T. Nguyen, M. Yun & J. Zmuidzinas, eds., *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, vol. 5498 of *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, 556–566. (Not cited.)
- FALCETA-GONÇALVES, D., VIDOTTO, A.A. & JATENCO-PEREIRA, V. (2006). On the magnetic structure and wind parameter profiles of Alfvén wave driven winds in late-type supergiant stars. *Monthly Notices of the Royal Astronomical Society*, **368**, 1145–1150. (Not cited.)
- FAMAHEY, B., JORISSEN, A., LURI, X., MAYOR, M., UDRY, S., DEJONGHE, H. & TURON, C. (2005). Local kinematics of K and M giants from CORAVEL/Hipparcos/Tycho-2 data. Revisiting the concept of superclusters. *Astronomy & Astrophysics*, **430**, 165–186. (Not cited.)
- FRACASSINI, M., PASINETTI, L.E. & MANZOLINI, F. (1981). Catalogue of apparent diameters and absolute radii of stars /CADARS/. *Astronomy & Astrophysics Supplemental*, **45**, 145–174. (Not cited.)
- GEHRZ, R.D. & WOOLF, N.J. (1971). Mass Loss from M Stars. *Astrophysical Journal*, **165**, 285. (Not cited.)
- GILLILAND, R.L. & DUPREE, A.K. (1996). First Image of the Surface of a Star with the Hubble Space Telescope. *Astrophysical Journal Letters*, **463**, L29. (Not cited.)
- GLASSGOLD, A.E. & HUGGINS, P.J. (1986). The ionization structure of the circumstellar envelope of Alpha Orionis. *Astrophysical Journal*, **306**, 605–617. (Not cited.)
- GOLDBERG, L. (1984). The variability of alpha Orionis. *Publications of the Astronomical Society of the Pacific*, **96**, 366–371. (Not cited.)
- GOLDBERG, L., RAMSEY, L., TESTERMAN, L. & CARBON, D. (1975). High-resolution profiles of sodium and potassium lines in alpha Orionis. *Astrophysical Journal*, **199**, 427–431. (Not cited.)
- GOLDREICH, P. & SCOVILLE, N. (1976). OH-IR stars. I - Physical properties of circumstellar envelopes. *Astrophysical Journal*, **205**, 144–154. (Not cited.)
- GRADSHTEYN, I.S. & RYZHIK, I.M. (1994). *Table of integrals, series and products*. (Not cited.)
- GRAY, D.F. (2008). Mass Motions in the Photosphere of Betelgeuse. *Astronomical Journal*, **135**, 1450–1458. (Not cited.)
- GRAY, R.O., CORBALLY, C.J., GARRISON, R.F., MCFADDEN, M.T., BUBAR, E.J., MCGAHEE, C.E., O'DONOGHUE, A.A. & KNOX, E.R. (2006). Contributions to the Nearby Stars (NStars) Project: Spectroscopy of Stars Earlier than M0 within 40 pc-The Southern Sample. *Astronomical Journal*, **132**, 161–170. (Not cited.)
- GRIFFIN, R.E.M. (1996). Arcturus and human evolution. *The Observatory*, **116**, 404–405. (Not cited.)

REFERENCES

- GRIFFIN, R.E.M. & LYNAS-GRAY, A.E. (1999). The Effective Temperature of Arcturus. *Astronomical Journal*, **117**, 2998–3006. (Not cited.)
- GUESTEN, R., HARTOGH, P., HUEBERS, H.W., GRAF, U.U., JACOBS, K., ROESER, H.P., SCHAEFER, F., SCHIEDER, R.T., STARK, R., STUTZKI, J., VAN DER WAL, P. & WUNSCH, A. (2000). GREAT: the first-generation German heterodyne receiver for SOFIA. In R. K. Melugin & H.-P. Röser, ed., *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, vol. 4014, 23–30. (Not cited.)
- HABING, H.J. (1996). Circumstellar envelopes and Asymptotic Giant Branch stars. *Astronomy & Astrophysics Review*, **7**, 97–207. (Not cited.)
- HAISCH, B.M., LINSKY, J.L. & BASRI, G.S. (1980). Outer atmospheres of cool stars. IV - A discussion of cool stellar wind models. *Astrophysical Journal*, **235**, 519–533. (Not cited.)
- HAMAKER, J.P., BREGMAN, J.D. & SAULT, R.J. (1996). Understanding radio polarimetry. I. Mathematical foundations. *Astronomy & Astrophysics Supplemental*, **117**, 137–147. (Not cited.)
- HARPER, G.M. (1988). *The outer atmospheres of hybrid giants*. Ph.D. thesis, Oxford Univ. (England). (Not cited.)
- HARPER, G.M. (1994). A Computer Program for Solving Multilevel Non-Lte Radiative Transfer Problems in Spherical Geometry in Moving or Static Atmospheres. *Monthly Notices of the Royal Astronomical Society*, **268**, 894. (Not cited.)
- HARPER, G.M. (2010). Betelgeuse: A Case Study of an Inhomogeneous Extended Atmosphere. In C. Leitherer, P.D. Bennett, P.W. Morris & J.T. Van Loon, eds., *Hot and Cool: Bridging Gaps in Massive Star Evolution*, vol. 425 of *Astronomical Society of the Pacific Conference Series*, 152. (Not cited.)
- HARPER, G.M. (2013). Atmospheric structure and dynamics: the spatial and temporal domains. In P. Kervella, T. Le Bertre & G. Perrin, eds., *EAS Publications Series*, vol. 60 of *EAS Publications Series*, 59–68. (Not cited.)
- HARPER, G.M., BROWN, A. & LIM, J. (2001). A Spatially Resolved, Semiempirical Model for the Extended Atmosphere of α Orionis (M2 Iab). *Astrophysical Journal*, **551**, 1073–1098. (Not cited.)
- HARPER, G.M., BROWN, A., BENNETT, P.D., BAADE, R., WALDER, R. & HUMMEL, C.A. (2005). VLA Observations of ζ Aurigae: Confirmation of the Slow Acceleration Wind Density Structure. *Astronomical Journal*, **129**, 1018–1034. (Not cited.)
- HARPER, G.M., BROWN, A. & GUINAN, E.F. (2008). A New VLA-Hipparcos Distance to Betelgeuse and its Implications. *Astronomical Journal*, **135**, 1430–1440. (Not cited.)
- HARPER, G.M., RICHTER, M.J., RYDE, N., BROWN, A., BROWN, J., GREATHOUSE, T.K. & STRONG, S. (2009). Texas Observations of M Supergiants: Dynamics and Thermodynamics of Wind Acceleration. *Astrophysical Journal*, **701**, 1464–1483. (Not cited.)

REFERENCES

- HARPER, G.M., O’RIAIN, N. & AYRES, T.R. (2013). Chromospheric thermal continuum millimetre emission from non-dusty K and M red giants. *Monthly Notices of the Royal Astronomical Society*, **428**, 2064–2073. (Not cited.)
- HARRIS, M.J. & LAMBERT, D.L. (1984). Oxygen isotopes in the atmospheres of Betelgeuse and Antares. *Astrophysical Journal*, **281**, 739–745. (Not cited.)
- HARTMANN, L. & AVRETT, E.H. (1984). On the extended chromosphere of Alpha Orionis. *Astrophysical Journal*, **284**, 238–249. (Not cited.)
- HARTMANN, L. & MACGREGOR, K.B. (1980). Momentum and energy deposition in late-type stellar atmospheres and winds. *Astrophysical Journal*, **242**, 260–282. (Not cited.)
- HARTMANN, L., AVRETT, E. & EDWARDS, S. (1982). Wave-driven winds from cool stars. II - Models for T Tauri stars. *Astrophysical Journal*, **261**, 279–292. (Not cited.)
- HATZES, A.P. & COCHRAN, W.D. (1993). Long-period radial velocity variations in three K giants. *Astrophysical Journal*, **413**, 339–348. (Not cited.)
- HATZES, A.P. & COCHRAN, W.D. (1998). On the nature of the radial velocity variability of Aldebaran - A search for spectral line bisector variations. *Monthly Notices of the Royal Astronomical Society*, **293**, 469. (Not cited.)
- HAUBOIS, X., PERRIN, G., LACOUR, S., VERHOELST, T., MEIMON, S., MUGNIER, L., THIÉBAUT, E., BERGER, J.P., RIDGWAY, S.T., MONNIER, J.D., MILLAN-GABET, R. & TRAUB, W. (2009). Imaging the spotty surface of μ ASTROBJ₁Betelgeuse/ μ ASTROBJ₂ in the H band. *Astronomy & Astrophysics*, **508**, 923–932. (Not cited.)
- HINKLE, K.H., LAMBERT, D.L. & SNELL, R.L. (1976). The C-12/C-13 ratio in stellar atmospheres. VI - Five luminous cool stars. *Astrophysical Journal*, **210**, 684–693. (Not cited.)
- HJELLMING, R.M. & NEWELL, R.T. (1983). Radio emission from Antares and an ionized cavity in its wind. *Astrophysical Journal*, **275**, 704–708. (Not cited.)
- HÖGBOM, J.A. (1974). Aperture Synthesis with a Non-Regular Distribution of Interferometer Baselines. *Astronomy & Astrophysics Supplemental*, **15**, 417. (Not cited.)
- HOLLENBACH, D. & MCKEE, C.F. (1989). Molecule formation and infrared emission in fast interstellar shocks. III - Results for J shocks in molecular clouds. *Astrophysical Journal*, **342**, 306–336. (Not cited.)
- HOLLWEG, J.V. (1973). ALFVtN Waves in a Two-Fluid Model of the Solar Wind. *Astrophysical Journal*, **181**, 547–566. (Not cited.)
- HOLZER, T.E. & MACGREGOR, K.B. (1985). Mass loss mechanisms for cool, low-gravity stars. In M. Morris & B. Zuckerman, eds., *Mass Loss from Red Giants*, vol. 117 of *Astrophysics and Space Science Library*, 229–255. (Not cited.)
- HOLZER, T.E., FLA, T. & LEER, E. (1983). Alfven waves in stellar winds. *Astrophysical Journal*, **275**, 808–835. (Not cited.)

REFERENCES

- HOogerwerf, R., de Bruijne, J.H.J. & de Zeeuw, P.T. (2000). The Origin of Runaway Stars. *Astrophysical Journal Letters*, **544**, L133–L136. (Not cited.)
- HUBER, K.P. & HERZBERG, G. (1979). *Molecular Spectra and Molecular Structure IV. Constants of Diatomic Molecules*. New York: Van Nostrand Reinhold. (Not cited.)
- HUGGINS, P.J. (1987). CO in the circumstellar envelope of Alpha Orionis. *Astrophysical Journal*, **313**, 400–407. (Not cited.)
- HUGGINS, P.J., BACHILLER, R., COX, P. & FORVEILLE, T. (1994). Neutral carbon in the circumstellar envelope of alpha Orionis. *Astrophysical Journal Letters*, **424**, L127–L130. (Not cited.)
- HUMMER, D.G. (1988). A fast and accurate method for evaluating the nonrelativistic free-free Gaunt factor for hydrogenic ions. *Astrophysical Journal*, **327**, 477–484. (Not cited.)
- IBEN, I., JR. (1967). Stellar Evolution Within and off the Main Sequence. *Annual Review of Astronomy & Astrophysics*, **5**, 571. (Not cited.)
- IBEN, I., JR. (1993). On Why Intermediate-Mass Stars Become Giants after the Exhaustion of Hydrogen in Their Cores. *Astrophysical Journal*, **415**, 767. (Not cited.)
- IROSHNIKOV, P.S. (1964). Turbulence of a Conducting Fluid in a Strong Magnetic Field. *Soviet Astronomy*, **7**, 566. (Not cited.)
- JACKSON, N. (2008). Principles of Interferometry. In F. Bacciotti, L. Testi & E. Whelan, eds., *Jets from Young Stars II*, vol. 742 of *Lecture Notes in Physics*, Berlin Springer Verlag, 193. (Not cited.)
- JATENCO-PEREIRA, V. & O'PHER, R. (1989). Effect of diverging magnetic fields on mass loss in late-type giant stars. *Astronomy & Astrophysics*, **209**, 327–336. (Not cited.)
- JEWELL, P.R., SNYDER, L.E., WALMSLEY, C.M., WILSON, T.L. & GENSHEIMER, P.D. (1991). Observational properties of $V = 1$, $J = 1 - 0$ SiO masers. *Astronomy & Astrophysics*, **242**, 211–234. (Not cited.)
- JOHNSON, H.R. (1973). Cooling effect of CO in stellar atmospheres. *Astrophysical Journal*, **180**, 81–89. (Not cited.)
- JONES, M.H. (2008). The incidence of mid-infrared excesses in G and K giants. *Monthly Notices of the Royal Astronomical Society*, **387**, 845–855. (Not cited.)
- JOSSELIN, E. & PLEZ, B. (2007). Atmospheric dynamics and the mass loss process in red supergiant stars. *Astronomy & Astrophysics*, **469**, 671–680. (Not cited.)
- JUDGE, P.G. (1986). Constraints on the outer atmospheric structure of late-type giant stars with IUE - Methods and application to Arcturus (Alpha Boo K2III). *Monthly Notices of the Royal Astronomical Society*, **221**, 119–153. (Not cited.)
- JUDGE, P.G. & CARPENTER, K.G. (1998). On Chromospheric Heating Mechanisms of “Basal Flux” Stars. *Astrophysical Journal*, **494**, 828. (Not cited.)

REFERENCES

- KALER, J.B., IBEN, I., JR. & BECKER, S.A. (1978). On the enhancement of helium and nitrogen in planetary nebulae. *Astrophysical Journal Letters*, **224**, L63–L66. (Not cited.)
- KALLINGER, T., WEISS, W.W., BARBAN, C., BAUDIN, F., CAMERON, C., CARRIER, F., DE RIDDER, J., GOUPIL, M.J., GRUBERBAUER, M., HATZES, A., HEKKER, S., SAMADI, R. & DELEUIL, M. (2010). Oscillating red giants in the CoRoT exofield: asteroseismic mass and radius determination. *Astronomy & Astrophysics*, **509**, A77. (Not cited.)
- KAMIŃSKI, T., GOTTLIEB, C.A., SCHMIDT, M.R., PATEL, N.A., YOUNG, K.H., MENTEN, K.M., BRÜNKEN, S., MÜLLER, H.S.P., WINTERS, J.M. & MCCARTHY, M.C. (2013). Dust-forming molecules in VY Canis Majoris (and Betelgeuse). In *EAS Publications Series*, vol. 60 of *EAS Publications Series*, 191–198. (Not cited.)
- KARZAS, W.J. & LATTER, R. (1961). Electron Radiative Transitions in a Coulomb Field. *Astrophysical Journal Supplemental Series*, **6**, 167. (Not cited.)
- KELCH, W.L., CHANG, S.H., FURENLID, I., LINSKY, J.L., BASRI, G.S., CHIU, H.Y. & MARAN, S.P. (1978). Stellar model chromospheres. VII - Capella /G5 III +/, Pollux /K0 III/, and Aldebaran /K5 III/. *Astrophysical Journal*, **220**, 962–979. (Not cited.)
- KERVELLA, P., VERHOELST, T., RIDGWAY, S.T., PERRIN, G., LACOUR, S., CAMI, J. & HAUBOIS, X. (2009). The close circumstellar environment of Betelgeuse. Adaptive optics spectro-imaging in the near-IR with VLT/NACO. *Astronomy & Astrophysics*, **504**, 115–125. (Not cited.)
- KERVELLA, P., PERRIN, G., CHIAVASSA, A., RIDGWAY, S.T., CAMI, J., HAUBOIS, X. & VERHOELST, T. (2011). The close circumstellar environment of Betelgeuse. II. Diffraction-limited spectro-imaging from 7.76 to 19.50 μm with VLT/VISIR. *Astronomy & Astrophysics*, **531**, A117. (Not cited.)
- KNAPP, G.R. & MORRIS, M. (1985). Mass loss from evolved stars. III - Mass loss rates for fifty stars from CO J = 1-0 observations. *Astrophysical Journal*, **292**, 640–669. (Not cited.)
- KNAPP, G.R., PHILLIPS, T.G. & HUGGINS, P.J. (1980). Detection of CO Emission at 1.3-MILLIMETERS from the Betelgeuse Circumstellar Shell. *Astrophysical Journal Letters*, **242**, L25+. (Not cited.)
- KOLMOGOROV, A. (1941). The Local Structure of Turbulence in Incompressible Viscous Fluid for Very Large Reynolds' Numbers. *Akademiia Nauk SSSR Doklady*, **30**, 301–305. (Not cited.)
- KRAUS, J.D., TIURI, M., RAISANEN, A.V. & CARR, T.D. (1986). *Radio astronomy receivers*. (Not cited.)
- KROGULEC, M. (1988). Alfven wave driven wind - Models of Alpha Boo atmosphere. *Acta Astronomica*, **38**, 107–126. (Not cited.)
- KROGULEC, M. (1989). Alfven wave driven wind. II - Models of cool giant and supergiant stars. *Acta Astronomica*, **39**, 51–67. (Not cited.)

REFERENCES

- LACOUR, S., MEIMON, S., THIÉBAUT, E., PERRIN, G., VERHOELST, T., PEDRETTI, E., SCHULLER, P.A., MUGNIER, L., MONNIER, J., BERGER, J.P., HAUBOIS, X., PONCELET, A., LE BESNERAIS, G., ERIKSSON, K., MILLAN-GABET, R., RAGLAND, S., LACASSE, M. & TRAUB, W. (2008). The limb-darkened Arcturus: imaging with the IOTA/IONIC interferometer. *Astronomy & Astrophysics*, **485**, 561–570. (Not cited.)
- LAMB, S.A., IBEN, I., JR. & HOWARD, W.M. (1976). On the evolution of massive stars through the core carbon-burning phase. *Astrophysical Journal*, **207**, 209–232. (Not cited.)
- LAMBERT, D.L. & VANDEN BOUT, P.A. (1978). Constraints on the properties of circumstellar shells from observations of thermal CO and SiO millimeter line emission. *Astrophysical Journal*, **221**, 854–860. (Not cited.)
- LAMBERT, D.L., DEARBORN, D.S. & SNEDEN, C. (1974). The C-12/C-13 ratio in stellar atmospheres. II - CN and CO in alpha Orionis. *Astrophysical Journal*, **193**, 621–630. (Not cited.)
- LAMBERT, D.L., BROWN, J.A., HINKLE, K.H. & JOHNSON, H.R. (1984). Carbon, nitrogen, and oxygen abundances in Betelgeuse. *Astrophysical Journal*, **284**, 223–237. (Not cited.)
- LAMERS, H.J.G.L.M. (1998). Stellar Wind Theories. *Astrophysics & Space Science*, **260**, 81–100. (Not cited.)
- LAMERS, H.J.G.L.M. & CASSINELLI, J.P. (1999). *Introduction to Stellar Winds*. (Not cited.)
- LAMERS, H.J.G.L.M., CERRUTI-SOLA, M. & PERINOTTO, M. (1987). The 'SEI' method for accurate and efficient calculations of line profiles in spherically symmetric stellar winds. *Astrophysical Journal*, **314**, 726–738. (Not cited.)
- LEBZELTER, T., HEITER, U., ABIA, C., ERIKSSON, K., IRELAND, M., NEILSON, H., NOWOTNY, W., MALDONADO, J., MERLE, T., PETERSON, R., PLEZ, B., SHORT, C.I., WAHLGREN, G.M., WORLEY, C., ARINGER, B., BLADH, S., DE LAVERNY, P., GOSWAMI, A., MORA, A., NORRIS, R.P., RECIO-BLANCO, A., SCHOLZ, M., THÉVENIN, F., TSUJI, T., KORDOPATIS, G., MONTESINOS, B. & WING, R.F. (2012). Comparative modelling of the spectra of cool giants. *Astronomy & Astrophysics*, **547**, A108. (Not cited.)
- LEHE, R., PARRISH, I.J. & QUATAERT, E. (2009). The Heating of Test Particles in Numerical Simulations of Alfvénic Turbulence. *Astrophysical Journal*, **707**, 404–419. (Not cited.)
- LEVESQUE, E.M. (2010). The Physical Properties of Red Supergiants. In C. Leitherer, P.D. Bennett, P.W. Morris & J.T. Van Loon, eds., *Hot and Cool: Bridging Gaps in Massive Star Evolution*, vol. 425 of *Astronomical Society of the Pacific Conference Series*, 103. (Not cited.)
- LEVESQUE, E.M., MASSEY, P., OLSEN, K.A.G., PLEZ, B., JOSSELIN, E., MAEDER, A. & MEYNET, G. (2005). The Effective Temperature Scale of Galactic Red Supergiants: Cool, but Not As Cool As We Thought. *Astrophysical Journal*, **628**, 973–985. (Not cited.)
- LIGHTHILL, M.J. (1952). On Sound Generated Aerodynamically. I. General Theory. *Royal Society of London Proceedings Series A*, **211**, 564–587. (Not cited.)

REFERENCES

- LIM, J., CARILLI, C.L., WHITE, S.M., BEASLEY, A.J. & MARSON, R.G. (1998). Large convection cells as the source of Betelgeuse’s extended atmosphere. *Nature*, **392**, 575–577. (Not cited.)
- LINSKY, J.L. & HAISCH, B.M. (1979). Outer atmospheres of cool stars. I - The sharp division into solar-type and non-solar-type stars. *Astrophysical Journal Letters*, **229**, L27–L32. (Not cited.)
- LUCK, R.E. & CHALLENGER, S.L. (1995). Chemical Abundances for Very Strong-Lined Giants. *Astronomical Journal*, **110**, 2968. (Not cited.)
- LUTTERMOSER, D.G., JOHNSON, H.R. & EATON, J. (1994). The chromospheric structure of the cool giant star G Herculis. *Astrophysical Journal*, **422**, 351–365. (Not cited.)
- MACCHETTO, F. & PENSTON, M.V. (1978). The International Ultraviolet Explorer. *ESA Bulletin*, **13**, 9–14. (Not cited.)
- MAEDER, A. & MEYNET, G. (1987). Grids of evolutionary models of massive stars with mass loss and overshooting - Properties of Wolf-Rayet stars sensitive to overshooting. *Astronomy & Astrophysics*, **182**, 243–263. (Not cited.)
- MASSAROTTI, A., LATHAM, D.W., STEFANIK, R.P. & FOGEL, J. (2008). Rotational and Radial Velocities for a Sample of 761 HIPPARCOS Giants and the Role of Binarity. *Astronomical Journal*, **135**, 209–231. (Not cited.)
- MATHISEN, R. (1984). Photo cross-sections for stellar atmosphere calculations – compilation of references and data. Inst. Theor. Astrophys. Univ. Oslo, publ. Series No. 1. (Not cited.)
- MATTHAEUS, W.H., ZANK, G.P., OUGHTON, S., MULLAN, D.J. & DMITRUK, P. (1999). Coronal Heating by Magnetohydrodynamic Turbulence Driven by Reflected Low-Frequency Waves. *Astrophysical Journal Letters*, **523**, L93–L96. (Not cited.)
- McMULLIN, J.P., WATERS, B., SCHIEBEL, D., YOUNG, W. & GOLAP, K. (2007). CASA Architecture and Applications. In R.A. Shaw, F. Hill & D.J. Bell, eds., *Astronomical Data Analysis Software and Systems XVI*, vol. 376 of *Astronomical Society of the Pacific Conference Series*, 127. (Not cited.)
- McMURRY, A.D. (1999). The outer atmosphere of Tau - I. A new chromospheric model. *Monthly Notices of the Royal Astronomical Society*, **302**, 37–47. (Not cited.)
- McWILLIAM, A. (1990). High-resolution spectroscopic survey of 671 GK giants. I - Stellar atmosphere parameters and abundances. *Astrophysical Journal Supplemental Series*, **74**, 1075–1128. (Not cited.)
- MERLINE, W.J. (1999). Precise Velocity Observation of K-Giants: Evidence for Solar-Like Oscillations in Arcturus. In J.B. Hearnshaw & C.D. Scarfe, eds., *IAU Colloq. 170: Precise Stellar Radial Velocities*, vol. 185 of *Astronomical Society of the Pacific Conference Series*, 187. (Not cited.)
- MEYNET, G. & MAEDER, A. (2003). Stellar evolution with rotation. X. Wolf-Rayet star populations at solar metallicity. *Astronomy & Astrophysics*, **404**, 975–990. (Not cited.)

REFERENCES

- MICHELSON, A.A. (1890). . *Phil. Mag.*, **30**, 1–21. (Not cited.)
- MIHALAS, D. (1978). *Stellar atmospheres /2nd edition/*. (Not cited.)
- MORRIS, R.A.H. (2001). Ph.D. thesis, University of Wales, Cardiff (Wales). (Not cited.)
- MORTON, D.C. (1967). Mass Loss from Three OB Supergiants in Orion. *Astrophysical Journal*, **150**, 535. (Not cited.)
- MUXLOW, T.W.B. (2003). e-MERLIN - a Real-Time VLBI Array. In Y.C. Minh, ed., *New technologies in VLBI*, vol. 306 of *Astronomical Society of the Pacific Conference Series*, 245. (Not cited.)
- NAVARRO, J.F., HELMI, A. & FREEMAN, K.C. (2004). The Extragalactic Origin of the Arcturus Group. *Astrophysical Journal Letters*, **601**, L43–L46. (Not cited.)
- NEUGEBAUER, M. & SNYDER, C.W. (1962). Solar Plasma Experiment. *Science*, **138**, 1095–1097. (Not cited.)
- NEWELL, R.T. & HJELLMING, R.M. (1982). Radio emission from the extended chromosphere of Alpha Orionis. *Astrophysical Journal Letters*, **263**, L85–L87. (Not cited.)
- O’GORMAN, E. & HARPER, G.M. (2011). What is Heating Arcturus’ Wind? In C. Johns-Krull, M.K. Browning & A.A. West, eds., *16th Cambridge Workshop on Cool Stars, Stellar Systems, and the Sun*, vol. 448 of *Astronomical Society of the Pacific Conference Series*, 691. (Not cited.)
- O’GORMAN, E., HARPER, G.M., BROWN, J.M., BROWN, A., REDFIELD, S., RICHTER, M.J. & REQUENA-TORRES, M.A. (2012). CARMA CO(J = 2 - 1) Observations of the Circumstellar Envelope of Betelgeuse. *Astronomical Journal*, **144**, 36. (Not cited.)
- O’GORMAN, E., HARPER, G.M., BROWN, A., DRAKE, S. & RICHARDS, A.M.S. (2013). Multi-wavelength Radio Continuum Emission Studies of Dust-free Red Giants. *Astronomical Journal*, **144**, 36. (Not cited.)
- OHNAKA, K. (2013). Spatially resolved, high-spectral resolution observation of the K giant Aldebaran in the CO first overtone lines with VLTI/AMBER. *Astronomy & Astrophysics*, **553**, A3. (Not cited.)
- OHNAKA, K., WEIGELT, G., MILLOUR, F., HOFMANN, K.H., DRIEBE, T., SCHERTL, D., CHELLI, A., MASSI, F., PETROV, R. & STEE, P. (2011). Imaging the dynamical atmosphere of the red supergiant Betelgeuse in the CO first overtone lines with VLTI/AMBER. *Astronomy & Astrophysics*, **529**, A163. (Not cited.)
- OLNON, F.M. (1975). Thermal bremsstrahlung radiospectra for inhomogeneous objects, with an application to MWC 349. *Astronomy & Astrophysics*, **39**, 217–223. (Not cited.)
- OWOCKI, S. (2004). Stellar wind mechanisms and instabilities. In M. Heydari-Malayeri, P. Stee & J.P. Zahn, eds., *EAS Publications Series*, vol. 13 of *EAS Publications Series*, 163–250. (Not cited.)

REFERENCES

- PANAGIA, N. & FELLI, M. (1975). The spectrum of the free-free radiation from extended envelopes. *Astronomy & Astrophysics*, **39**, 1–5. (Not cited.)
- PARKER, E.N. (1958). Dynamics of the Interplanetary Gas and Magnetic Fields. *Astrophysical Journal*, **128**, 664. (Not cited.)
- PARKER, E.N. (1983). Magnetic Neutral Sheets in Evolving Fields - Part Two - Formation of the Solar Corona. *Astrophysical Journal*, **264**, 642. (Not cited.)
- PARKER, E.N. (1988). Nanoflares and the solar X-ray corona. *Astrophysical Journal*, **330**, 474–479. (Not cited.)
- PAVLENKO, Y.V., JONES, H.R.A. & LONGMORE, A.J. (2003). Carbon abundances and $^{12}\text{C}/^{13}\text{C}$ from globular cluster giants. *Monthly Notices of the Royal Astronomical Society*, **345**, 311–324. (Not cited.)
- PERLEY, R.A. & BUTLER, B.J. (2013). An Accurate Flux Density Scale from 1 to 50 GHz. *Astrophysical Journal Supplemental Series*, **204**, 19. (Not cited.)
- PERRIN, G., RIDGWAY, S.T., COUDÉ DU FORESTO, V., MENNESSON, B., TRAUB, W.A. & LACASSE, M.G. (2004). Interferometric observations of the supergiant stars α Orionis and α Herculis with FLUOR at IOTA. *Astronomy & Astrophysics*, **418**, 675–685. (Not cited.)
- PERRIN, G., VERHOELST, T., RIDGWAY, S.T., CAMI, J., NGUYEN, Q.N., CHESNEAU, O., LOPEZ, B., LEINERT, C. & RICHICHI, A. (2007). The molecular and dusty composition of Betelgeuse’s inner circumstellar environment. *Astronomy & Astrophysics*, **474**, 599–608. (Not cited.)
- PERRYMAN, M.A.C., LINDEGREN, L., KOVALEVSKY, J., HOEG, E., BASTIAN, U., BERNACCA, P.L., CRÉZÉ, M., DONATI, F., GRENON, M., GREWING, M., VAN LEEUWEN, F., VAN DER MAREL, H., MIGNARD, F., MURRAY, C.A., LE POOLE, R.S., SCHRIJVER, H., TURON, C., ARENOU, F., FROESCHLÉ, M. & PETERSEN, C.S. (1997). The HIPPARCOS Catalogue. *Astronomy & Astrophysics*, **323**, L49–L52. (Not cited.)
- PILBRATT, G.L., RIEDINGER, J.R., PASSVOGEL, T., CRONE, G., DOYLE, D., GAGEUR, U., HERAS, A.M., JEWELL, C., METCALFE, L., OTT, S. & SCHMIDT, M. (2010). Herschel Space Observatory. An ESA facility for far-infrared and submillimetre astronomy. *Astronomy & Astrophysics*, **518**, L1. (Not cited.)
- PLEZ, B. & LAMBERT, D.L. (2002). The outer atmosphere of the M-type supergiant alpha Orionis: K I 7699 Å emission. *Astronomy & Astrophysics*, **386**, 1009–1018. (Not cited.)
- RAMÍREZ, I. & ALLENDE PRIETO, C. (2011). Fundamental Parameters and Chemical Composition of Arcturus. *Astrophysical Journal*, **743**, 135. (Not cited.)
- RAMÍREZ, S.V., SELLGREN, K., CARR, J.S., BALACHANDRAN, S.C., BLUM, R., TERNDROP, D.M. & STEED, A. (2000). Stellar Iron Abundances at the Galactic Center. *Astrophysical Journal*, **537**, 205–220. (Not cited.)
- REIMERS, D. (1975). Circumstellar absorption lines and mass loss from red giants. *Memoires of the Societe Royale des Sciences de Liege*, **8**, 369–382. (Not cited.)

REFERENCES

- REIMERS, D. (1977). On the absolute scale of mass-loss in red giants. I - Circumstellar absorption lines in the spectrum of the visual companion of Alpha-1 HER. *Astronomy & Astrophysics*, **61**, 217–224. (Not cited.)
- REIMERS, D. (1982). Detection of further red giants with 'hybrid' atmospheres and a possible correlation with double circumstellar MG II and CA II lines. *Astronomy & Astrophysics*, **107**, 292–299. (Not cited.)
- RENZINI, A., GREGGIO, L., RITOSSA, C. & FERRARIO, L. (1992). Why stars inflate to and deflate from red giant dimensions. *Astrophysical Journal*, **400**, 280–303. (Not cited.)
- RETTET, A., BEDDING, T.R., BUZASI, D.L., KJELDSSEN, H. & KISS, L.L. (2003). Oscillations in Arcturus from WIRE Photometry. *Astrophysical Journal Letters*, **591**, L151–L154. (Not cited.)
- RICH, J.W., DE BLOK, W.J.G., CORNWELL, T.J., BRINKS, E., WALTER, F., BAGETAKOS, I. & KENNICUTT, R.C., JR. (2008). Multi-Scale CLEAN: A Comparison of its Performance Against Classical CLEAN on Galaxies Using THINGS. *Astronomical Journal*, **136**, 2897–2920. (Not cited.)
- RICHARDS, A.M.S., DAVIS, R.J., DECIN, L., ETOKA, S., HARPER, G.M., LIM, J.J., GARRINGTON, S.T., GRAY, M.D., McDONALD, I., O'GORMAN, E. & WITTKOWSKI, M. (2013). e-MERLIN resolves Betelgeuse at λ 5 cm: hotspots at 5 R. *Monthly Notices of the Royal Astronomical Society*, **432**, L61. (Not cited.)
- RICHICHI, A. & ROCCATAGLIATA, V. (2005). Aldebaran's angular diameter: How well do we know it? *Astronomy & Astrophysics*, **433**, 305–312. (Not cited.)
- ROBINSON, R.D., CARPENTER, K.G. & BROWN, A. (1998). Goddard High-Resolution Spectrograph Observations of Cool Low-Gravity Stars. IV. A Comparison of the K5 III Stars alpha Tauri and gamma Draconis. *Astrophysical Journal*, **503**, 396. (Not cited.)
- RODGERS, B. & GLASSGOLD, A.E. (1991). The temperature of the circumstellar envelope of Alpha Orionis. *Astrophysical Journal*, **382**, 606–616. (Not cited.)
- ROHLFS, K. & WILSON, T.L. (1996). *Tools of Radio Astronomy*. (Not cited.)
- RYAN, S.G. & NORTON, A.J. (2010). *Stellar Evolution and Nucleosynthesis*. (Not cited.)
- RYBICKI, G.B. & LIGHTMAN, A.P. (1979). *Radiative processes in astrophysics*. (Not cited.)
- RYDE, N., LAMBERT, D.L., RICHTER, M.J. & LACY, J.H. (2002). Detection of Water Vapor in the Photosphere of Arcturus. *Astrophysical Journal*, **580**, 447–458. (Not cited.)
- SAHA, S.K. (2011). *Aperture Synthesis*. (Not cited.)
- SANFORD, R.F. (1933). No. 464. The variation in the radial velocity of alpha Orionis from 1923 to 1931. *Contributions from the Mount Wilson Observatory / Carnegie Institution of Washington*, **464**, 1–10. (Not cited.)

REFERENCES

- SAULT, R.J., TEUBEN, P.J. & WRIGHT, M.C.H. (1995). A Retrospective View of MIRIAD. In R.A. Shaw, H.E. Payne & J.J.E. Hayes, eds., *Astronomical Data Analysis Software and Systems IV*, vol. 77 of *Astronomical Society of the Pacific Conference Series*, 433. (Not cited.)
- SCHARMER, G.B. & CARLSSON, M. (1985). A new approach to multi-level non-LTE radiative transfer problems. *Journal of Computational Physics*, **59**, 56–80. (Not cited.)
- SCHRIJVER, C.J. & HARVEY, K.L. (1989). The distribution of solar magnetic fluxes and the nonlinearity of stellar flux-flux relations. *Astrophysical Journal*, **343**, 481–488. (Not cited.)
- SCHRÖDER, K.P. & SEDLMAYR, E. (2001). The galactic mass injection from cool stellar winds of the 1 to 2.5 M_{sun} stars in the solar neighbourhood. *Astronomy & Astrophysics*, **366**, 913–922. (Not cited.)
- SCHWARZ, U.J. (1978). Mathematical-statistical Description of the Iterative Beam Removing Technique (Method CLEAN). *Astronomy & Astrophysics*, **65**, 345. (Not cited.)
- SCHWARZSCHILD, M. (1975). On the scale of photospheric convection in red giants and supergiants. *Astrophysical Journal*, **195**, 137–144. (Not cited.)
- SCUDERI, S., PANAGIA, N., STANGHELLINI, C., TRIGILIO, C. & UMANA, G. (1998). Radio observations of stellar winds from early type stars. *Astronomy & Astrophysics*, **332**, 251–267. (Not cited.)
- SEAQUIST, E.R. & TAYLOR, A.R. (1987). A detailed analysis of the radio emission from the symbiotic star RX Puppis. *Astrophysical Journal*, **312**, 813–821. (Not cited.)
- SENNHAUSER, C. & BERDYUGINA, S.V. (2011). First detection of a weak magnetic field on the giant Arcturus: remnants of a solar dynamo? *Astronomy & Astrophysics*, **529**, A100. (Not cited.)
- SHANG, H., GLASSGOLD, A.E., SHU, F.H. & LIZANO, S. (2002). Heating and Ionization of X-Winds. *Astrophysical Journal*, **564**, 853–876. (Cited on page 7.)
- SHU, F.H. (1991). *The physics of astrophysics. Volume 1: Radiation..* (Not cited.)
- SIMON, T., LINSKY, J.L. & STENCEL, R.E. (1982). On the reality of a boundary in the H-R diagram between late-type stars with and without high temperature outer atmospheres. *Astrophysical Journal*, **257**, 225–246. (Not cited.)
- SKINNER, C.J., DOUGHERTY, S.M., MEIXNER, M., BODE, M.F., DAVIS, R.J., DRAKE, S.A., ARENS, J.F. & JERNIGAN, J.G. (1997). Circumstellar environments - V. The asymmetric chromosphere and dust shell of alpha Orionis. *Monthly Notices of the Royal Astronomical Society*, **288**, 295–306. (Not cited.)
- SLEE, O.B., STEWART, R.T., BUNTON, J.D., BEASLEY, A.J., CARTER, B.D. & NELSON, G.J. (1989). A microwave survey of southern red giants. *Monthly Notices of the Royal Astronomical Society*, **239**, 913–922. (Not cited.)
- SMITH, M.A., PATTEN, B.M. & GOLDBERG, L. (1989). Radial-velocity variations in Alpha Ori, Alpha Sco, and Alpha HER. *Astronomical Journal*, **98**, 2233–2248. (Not cited.)

REFERENCES

- SMITH, N., HINKLE, K.H. & RYDE, N. (2009). Red Supergiants as Potential Type II_n Supernova Progenitors: Spatially Resolved 4.6 μ m CO Emission Around VY CMa and Betelgeuse. *Astronomical Journal*, **137**, 3558–3573. (Not cited.)
- SODERHJELM, S. & MIGNARD, F. (1998). Arcturus as a double star. *The Observatory*, **118**, 365–366. (Not cited.)
- SPENCER JONES, H. (1928). The radial velocity variations of a Orionis and a Scorpii. *Monthly Notices of the Royal Astronomical Society*, **88**, 660. (Not cited.)
- SPINRAD, H. & VARDYA, M.S. (1966). Approximate Abundances of the Light Elements from the Molecular Spectra of M and S Stars. *Astrophysical Journal*, **146**, 399. (Not cited.)
- SPITZER, L. (1978). *Physical processes in the interstellar medium*. New York Wiley-Interscience, 1978. 333 p. (Not cited.)
- SPITZER, L., JR. (1939). Spectra of M Supergiant Stars. *Astrophysical Journal*, **90**, 494. (Not cited.)
- STANCLIFFE, R.J., CHIEFFI, A., LATTANZIO, J.C. & CHURCH, R.P. (2009). Why Do Low-Mass Stars Become Red Giants. **26**, 203–208. (Not cited.)
- STEBBINS, J. & HUFFER, C.M. (1931). Photo-electric studies of five variable stars. *Publications of the Washburn Observatory*, **15**, 178–213. (Not cited.)
- STEIN, R.F. (1967). Generation of Acoustic and Gravity Waves by Turbulence in an Isothermal Stratified Atmosphere. *Solar Physics*, **2**, 385–432. (Not cited.)
- STOTHERS, R. & CHIN, C.W. (1969). Advanced Phases of Evolution in Massive Red Supergiants. *Astrophysical Journal*, **158**, 1039. (Not cited.)
- STOTHERS, R. & LEUNG, K.C. (1971). Luminosities, masses and periodicities of massive red supergiants. *Astronomy & Astrophysics*, **10**, 290–300. (Not cited.)
- STRASSMEIER, K.G., HALL, D.S., FEKEL, F.C. & SCHECK, M. (1993). A catalog of chromospherically active binary stars (second edition). *Astronomy & Astrophysics Supplemental*, **100**, 173–225. (Not cited.)
- SUGIMOTO, D. & FUJIMOTO, M.Y. (2000). Why Stars Become Red Giants. *Astrophysical Journal*, **538**, 837–853. (Not cited.)
- SUTMANN, G. & CUNTZ, M. (1995). Generation of mass loss in K giants: The failure of global oscillation modes and possible implications. *Astrophysical Journal Letters*, **442**, L61–L64. (Not cited.)
- SUZUKI, T.K. (2007). Structured Red Giant Winds with Magnetized Hot Bubbles and the Corona/Cool Wind Dividing Line. *Astrophysical Journal*, **659**, 1592–1610. (Not cited.)
- TATEBE, K., CHANDLER, A.A., WISHNOW, E.H., HALE, D.D.S. & TOWNES, C.H. (2007). The Nonspherical Shape of Betelgeuse in the Mid-Infrared. *Astrophysical Journal Letters*, **670**, L21–L24. (Not cited.)

REFERENCES

- TAYLOR, G.B., CARILLI, C.L. & PERLEY, R.A., eds. (1999). *Synthesis Imaging in Radio Astronomy II*, vol. 180 of *Astronomical Society of the Pacific Conference Series*. (Not cited.)
- TSUJI, T. (1979). A probe of optically thin dust shells around late-type stars. *Publications of the Astronomical Society of Japan*, **31**, 43–60. (Not cited.)
- TSUJI, T. (2000). Water on the Early M Supergiant Stars α Orionis and μ Cephei. *Astrophysical Journal*, **538**, 801–807. (Not cited.)
- TSUJI, T. (2008). Cool luminous stars: the hybrid nature of their infrared spectra. *Astronomy & Astrophysics*, **489**, 1271–1289. (Not cited.)
- TSUJI, T. (2009). The K giant star Arcturus: the hybrid nature of its infrared spectrum. *Astronomy & Astrophysics*, **504**, 543–559. (Not cited.)
- TURNER, N.H., TEN BRUMMELAAR, T.A. & MASON, B.D. (1999). Adaptive Optics Observations of Arcturus using the Mount Wilson 100 Inch Telescope. *Publications of the Astronomical Society of the Pacific*, **111**, 556–558. (Not cited.)
- TUTHILL, P.G., HANIFF, C.A. & BALDWIN, J.E. (1997). Hotspots on late-type supergiants. *Monthly Notices of the Royal Astronomical Society*, **285**, 529–539. (Not cited.)
- UITENBROEK, H., DUPREE, A.K. & GILLILAND, R.L. (1998). Spatially Resolved Hubble Space Telescope Spectra of the Chromosphere of α Orionis. *Astronomical Journal*, **116**, 2501–2512. (Not cited.)
- VAN CITTERT, P.H. (1934). Die Wahrscheinliche Schwingungsverteilung in Einer von Einer Lichtquelle Direkt Oder Mittels Einer Linse Beleuchteten Ebene. *Physica*, **1**, 201–210. (Not cited.)
- VAN LEEUWEN, F. (2007). Validation of the new Hipparcos reduction. *Astronomy & Astrophysics*, **474**, 653–664. (Not cited.)
- VAN LOON, J.T. (2010). The Effects of Red Supergiant Mass Loss on Supernova Ejecta and the Circumburst Medium. In C. Leitherer, P.D. Bennett, P.W. Morris & J.T. Van Loon, eds., *Hot and Cool: Bridging Gaps in Massive Star Evolution*, vol. 425 of *Astronomical Society of the Pacific Conference Series*, 279. (Not cited.)
- VAN LOON, J.T., CIONI, M.R.L., ZIJLSTRA, A.A. & LOUP, C. (2005). An empirical formula for the mass-loss rates of dust-enshrouded red supergiants and oxygen-rich Asymptotic Giant Branch stars. *Astronomy & Astrophysics*, **438**, 273–289. (Not cited.)
- VERDINI, A. & VELLI, M. (2007). Alfvén Waves and Turbulence in the Solar Atmosphere and Solar Wind. *Astrophysical Journal*, **662**, 669–676. (Not cited.)
- VERHOELST, T., DECIN, L., VAN MALDEREN, R., HONY, S., CAMI, J., ERIKSSON, K., PERRIN, G., DEROO, P., VANDENBUSSCHE, B. & WATERS, L.B.F.M. (2006). Amorphous alumina in the extended atmosphere of α Orionis. *Astronomy & Astrophysics*, **447**, 311–324. (Not cited.)

REFERENCES

- VIDOTTO, A.A., FALCETA-GONÇALVES, D. & JATENCO-PEREIRA, V. (2006). A Self-Consistent Determination of the Temperature Profile and The Magnetic Field Geometry in Winds of Late-Type Stars. *Space Science Reviews*, **122**, 181–188. (Not cited.)
- VÖGLER, A. & SCHÜSSLER, M. (2007). A solar surface dynamo. *Astronomy & Astrophysics*, **465**, L43–L46. (Not cited.)
- WEI, L., WOODY, D., TEUBEN, P., LA VIGNE, M. & VOGEL, S. (2008). Bandpass Calibration for CARMA. In *CARMA Memorandum Series*, vol. 45. (Not cited.)
- WEYMANN, R. (1962). Physical Conditions in the Circumstellar Envelope of α Orionis. *Astrophysical Journal*, **136**, 844–+. (Not cited.)
- WHITE, S.M. (2000). The Contributions of the VLA to the Study of Radio Stars. In D.G. Finley & W.M. Goss, eds., *Radio interferometry : the saga and the science*, 86. (Not cited.)
- WHITE, S.M. (2009). Single Dish Aperture Efficiency Measurements at CARMA. In *CARMA Memorandum Series*, vol. 49 of *CARMA Memorandum Series*. (Not cited.)
- WIEDEMANN, G., AYRES, T.R., JENNINGS, D.E. & SAAR, S.H. (1994). Carbon Monoxide Fundamental Bands in Late-Type Stars. III. Chromosphere or CO-mosphere? *Astrophysical Journal*, **423**, 806. (Not cited.)
- WILLIAMS, M.E.K., FREEMAN, K.C., HELMI, A. & RAVE COLLABORATION (2009). The Arcturus Moving Group: Its Place in the Galaxy. In J. Andersen, Nordströara, B. m & J. Bland-Hawthorn, eds., *IAU Symposium*, vol. 254 of *IAU Symposium*, 139–144. (Not cited.)
- WILSON, R.W., BALDWIN, J.E., BUSCHER, D.F. & WARNER, P.J. (1992). High-resolution imaging of Betelgeuse and Mira. *Monthly Notices of the Royal Astronomical Society*, **257**, 369–376. (Not cited.)
- WOOD, B.E., HARPER, G.M., MÜLLER, H.R., HEERIKHUISEN, J. & ZANK, G.P. (2007). The Wind-ISM Interaction of α Tauri. *Astrophysical Journal*, **655**, 946–957. (Not cited.)
- WRIGHT, A.E. & BARLOW, M.J. (1975). The radio and infrared spectrum of early-type stars undergoing mass loss. *Monthly Notices of the Royal Astronomical Society*, **170**, 41–51. (Not cited.)
- WRIGHT, K.O. (1970). The Zeta Aurigae stars. *Vistas in Astronomy*, **12**, 147–182. (Not cited.)
- WRIGHT, M. (2009). Bandpass Calibration and Stability. In *CARMA Memorandum Series*, vol. 55. (Not cited.)
- WRIGHT, M.C.H. (1999). Image Fidelity. In *BIMA memo 73: Image Fidelity*, vol. 73 of *BIMA memos*. (Not cited.)
- ZUCKERMAN, B., KIM, S.S. & LIU, T. (1995). Luminosity Class III Stars with Excess Far-Infrared Emission. *Astrophysical Journal Letters*, **446**, L79. (Not cited.)