



Radio Interferometric Studies of Cool Evolved Stellar Winds

Eamon O'Gorman

Trinity College Dublin
November 15, 2013

Supervisor: Prof. Graham Harper

Overview

- Stellar Winds Across the HR Diagram
- Stellar Radio Emission
 - 1) Winds of Red Supergiants (CSE and inner wind)
 - 2) Winds of Red Giants (inner wind)
- Summary

Stellar Winds Across the H-R Diagram





Image Credit: S. R. Cranmer

Red Supergiants & Red Giants

	Red Giants	Red Supergiants
Mass (M_{\odot})	$\sim 0.4 \rightarrow \sim 8$	$\sim 8 \rightarrow \sim 40$
Radius (R_{\odot})	~ 50	~ 500
Lifetime (yr)	$\sim 10^9$	$\sim 5 \times 10^5$
dM/dt ($M_{\odot} \text{ yr}^{-1}$)	$\sim 10^{-10}$	$\sim 10^{-5}$

Red Supergiants & Red Giants

	Red Giants	Red Supergiants
Mass (M_{\odot})	$\sim 0.4 \rightarrow \sim 8$	$\sim 8 \rightarrow \sim 40$
Radius (R_{\odot})	~ 50	~ 500
Lifetime (yr)	$\sim 10^9$	$\sim 5 \times 10^5$
dM/dt ($M_{\odot} \text{ yr}^{-1}$)	$\sim 10^{-10}$	$\sim 10^{-5}$



■ $H_p/R_* \sim 10^{-2}$

■ $V_{\infty} < V_{\text{escape}}(R_*)$

■ $T_{\text{wind}} < 10^4 \text{ K}$

Red Supergiants & Red Giants

- No coronae
- Radiation field peaks in the IR
- Small pulsation amplitudes and little dust
- Unknown mass-loss mechanism




Image Credit: L. Decin

Red Supergiants & Red Giants

- No coronae
- Radiation field peaks in the IR
- Small pulsation amplitudes and little dust
- Unknown mass-loss mechanism
- Poor knowledge of outflow conditions
 r , $v(r)$, $T(r)$
- Disk averaged observations
- No wind emission features at UV or optical λ .




Image Credit: L. Decin

Red Supergiants & Red Giants

- No coronae
- Radiation field peaks in the IR
- Small pulsation amplitudes and little dust
- Unknown mass-loss mechanism
- Poor knowledge of outflow conditions
 r , $v(r)$, $T(r)$
- Disk averaged observations
- No wind emission features at UV or optical λ .





Image Credit: L. Decin

Thesis Goal: Improve understanding of outflow conditions to gain insight into mass-loss mechanism. → Radio

Aperture Synthesis




- No need to measure all the incoming rays at one time.

- Break aperture into N sub-apertures:
 $N(N-1)/2$ pairs


- FT of each measurement is a sinusoid with spacing λ/B between successive peaks.

- Build up image of sky by summing many such sinusoids.


Aperture Synthesis




Aperture Synthesis




Aperture Synthesis




Aperture Synthesis



Aperture Synthesis




Aperture Synthesis




1) Field of View: λ/D (D = antenna diameter)

Aperture Synthesis



- 1) Field of View: λ/D (D = antenna diameter)
- 2) Resolution: λ/B_{\max} (B_{\max} = longest projected baseline)

Aperture Synthesis



- 1) Field of View: λ/D (D = antenna diameter)
- 2) **Resolution:** λ/B_{\max} (B_{\max} = longest projected baseline)
- 3) **Resolving out scale:** λ/B_{\min} (B_{\min} = shortest projected baseline)

Stellar Radio Emission

Radio Sky at 4.85 GHz (300ft Green Bank)



Credit: NRAO/AUI

$$(P_{\text{radio}}/P_{\text{optical}})_{\odot} \sim 10^{-15}$$

$S_{\nu=4.6 \text{ GHz}} \sim 40 \mu\text{Jy}$ at $\alpha \text{ Cen}$
(not detectable with 'old' VLA!)

Stellar Radio Emission


Radio Sky at 4.85 GHz (300ft Green Bank)



Credit: NRAO/AUI

$$(P_{\text{radio}}/P_{\text{optical}})_{\odot} \sim 10^{-15}$$


$S_{\nu=4.6 \text{ GHz}} \sim 40 \mu\text{Jy}$ at $\alpha \text{ Cen}$
(not detectable with 'old' VLA!)



1)

Winds of Red Supergiants


Betelgeuse (M2 Iab)

(Decin *et al.*, 2013)

Log(L/L_{\odot})	5
Distance	197 ± 45 parsec
Photospheric Radius	22.5 mas ($950 R_{\odot}$)
Mass	$\sim 15 M_{\odot}$
Mass loss rate	$3 \times 10^{-6} M_{\odot} \text{ yr}^{-1}$
Photospheric Temperature	3,600 K (cool star)
Origin	O-type main sequence
$^{12}\text{C}/^{13}\text{C}$	6 ± 1



Betelgeuse: Circumstellar Environment

- At least two *recent* mass loss phases
- Two distinct shells spectrally resolved at 4.6 μm :
 - S2, moving at 17 km s^{-1}
 - S1, moving at 10 km s^{-1}
 - Spatial extent not directly determined



Betelgeuse: Circumstellar Environment



- At least two *recent* mass loss phases
- Two distinct shells spectrally resolved at 4.6 μm :
 - S2, moving at 17 km s^{-1}
 - S1, moving at 10 km s^{-1}
 - Spatial extent not directly determined
- Plez & Lambert (2002) appear to detect S2 out to $50''$
- IRAM 30 m ($\theta_{\text{HPBW}} \sim 12''$) fails to resolve S2 shell at 1.3 mm
- Single dish $^{12}\text{C}^{16}\text{O}$ mm-observations reveal only high velocity S2 shell.



Betelgeuse: Circumstellar Environment

- At least two *recent* mass loss phases
- Two distinct shells spectrally resolved at 4.6 μm :
 - S2, moving at 17 km s^{-1}
 - S1, moving at 10 km s^{-1}
 - Spatial extent not directly determined
- Plez & Lambert (2002) appear to detect S2 out to $50''$
- IRAM 30 m ($\theta_{\text{HPBW}} \sim 12''$) fails to resolve S2 shell at 1.3 mm
- Single dish $^{12}\text{C}^{16}\text{O}$ mm-observations reveal only high velocity S2 shell.

Goal: Measure both the spatial scales and the velocities of Betelgeuse's outflow region using $^{12}\text{C}^{16}\text{O}$ $J = 2-1$ line as a tracer to sort out puzzling evidence.



CARMA


- **C**ombined **A**rray for **R**esearch in **M**illimeter-wave **A**stronomy
- Cedar Flat, eastern California (~ 2,200 m)
- 15 element interferometer (9 x 6.1 m + 6 x 10.4 m antennas)
- Merger of two independent arrays: BIMA + OVRO (2007)
- 105 baselines ($n(n-1)/2$) with 5 configurations
- Three bands: 7 mm, 3 mm and 1.3 mm




Credit:
John
Carlstrom

mm Observations


Date	Config	Time (hr)	Resolution (")	Max Scale (")
Jun 07	D	8.4	2.1	15
Jul 09	E	3.2	4.4	19
Nov 09	C	8.4	0.9	6



Individual Configurations





(O'Gorman *et al.*, 2012)





No shells in individual configuration image cubes.



Spherically Symmetric Shell





Spherically Symmetric Shell





Spherically Symmetric Shell





Spherically Symmetric Shell





Spherically Symmetric Shell





Spherically Symmetric Shell





Spherically Symmetric Shell





Spherically Symmetric Shell





Spherically Symmetric Shell





Spherically Symmetric Shell




Spherically Symmetric Shell




Spherically Symmetric Shell



Combined Configuration




Combined Configurations




S2 Flow

S2 flow not present at low absolute velocities.



(O'Gorman *et al.*, 2012)



$$r_{\text{chan}} = R_{S2} \sin \left[\cos^{-1} \left(\frac{v_{\text{chan}}}{V_{S2}} \right) \right]$$



S1 Flow



Conclusions

- High spatial resolution configuration resolves out S2 emission providing S1 profile.
- Multiple CARMA configurations provide the high spatial resolution needed to study the inner S1 flow, while ensuring the extended S2 flow is not resolved out.




	S1	S2
Outflow Velocities (km s^{-1})	9.8	14.3
Maximum Spatial Extent (")	4 \rightarrow 6	17
Age (yr)	400 \rightarrow 600	1100

- Previous line profiles not horned shaped because:
 - S1 emission is also present
 - S2 emission weak at low absolute velocities


Betelgeuse's Wind Acceleration Region

(Newell & Hjellming, 1982)




Betelgeuse's Wind Acceleration Region

(Newell & Hjellming, 1982)



(Lim *et al.*, 1998)




$$F_\nu = \frac{\pi k \phi^2 T_b}{2\lambda^2}$$

Brightness Temperature


Size

Betelgeuse with e-MERLIN (5.2 cm)




(Credit: MERLIN/VLBI national facility)

Betelgeuse with e-MERLIN (5.2 cm)



(Credit: MERLIN/VLBI national facility)



Two unresolved *hot spots*:


$$T_e > 3800 \text{ K}$$

$$T_e > 5400 \text{ K}$$

Where is the photosphere?

Betelgeuse with e-MERLIN (5.2 cm)


Astrometric solution of Harper *et al.*, (2008)




- Hottest source at $4 R_*$.
- At least ~ 3 times the predicted T_e .

Betelgeuse with e-MERLIN (5.2 cm)

Astrometric solution of Harper *et al.*, (2008)




- Hottest source at $4 R_*$.
- At least ~ 3 times the predicted T_e .



Spherically symmetric semi-empirical model of Harper *et al.*, (2001)


Betelgeuse with e-MERLIN (5.2 cm)

Astrometric solution of Harper *et al.*, (2008)



- Hottest source at $4 R_*$.
- At least ~ 3 times the predicted T_e .

Goal: Analyse high resolution archival cm data to search for signatures of hotspots.



Spherically symmetric semi-empirical model of Harper *et al.*, (2001)

Betelgeuse with VLA – Pie Town

VLA



+

Pie Town Antenna



=

e-MERLIN

w/l (cm)	Resolution (")
5.2	80 x 60

VLA + Pie Town


w/l (cm)	Resolution (")
0.7	40 x 25
1.3	80 x 40
2.0	120 x 90
3.5	200 x 130
6.2	380 x 270

Betelgeuse with VLA – Pie Town

Variability 1998-2004

w/l (cm)	Variability
0.7	23%
1.3	27%
2.0	32%
3.5	21%
6.2	35%

Thermal Profile




Betelgeuse with VLA – Pie Town


Variability 1998-2004

w/l (cm)	Variability
0.7	23%
1.3	27%
2.0	32%
3.5	21%
6.2	35%

Thermal Profile



Q band (0.7 cm)




Betelgeuse with VLA – Pie Town


Variability 1998-2004

w/l (cm)	Variability
0.7	23%
1.3	27%
2.0	32%
3.5	21%
6.2	35%


Thermal Profile



Q band (0.7 cm)



Betelgeuse with VLA – Pie Town



- **No clear signature of hot spots any in any maps.**

- Time dependence?

- Opacity?

$$\frac{\tau_{5.2 \text{ cm}}}{\tau_{1.3 \text{ cm}}} \simeq 20$$


Conclusions

- e-MERLIN has revealed two unresolved *hotspots* separated by $4 R_*$
- One may be at the position of photosphere
- VLA + Pie Town data in good agreement with Lim *et al.* (1998)
- Flux and structure variability present in data
- No clear signature of e-MERLIN hotspots in any of the VLA + Pie Town data

2)

Winds of Red Giants

- Atmospheres cannot be spatially resolved at radio wavelengths.
- Wind properties generally traditionally determined by analysing strong UV and optical resonance lines.
- At cm/mm the thermal continuum Planck function depends linearly on T , unlike the UV.
- Continuum flux measurements at cm/mm: opacity is proportional to $\sim \lambda^{2.1} n_e n_{\text{ion}}$.



Goal: *Observe two 'standard' red giants at all possible cm wavelengths to test and improve existing models.*

Red Giant Targets

	Arcturus (α Boo: K2 III)	Aldebaran (α Tau: K5 III)
Distance (pc)	11.3	20.4
Photospheric Radius (R_{\odot})	25.4	44.2
Mass (M_{\odot})	0.8	1.3
Mass loss rate ($M_{\odot} \text{ yr}^{-1}$)	2×10^{-10}	1.6×10^{-11}
T_{eff} (K)	4290	3970

- Single, non-dusty and non-pulsating
- Nearby, with well known stellar parameters
- Semi-empirical 1-D chromospheric and wind models that can be directly tested

Karl G. Jansky Very Large Array

Credit:
NRAO



- 27 25 m antennas, New Mexico. Max baseline ~ 36 km
- Full frequency coverage between 1.0 and 50 GHz
- Continuum sensitivity improvement over the VLA by factors of 5 to 20
- Spectral Capability: A minimum of 16,384 and a maximum of 4,194,304 channels

New Observations

Open Shared Risk Observing (OSRO)

- B config (128 MHz)
- α Boo: S \rightarrow Q band (13th - 22nd Feb 2011)
- α Tau: S \rightarrow Q band (11th - 13th Feb 2011)
- A unique dataset


Directors Discretionary Time (DDT)

- B config (2 GHz)
- α Boo: S & L band (July 2012)

Red Giant Radio Maps


Example: Aldebaran X (3 cm), C (6 cm), and S (10 cm) band

X-band (8 GHz)
 $S_{\nu} = 0.3$ mJy




Contours = $(-2, 2, 4, \dots, 16) \times \sigma$
 $\sigma = 16$ μ Jy

C-band (5 GHz)
 $S_{\nu} = 0.15$ mJy




Contours = $(-2, 2, 4, \dots, 14) \times \sigma$
 $\sigma = 10$ μ Jy

S-band (3 GHz)
 $S_{\nu} = 0.06$ mJy




Contours = $(-3, -2, -1, 1, 2, 3) \times \sigma$
 $\sigma = 18$ μ Jy


Spectral Energy Distribution – α Boo



Spectral Energy Distribution – α Boo



Spectral Indices



■ If, $n_e \propto r^{-p}$

$$T_e \propto r^{-n}$$

then,

$$\alpha = f(p, n)$$


■ α Tau: wind ~ wind optically thin

■ α Boo:


Assume constant velocity wind,

$$T_e \propto r^{-1.65}$$

Spectral Energy Distribution – α Boo



Spectral Energy Distribution – α Boo



2)

Conclusions

- Most comprehensive set of multi- λ radio observations of two *standard* red giants.
- Tested theoretical and semi-empirical atmospheric models.
- Radio spectral indices allow flow properties to be investigated.
 - Optically thin wind for α Tau.
 - Rapidly cooling wind for α Boo.
- New analytical advection wind model for α Boo.

Summary

- Established spatial scales for the two flows in CSE of Betelgeuse
- Evidence for asymmetries in both S1 and S2 flows
- e-MERLIN results are surprising. Episodic mass-loss mechanism in RSGs?
- 1st multi-wavelength radio study of red giants
- Provide wind diagnostics and updated outflow models.
- Understanding of mass-loss is improving and radio interferometry will continue to play a major role in future developments.

Thermal Energy Balance

- Close to photosphere: $v^2(r) \ll v_{\text{esc}}^2(r)$
- To escape star: $v^2(r) > v_{\text{esc}}^2(r)$
- Energy added in the form of either heat or momentum

Goal: *Create a wind thermal energy balance to investigate possible heating mechanisms in Arcturus' inner outflow region.*

Thermal Energy Balance

- Close to photosphere: $v^2(r) \ll v_{\text{esc}}^2(r)$
- To escape star: $v^2(r) > v_{\text{esc}}^2(r)$
- Energy added in the form of either heat or momentum

Goal: Carry out a thermal energy balance to investigate possible heating mechanisms in Arcturus' inner outflow region.


Notation:

$$\sum \lambda_i = \overbrace{-\frac{4}{3} - \frac{2}{3} \frac{d(\ln v)}{d(\ln r)}}^{\text{Adiabatic Expansion}} + \overbrace{\sum_{i=1} \mathcal{H}_i}^{\text{Heating}} - \overbrace{\sum_{j=1} \mathcal{L}_j}^{\text{Cooling}}$$


Gas kinetic power law slope

Cooling and Heating Processes


Cooling




Heating




Cooling and Heating Processes



Cooling and Heating Processes




Cooling and Heating Processes



Radio Emission Mechanisms

1) Molecular Emission lines

- Line profiles: flat-topped or horned shaped (optically thin)
- Line full widths typically between 20 and 50 km s⁻¹
- CO: Both O-rich and C-rich stars. Stable.
- Excitation processes for CO: H and H₂ collisions. Photo-excitation of vibrational levels by IR photons.



1) Thermal Bremsstrahlung Emission

- Opacity is proportional to $\sim \lambda^{2.1}$

$$F_\nu = \frac{\pi k \phi^2 T_b}{2\lambda^2}$$

Source or beam size

