



Radio Interferometric Studies of Cool Evolved Stellar Winds

Eamon O'Gorman

Trinity College Dublin
November 15, 2013

Supervisor: Prof. Graham Harper

Overview

- Stellar Winds Across the HR Diagram
- Stellar Radio Emission
 - 1) Winds of Red Supergiants (CSE and inner wind)
 - 2) Winds of Red Giants (inner wind)
- Summary

Stellar Winds Across the H-R Diagram




Image Credit: S. R. Cranmer

Red Supergiants & Red Giants


	Red Supergiants	Red Giants
Mass (M_{\odot})	$\sim 8 \rightarrow \sim 40$	$\sim 0.4 \rightarrow \sim 8$
Radius (R_{\odot})	~ 500	~ 50
Lifetime (yr)	$\sim 10^6$	$\sim 10^9$
dM/dt ($M_{\odot} \text{ yr}^{-1}$)	$\sim 10^{-5}$	$\sim 10^{-10}$

Red Supergiants & Red Giants

	Red Supergiants	Red Giants
Mass (M_{\odot})	$\sim 8 \rightarrow \sim 40$	$\sim 0.4 \rightarrow \sim 8$
Radius (R_{\odot})	~ 500	~ 50
Lifetime (yr)	$\sim 10^6$	$\sim 10^9$
dM/dt ($M_{\odot} \text{ yr}^{-1}$)	$\sim 10^{-5}$	$\sim 10^{-10}$


$$H_p/R_* \sim 10^{-2}$$


$$V_{\infty} < V_{\text{escape}}(R_*)$$


$$T_{\text{wind}} < 10^4 \text{ K}$$

Red Supergiants & Red Giants

	Red Supergiants	Red Giants
Mass (M_{\odot})	$\sim 8 \rightarrow \sim 40$	$\sim 0.4 \rightarrow \sim 8$
Radius (R_{\odot})	~ 500	~ 50
Lifetime (yr)	$\sim 10^6$	$\sim 10^9$
dM/dt ($M_{\odot} \text{ yr}^{-1}$)	$\sim 10^{-5}$	$\sim 10^{-10}$

$$\blacksquare H_p/R_* \sim 10^{-2}$$

$$\blacksquare V_{\infty} < V_{\text{escape}}(R_*)$$

$$\blacksquare T_{\text{wind}} < 10^4 \text{ K}$$

\blacksquare No detected coronae, radiation field peaks in IR, small pulsation amplitudes, and little dust

Red Supergiants & Red Giants

	Red Supergiants	Red Giants
Mass (M_{\odot})	$\sim 8 \rightarrow \sim 40$	$\sim 0.4 \rightarrow \sim 8$
Radius (R_{\odot})	~ 500	~ 50
Lifetime (yr)	$\sim 10^6$	$\sim 10^9$
dM/dt ($M_{\odot} \text{ yr}^{-1}$)	$\sim 10^{-5}$	$\sim 10^{-10}$

$$\blacksquare H_p/R_* \sim 10^{-2}$$

$$\blacksquare V_{\infty} < V_{\text{escape}}(R_*)$$

$$\blacksquare T_{\text{wind}} < 10^4 \text{ K}$$

- No detected coronae, radiation field peaks in IR, small pulsation amplitudes, and little dust
- Disk averaged observations, few wind emission features in the optical and UV

Red Supergiants & Red Giants

	Red Supergiants	Red Giants
Mass (M_{\odot})	$\sim 8 \rightarrow \sim 40$	$\sim 0.4 \rightarrow \sim 8$
Radius (R_{\odot})	~ 500	~ 50
Lifetime (yr)	$\sim 10^6$	$\sim 10^9$
dM/dt ($M_{\odot} \text{ yr}^{-1}$)	$\sim 10^{-5}$	$\sim 10^{-10}$

$$\blacksquare H_p/R_* \sim 10^{-2}$$

$$\blacksquare V_{\infty} < V_{\text{escape}}(R_*)$$

$$\blacksquare T_{\text{wind}} < 10^4 \text{ K}$$

■ No detected coronae, radiation field peaks in IR, small pulsation amplitudes, and little dust


■ Disk averaged observations, few wind emission features in the optical and UV

■ **Thesis Goal:** Improve understanding of outflow conditions → Radio

Aperture Synthesis




Aperture Synthesis




- 1) **Spatial resolution:** λ/B_{\max} (B_{\max} = longest projected baseline)

Aperture Synthesis



- 1) **Spatial resolution:** λ/B_{max} (B_{max} = longest projected baseline)
- 2) **Field of View:** λ/D (D = individual antenna diameter)

Aperture Synthesis



- 1) **Spatial resolution:** λ/B_{\max} (B_{\max} = longest projected baseline)
- 2) Field of View: λ/D (D = individual antenna diameter)
- 3) **Resolving out scale:** λ/B_{\min} (B_{\min} = shortest projected baseline)

Stellar Radio Emission

Radio Sky at 4.85 GHz (300ft Green Bank)



Credit: NRAO/AUI

$$(P_{\text{radio}}/P_{\text{optical}})_{\odot} \sim 10^{-15}$$

$S_{\nu=4.6 \text{ GHz}} \sim 40 \mu\text{Jy}$ at $\alpha \text{ Cen}$
(not detectable with 'old' VLA!)

Stellar Radio Emission


Radio Sky at 4.85 GHz (300ft Green Bank)



Credit: NRAO/AUI

$$(P_{\text{radio}}/P_{\text{optical}})_{\odot} \sim 10^{-15}$$


$S_{\nu=4.6 \text{ GHz}} \sim 40 \mu\text{Jy}$ at $\alpha \text{ Cen}$
(not detectable with 'old' VLA!)



1)

Winds of Red Supergiants

Betelgeuse (M2 Iab)




(Decin *et al.*, 2013)

Distance	197 ± 45 parsec
Photospheric Radius	22.5 mas ($950 R_{\odot}$)
Mass	$\sim 15 M_{\odot}$
Origin	O-type main sequence
Mass loss rate	$3 \times 10^{-6} M_{\odot} \text{ yr}^{-1}$
$^{12}\text{C}/^{13}\text{C}$	6 ± 1



Betelgeuse: Circumstellar Environment

- At least two *recent* mass loss phases
- Two distinct shells spectrally resolved at 4.6 μm :
 - S2, moving at 17 km s^{-1}
 - S1, moving at 10 km s^{-1}
 - Spatial extent not directly determined



Betelgeuse: Circumstellar Environment



- At least two *recent* mass loss phases
- Two distinct shells spectrally resolved at 4.6 μm :
 - S2, moving at 17 km s^{-1}
 - S1, moving at 10 km s^{-1}
 - Spatial extent not directly determined
- Plez & Lambert (2002) appear to detect S2 out to $50''$
- IRAM 30 m ($\theta_{\text{HPBW}} \sim 12''$) fails to resolve S2 shell at 1.3 mm
- Single dish $^{12}\text{C}^{16}\text{O}$ mm-observations reveal only high velocity S2 shell.



Betelgeuse: Circumstellar Environment

- At least two *recent* mass loss phases
- Two distinct shells spectrally resolved at 4.6 μm :
 - S2, moving at 17 km s^{-1}
 - S1, moving at 10 km s^{-1}
 - Spatial extent not directly determined
- Plez & Lambert (2002) appear to detect S2 out to $50''$
- IRAM 30 m ($\theta_{\text{HPBW}} \sim 12''$) fails to resolve S2 shell at 1.3 mm
- Single dish $^{12}\text{C}^{16}\text{O}$ mm-observations reveal only high velocity S2 shell.

Goal: Measure both the spatial scales and the velocities of Betelgeuse's outflow region using $^{12}\text{C}^{16}\text{O}$ $J = 2-1$ line as a tracer to sort out puzzling evidence.




CARMA

- **C**ombined **A**rray for **R**esearch in **M**illimeter-wave **A**stronomy
- 15 element interferometer (9 x 6.1 m + 6 x 10.4 m antennas)
- 105 baselines ($n(n-1)/2$) with 5 configurations
- Three bands: 7 mm, 3 mm and 1.3 mm



Credit:
John
Carlstrom


mm Observations




Resolution (") $0.9'' \rightarrow (\sim 40 R_*)$

Resolving out scale (") $20'' \rightarrow (\sim 800 R_*)$

Individual Configurations





(O'Gorman *et al.*, 2012)





No shells in individual configuration image cubes.



Spherically Symmetric Shell





Spherically Symmetric Shell





Spherically Symmetric Shell





Spherically Symmetric Shell





Spherically Symmetric Shell





Spherically Symmetric Shell





Spherically Symmetric Shell





Spherically Symmetric Shell





Spherically Symmetric Shell





Spherically Symmetric Shell




Spherically Symmetric Shell




Spherically Symmetric Shell



Combined Configuration




Combined Configurations



Spatial Extent of Flows


S2 flow not present at low absolute velocities.



$$r_{\text{chan}} = R_{\text{S2}} \sin \left[\cos^{-1} \left(\frac{v_{\text{chan}}}{V_{\text{S2}}} \right) \right]$$


Spatial Extent of Flows

S2 flow not present at low absolute velocities.



$$r_{\text{chan}} = R_{\text{S2}} \sin \left[\cos^{-1} \left(\frac{v_{\text{chan}}}{V_{\text{S2}}} \right) \right]$$

S1 flow



S1 flow: Density consistent with R^{-2} .
Also clumping.

Conclusions


- High spatial resolution configuration resolves out S2 emission providing S1 profile.
- Multiple CARMA configurations provide the high spatial resolution needed to study the inner S1 flow, while ensuring the extended S2 flow is not resolved out.



	S1	S2
Outflow Velocities (km s^{-1})	9.8	14.3
Maximum Spatial Extent (")	4 \rightarrow 6	17
Age (yr)	400 \rightarrow 600	1100

Betelgeuse's Wind Acceleration Region

(Lim *et al.*, 1998)




$$F_\nu = \frac{\pi k \phi^2 T_b}{2\lambda^2}$$

Angular Diameter


Brightness Temperature

Betelgeuse with e-MERLIN (5.2 cm)




(Credit: MERLIN/VLBI national facility)

Betelgeuse with e-MERLIN (5.2 cm)



(Credit: MERLIN/VLBI national facility)



Two unresolved *hot spots*:


$$T_e > 3800 \text{ K}$$

$$T_e > 5400 \text{ K}$$

Where is the photosphere?

Betelgeuse with e-MERLIN (5.2 cm)


Astrometric solution of Harper *et al.*, (2008)




- Hottest source at $4 R_*$.
- At least ~ 3 times the predicted T_e .

Betelgeuse with e-MERLIN (5.2 cm)

Astrometric solution of Harper *et al.*, (2008)




- Hottest source at $4 R_*$.
- At least ~ 3 times the predicted T_e .



Spherically symmetric semi-empirical model of Harper *et al.*, (2001)


Betelgeuse with e-MERLIN (5.2 cm)

Astrometric solution of Harper *et al.*, (2008)



- Hottest source at $4 R_*$.
- At least ~ 3 times the predicted T_e .

Goal: Analyse high resolution archival cm data to search for signatures of hotspots.



Spherically symmetric semi-empirical model of Harper *et al.*, (2001)

Betelgeuse with VLA – Pie Town

VLA



+

Pie Town Antenna



=

e-MERLIN

w/l (cm)	Resolution (")
5.2	80 x 60

VLA + Pie Town


w/l (cm)	Resolution (")
0.7	40 x 25
1.3	80 x 40
2.0	120 x 90
3.5	200 x 130
6.2	380 x 270

Betelgeuse with VLA – Pie Town

Variability 1998-2004

w/l (cm)	Variability
0.7	23%
1.3	27%
2.0	32%
3.5	21%
6.2	35%

Thermal Profile




Betelgeuse with VLA – Pie Town


Variability 1998-2004

w/l (cm)	Variability
0.7	23%
1.3	27%
2.0	32%
3.5	21%
6.2	35%

Thermal Profile



Q band (0.7 cm)




Betelgeuse with VLA – Pie Town


Variability 1998-2004

w/l (cm)	Variability
0.7	23%
1.3	27%
2.0	32%
3.5	21%
6.2	35%


Thermal Profile



Q band (0.7 cm)



Betelgeuse with VLA – Pie Town



- **No clear signature of hot spots any in any maps.**
 - Time dependence?
 - Opacity?


Conclusions

- e-MERLIN has revealed two unresolved *hotspots* separated by $4 R_*$
- One may be at the position of photosphere
- VLA + Pie Town data in good agreement with Lim *et al.* (1998)
- No clear signature of e-MERLIN hotspots in any of the VLA + Pie Town data

2)

Winds of Red Giants

- Atmospheres cannot be spatially resolved at radio wavelengths.
- Wind properties generally traditionally determined by analysing strong UV and optical resonance lines.
- At cm/mm the thermal continuum Planck function depends linearly on T , unlike the UV.
- Continuum flux measurements at cm/mm: opacity is proportional to $\sim \lambda^{2.1} n_e n_{\text{ion}}$.



Goal: *Observe two 'standard' red giants at all possible cm wavelengths to test and improve existing models.*

Red Giant Targets

	Arcturus (α Boo: K2 III)	Aldebaran (α Tau: K5 III)
Distance (pc)	11.3	20.4
Photospheric Radius (R_{\odot})	25.4	44.2
Mass (M_{\odot})	0.8	1.3
Mass loss rate ($M_{\odot} \text{ yr}^{-1}$)	2×10^{-10}	1.6×10^{-11}

- Single, non-dusty, low molecular abundances, and non-pulsating
- Nearby with large angular diameters
- Semi-empirical 1-D chromospheric and wind models that can be directly tested

Karl G. Jansky Very Large Array

Credit:
NRAO




- Full frequency coverage between 1.0 and 50 GHz
- Observations:
 - B config (11 km) (128 MHz bandwidth)
 - S (13 cm) → Q (0.7 cm) band (13th - 22nd Feb 2011)
 - A unique data set

Red Giant Radio Maps


Example: Aldebaran X (3 cm), C (6 cm), and S (10 cm) band

X-band (8 GHz)
 $S_{\nu} = 0.3$ mJy




Contours = $(-2, 2, 4, \dots, 16) \times \sigma$
 $\sigma = 16$ μ Jy

C-band (5 GHz)
 $S_{\nu} = 0.15$ mJy




Contours = $(-2, 2, 4, \dots, 14) \times \sigma$
 $\sigma = 10$ μ Jy

S-band (3 GHz)
 $S_{\nu} = 0.06$ mJy




Contours = $(-3, -2, -1, 1, 2, 3) \times \sigma$
 $\sigma = 18$ μ Jy

Spectral Energy Distribution – α Boo




Spectral Energy Distribution – α Boo



Spectral Indices and Power Laws

$$F_\nu \propto \nu^\alpha$$



■ If, $n_e \propto r^{-p}$


$$T_e \propto r^{-n}$$

then,

$$\alpha = f(p, n)$$

Spectral Indices and Power Laws

$$F_\nu \propto \nu^\alpha$$



■ If, $n_e \propto r^{-p}$

$$T_e \propto r^{-n}$$

then,

$$\alpha = f(p, n)$$


■ α Tau: wind ~ wind optically thin

■ α Boo:


Assume constant velocity wind,

$$T_e \propto r^{-1.65}$$

Spectral Energy Distribution – α Boo



Spectral Energy Distribution – α Boo



2)

Conclusions

- Most comprehensive set of multi- λ radio observations of two *standard* red giants.
- Tested theoretical and semi-empirical atmospheric models.
- Radio spectral indices allow flow properties to be investigated.
 - Optically thin wind for α Tau.
 - Rapidly cooling wind for α Boo.
- New analytical advection wind model for α Boo.

Summary

- Established spatial scales for the two flows in CSE of Betelgeuse
- e-MERLIN results are surprising. Episodic mass-loss mechanism in RSGs?
- 1st multi-wavelength radio study of red giants
- Provide wind diagnostics and updated outflow models.
- Understanding of mass-loss is improving and radio interferometry will continue to play a major role in future developments.

Summary

- Established spatial scales for the two flows in CSE of Betelgeuse
- e-MERLIN results are surprising. Episodic mass-loss mechanism in RSGs?
- 1st multi-wavelength radio study of red giants
- Provide wind diagnostics and updated outflow models.
- Understanding of mass-loss is improving and radio interferometry will continue to play a major role in future developments.

Thank you.