BOOK CHAPTERS

- 1. <u>Snow A.D.</u> and Kisilevsky R. Time relation between glycosaminoglycan accumulation and amyloid deposition during experimental amyloidosis. In: "Amyloidosis", edited by Glenner G.G., Osserman E.F., Benditt E.P., Calkins E., Cohen A.S. and Zucker-Franklin D., p. 149-156, Plenum Press, 1986.
- 2. Kisilevsky R., <u>Snow A.D.</u>, Subrahmanyan L., Boudreau L. and Tan R. What factors are necessary for the induction of AA amyloidosis? (AA amyloid, glycosaminoglycans, amyloid enhancing factor, apo-SAA). In: "Amyloidosis", edited by Marrink J. and Van Rijswijk M.H., p. 301-310, Kluwer Academic Publishers,1986.
- 3. <u>Snow A.D.</u>, Kisilevsky R. and Wight, T.N. Immunolocalization of heparan sulfate proteoglycans to AA amyloid deposition sites in spleen and liver during experimental amyloidosis. In: "Amyloid and Amyloidosis", edited by T. Isobe, S. Araki, F. Uchino, S. Kito and E. Tsubura, Plenum Press, New York, 1988, pp. 87-93.
- 4. <u>Snow A.D.</u>, Mar H., Nochlin D., Raskind M. and Wight T.N. Corpora amylacea in aging and Alzheimer's brain: immunolocalization of chondroitin sulfate and heparan sulfate proteoglycans. In "Amyloid and Amyloidosis", edited by T. Isobe, S. Araki, F. Uchino, S. Kito and E. Tsubura, Plenum Press, New York, 1988, pp. 561-566.
- 5. <u>Snow A.D.</u>, Mar H., Nochlin D. and Wight T.N. Congo red staining on 1 micron de-plasticized sections for detection of lesions in Alzheimer's disease and related disorders. In " Alzheimer's Disease and Related Disorders", edited by K. Iqbal, H.Wisniewski, and B. Winblad, Alan R. Liss inc., New York, 1989, pp. 383-391.
- 6. <u>Snow A.D.</u> The possible involvement of specific proteoglycans in the pathogenesis of Alzheimer's disease, Down's syndrome and other amyloidoses. In "Aging of the Brain. Cellular and Molecular Aspects of Brain Aging and Alzheimer's Disease", edited by T. Nagatsu and O. Hayaishi, Karger Press, New York, 1991, pp. 185-204.
- 7. <u>Snow A.D.</u> Heparan sulfate accumulation in the beta-amyloid protein containing lesions of Down's syndrome occurs prior to the appearance of fibrillar amyloid. In "Amyloidosis", edited by J.B. Natvig,O. Forre, G. Husby, A. Husebekk, B. Skogen, K. Sletten and P. Westermark, Kluwer Academic Publishers. 1991.
- 8. <u>Snow A.D.</u> Peripheral distribution of a dermatan sulfate proteoglycan (decorin) in amyloid plaques and their presence in neurofibrillary tangles in Alzheimer's disease. In "Amyloidosis", edited by J.B. Natvig, O. Forre, G. Husby, A. Husebekk, B. Skogen, K. Sletten and P. Westermark, Kluwer Academic Publishers. 1991.
- 9. Fraser P.E., McLachlan D.R., Surewicz W.K., Mizzen C.A., <u>Snow A.D.</u>, Nguyen J.T., and Kirschner D.A. Conformation and fibrillogenesis of Alzheimer β-amyloid peptides with selected substitution of charged residues. In "Research Advances in Alzheimer's Disease and Related Disorders", edited by K. Iqbal, J.A. Mortimer, B.Winblad, and H.M. Wisniewski, John Wiley & Sons Ltd. 1995.

10. Castillo G.M., Kirschner D.A., Yee A.G., and <u>Snow A.D.</u> Electron microscopy and x-ray diffraction studies further confirm the efficacy of PTI-00703® (Cat's claw derivative) as a potential inhibitor of Alzheimer's beta-amyloid protein fibrillogenesis. In "Alzheimer's Disease: Advances in Etiology, Pathogenesis and Therapeutics. The Proceedings of the 7th International Conference on Alzheimer's Disease and Related Disorders", edited by K. Iqbal, S.S. Sisodia, and B. Winblad, John Wiley & Sons, pp. 450-460, 2001.