

# A Cubical Path from Algebra to Analysis

Lorenzo Molena<sup>1,2</sup>, Marcin Jan Turek-Grzybowski<sup>3</sup>, and Riccardo Borsetto<sup>1</sup>

<sup>1</sup> University of Verona

`lorenzo.molena@studenti.univr.it`, `riccardo.borsetto@univr.it`

<sup>2</sup> University of Stuttgart

<sup>3</sup> Formal Foundry

`marcinjangrzybowski@gmail.com`

## Abstract

We present contributions to the `agda/cubical` library for constructive analysis. Starting from *pseudolattices* and *ordered commutative rings*—algebraic structures adapted from the HoTT Book’s ordered Heyting fields—we build a unified framework that applies uniformly to integers, dyadic rationals, rationals, and reals. We also outline ongoing work on *premetric spaces* following Gilbert’s approach, generalizing the closeness relation to use the positive cone of an ordered commutative ring rather than just the positive rationals.

## 1 Introduction

Constructive analysis in type theory requires careful attention to the algebraic and order-theoretic structures underlying the real numbers. Different constructions of the reals (Cauchy sequences [5], Dedekind cuts [9, 3], Eudoxus reals [2, 11], or HIITs [6]) share common properties: any reasonable definition should be an Archimedean field where all Cauchy sequences of rationals converge [7]. Cubical Agda provides computational univalence and higher inductive types, enabling formalizations where equality has computational content.

We contribute to the `agda/cubical` library [23] with two main developments: (1) algebraic structures for ordered commutative rings and pseudolattices [17, 16, 15], providing a unified framework for reasoning about ordered number types; and (2) ongoing work on premetric spaces following Gilbert [12], with a generalization that allows closeness relations valued in arbitrary ordered commutative rings rather than just the positive rationals. In particular, the HIIT construction of the reals yields Cauchy-completeness directly, unlike naive Cauchy sequence constructions which may fail to be complete [14].

## 2 Ordered Algebraic Structures

We begin with order structure. A *pseudolattice* combines a poset with binary meet ( $\wedge$ ) and join ( $\vee$ ) operations. (The `agda/cubical` library [23] reserves “lattice” for bounded pseudolattices.) We define `IsPseudolattice` with a `Properties` module establishing standard lattice laws and connection lemmas such as  $(a \leq b) \simeq (a = a \wedge b)$ .

Building on this, an *ordered commutative ring* (OCR) combines commutative ring structure with compatible order structure, adapting the HoTT Book’s ordered Heyting fields [19] by dropping invertibility axioms. The record `IsOrderedCommRing` bundles commutative ring and pseudolattice structures with a compatible strict ordering  $<$  satisfying weak linearity ( $x < z \rightarrow (x < y) \vee (y < z)$ ), mixed transitivity, monotonicity of operations, and positivity conditions including  $0 < 1$  and closure of positives under multiplication. These axioms are not all independent [7]; when an OCR additionally forms a Heyting field with apartness, we obtain an *ordered Heyting field* as in the HoTT Book.

We use rings rather than fields for generality: the same structure applies to integers, dyadic rationals [13], rationals, and reals.

Finally, an *Archimedean ring* is an OCR with a cancellability property and the Archimedean property: for all  $x, y$  such that  $0 < x$  and  $0 < y$ , there exists  $n : \mathbb{N}^+$  such that  $x < n \cdot y$ .

### 3 Premetric Spaces

With the algebraic foundations in place, we turn to analysis. Following Gilbert [12], a *premetric space* is a type  $A$  equipped with a closeness relation  $\approx : \mathbb{Q}^+ \rightarrow A \rightarrow A \rightarrow \mathbf{Prop}$ , interpreted as “distance( $x, y$ )  $< \varepsilon$ ”, satisfying reflexivity, symmetry, separatedness, triangularity, and roundness. The HIIT (higher inductive-inductive type) completion  $\mathcal{C}(T)$  of a premetric space  $T$  mutually defines a closeness relation on  $\mathcal{C}(T)$ , with point constructors  $\eta : T \rightarrow \mathcal{C}(T)$  (embedding) and  $\text{lim}$  (limits of Cauchy approximations), and a path constructor identifying arbitrarily close points. This forms an idempotent monad on premetric spaces with Lipschitz functions, yielding the reals as completion of rationals or dyadics.

Our algebraic structures enable a generalization:  $\mathbb{Q}^+$  can be replaced with the positive cone of an arbitrary OCR. We show that any OCR in which  $2 := 1 + 1$  has a multiplicative inverse admits a premetric structure valued in its own positive cone. In particular,  $\mathbb{Q}$  is a standard premetric space. Moreover, any Archimedean ring can be turned into a premetric space: for such a ring, we would like to define a closeness relation by  $|x - y| < \varepsilon$  (where  $|z| := z \vee -z$ ). However, since the rationals do not embed into every Archimedean ring, we multiply both sides by the denominator of  $\varepsilon = \frac{a}{b}$ , obtaining  $b \cdot |x - y| < a$ , which can be expressed using only the Archimedean structure.

In parallel with this generalization, a comprehensive development of Cauchy reals following HoTT Book Chapter 11.3 [22] includes the Riemann integral, the fundamental theorem of calculus, the mean value theorem,  $n$ th roots, and trigonometric functions.

Current work includes adapting the Closeness module [22] to the generalized framework and proving that the HoTT Book reals are initial among Cauchy-complete Archimedean ordered fields [19, 10, 4].

### 4 Conclusion and Future Work

From pseudolattices through ordered commutative rings and Archimedean rings to premetric spaces, we trace a path from algebra to analysis. Unlike approaches that rely on setoids [8, 20, 18], we leverage computational univalence and higher inductive types to work directly with identity types.

For the development of the theory, we follow Bishop [5] and Booi [7], adapting the proofs for arbitrary Archimedean ordered fields with appropriate completeness conditions.

Future work includes implementing the premetric generalization, proving monadicity of the HIIT completion, and comparison with the Agda-UniMath library’s approach to metric spaces [1]. Our long-term goal is to formalize a constructive version of the Picard-Lindelöf theorem, establishing existence and uniqueness of solutions to ordinary differential equations, adapting the proof in Booi’s thesis [7, 21].

## References

- [1] Agda-UniMath Contributors. Agda-unimath: A library of formalized mathematics in Agda. <https://unimath.github.io/agda-unimath/>, 2024. Metric spaces module.
- [2] R. D. Arthan. The Eudoxus Real Numbers, May 2004.
- [3] Andrej Bauer and contributors. A formalization of the Dedekind reals in Coq. <https://github.com/andrejbauer/dedekind-reals>, 2019.
- [4] Andrej Bauer, Jason Gross, Peter LeFanu Lumsdaine, Mike Shulman, Matthieu Sozeau, and Bas Spitters. The HoTT Library: A formalization of homotopy type theory in Coq, December 2016.
- [5] Errett Bishop and Douglas Bridges. *Constructive Analysis*, volume 279 of *Grundlehren Der Mathematischen Wissenschaften*. Springer, Berlin, Heidelberg, 1985.
- [6] Auke Bart Booi. The HoTT reals coincide with the Escardó-Simpson reals, June 2017.
- [7] Auke Bart Booi. *Analysis in Univalent Type Theory*. PhD thesis, University of Birmingham, July 2020.
- [8] D. Bridges and F. Richman. *Varieties of Constructive Mathematics*. Cambridge University Press, 1987.
- [9] R. Dedekind. *Stetigkeit und irrationale Zahlen*. 1872.
- [10] M.H. Escardo and A.K. Simpson. A universal characterization of the closed Euclidean interval. In *Proceedings 16th Annual IEEE Symposium on Logic in Computer Science*, pages 115–125, June 2001.
- [11] S. J. Anna Fokma. The eudoxus reals constructed in homotopy type theory. Master’s thesis, Utrecht University, 2020.
- [12] Gaëtan Gilbert. Formalising real numbers in homotopy type theory. In *Proceedings of the 6th ACM SIGPLAN Conference on Certified Programs and Proofs*, CPP 2017, pages 112–124, New York, NY, USA, January 2017. Association for Computing Machinery.
- [13] David R. Lester. The world’s shortest correct exact real arithmetic program? *Information and Computation*, 216:39–46, July 2012.
- [14] Robert S. Lubarsky. On the Cauchy completeness of the constructive Cauchy reals. *Mathematical Logic Quarterly*, 53(4-5):396–414, 2007.
- [15] Lorenzo Molena. Properties of `Pseudolattice`. Pull Request #1272, <https://github.com/agda/cubical/pull/1272>, 2025. agda/cubical library.
- [16] Lorenzo Molena and Ettore Forigo. Instances for `Pseudolattice` and `OrderedCommRing`. Pull Request #1271, <https://github.com/agda/cubical/pull/1271>, 2025. agda/cubical library.
- [17] Lorenzo Molena, Ettore Forigo, and Marcin Jan Turek-Grzybowski. Definition of `OrderedCommRing`. Pull Request #1246, <https://github.com/agda/cubical/pull/1246>, 2025. agda/cubical library.
- [18] R. S. S. O’Connor. *Incompleteness & Completeness : Formalizing Logic and Analysis in Type Theory*. S.l : s.n., 2009.
- [19] The Univalent Foundations Program. Homotopy Type Theory: Univalent Foundations of Mathematics, August 2013.
- [20] Fred Richman. Real numbers and other completions. *Mathematical Logic Quarterly*, 54(1):98–108, 2008.
- [21] H. L. Royden and P. M. Fitzpatrick. *Real Analysis*. Pearson, 4th edition, 2010.
- [22] Marcin Jan Turek-Grzybowski. Cauchy reals following HoTT book 11.3. Pull Request #1182, <https://github.com/agda/cubical/pull/1182>, 2024. agda/cubical library, draft.
- [23] Andrea Vezzosi, Anders Mörtberg, and Andreas Abel. Cubical Agda: A dependently typed programming language with univalence and higher inductive types. <https://github.com/agda/cubical>, 2019. Cubical Agda library.