

Real Number Definitions, Equivalence Classes, and Embeddings (No Countable Choice)

This document lists the first twenty-three constructive definitions of real numbers, partitions them into equivalence classes, and records the canonical embeddings. Note that while many equivalences hold in standard constructive mathematics (assuming Countable Choice, AC_ω), in plain Cubical Agda without AC_ω , the hierarchy is more fractured (e.g., Cauchy reals are not provably Dedekind reals). General surveys and formalization resources include [95, 109, 5].

1. Definitions (1–33)

1. \mathbb{R}_D : Dedekind reals (located cuts of \mathbb{Q}) [34, 21, 22, 28, 65, 108, 122, 75, 101, 84, 50, 35].
2. \mathbb{R}_C : Cauchy reals (modulated Cauchy sequences of rationals, quotiented) [21, 22, 63, 76, 73, 71, 79].
3. \mathbb{R}_E : Eudoxus reals (almost-homomorphisms $\mathbb{Z} \rightarrow \mathbb{Z}$) [9, 98, 102, 93, 45, 56, 74].
4. $\mathbb{R}_{FC} / \mathbb{R}_I$: fast Cauchy reals / interval reals (Cauchy sequences with explicit moduli or nested rational intervals) [31, 3, 114, 119].
5. \mathbb{R}_{CF} : continued fraction reals (streams of partial quotients) [30, 118, 112].
6. \mathbb{R}_b : coinductive base- b reals (digit streams, e.g., binary/decimal) [115].
7. \mathbb{R}_{SD} : signed-digit reals (streams over $\{-1, 0, 1\}$) [8, 42, 115, 17, 77, 54, 59, 24, 66, 27, 99, 20, 19, 41, 120].
8. \mathbb{R}_{ID} : interval domain reals (maximal elements of the interval domain) [113, 87, 53, 13, 15, 38, 37, 36, 106, 39, 14].
9. \mathbb{R}_L : lower reals (rounded lower sets of \mathbb{Q}) [83, 23, 33].
10. \mathbb{R}_U : upper reals (rounded upper sets of \mathbb{Q}) [83, 23, 33].
11. \mathbb{R}_M : MacNeille reals (double-negation closed cuts) [68, 82, 46, 100].
12. \mathbb{R}_H : HIT/HoTT-book reals (higher inductive type with universal property) [107, 81, 26, 25, 12, 92].
13. \mathbb{R}_{ES} : Escardó–Simpson reals (least Cauchy-complete subobject of \mathbb{R}_D containing \mathbb{Q}) [40, 25].
14. $\mathbb{R}_{\text{formal}}$: formal/locale reals (points of the locale of reals) [67, 51, 52, 91, 49, 84].
15. \mathbb{R}_{init} : initial sequentially modulated Cauchy-complete Archimedean ordered field [40, 25, 78].
16. \mathbb{R}_{term} : terminal Archimedean ordered field [40, 67, 78]. Items 34 and 35 are distinct definitions with different construction proofs, but they result in the same object if the category is well-behaved.

17. $\mathbb{R}_{\text{DedComp}}$: Dedekind-complete ordered field (axiomatic characterization) [67, 52, 11].
18. $\mathbb{R}_{\text{CauComp}}$: Cauchy-complete ordered field (axiomatic characterization of the Cauchy completion) [22, 109].
19. $\mathbb{R}_{\text{Tarski}}$: Archimedean Tarski group reals (characterization via Tarski’s axioms) [104, 35].
20. $[0, 1]_{\text{coalg}}$: unit interval as a terminal coalgebra [40, 4, 80, 89].
21. $\mathbb{R}_{\text{coalg}}^+$: positive reals as a terminal coalgebra [40, 4, 80].
22. Sheaf-theoretic reals: the internal real numbers object in a topos [67, 52, 101, 49, 84].
23. Real numbers object (RNO) in a topos [67, 52, 101, 49, 84]. Items 41 and 42 are essentially the same mathematical object described from two different points of view (internal language vs. category theory).
24. \mathbb{R}_{SDG} : Smooth Reals (synthetic differential geometry). In Synthetic Differential Geometry, the reals are defined to include “nilpotent” infinitesimals (elements d where $d^2 = 0$ but $d \neq 0$). These are distinct from standard Dedekind/Cauchy reals because they violate the field axiom $x \neq 0 \implies x$ is invertible (nilpotents are not invertible). They are a distinct mathematical object internal to a smooth topos, not isomorphic to the usual Cauchy/Dedekind reals [58, 60, 72, 85].
25. $^*\mathbb{R}$: Hyperreals (non-standard analysis). These include infinite and infinitesimal numbers. While usually constructed classically (using ultrafilters), there are constructive approaches (e.g., Palmgren’s constructive non-standard analysis) that result in a structure distinct from \mathbb{R}_D or \mathbb{R}_C . They are strict extensions of the ordinary reals [96, 55, 86].
26. Predicative Reals: In systems stricter than Agda (like those prohibiting impredicativity), Dedekind cuts must be restricted (e.g., to “generalized” or “weak” cuts) to avoid circular definitions. The document hints at this with lower/upper reals (items 9, 10), but specific predicative formalizations often stand alone [44, 29, 88].
27. **No**: Surreal Numbers (Conway’s construction). While the Surreals contain the Reals, the “Real subset” of the Surreals is a valid constructive definition of the reals. Inside **No** there is a canonical embedded copy of \mathbb{R} ; this embedding can be used as yet another definition of the real line [32, 48, 69, 57, 121].
28. Geometric Reals: Defined synthetically in Euclidean Geometry (e.g., Tarski’s axioms for geometry, or Hilbert’s axioms). Defined as “points on a line” rather than arithmetically. Constructively, relating “points on a line” to “Dedekind cuts” is a non-trivial project (requires the Cantor-Dedekind axiom) [104, 105, 16, 90, 116].
29. Computable Reals (Turing): Specifically defined as “Turing machines that output digits”. This is distinct from \mathbb{R}_C because \mathbb{R}_C allows *any* function, whereas Computable Reals restrict the functions to computable ones. In strongly normalizing type theories, every *definable* function is computable (meta-theoretically), so formalising computable reals inside such a system is natural. But this does not by itself make \mathbb{R}_C “the same” as the usual Cauchy reals object; you still have to choose a semantic setting (e.g. an effective topos) where every function in the space is interpreted computably [110, 1, 114, 47].
30. Decimal / Base-10 Cauchy Reals: Reals as equivalence classes of decimal expansions; classically standard, but constructively they are just another representation type akin to digit-based reals [103, 10, 117].

31. Apartness / Located Reals (Bishop Style): Reals as located, rounded lower cuts (or Cauchy sequences with an apartness relation). Bishop’s “Constructive Analysis” uses a specific flavor of Cauchy reals (regular sequences with a fixed modulus of convergence, usually $1/n$). While often isomorphic to standard Cauchy reals, in a strict intensional type theory, the specific choice of modulus makes the type definition distinct [21, 22, 29, 94].
32. Filter-based Completions: Reals as equivalence classes of Cauchy filters (or regular Cauchy filters) on \mathbb{Q} ; conceptually close to Dedekind/Cauchy but more topological [43].
33. Locale-of-Reals Variants: Several flavours exist internally: lower reals, upper reals, rounded reals, etc. Some topos texts distinguish a few different “real objects” as default [111, 46, 51, 84].

2. Equivalence Classes (Provable without Countable Choice)

Each class below consists of definitions that are often equivalent in constructive mathematics with AC_ω . In plain Cubical Agda, equivalences between classes (e.g., A and B) may fail.

A. Dedekind-Type Completions

\mathbb{R}_D , $\mathbb{R}_{\text{formal}}$, Sheaf-theoretic reals, and the real numbers object in a topos all present the Dedekind completion of \mathbb{Q} via localized/topos-theoretic perspectives [67, 51, 52, 84, 49].

B. Cauchy/HIT-Type Completions

\mathbb{R}_C , \mathbb{R}_{FC} , \mathbb{R}_I , \mathbb{R}_H , \mathbb{R}_{init} , \mathbb{R}_{ES} , and (axiomatically) $\mathbb{R}_{\text{CauComp}}$ represent the Cauchy completion, differing only in presentation (explicit modulus, higher inductive, universal property, or internal closure of \mathbb{Q} within \mathbb{R}_D) [107, 26, 25, 76, 99].

C. Representation (Digit/Continued Fraction) Pre-Reals

\mathbb{R}_{CF} , \mathbb{R}_b , \mathbb{R}_{SD} , Decimal/Base-10 reals give concrete digit- or fraction-based streams. These are not literally “the reals” until quotiented; they are “presentations of \mathbb{R} ”. Raw types are not fields because of non-unique encodings, but their quotients by the appropriate equivalence relation coincide with Class B [114, 115, 70, 18, 17, 77, 54, 59, 24, 66, 27, 99, 20, 19].

D. Coalgebraic Subspaces

$[0, 1]_{\text{coalg}}$ and $\mathbb{R}_{\text{coalg}}^+$ describe the unit interval and positive reals as terminal coalgebras. Constructively they model subspaces of \mathbb{R}_D but do not deliver the entire field without additional principles [40, 4, 80, 97, 41].

E. Generalized Cuts

\mathbb{R}_L , \mathbb{R}_U , and \mathbb{R}_M relax locatedness/density requirements. They contain \mathbb{R} as a canonical subobject (or as maximal elements) but are bigger structures and not isomorphic to \mathbb{R} as an ordered field [111, 68, 23, 46, 100].

F. Domain-Theoretic

\mathbb{R}_D sits in domain/locale theory. Its equivalence to Dedekind reals is not provable in plain Cubical Agda, so it remains a separate class. Note that in frameworks like Abstract Stone Duality or general Topos Theory, these often collapse into Class A (Dedekind-type) via duality

results, but internally to Agda without extra axioms, the distinction is maintained [2, 38, 113, 87, 13, 36, 106].

G. Axiomatic/Universal Characterizations

\mathbb{R}_{term} , $\mathbb{R}_{\text{DedComp}}$, and $\mathbb{R}_{\text{Tarski}}$ capture Dedekind-like structures via universal properties or axioms. These are abstract characterizations; one still needs to show they are realized by some concrete construction. They coincide with the usual reals only once classical principles (e.g., choice) are assumed [67, 52, 11].

H. Isolated/Unresolved

\mathbb{R}_{E} (Eudoxus reals) currently lacks a constructive proof of equivalence with either Dedekind or Cauchy completions. We therefore mark it as isolated [9, 98, 93, 45, 56].

2b. Diagrams

Legend

- Solid arrows: provable embeddings in plain Cubical Agda (no countable choice, no LEM).
- Dashed arrows: relationships that typically require countable choice or classical principles.
- “quotient” labels: representations coincide with Cauchy reals after quotienting the appropriate equivalence.

Diagram: Plain Cubical Agda (no CC/LEM)





Diagram: With Countable Choice (illustrative collapse)



ASCII Fallback

```

Representations (raw) --quotient--> RC --embed--> RD --embed--> RL, RU, RM
      |                                     \
      |                                     \-- (subspaces) [0,1], R+ (via coalgebras; caveats)
      +-- R_CF, R_b, R_SD

Cauchy/HIT-type: RC, RFC, RI, [RH, R_init (distinct from RD constructively)], RES

Axiomatic: R_term, R_DedComp, R_Tarski, R_CauComp ... (classically collapse to mainstream)

Domain-theoretic: R_ID ... (related to RD; not provably equivalent)

```

Eudoxus: \mathbb{R}_E ... (isolated; classically related to Cauchy/Dedekind)

2c. Corrections and Cautions

- Embeddings go $\mathbb{R}_D \rightarrow \mathbb{R}_L, \mathbb{R}_U$ by taking lower/upper cuts; the reverse is not constructive.
- Place \mathbb{R}_{ES} with Cauchy-type (the Cauchy closure inside \mathbb{R}_D), not with Dedekind-type.
- Keep \mathbb{R}_H and \mathbb{R}_{init} together and distinct from \mathbb{R}_D (they coincide only with extra principles).
- Treat \mathbb{R}_{ID} as domain-theoretic and distinct in plain Cubical Agda; do not assert equivalence to \mathbb{R}_D without additional structure.
- Signed-digit/base- b /continued-fraction representations coincide with Cauchy only after quotienting and with suitable moduli.

3. Canonical Embeddings (No Countable Choice)

- Cauchy into Dedekind: there is a canonical field embedding $\mathbb{R}_C \hookrightarrow \mathbb{R}_D$ that sends each Cauchy real to the located cut defined by its values [22, 63, 64, 73].
- Cauchy/HIT equivalences: $\mathbb{R}_{FC}, \mathbb{R}_I, \mathbb{R}_H, \mathbb{R}_{init}, \mathbb{R}_{ES}$, and $\mathbb{R}_{CauComp}$ are inter-definable and embed into \mathbb{R}_C (hence into \mathbb{R}_D) [107, 26, 25, 40].
- Representation quotients: the quotient of $\mathbb{R}_{CF}, \mathbb{R}_b$, or \mathbb{R}_{SD} by the digit-equivalence relation embeds into \mathbb{R}_C . Without quotienting there is a surjection obstruction due to non-unique encodings (Booij [27] shows locators provide a choice-free signed-digit conversion for Dedekind reals with extra structure) [114, 18, 70, 115, 17, 77, 54, 59, 24, 66, 99, 20, 19].
- Dedekind to generalized cuts: taking lower (resp. upper) shadows gives embeddings $\mathbb{R}_D \rightarrow \mathbb{R}_L$ and $\mathbb{R}_D \rightarrow \mathbb{R}_U$. Composing with double-negation closure embeds into \mathbb{R}_M [111, 83, 82, 68].
- Coalgebraic subspaces: maps $[0, 1]_{\text{coalg}} \rightarrow [0, 1] \subseteq \mathbb{R}_D$ and $\mathbb{R}_{\text{coalg}}^+ \rightarrow \mathbb{R}^+ \subseteq \mathbb{R}_D$ exist, but surjectivity constructs require additional principles [40, 4, 80].
- Axiomatic to Dedekind/Cauchy: the objects $\mathbb{R}_{\text{term}}, \mathbb{R}_{\text{DedComp}},$ and $\mathbb{R}_{\text{Tarski}}$ admit maps into \mathbb{R}_D matching their universal properties, yet converses rely on classical axioms and are not derivable constructively [67, 52, 11, 35].
- Domain to Dedekind: \mathbb{R}_{ID} maps to \mathbb{R}_D via evaluation at maximal elements, but equivalence is unproven constructively [2, 38, 13, 37].
- Eudoxus: natural maps from \mathbb{R}_E into either \mathbb{R}_C or \mathbb{R}_D require countable choice for surjectivity and therefore remain dashed (non-provable) in this setting [9, 93, 45, 56, 74].

4. Higher Coinductive Types (HCITs) for Signed-Digit Reals

Following Altenkirch [7], we explore defining the signed-digit reals as a *Higher Coinductive Type* (HCIT)—the coinductive dual of HITs/QIITs. An HCIT has constructors (operations) and path constructors (equations built into the type), with its universal property being *terminality* rather than *initiality*.

4.1 Signature

An HCIT for the signed-digit interval \mathbb{I} is specified by:

- **Operations:** $\text{cons} : \text{Digit} \rightarrow \mathbb{I} \rightarrow \mathbb{I}$, $\text{inc}, \text{dec} : \mathbb{I} \rightarrow \mathbb{I}$
- **Inc/dec equations** (carry/borrow propagation):

$$\begin{aligned}\text{inc}(\text{cons } (-1) x) &\equiv \text{cons } 0 (\text{inc } x) \\ \text{inc}(\text{cons } 0 x) &\equiv \text{cons } (+1) (\text{cons } 0 x) \\ \text{inc}(\text{cons } (+1) x) &\equiv \text{cons } (+1) (\text{inc } x)\end{aligned}$$

(symmetric for dec)

- **Generation:** $\forall y. \exists d x. y \equiv \text{cons } d x$
- **Completeness:** $\text{cons } 0 x \equiv \text{inc } y \Rightarrow \text{cons } (-1) x \equiv \text{cons } 0 y$
- **Separation:** $\text{cons } (-1) x \equiv \text{cons } 0 y \Rightarrow \text{cons } 0 x \equiv \text{inc } y$
- **Set truncation:** $\text{isSet}(\mathbb{I})$

The carry/borrow equations $\text{cons } (+1) (\text{cons } (-1) x) \equiv \text{cons } 0 (\text{inc } x)$ are derivable from completeness + separation.

Note: The “no-confusion” axiom $\text{cons } (-1) x \equiv \text{cons } (+1) y \rightarrow \perp$ from Altenkirch’s first attempt (slide 10) is *false* in the quotient model: $[-1 :: 1^\omega]$ and $[+1 :: (-1)^\omega]$ both represent 0.

4.2 Type-theoretic rules

Formation. Given an HCIT signature $\Sigma = (\text{Ops}, \text{Eqs})$:

$$\frac{\Sigma : \text{HCIT-Sig}}{\text{HCIT}(\Sigma) : \text{Type}}$$

Introduction. Operations become point constructors; equations become path constructors:

$$\frac{d : \text{Digit} \quad x : \text{HCIT}(\Sigma)}{\text{cons } d x : \text{HCIT}(\Sigma)} \quad \frac{}{\text{carry} : \text{cons } (+1) (\text{cons } (-1) x) \equiv \text{cons } 0 (\text{inc } x)}$$

Elimination (terminality). For any Σ -algebra A , there is a unique morphism into $\text{HCIT}(\Sigma)$:

$$\frac{A : \Sigma\text{-Alg}}{\text{corec}_A : A.\text{Carrier} \rightarrow \text{HCIT}(\Sigma)}$$

Computation (β). $\text{corec}_A(\text{cons}_A d x) \equiv \text{cons } d (\text{corec}_A x)$

Uniqueness (η). $f : \Sigma\text{-Hom } A \text{ HCIT}(\Sigma) \Rightarrow f = \text{corec}_A$

4.3 Encoding in Cubical Agda

Native HCITs are unavailable in Cubical Agda. The quotient-of-codata encoding $\mathbb{I}_{\text{sd}} = \mathbb{N}^\mathbb{N} / \approx_{\text{sd}}$ recovers the equations but the corecursion principle (terminality) requires lifting through the quotient, which needs AC_ω (countable dependent choice). This matches the obstruction for $\lim_{\mathbb{I}_{\text{sd}}}$ documented in the codebase.

4.4 Comparison with existing approaches

| Approach | Type theory | Equalities | Corecursion | AC _ω -free |
|--------------------|---------------------|--------------------|----------------|-----------------------|
| Quotient of codata | Cubical Agda | via quotient | blocked | No |
| Native HCIT | Cubical + primitive | native | native | Yes |
| Coconditions | dTT/Narya | via coconditions | via comatching | Yes |
| Greatest HITs | Clocked Cubical | guarded rec + HITs | guarded rec | Yes |

References: Altenkirch [7] (HCIT proposal); Lorenzen–Shulman [62] (coconditions); Kristensen–Møgelberg [61] (Greatest HITs); Ahrens–Capriotti–Spadotti [6] (M-types in HoTT); Shulman et al. (Narya, narya.readthedocs.io) (displayed coinductive types).

References

- [1] O. Aberth. *Computable Analysis*. McGraw-Hill, 1980. ISBN: 978-0070000797.
- [2] S. Abramsky and A. Jung. “Domain Theory”. In: *Handbook of Logic in Computer Science*. Vol. 3. Oxford University Press, 1994, pp. 1–168. URL: <http://www.cs.bham.ac.uk/~axj/>.
- [3] J.-R. Abrial and D. Cansell. *Constructing the Reals from the Integers*. EBRP Project Notes. 2021.
- [4] J. Adámek, S. Milius, and L. Moss. *Initial Algebras and Terminal Coalgebras*. Cambridge Tracts in Theoretical Computer Science. Cambridge University Press, 2025. DOI: 10.1017/9781108884112.
- [5] Agda Wiki. *Agda Wiki Portal*. 2024. URL: <https://wiki.portal.chalmers.se/agda/pmwiki.php>.
- [6] Benedikt Ahrens, Paolo Capriotti, and Régis Spadotti. “Non-Wellfounded Trees in Homotopy Type Theory”. In: *Proc. 13th International Conference on Typed Lambda Calculi and Applications (TLCA)*. 2015. arXiv: 1504.02949.
- [7] Thorsten Altenkirch. *The Reals as a Higher Coinductive Type?* Talk at Martín Escardó 60th Birthday Seminar. 2024.
- [8] Todd Waugh Ambridge. “Exact Real Search: Formalised Optimisation and Regression in Constructive Univalent Mathematics”. Implements ternary signed-digit encodings in Agda with verified arithmetic operations. PhD thesis. University of Birmingham, 2024. DOI: 10.48550/arXiv.2401.09270. arXiv: 2401.09270. URL: <https://arxiv.org/abs/2401.09270>.
- [9] R. D. Arthan. “The Eudoxus Real Numbers”. In: *arXiv preprint arXiv:math/0405454* (2004). arXiv: math/0405454. URL: <https://arxiv.org/abs/math/0405454>.
- [10] R. G. Bartle and D. R. Sherbert. *Introduction to Real Analysis*. 4th. Wiley, 2011. ISBN: 978-0-471-43331-6.
- [11] A. Bauer. *On Complete Ordered Fields*. 2019. URL: <https://math.andrej.com/2019/09/09/on-complete-ordered-fields/>.
- [12] A. Bauer. *The Real Numbers in Homotopy Type Theory*. CCA 2016 slides. 2016. URL: <https://math.andrej.com/wp-content/uploads/2016/06/hott-reals-cca2016.pdf>.
- [13] A. Bauer. *The Role of the Interval Domain in Modern Exact Real Arithmetic*. 2007. URL: <https://math.andrej.com/2007/09/18/the-role-of-the-interval-domain-in-modern-exact-real-arithmetic/>.

- [14] A. Bauer and A. Kavkler. “Implementing Real Numbers With RZ”. In: *Electronic Notes in Theoretical Computer Science* 202 (2008), pp. 365–384. DOI: 10.1016/j.entcs.2008.03.027.
- [15] A. Bauer and P. Taylor. “The Dedekind Reals in Abstract Stone Duality”. In: *Electronic Notes in Theoretical Computer Science* 203 (2009), pp. 69–90. DOI: 10.1016/j.entcs.2008.05.018.
- [16] M. Beeson. “A Constructive Version of Tarski’s Geometry”. In: *Annals of Pure and Applied Logic* 166.11 (2015), pp. 1199–1273. DOI: 10.1016/j.apal.2015.07.006.
- [17] Ulrich Berger. “From Coinductive Proofs to Exact Real Arithmetic: Theory and Applications”. In: *Logical Methods in Computer Science* 7.1 (2011). DOI: 10.2168/LMCS-7(1:8)2011.
- [18] Ulrich Berger and Tie Hou. “Coinduction for Exact Real Number Computation”. In: *Theory of Computing Systems* 43 (2008), pp. 38–51. DOI: 10.1007/s00224-007-9017-6.
- [19] Ulrich Berger and Sion Lloyd. “A Coinductive Approach to Verified Exact Real Number Computation”. In: *Electronic Communications of the EASST* 23 (2009). DOI: 10.14279/tuj.eceasst.23.331. URL: <https://eceasst.org/index.php/eceasst/article/view/1598>.
- [20] Ulrich Berger and Monika Seisenberger. “From Coinductive Proofs to Exact Real Arithmetic: Theory and Applications”. In: *Journal of Logic and Analysis* 4 (2012), pp. 1–44. eprint: 1101.2162. URL: <https://arxiv.org/abs/1101.2162>.
- [21] E. Bishop. *Foundations of Constructive Analysis*. McGraw-Hill, 1967. URL: https://openlibrary.org/books/OL5546656M/Foundations_of_constructive_analysis.
- [22] E. Bishop and D. Bridges. *Constructive Analysis*. Springer, 1985. DOI: 10.1007/978-3-642-61667-9.
- [23] I. Blechschmidt and M. Hutzler. *A Constructive Knaster–Tarski Proof of the Uncountability of the Reals*. 2019. arXiv: 1902.07366. URL: <https://arxiv.org/abs/1902.07366>.
- [24] Hans-Juergen Boehm et al. “Exact Real Arithmetic: A Case Study in Higher Order Programming”. In: *Proceedings of the 1986 ACM Conference on LISP and Functional Programming (LFP ’86)*. ACM, 1986, pp. 162–173. DOI: 10.1145/319838.319860.
- [25] A. Booiij. *The HoTT Reals are the Cauchy Completion of the Rationals*. 2020. arXiv: 2004.04582. URL: <https://arxiv.org/abs/2004.04582>.
- [26] A. Booiij. *The HoTT Reals Coincide with the Escardó–Simpson Reals*. 2017. arXiv: 1706.05956. URL: <https://arxiv.org/abs/1706.05956>.
- [27] Auke Booiij. “Extensional Constructive Real Analysis via Locators”. In: *Mathematical Structures in Computer Science* 31 (2021). Choice-free signed-digit conversion via locators, pp. 64–88. eprint: 1805.06781. URL: <https://arxiv.org/abs/1805.06781>.
- [28] D. Bridges and F. Richman. *Varieties of Constructive Mathematics*. Cambridge University Press, 1987. DOI: 10.1017/CB09780511565663.
- [29] D. Bridges et al., eds. *Handbook of Constructive Mathematics*. Cambridge University Press, 2023. DOI: 10.1017/9781009039888.
- [30] Brilliant.org. *Continued Fractions*. 2024. URL: <https://brilliant.org/wiki/continued-fractions/>.
- [31] V. Chernov. *Constructive Real Numbers and Connectedness*. 2021. arXiv: 2108.11189. URL: <https://arxiv.org/abs/2108.11189>.
- [32] J. H. Conway. *On Numbers and Games*. Academic Press, 1976. ISBN: 978-0121863500.

- [33] Coq Development Team. *Constructive Reals Interface*. 2024. URL: <https://coq.inria.fr/stdlib/>.
- [34] R. Dedekind. *Stetigkeit und irrationale Zahlen*. 1872. URL: <https://gdz.sub.uni-goettingen.de/id/PPN23569441X>.
- [35] G. Devillanova. “The Fabulous Destiny of Richard Dedekind”. In: *Atti della Accademia Peloritana* 99 (2021). DOI: 10.1478/AAPP.99S1A17.
- [36] P. Di Gianantonio. “Real Number Computability and Domain Theory”. In: *Information and Computation* 127 (1996), pp. 11–25. DOI: 10.1006/inco.1996.0048.
- [37] A. Edalat. “A Domain-Theoretic Approach to Computability on the Real Line”. In: *Theoretical Computer Science* 210 (1999), pp. 73–98. DOI: 10.1016/S0304-3975(98)00097-8.
- [38] A. Edalat and R. Heckmann. “A Computational Model for Metric Spaces”. In: *Theoretical Computer Science* 193 (1998), pp. 53–73. DOI: 10.1016/S0304-3975(97)00050-9.
- [39] M. Escardó. “PCF Extended with Real Numbers”. In: *Theoretical Computer Science* 162 (1996), pp. 79–115. DOI: 10.1016/0304-3975(95)00250-2.
- [40] M. Escardó and A. Simpson. “A Universal Characterization of the Closed Real Interval”. In: *Theoretical Computer Science* 287 (2002), pp. 513–546. DOI: 10.1016/S0304-3975(01)00178-5.
- [41] Martín Escardó and Alex Simpson. *Abstract Datatypes for Real Numbers in Type Theory*. Representation-independent approach via abstract datatypes for $[-1,1]$; midpoint algebra. 2014. URL: <https://martinescardo.github.io/papers/realadt.pdf>.
- [42] Martín H. Escardó and contributors. *TypeTopology: Various new theorems in constructive univalent mathematics written in Agda*. Includes signed-digit real encodings in TWA. Thesis. Chapter 5. Sig. 2024. URL: <https://github.com/martinescardo/TypeTopology>.
- [43] I. Farah and I. Weiss. “The Reals as Rational Cauchy Filters”. In: *The Journal of Symbolic Logic* (2015). DOI: 10.48550/arXiv.1503.04348. URL: <https://arxiv.org/abs/1503.04348>.
- [44] S. Feferman. “Predicative Foundations of Arithmetic”. In: *Essays on the Foundations of Mathematics*. 1962. URL: <https://math.stanford.edu/~feferman/papers.html>.
- [45] A. Fokma. “The Eudoxus Reals Constructed in Homotopy Type Theory”. MA thesis. TU Eindhoven, 2021. URL: <https://research.tue.nl/en/studentTheses/the-eudoxus-reals-constructed-in-homotopy-type-theory>.
- [46] M. Fourman and D. Grayson. “Formal Spaces”. In: *Studies in Logic*. Vol. 31. North-Holland, 1982, pp. 107–122. URL: <https://www.sciencedirect.com/science/article/pii/S0049237X0800160X>.
- [47] H. Geuvers and M. Niqui. “Constructive Analysis, Types and Exact Real Numbers”. In: *Mathematical Structures in Computer Science* 17 (2007), pp. 879–916. DOI: 10.1017/S0960129506005834.
- [48] H. Gonshor. *An Introduction to the Theory of Surreal Numbers*. Cambridge University Press, 1986. DOI: 10.1017/CB09780511629143.
- [49] C. Grossack. *Life in Johnstone’s Topological Topos*. 2024. URL: <https://grossack.site/2024/07/03/life-in-johnstones-topological-topos.html>.
- [50] J. Hall. *Completeness of Ordered Fields*. Tech. rep. Cal Poly, 2010. URL: <https://digitalcommons.calpoly.edu/mathsp/3>.
- [51] P. T. Johnstone. “Metric Spaces in Topoi”. In: *Category Theory Conference*. Vol. 871. Springer LNM. 1981, pp. 247–269. DOI: 10.1007/BFb0090024.

- [52] P. T. Johnstone. *Sketches of an Elephant: A Topos Theory Compendium*. Oxford University Press, 2002. URL: <https://global.oup.com/academic/product/sketches-of-an-elephant-a-topos-theory-compendium-9780198534259>.
- [53] T. de Jong. “Domain Theory in Constructive and Predicative Univalent Foundations”. PhD thesis. 2023. arXiv: 2301.12405. URL: <https://arxiv.org/abs/2301.12405>.
- [54] Abraham Kaganovsky. “Computing with Exact Real Numbers in a Radix-r System”. In: *Electronic Notes in Theoretical Computer Science* 13 (1998), pp. 92–118. DOI: 10.1016/S1571-0661(05)80217-8.
- [55] H. J. Keisler. *Elementary Calculus: An Infinitesimal Approach*. Prindle, Weber & Schmidt, 1976. URL: <https://www.math.wisc.edu/~keisler/calc.html>.
- [56] A. Keskin. *Eudoxus Reals*. Archive of Formal Proofs. 2025. URL: https://www.isa-afp.org/entries/Eudoxus_Reals.html.
- [57] D. E. Knuth. *Surreal Numbers*. Addison-Wesley, 1974. ISBN: 978-0201038125.
- [58] A. Kock. *Synthetic Differential Geometry*. 2nd. Cambridge University Press, 2006. DOI: 10.1017/CB09780511550812.
- [59] Ulrich Köpp and Helmut Schwichtenberg. “Lookahead Analysis in Exact Real Arithmetic with Logical Methods”. In: *Theoretical Computer Science* 943 (2022), pp. 171–186. DOI: 10.1016/j.tcs.2022.10.003.
- [60] R. Kostecki. *An Introduction to Synthetic Differential Geometry*. 2009. URL: <https://www.fuw.edu.pl/~kostecki/sdg.pdf>.
- [61] Magnus Baunsgaard Kristensen and Rasmus Ejlers Møgelberg. “Greatest HITs: Higher inductive types in coinductive definitions via induction under clocks”. In: *Proc. 37th Annual ACM/IEEE Symposium on Logic in Computer Science (LICS)*. 2022. arXiv: 2102.01969.
- [62] Jarl G. Taxerås Lorenzen and Michael Shulman. *(Co)condition hits the Path*. 2024. arXiv: 2405.12994 [cs.LO].
- [63] R. Lubarsky. “On the Cauchy Completeness of the Constructive Cauchy Reals”. In: *Electronic Notes in Theoretical Computer Science* 167 (2007), pp. 307–323. DOI: 10.1016/j.entcs.2006.05.018.
- [64] R. Lubarsky. “On the Cauchy Completeness of the Constructive Dedekind Reals”. In: *Mathematical Logic Quarterly* 53 (2007), pp. 396–414. DOI: 10.1002/malq.200710009.
- [65] R. Lubarsky and M. Rathjen. “On the Constructive Dedekind Reals”. In: *Logic Colloquium 2004*. Vol. 4514. LNCS. Springer, 2007, pp. 129–142. DOI: 10.1007/978-3-540-72734-7_9.
- [66] Robert S. Lubarsky and Fred Richman. *Signed-Bit Representations of Real Numbers*. 2015. arXiv: 1510.00648. URL: <https://arxiv.org/abs/1510.00648>.
- [67] S. Mac Lane and I. Moerdijk. *Sheaves in Geometry and Logic*. Springer, 1992. DOI: 10.1007/978-1-4612-0927-0.
- [68] H. M. MacNeille. “Partially Ordered Sets”. In: *Transactions of the American Mathematical Society* 42 (1937), pp. 416–460. DOI: 10.1090/S0002-9947-1937-1501929-X.
- [69] L. E. Mamane. “Surreal Numbers in Coq”. In: *Types for Proofs and Programs*. Vol. 3839. LNCS. Springer, 2004, pp. 170–185. DOI: 10.1007/11617990_11.
- [70] Math StackExchange. *Constructive Representation of Real Numbers*. 2024. URL: <https://math.stackexchange.com/questions/5093384>.
- [71] MathOverflow. *Cauchy Real Numbers with and without Modulus*. 2024. URL: <https://mathoverflow.net/questions/289900>.

- [72] MathOverflow. *Constructive Analysis and Synthetic Differential Geometry*. 2024. URL: <https://mathoverflow.net/questions/286187>.
- [73] MathOverflow. *Difference Between Constructive Dedekind and Cauchy Reals in Computation*. 2024. URL: <https://mathoverflow.net/questions/236483>.
- [74] MathOverflow. *Is Bauer–Hanson’s Result There is a Topos Where the Dedekind Reals are Countable...* 2024. URL: <https://mathoverflow.net/questions/453312>.
- [75] MathOverflow. *Locales in Constructive Mathematics*. 2024. URL: <https://mathoverflow.net/questions/275548>.
- [76] Z. Murray. *Constructive Analysis in the Agda Proof Assistant*. 2022. arXiv: 2205.08354. URL: <https://arxiv.org/abs/2205.08354>.
- [77] Milad Niqui. “Coinductive Formal Reasoning in Exact Real Arithmetic”. In: *Logical Methods in Computer Science* 4.3 (2008). DOI: 10.2168/LMCS-4(3:6)2008.
- [78] nLab. *Archimedean Ordered Field*. 2024. URL: <https://ncatlab.org/nlab/show/Archimedean+ordered+field>.
- [79] nLab. *Cauchy Real Number*. 2024. URL: <https://ncatlab.org/nlab/show/Cauchy+real+number>.
- [80] nLab. *Coalgebra of the Real Interval*. 2024. URL: <https://ncatlab.org/nlab/show/coalgebra+of+the+real+interval>.
- [81] nLab. *HoTT Book Real Number*. 2024. URL: <https://ncatlab.org/nlab/show/HoTT+book+real+number>.
- [82] nLab. *MacNeille Real Number*. 2024. URL: <https://ncatlab.org/nlab/show/MacNeille+real+number>.
- [83] nLab. *One-Sided Real Number*. 2024. URL: <https://ncatlab.org/nlab/show/one-sided+real+number>.
- [84] nLab. *Real Numbers Object*. 2024. URL: <https://ncatlab.org/nlab/show/real+numbers+object>.
- [85] nLab. *Smooth Topos*. 2024. URL: <https://ncatlab.org/nlab/show/smooth+topos>.
- [86] E. Palmgren. “Constructive Nonstandard Analysis”. In: *Mathematical Logic Quarterly* 44 (1998), pp. 71–103. DOI: 10.1002/malq.19980440105.
- [87] D. Pattinson and A. Mohammadian. “Constructive Domains with Classical Witnesses”. In: *Logical Methods in Computer Science* 17 (2021). DOI: 10.23638/LMCS-17(3:17)2021.
- [88] C. Paulin-Mohring. “Inductive Definitions in the System Coq - Rules and Properties”. In: *Typed Lambda Calculi and Applications*. Vol. 664. LNCS. Springer, 1993, pp. 328–345. DOI: 10.1007/BFb0037116.
- [89] D. Pavlović and D. Prange. “The Continuum as a Final Coalgebra”. In: *Theoretical Computer Science* 280 (2002), pp. 285–301. DOI: 10.1016/S0304-3975(01)00027-5.
- [90] D. Perout. “Tarski’s Axioms of Euclidean Geometry”. MA thesis. Charles University in Prague, 2013. URL: <http://hdl.handle.net/20.500.11956/57343>.
- [91] J. Picado and A. Pultr. *Frames and Locales*. Springer, 2012. DOI: 10.1007/978-3-0348-0154-6.
- [92] F. Pratali. “The Construction of Real Numbers in Homotopy Type Theory”. MA thesis. University of Pisa, 2016. URL: <https://etd.adm.unipi.it/theses/available/etd-06082016-113314/>.
- [93] PROMYS Research Lab. *The Eudoxus Reals*. 2023. arXiv: 2310.04534. URL: <https://arxiv.org/abs/2310.04534>.

- [94] ResearchGate. *Mathematical Foundations of Real Numbers and its Application in Computation*. 2024. URL: <https://www.researchgate.net/publication/385359856>.
- [95] F. Richman. “Real Numbers in Constructive Mathematics”. In: *American Mathematical Monthly* 104 (1997), pp. 546–551. DOI: 10.1080/00029890.1997.11990683.
- [96] A. Robinson. *Non-standard Analysis*. North-Holland, 1966. ISBN: 978-0691044903.
- [97] J. J. M. M. Rutten. “Universal Coalgebra: A Theory of Systems”. In: *Theoretical Computer Science* 249 (2000), pp. 3–80. DOI: 10.1016/S0304-3975(00)00056-6.
- [98] S. H. Schanuel. “The Eudoxus Real Numbers”. In: *Proceedings of the American Mathematical Society* 114 (1992), pp. 543–552. DOI: 10.1090/S0002-9939-1992-1076586-6.
- [99] Helmut Schwichtenberg and Franziskus Wiesnet. “Limits in the Signed Digit Representation”. In: *Logical Methods in Computer Science* 17.2 (2021). SdLim theorem; explicitly uses CC and DC, 7:1–7:27. eprint: 2103.15702. URL: <https://arxiv.org/abs/2103.15702>.
- [100] M. Shulman. *MacNeille Reals*. 2022. URL: <https://ncatlab.org/nlab/show/MacNeille+completion>.
- [101] L. N. Stout. “Topological Properties of the Real Numbers Object in a Topos”. In: *Cahiers de Topologie et Géométrie Différentielles Catégoriques* 17 (1976), pp. 295–326. URL: http://www.numdam.org/item/CTGDC_1976__17_3_295_0/.
- [102] R. Street. *The Most General Real Numbers*. 2003. arXiv: math/0309030. URL: <https://arxiv.org/abs/math/0309030>.
- [103] T. Tao. *Analysis I*. Springer, 2016. DOI: 10.1007/978-981-10-1789-6.
- [104] A. Tarski. 2nd. University of California Press, 1951. URL: <https://www.rand.org/pubs/reports/R109.html>.
- [105] A. Tarski. “What is Elementary Geometry?” In: *The Axiomatic Method*. North-Holland, 1959, pp. 16–29. DOI: 10.1016/S0049-237X(08)71527-4.
- [106] P. Taylor. *The Dedekind Reals in ASD*. Slides. 2005. URL: <https://paultaylor.eu/slides/05-CCA-Kyoto1.pdf>.
- [107] The Univalent Foundations Program. *Homotopy Type Theory: Univalent Foundations of Mathematics*. 2013. URL: <https://homotopytypetheory.org/book>.
- [108] L. Tomasi. *Fondamenti della Matematica: Sezioni di Dedekind*. University of Padova lecture notes. 2012.
- [109] A. S. Troelstra and D. van Dalen. *Constructivism in Mathematics*. Vol. I–II. North-Holland, 1988. URL: <https://www.sciencedirect.com/bookseries/studies-in-logic-and-the-foundations-of-mathematics/vol/121>.
- [110] A. M. Turing. “On Computable Numbers, with an Application to the Entscheidungsproblem”. In: *Proceedings of the London Mathematical Society* 42 (1937), pp. 230–265. DOI: 10.1112/plms/s2-42.1.230.
- [111] S. Vickers. *Topology via Logic*. Cambridge University Press, 1996. DOI: 10.1017/CB09780511569258.
- [112] J. Vuillemin. “Two Constructive Results in Continued Fractions”. In: *SIAM Journal on Computing* 5.2 (1976), pp. 231–248. DOI: 10.1137/0205019.
- [113] N. van der Weide and D. Frumin. “The Interval Domain in Homotopy Type Theory”. In: *TYPES 2023*. Vol. 14560. LNCS. Springer, 2024, pp. 381–399. DOI: 10.1007/978-3-031-58367-4_18.
- [114] K. Weihrauch. *Computable Analysis*. Springer, 2000. DOI: 10.1007/978-3-642-56999-9.

- [115] T. Wiesnet and T. Kopp. “Limits of Real Numbers in the Binary Signed Digit Representation”. In: *Logical Methods in Computer Science* 18 (2022). DOI: 10.46298/lmcs-18(3:24)2022.
- [116] Wikipedia. *Cantor–Dedekind Axiom*. 2024. URL: https://en.wikipedia.org/wiki/Cantor%E2%80%93Dedekind_axiom.
- [117] Wikipedia. *Construction of the Real Numbers*. 2024. URL: https://en.wikipedia.org/wiki/Construction_of_the_real_numbers.
- [118] Wikipedia. *Continued Fraction*. 2024. URL: https://en.wikipedia.org/wiki/Continued_fraction.
- [119] Wikipedia. *Nested Intervals*. 2024. URL: https://en.wikipedia.org/wiki/Nested_intervals.
- [120] Wikipedia. *Signed-digit representation*. 2024. URL: https://en.wikipedia.org/wiki/Signed-digit_representation.
- [121] Wikipedia. *Surreal Number*. 2024. URL: https://en.wikipedia.org/wiki/Surreal_number.
- [122] A. Zanardo. *La Struttura dei Numeri Reali: Costruzione e Proprietà*. Dept. of Mathematics, Univ. of Padova. 2012.