
a
b
c
d
Acoustics Indices
nROI
aROI
EPS
BIO
MFC
EVNspFract
EAS
TFSD
ACTspMean
EVNspCount
AGI
NDSI
rBA
ACTspFract
ACTspCount
NP
EVNspMean
H_Havrda
H_Renyi
BioEnergy
H_GiniSimpson
ECU
LFC
ECV
H
H_pairedShannon
Hf
ZCR
SNRf
EVNtMean
MEANf
H_gamma
RAOQ
EVNtFraction
M
BGNt
AEI
Ht
ACI
ROU
LEQf
ACTtFraction
ACTtCount
VARf
EVNtCount
EPS_SKEW
BGNf
SKEWt
EPS_KURT
LEQt
VARt
SNRt
MEANt
KURTt
ACTtMean
ADI
HFC
AnthroEnergy
SKEWf
KURTf
all habitats (unbalanced)
all habitats (balanced)
Corr. with species richnesss (Spearman's R)
Coefficient importance
Species richness
nROI
CCC=0.49
MAE=1.02
p=0.01
Arable land
Forest - Subtropical-tropical moist lowland
Forest - Subtropical-tropical moist montane
Forest - Temperate
Plantations
Shrubland - Subtropical-tropical high altitude
Shrubland - Temperate
5 samples/day/site
15 samples/day/site
30 samples/day/site
60 samples/day/site
Fit with 95% Confidence Limits
95% Prediction Limits