

Now. 10 am.


Lets meet 3 hours later. When will we meet? $10+3=\overline{13}(a.m.)$?

Lets meet 15 hours later. 10+15 = 25]

10 am -> 10 pm -> 12 am 2 hrs 1 am

Lets meet 27 hours later $10+27=\overline{37}$ $10 \text{ am} \xrightarrow{24 \text{ hrs}} 10 \text{ am} \xrightarrow{3 \text{ hrs}} 1 \text{ pm}.$

1 am. 1 p.m. are like this in clock.



$$|3 = |2 \times 1 + 1|$$
 $25 = |2 \times 2 + 1|$
 $37 = |2 \times 3 + 1|$

Same remainder.

$$13 = 25 \pmod{12}$$

$$25 \equiv 1 \pmod{12}$$

Classify numbers in congruence class

(have same expression in clock).

Question: Where do we put hegative integers?

Conquence class

	J							0.000			4.		
	1	;	:	:	:		:	:	:	:			
	-24	-23	-22	-21	-20	-19	-18	-17	-16	-15	-14	-	5-12
representation	-12	-11	-10	-9	-8	-7	-6	-5	-4	-3	-2	-	7-12
	0.	1	2	2	4	5	6	7	8	9	10	11	/
	12	13	= 14	15	16	17	18	19	20	21	22	23	112
	12	17		()	10	. ,	10	' ')+12
	24	15	26	27	28	29	30	31	32	33	34	25	V
	~ /	2.7	50	50	40	41	42	43	11	45	46	127	2+12
	36	37	38	39	40	41	12	77	44	73	40	יין	
	v			1	;	1 ,	•	1	;	1	;	1	
	3 March 2 March	-		-		•		•	•		12		
				¥.									
				1				- 1					
				. 1									

12 classes all together (mod 12) Generally. In classes (mod 17) Practice:

- 11) $5 \neq 100 \pmod{12}$ No! $5 \stackrel{\text{H}^2}{\to} 17 \stackrel{\text{H}^2}{\to} 29 \stackrel{\text{H}^2}{\to} 41 \stackrel{\text{H}^2}{\to} 53$ $\stackrel{\text{H}^2}{\to} 65 \stackrel{\text{H}^2}{\to} 77 \stackrel{\text{H}^2}{\to} 89 \stackrel{\text{H}^2}{\to} 103$ (pass 100)
- (2) $5 \not\equiv -7 \pmod{12}$ Yes: $-7 \stackrel{?}{=} 5$
- (3) 2 ≠ 14 (mod 12) Yes!
- (4) 26 ≠ 14 (mod 12) Yes!
- (5) 1 ≠ -1 (mod 12)

Faster method

Q 单 b (mod 12)

Check whether a-b is multiple of 12

∫ Yes → congruent

No → not congruent.