

Loading a Package

```
library(PACKAGE NAME)
```

Reading in Data

```
NAME OF DATASET <- read_csv("PATH & NAME OF DATASET.csv")
```

Note: The name of the dataset will change, but it will always need to have the .csv at the end of its name!

Note: Do not put spaces in the name you give the data set.

Preview a Dataset

```
glimpse(NAME OF DATASET)
```

```
head(NAME OF DATASET) – shows first 6 rows
```

```
names(NAME OF DATASET) – outputs the names of the columns/variables
```

Plotting a One Categorical Variable Bar Plot with Counts

```
ggplot(data = NAME OF DATASET,  
       mapping = aes(x = NAME OF VARIABLE)) +  
  geom_bar(stat = "count") +  
  labs(title = "TITLE FOR GRAPH",  
        x = "TITLE FOR THE X-AXIS",  
        y = "TITLE FOR THE Y-AXIS")
```

Note: This bar plot has the variable names on the x-axis. If the names are squished, then you should use **y = NAME OF VARIABLE** instead of **x = NAME OF VARIABLE**.

Plotting a One Categorical Variable Bar Plot with Proportions

```
ggplot(data = NAME OF DATASET,  
       mapping = aes(x = NAME OF VARIABLE)) +  
  geom_bar(stat = "count", aes(y = ..prop.., group = 1)) +  
  labs(title = "TITLE FOR GRAPH",  
        x = "TITLE FOR THE X-AXIS",  
        y = "TITLE FOR THE Y-AXIS")
```

Note: This bar plot has the variable names on the x-axis. If the names are squished, then you should use **y = NAME OF VARIABLE** instead of **x = NAME OF VARIABLE**.

Creating a Summary Table of Observations of One Categorical Variable

```
NAME OF DATASET |>  
count(NAME OF VARIABLE)
```

Conducting an Exact Binomial Hypothesis Test for One Proportion

```
binom.test(x = NUMBER OF SUCCESSES, n = SAMPLE SIZE, p = NULL VALUE, alternative = "DIRECTION")
```

Note: The alternative direction can be "greater", "less", or "two.sided"

Performing a Chi-Squared Goodness-of-Fit Test (One Categorical Variable)

```
chisq_test(x = NAME OF DATASET,  
  response = NAME OF VARIABLE,  
  p = c("CATEGORY 1" = EXPECTED PROP 1,  
        "CATEGORY 2" = EXPECTED PROP 2,  
        "CATEGORY LAST" = EXPECTED PROP LAST  
  )  
)
```

*Make sure to check conditions first!

Plotting a Two Categorical Variable Bar Plot

```
ggplot(data = NAME OF DATASET,  
  mapping = aes(x = EXPLANATORY VARIABLE,  
                fill = RESPONSE VARIABLE)  
  ) +  
  geom_bar(stat = "count",  
    position = "fill") +  
  labs(title = "TITLE FOR GRAPH",  
    x = "TITLE FOR THE X-AXIS",  
    y = "TITLE FOR THE Y-AXIS")
```

Note: If you want a side-by-side bar plot you need to change position to "dodge". If you want a stacked bar plot, you need change position to "stack".

Creating a Summary Table of Observations from Two Categorical Variables

```
NAME OF DATASET |>  
count(NAME OF VARIABLE 1, NAME OF VARIABLE 2)
```

Creating a Contingency Table of Observed Counts from Two Categorical Variables

```
NAME OF DATASET |>  
  count(EXPLANATORY VARIABLE, RESPONSE VARIABLE) |>  
  pivot_wider(names_from = RESPONSE VARIABLE,  
              values_from = n) |>  
  adorn_totals(where = c("row", "col"))
```

Note: Your explanatory variable should be in the rows and your response variable should be in the columns. So, the variable you insert into names_from should be the response variable you are interested in.

Creating a Contingency Table of Observed Proportions from Two Categorical Variables

```
NAME OF DATASET |>  
  count(EXPLANATORY VARIABLE, RESPONSE VARIABLE) |>  
  pivot_wider(names_from = RESPONSE VARIABLE,  
              values_from = n) |>  
  adorn_totals(where = c("row", "col")) |>  
  adorn_percentages(denominator = "row")
```

Note: Since your explanatory variable (groups) should be in your rows from above, we want to calculate our proportions in respect to the group totals.

Performing a Chi-Square Test (Two Categorical Variables)

```
chisq_test(x = NAME OF DATASET,  
           response = RESPONSE VARIABLE,  
           explanatory = EXPLANATORY VARIABLE)
```

*Make sure to check conditions first!

Calculating Summary Statistics for One Numeric Variable

```
favstats(~ NAME OF VARIABLE, data = NAME OF DATASET)
```

Note: The ~ (top left keyboard) **must** be included *before* the variable's name!

Histogram for One Numeric Variable

```
ggplot(data = NAME OF DATASET,  
  mapping = aes(x = NAME OF VARIABLE)) +  
geom_histogram(binwidth = WIDTH OF BINS, color = "white") +  
  labs(title = "TITLE FOR GRAPH",  
    x = "TITLE FOR THE X-AXIS",  
    y = "TITLE FOR THE Y-AXIS")
```

Note: A histogram **must** have a numeric variable on the x-axis! If your variable has a space in it, you will need to use tick marks.

Dotplot for One Numeric Variable

```
ggplot(data = NAME OF DATASET,  
  mapping = aes(x = NAME OF VARIABLE)) +  
geom_dotplot() +  
  labs(title = "TITLE FOR GRAPH",  
    x = "TITLE FOR THE X-AXIS",  
    y = "TITLE FOR THE Y-AXIS")
```

Note: A dotplot **must** have the variable on the x-axis!

Boxplot for One Numeric Variable

```
ggplot(data = NAME OF DATASET,  
  mapping = aes(x = NAME OF VARIABLE)) +  
geom_boxplot() +  
  labs(title = "TITLE FOR GRAPH",  
    x = "TITLE FOR THE X-AXIS",  
    y = "")
```

Note: This boxplot is horizontal. If you want for your boxplot to be vertical, in the mapping aes(), you use **y =** instead of **x =**. Keep in mind you will need to change the location of you axis label, too!

Performing a t-test for One Mean (and Confidence Interval)

```
t_test(x = NAME OF DATASET,  
      response = NAME OF VARIABLE,  
      mu = VALUE FROM NULL HYPOTHESIS FOR Mu,  
      alternative = "two-sided",  
      conf_level = 0.95)
```

Note: If you want a 90% confidence interval, you change `conf_level` to 0.90. If you want a 99% confidence interval, you change `conf_level` to 0.99

Note: If you are doing a one-sided hypothesis test, you change `alternative` to either “greater” or “less”

Calculating Summary Statistics for One Numerical Variable and One Categorical Variable

```
favstats(NAME OF NUMERICAL VARIABLE ~ NAME OF CATEGORICAL VARIABLE,  
        data = NAME OF DATASET)
```

Note: The ~ **must** be included! This is from the mosaic plot.

Faceted Histograms

```
ggplot(data = NAME OF DATASET,  
      mapping = aes(x = NAME OF NUMERICAL VARIABLE)) +  
  geom_histogram(binwidth = WIDTH OF BINS, color = "white") +  
  facet_wrap(~NAME OF CATEGORICAL VARIABLE) +  
  labs(title = "TITLE FOR GRAPH",  
        x = "TITLE FOR THE X-AXIS",  
        y = "TITLE FOR THE Y-AXIS")
```

Note: A histogram **must** have the variable on the x-axis!

Side-by-Side Boxplots

```
ggplot(data = NAME OF DATASET,  
      mapping = aes(x = NAME OF NUMERICAL VARIABLE,  
                    y = NAME OF CATEGORICAL VARIABLE)  
      ) +  
  geom_boxplot()+  
  labs(title = "TITLE FOR GRAPH",  
        x = "TITLE FOR THE X-AXIS",  
        y = "TITLE FOR THE Y-AXIS")
```

Note: For vertically stacked boxplots, the categorical variable should be on the y-axis. For horizontally stacked boxplots, the categorical variable should be on the x-axis.

Performing a Two-Sample Independent t-test (Difference in Means)

```
t_test(x = NAME OF DATASET,  
      response = NAME OF NUMERICAL VARIABLE,  
      explanatory = NAME OF CATEGORICAL VARIABLE,  
      mu = 0,  
      conf_int = TRUE,  
      conf_level = 0.95,  
      alternative = "two-sided")
```

Note: If you want a 90% or 99% confidence interval, you change `conf_level` to 0.90 or 0.99

Note: If you are doing a one-sided hypothesis test, you change `alternative` to either "greater" or "less"