

Documentation for Estimation of PM_{2.5} in western US: Total and Attributed to Wildfires and Prescribed Fires

C.E. Reid¹, M.M. Maestas¹, E. Considine¹, G. Li¹,
N.H.F. French², M. Billmire², M. Jerrett³

¹University of Colorado Boulder and ²Michigan Technological University
and ³University of California, Los Angeles

December 11, 2018

Abstract

The purpose of this document is to provide detailed information about the estimation of PM_{2.5} (total and attributed to prescribed fires and wildfires) that our work could be reproduced. Figure 1 shows the study area of interest.

Contents

1 Ideas, To Do, Resources, etc	3
2 PM2.5 Surface Paper Notes	4
2.1 Papers published in Atmospheric Environment - use as style example	4
3 Papers to cite/discuss in Introduction and/or Discussion	4
3.1 Notes on Papers	4
4 Fire attribution paper	5
4.1 text written for the COPD paper - variation of this may be useful	5
5 Data Sources for Machine Learning	6
5.1 PM2.5 Monitor data from US EPA AQS Air Data Query Tool	11
5.2 EPA PM2.5 Plots	14
5.3 PM _{2.5} data from the Federal Land Manager Environmental Database	15
5.4 PM _{2.5} data from the Fire Cache Smoke Monitor Archive	16
5.5 Fire Cache Smoke Monitor (DRI) Plots	19
5.6 California State Air Quality and Meteorological Information System (AQMS) . .	21
5.7 CARB Plots	22
5.8 PM _{2.5} Monitor data from Uintah Basin	23
5.9 PM _{2.5} data from PCAPS in the Salt Lake Valley	25
5.10 Utah Department of Environmental Quality	26

5.11	Utah DEQ Plots	27
5.12	MODIS AOD	28
5.13	GASP-West AOD	30
5.14	MERRA-2	32
5.15	MAIAC	33
5.16	MODIS Thermal Anomalies/Fire Daily L3 Global 1km (MCD14DL)	34
5.17	Landsat-derived burned area essential climate variable (BAECV) fire activity data	35
5.18	MODIS/Terra and Aqua Burned Area Monthly L3 Global 500 m SIN Grid V006 (MCD64A1)	36
5.19	Visible Infrared Imaging Radiometer Suite (VIIRS) (VNP14IMGTDL_NRT)	37
5.20	Classified land cover information from the Landsat-derived NLCD 2011	39
5.21	MODIS Snow Cover Daily L3 Global 500m Grid, Version 6 (MOD10A1 and MYD10A1)	40
5.22	Elevation	41
5.23	Meteorological Data	42
5.24	Dust Storms	45
6	Data Sources for CAMx Modeling of Source-Attributed Air Quality Modeling	46
7	CAMx Modeling	47
8	Compiling Data	48
8.1	Processing PM2.5 data	48
8.1.1	Notes about very high data points	49
8.2	Compare 88101 to 88502 PM2.5	50
9	Machine Learning Methods	51
9.1	ML Scripts	51
10	Machine Learning Results	52
10.1	Geometric Centroids of Counties Images	53
10.2	ML Inputs Time Series Images	54
10.3	ML Inputs Map subset of days Images	87
10.4	ML Inputs Plot against PM2.5 Images	267
10.5	ML Inputs Map monthly medians Images	299
References		659

1 Ideas, To Do, Resources, etc

code up fires by type of land coverage

Consider using the work of Westerling et al for a comprehensive fire history (up through 2012)

<http://science.sciencemag.org/content/313/5789/940>, <http://www.pnas.org/content/108/32/13165>,
<http://rstb.royalsocietypublishing.org/content/371/1696/20150178> Westerling (2016b,a) Also look into the fire histories referenced in Westerling Westerling (2016b,a): http://fam.nwrg.gov/fam-web/weatherfirecd/fire_files.htm and <http://fam.nwrg.gov/fam-web/kcfast/mnmenu.htm> See also <http://www.nifc.gov>

look into the Fire and Smoke Model Evaluation Experiment (FASMEE) <http://www.fasmee.net>

Compare our results with EPA Downscaler <https://www.epa.gov/air-research/downscaler-model-predicting->

Look at Kollanus et al. (2016) again for references for PM2.5 paper, especially the introduction.

Consider using NAAPS in our study.

read ?

read ?

see also <https://www.5280.com/2018/09/can-colorado-burn-its-way-out-of-a-wildfire-crisis/>

Could we use inciweb to distinguish prescribed fires?

look up Global Fire Emissions Database (GFED3) - maybe it would be useful for our study as an input to the machine learning? see Liu et al. (2016)

see ? for potential data sources for ML project

emissions vary by temperature <https://cires.colorado.edu/news/wildfire-temperatures-key-better-understanding> and <https://www.atmos-chem-phys.net/18/9263/2018/>

read Monitoring Trends in Burn Severity MTBS, 2014. Data Access: Fire Level Geospatial Data. US Department of Agriculture, Forest Service and US Department of Interior, Geological Survey. [http://mtbs.gov/data/individualfiredata.html/](http://mtbs.gov/data/individualfiredata.html).

Idea: look at ambulance calls and PM2.5, similar to what Salimi et al. (2016) did in Australia.

read ?

Database of planned/proposed prescribed burns: WRAP's Fire Emissions Tracking System: <http://wrapfets.org/index.cfm>

See Di et al., 2016 and Johnston et al., 2012, Rappold et al., 2014 in ? - combine modelled and monitored/satellite data to estimate PM2.5

See page 11 of ? for discussion of discrepancies related to burned area estimates

<http://www.ptep-online.com/ctan/symbols-a4.pdf>

US National Atlas http://nationalmap.gov/small_scale/atlasftp.html

Thought: Using DigitalGlobe for fire data compared to NASA: would have higher spatial resolution, but not consistently viewing all areas (no cost to CU people)

Look up Openair R package

Papers/resources to look into: https://daac.ornl.gov/cgi-bin/dsviewer.pl?ds_id=1293

https://www.fs.fed.us/psw/publications/4451/psw_2009_4451-001.pdf

<https://labcit.ligo.caltech.edu/~ethrane/Resources/UNIX/>

<https://community.tableau.com/thread/141548>

According to ?, GEOS-Chem “can be classified according to emission source”, that implies that we could tag the emissions as wildfire vs prescribed fire vs urban. Would there be any advantages of this model over CAMx?

could analyze data with NAAQS and WHO PM2.5 standards
projection/datum info: <https://gis.stackexchange.com/questions/664/whats-the-difference-between-a-projected-coordinate-system-and-a-datum>
<http://resources.esri.com/help/9.3/arcgisengine/dotnet/89b720a5-7339-44b0-8b58-0f5bf2843393.htm>
<http://grindgis.com/blog/wgs84-vs-nad83>

Monitoring Trends in Burn Severity (MTBS) MTBS, 2016: Data Access: Fire Level Geospatial Data. USDA Forest Service/U.S. Geological Survey, accessed 8 October 2016, <https://mtbs.gov/direct-download>. Eidenshink, J., B. Schwind, K. Brewer, Z.-L. Zhu, B. Quayle, and S. Howard, 2007: A project for monitoring trends in burn severity. *Fire Ecol.*, 3, 3–21, <https://doi.org/10.4996/fireecology.0301003>.

Idea: Maybe instead of just distance to closest fire, we should follow the example of [Baek2016] and do distributed lags with concentric circles with information about fires in each concentric circle... also, instead of just distance to fire, maybe we could come up with a variable that is something like [distance*size of fire] since both are important.

Fire stats/records: https://www.nifc.gov/fireInfo/fireInfo_statistics.html

See ? for description of fire perimeter data that perhaps we could use (CA only)

See ? for info about MTBS and Active Fire Mapping Program and NWS smoke products. See also Lassman et al ? cited therein.

Read these papers cited in ?: Yao and Henderson, 2014; Henderson et al 2011; Liu et al 2015; Gan et al 2017; and look at their sources of PM2.5 data to see if we could add any of those to our project.

2 PM2.5 Surface Paper Notes

Discussion of trends in anthro PM2.5: ?

2.1 Papers published in Atmospheric Environment - use as style example

Need to go through these papers

- Brokamp et al. (2017) (partially done, done through intro)
- Sampson et al. (2013)
- Anyenda et al. (2016)
- Torvela et al. (2014)
- Whiteman et al. (2014)

Put in [Brokamp et al. \(2017\)](#); [Larsen et al. \(2017\)](#)

3 Papers to cite/discuss in Introduction and/or Discussion

[Westerling \(2016b,a\)](#)

try to find English version <http://80.24.165.149/webproduccion/PDFs/15CAP03.PDF>

For fire identification, consider using NOAA's Hazard Mapping System and BlueSky

3.1 Notes on Papers

See [J. et al. \(2016\)](#) for statistics about wildfires in western US, e.g., % started by humans, number of fires, etc.

4 Fire attribution paper

revisit ?

include ? - does a good job of summarizing the debate about more vs less prescribed burns
sources of fire data ?, ?

will need to compare our work to ?

include [Westerling \(2016b,a\)](#) and [Abatzoglou and Williams \(2016\)](#)

See ? for an alternative method of attributing PM2.5 to wildfire smoke (instead of CAMx)

See Le et al 2014 ?

See Huff et al ?

4.1 text written for the COPD paper - variation of this may be useful

Larsen et al., 2017 [Larsen et al. \(2017\)](#) found that, on average, ground-level PM_{2.5} concentrations increased by 2.9 $\mu\text{g}\cdot\text{m}^{-3}$ (2.8, 3.0) when there was a visible wildfire smoke plume overhead (from satellite imagery), as well as a 2.6 ppb (2.5-2.7) increase in O₃. Satellite data provides a wealth of data and can provide information about air quality where monitors are not present. However, satellite imagery inherently comes with a substantial uncertainty in that satellite data describes the entire atmospheric column and not specifically just air pollution at the ground level, where people are breathing.

5 Data Sources for Machine Learning

For the creation of the spatiotemporal daily exposure surface via machine learning, a large number of data sets will be collected as discussed below. The dependent variable will be daily 24-hour PM_{2.5} from monitoring data.




Figure 1: Map of 11-state study area.

All PM_{2.5} Monitor Locations

All PM2.5 Observation Locations



Figure 2: Map of locations of PM2.5 observations for entire study period, 2008 to 2014.

PM2.5 Observation Locations, 2008




Figure 3: Map of locations of PM2.5 observations during 2008.

PM2.5 Observation Locations, 2009




Figure 4: Map of locations of PM2.5 observations during 2009.

PM2.5 Observation Locations, 2010




Figure 5: Map of locations of PM2.5 observations during 2010.

PM2.5 Observation Locations, 2011




Figure 6: Map of locations of PM2.5 observations during 2011.

PM2.5 Observation Locations, 2012




Figure 7: Map of locations of PM2.5 observations during 2012.

PM2.5 Observation Locations, 2013




Figure 8: Map of locations of PM2.5 observations during 2013.

PM2.5 Observation Locations, 2014




Figure 9: Map of locations of PM2.5 observations during 2014.

5.1 PM2.5 Monitor data from US EPA AQS Air Data Query Tool

Data Source

- **Contact**

Can email the Air Quality Analysis Group (U.S. EPA Office of Air Quality Planning and Standards) on their website at <https://www.epa.gov/outdoor-air-quality-data/forms/contact-us-about-outdoor-air-quality-data>

- **Citation/Link**

United States Environmental Protection Agency. *Pre-Generated Data Files: Daily Summary Files, PM2.5 FRM/FEM Mass (88101) and PM2.5 non FRM/FEM Mass (88502), 2008-2014.* https://aqs.epa.gov/aqsweb/airdata/download_files.html#Daily

- **Data (local)**

- **Geographic Extent**

- **Temporal Extent** 2008 through 2014

- **Acknowledgment**

Download spreadsheet listing all AQS monitors with datums (https://aqs.epa.gov/aqsweb/airdata/aqs_monitors.zip) from "Monitor Listing" at https://aqs.epa.gov/aqsweb/airdata/download_files.html#Meta. The file name is aqs_monitors.csv in the AQS_Daily_Summaries folder in the S3 data.

Brief Description

We will download PM_{2.5} data from both the US EPA AQS Air Data Query Tool ([US EPA, 2017c](#)) and the IMPROVE monitors that capture air quality information in more rural areas ([US EPA, 2017e](#)) for the 11-state region (Figure 1) including any of the following parameter codes: 88101, 88500, 88502, 81104 ([US EPA, 2017a,d,f](#)).

Notes

File Format

Data Filtering and Processing

Final Variable(s)

Methods

1.

2.

Quality Control

Script Names

1.

Data File Names

1. daily_88101_2008.csv
2. daily_88101_2009.csv
3. daily_88101_2010.csv
4. daily_88101_2011.csv
5. daily_88101_2012.csv

6. daily_88101_2013.csv
7. daily_88101_2014.csv
8. daily_88502_2008.csv
9. daily_88502_2009.csv
10. daily_88502_2010.csv
11. daily_88502_2011.csv
12. daily_88502_2012.csv
13. daily_88502_2013.csv
14. daily_88502_2014.csv




Figure 10: Map of 88101 and 88502 PM_{2.5} Monitors.




Figure 11: EPA PM2.5 time series.

5.2 EPA PM2.5 Plots

5.3 PM_{2.5} data from the Federal Land Manager Environmental Database

Data Source

- Citation/Link <http://views.cira.colostate.edu/fed/DataWizard/Default.aspx>
- Download Date March 15, 2018
- Data (local) PM_{2.5} data from the Federal Land Manager Environmental Database
- Geographic Extent Nationwide
- Temporal Extent January 1, 2008 - December 31, 2014
- Acknowledgment - need to fill in

Downloading IMPROVE Aerosol, RHR II (New Equation) data (one parameter at a time):

1. Reports: Raw data
2. Datasets: "IMPROVE Aerosol, RHR II (New Equation)"
3. Sites: select all
4. Parameters:
 - (a) Mass, PM2.5 (Fine): Code MF, Type PM2.5, Units ug/m³ LC AQS ID 88101
 - (b) Mass, PM2.5 Reconstructed (Fine): Code RCFM, Type PM2.5 Units ug/m³ LC, AQS ID 88401
5. Select Dates: By Years and Months: 2008-2014; select all months
6. Aggregations: Non-aggregated
7. Fields: Select All
8. Options: Text File; Generate one file containing all the data; Comma delimited, Standard ("wide" format); Data & Metadata, Display Column Headers, Don't Display Section Titles, String Quotes: Double Quotes, Missing Values (blank); Date Format: 3/14/2002; Display Results: In a separate browser window; Show Report Log
9. Submit

Repeat the downloading steps above, except replace step #2 with these Datasets and parameters:

1. IMPROVE Aerosol, RHR III (DRAFT - Preliminary Most Impaired Days dataset)
 - (a) Mass, PM2.5 (Fine) is listed twice - these turned out to be the same data

File Formats

CSV

Original Data File Names

1. Federal_Land_Manager_IMPROVE_RHR_II_88101_20183151757452922Mvw0s.csv
2. Federal_Land_Manager_IMPROVE_RHR_II_88401_20185113533660420xLwJ.csv
3. Federal_Land_Manager_RHR_III_88101_first_param_201851152033932P22My0.csv

5.4 PM_{2.5} data from the Fire Cache Smoke Monitor Archive

Data Source

- **Contact** Josh Walston at 775-673-7624; Amber Ortega directed us to the website and Scott Landis suggested that a good person to contact about the page would be Mike Broughton from the US Forest Service (michaelbroughton@fs.fed.us)
- **Citation/Link** <https://wrcc.dri.edu/cgi-bin/smoke.pl>
- **Data (local)** PM_{2.5} data from the Fire Cache Smoke Monitor Archive
- **Geographic Extent**
- **Temporal Extent**
- **Acknowledgment**

Brief Description

Notes

Several of the files were password protected, so we contacted Josh and they were able to unlock most of them. As of March 20, 2018, only "Smoke NCFS E-BAM # 3" is still password protected. (Need to try calling Josh again.) Here are some comments that the system administrator passed along to us (via Josh): the data does not get quality controlled, so we should do our own qa/qc. The monitors/data were designed for the fire community to see data in real time, not for research purposes. If we want to speak with the people who ran the monitors, we should contact Josh and the director can probably put us in contact.

Update 2018-05-2018: sent email to Josh requesting the "Smoke NCFS E-BAM # 3" data with flags and with the other formatting settings we used on the other files. Also asked how to determine which datum is associated with the latitude/longitude data.

These monitors were not included because the website indicated that it did not have data during our study period (January 1, 2008 - December 31, 2014):

1. Smoke E-BAM 418
2. Smoke E-BAM 591
3. Smoke E-BAM 592
4. Smoke E-BAM 882
5. Smoke E-BAM 969
6. Smoke USFS R2-922
7. Smoke USFS R2-923
8. Smoke USFS R2-924
9. Smoke USFS R8-34
10. Smoke USFS R8-35
11. Smoke USFS R8-55
12. Smoke USFS R8-56
13. Smoke USFS 3015
14. Smoke USFS 3016
15. Smoke USFS R9-3017
16. Smoke USFS R9-3018
17. RSF Smoke Monitor 1
18. Lolo NF Smoke Monitor #1
19. Lolo NF Smoke Monitor #2

File Formats

.CSV

Data Filtering and Processing

Final Variable(s)

Methods

- 1.
- 2.

Quality Control

Script Names

- 1.

Original Data File Names

1. Fire_Cache_Smoke_DRI_Cache-NCFS-EBAM#3-with-flags.csv
2. Fire_Cache_Smoke_DRI_FWS_Smoke_N1.csv
3. Fire_Cache_Smoke_DRI_Smoke_N11.csv
4. Fire_Cache_Smoke_DRI_Smoke_N13.csv
5. Fire_Cache_Smoke_DRI_Smoke_N15.csv
6. Fire_Cache_Smoke_DRI_Smoke_N16.csv
7. Fire_Cache_Smoke_DRI_Smoke_N17.csv
8. Fire_Cache_Smoke_DRI_Smoke_N19.csv
9. Fire_Cache_Smoke_DRI_Smoke_N20.csv
10. Fire_Cache_Smoke_DRI_Smoke_N21.csv
11. Fire_Cache_Smoke_DRI_Smoke_N22.csv
12. Fire_Cache_Smoke_DRI_Smoke_N23.csv
13. Fire_Cache_Smoke_DRI_Smoke_N24.csv
14. Fire_Cache_Smoke_DRI_Smoke_N25.csv
15. Fire_Cache_Smoke_DRI_Smoke_N65.csv
16. Fire_Cache_Smoke_DRI_Smoke_N66.csv
17. Fire_Cache_Smoke_DRI_Smoke_N67.csv
18. Fire_Cache_Smoke_DRI_Smoke_N68.csv
19. Fire_Cache_Smoke_DRI_Smoke_N69.csv
20. Fire_Cache_Smoke_DRI_Smoke_N84.csv
21. Fire_Cache_Smoke_DRI_Smoke_N86.csv
22. Fire_Cache_Smoke_DRI_Smoke_N215.csv
23. Fire_Cache_Smoke_DRI_Smoke_N216.csv
24. Fire_Cache_Smoke_DRI_Smoke_N217.csv
25. Fire_Cache_Smoke_DRI_Smoke_E_BAM_52.csv
26. Fire_Cache_Smoke_DRI_Smoke_E_BAM_65.csv
27. Fire_Cache_Smoke_DRI_Smoke_E-BAM_231.csv
28. Fire_Cache_Smoke_DRI_Smoke_E-BAM_840.csv
29. Fire_Cache_Smoke_DRI_Smoke_E-BAM_866.csv
30. Fire_Cache_Smoke_DRI_Smoke_E-BAM_925.csv
31. Fire_Cache_Smoke_DRI_Smoke_NCFS_E-BAM_N1.csv

32. Fire_Cache_Smoke_DRI_Smoke_NCFS_E-BAM_N2.csv
33. Fire_Cache_Smoke_DRI_Smoke_USFS_R1-39.csv
34. Fire_Cache_Smoke_DRI_Smoke_USFS_R1-52.csv
35. Fire_Cache_Smoke_DRI_Smoke_USFS_R1-53.csv
36. Fire_Cache_Smoke_DRI_Smoke_USFS_R1-306.csv
37. Fire_Cache_Smoke_DRI_Smoke_USFS_R1-307.csv
38. Fire_Cache_Smoke_DRI_Smoke_USFS_R2-69.csv
39. Fire_Cache_Smoke_DRI_Smoke_USFS_R2-78.csv
40. Fire_Cache_Smoke_DRI_Smoke_USFS_R2-264.csv
41. Fire_Cache_Smoke_DRI_Smoke_USFS_R2-265.csv
42. Fire_Cache_Smoke_DRI_Smoke_USFS_R3-28.csv
43. Fire_Cache_Smoke_DRI_Smoke_USFS_R3-86.csv
44. Fire_Cache_Smoke_DRI_Smoke_USFS_R5-39.csv
45. Fire_Cache_Smoke_DRI_Smoke_USFS_R5-49.csv
46. Fire_Cache_Smoke_DRI_Smoke_USFS_R8-33.csv
47. Fire_Cache_Smoke_DRI_Smoke_USFS_R9-15.csv
48. Fire_Cache_Smoke_DRI_Smoke_USFS_R9-16.csv
49. Fire_Cache_Smoke_DRI_Smoke_USFS_R9-17.csv
50. Fire_Cache_Smoke_DRI_Smoke_USFS_R9-60.csv
51. Fire_Cache_Smoke_DRI_Smoke_NPS_Yosemite_01_California.csv

Processed/Cleaned Data File Names

- 1.
- 2.

Download instructions

1. <https://wrcc.dri.edu/cgi-bin/smoke.pl>
2. Hover over the appropriate drop-down menu and click on the monitor you want to download e.g., “Cache Monitors” then “Smoke #11”
3. On the left-side menu, click on “Data Details”
4. Set the starting date: January 1, 2008 (or as far back as it goes if it doesn’t go back to 2008)
5. Set the ending date: December 31, 2014 (or the last date possible if it ends before 2014)
6. Elements (ignore - default is to include all elements)
7. Options
8. Excel (.xls) (It had html code in the file if I chose other options.)
9. Data Source: Original
10. Represent missing data as: -9999.
11. Include data flags: Yes
12. Date format: MM/DD/YYYY hh:mm
13. Time format: LST 0-23
14. Table Header: Column header short descriptions
15. Field Delimiter: comma (,)
16. Select the Units: Metric
17. Leave Sub interval windows set to: January 01, December 31, Hours: 00 and 24
18. Submit Info




Figure 12: Fire Cache Smoke Monitor (DRI) time series.

19. Open in excel
20. Save as: Fire_Cache_Smoke_DRI_*.csv Where * is the monitor name with spaces replaced with underscore and # symbols replaced with the letter N, e.g., the file name for monitor “Smoke #11” is “Fire_Cache_Smoke_DRI_Smoke_N11.csv”
21. Upload file to S3 bucket: <https://732215511434.signin.aws.amazon.com/console>
22. Click on S3
23. Earthlab-reid-group
24. Fire_Cache_Smoke_DRI (folder)
25. Check the following:
26. The name of the monitor is in cell A1
27. The header is spread across rows 2-4
28. There are 34 columns of data (goes through columns “AH” in excel)
29. Concentration in the 11th (“K”) columns
30. List the file names in the overleaf documentation (PM25_Fire_Cache_Smoke_Monitor_Archive.tex)

5.5 Fire Cache Smoke Monitor (DRI) Plots

Fire Cache Smoke Monitor (DRI) Time Series, < 1000 ug/m³




Figure 13: Fire Cache Smoke Monitor (DRI) time series without data above 1000 ug/m³ so that the majority of data can be seen.

5.6 California State Air Quality and Meteorological Information System (AQMS)

Data Source

- **Contact** Denise Odenwalder, Denise.Odenwalder@arb.ca.gov
- **Citation/Link** To AQMIS: <https://www.arb.ca.gov/aqmis2/aqmis2.php>
- **Data (local)**
- **Geographic Extent** Whole state of California, wherever there are monitors
- **Temporal Extent** 2008-2014, daily averages
- **Acknowledgment** California Air Resources Board was very helpful in gathering and sending us this data.

Brief Description

- PM2.5 measurements at all monitoring stations in CA
- Some entries are 24-hour measurements while others are the average of hourly measurements
- One entry per 3 days

Notes

Reached out to aqmis@arb.ca.gov after determining that there was data being collected in CA that is not published on the EPA AQS website. They emailed us within a week, with a file of the data we requested.

File Formats

xlsx spreadsheet

Data Filtering and Processing

Final Variable(s)

Methods

1.

2.

Quality Control

Script Names

1.

Original Data File Names

1.

2.

Processed/Cleaned Data File Names

1.

2.




Figure 14: CARB time series.

5.7 CARB Plots

5.8 PM_{2.5} Monitor data from Uintah Basin

Data Source

- **Contact** Seth Lyman
- **Citation/Link** seth.lyman@usu.edu
- **Data (local)** PM_{2.5} measurements from 10 sites in Uintah Basin, Utah
- **Geographic Extent** Uintah Basin, Utah
- **Temporal Extent** October 2009 - March 2017
- **Acknowledgment** PM_{2.5} data from the Uintah Basin were provided by Seth Lyman at Utah State University.

Brief Description

PM_{2.5} data were provided by Seth Lyman at Utah State University via email on January 16, 2018. The .xlsx file has PM_{2.5} data from 10 stations during 2009-2017. The .png file has the longitude and latitude of each site.

Notes

Additional information from Seth's email:

"I've attached most of the PM2.5 observations that have ever been collected in the Uintah Basin. What are in the Excel file are 24-hr average data. Data from Roosevelt, Vernal, Ouray, Red Wash, Myton, and Rangely are from the EPA AQS database.

Data from Horsepool are from a BAM 1020 monitor that we operate every winter. Data in Ft. Duchesne and Randlett are 24-hr filter samples that were analyzed gravimetrically. Data from Rabbit Mountain are from a BAM 1020, and data through mid-2013 are in the AQS database.

I have hourly data from Horsepool and Rabbit Mountain if you'd rather have that.

Site locations are given in the list of monitoring stations for the Basin below."

The .png file is easier to read in some programs than others, e.g., it looks fine in "Paint," but not "Photos."

File Formats

Excel and png

Data Filtering and Processing

FinalPM2.5_multiyear_thruwint2017_sheet1.csv is the first sheet of FinalPM2.5_multiyear_thruwint2017.xlsx converted to .csv, and the second row of the header was merged into the first (24hr avg PM2.5).

FinalPM2.5_multiyear_thruwint2017_GISsheet.csv is the third sheet of FinalPM2.5_multiyear_thruwint2017.xlsx converted to .csv and gives the latitude and longitude of each site. This sheet originally did not have location information from the Rangely site, so this was filled in by hand with the numbers from UintahBasinSiteLocations.png.

Final Variable(s)

Methods

1.

2.

Quality Control

Script Names

1.

Original Data File Names

1. FinalPM2.5_multiyear_thruwinter2017.xlsx
2. UintahBasinSiteLocations.png

Processed/Cleaned Data File Names

1. FinalPM2.5_multiyear_thruwinter2017_sheet1.csv
2. UintahBasinSiteLocations.png

5.9 PM_{2.5} data from PCAPS in the Salt Lake Valley

Data Source

- **Contact** Dr. Geoff Silcox in Chemical Engineering at the University of Utah (geoff@chemeng.utah.edu)
- **Citation/Link** Publication: <https://www.sciencedirect.com/science/article/pii/S1352231011011204>
(Data was received from Dr. Silcox via email on February 6, 2018.)
- **Data (local)** PM_{2.5} data from the Persistent Cold Air Pool Study (PCAPS)
- **Geographic Extent** Salt Lake Valley
- **Temporal Extent** January - February, 2011
- **Acknowledgment** Dr. Geoff Silcox

Brief Description

Notes

File Formats

.xlsx

Data Filtering and Processing

PCAPS_Site_Locations.csv is the same data as Table 1 of final_publication.pdf, and has the site locations and elevation.

Final Variable(s)

Methods

1.

2.

Quality Control

Script Names

1.

Original Data File Names

1. final_publication.pdf (Publication of paper)
2. MiniVol_data.xlsx

Processed/Cleaned Data File Names

1. MiniVol_data.csv
2. PCAPS_Site_Locations.csv

5.10 Utah Department of Environmental Quality

Data Source

- Contact
- Citation/Link <http://www.airmonitoring.utah.gov/dataarchive/archpm25.htm>
- Data (local)
- Geographic Extent Varies...
- Temporal Extent Hourly Value CSVs
- Acknowledgment

Brief Description

PM2.5 data from all monitoring stations in Utah

Notes

There was a lot of overlap with the EPA AQS data, so we took data only from the PM2.5 stations not reported by the EPA. This ended up being one or more of three stations (NP, HC, and RS) for 2009, 2010, 2012, and 2013.

Information about the monitoring stations: <http://www.airmonitoring.utah.gov/network/Counties.htm>

Meta information about monitors obtained from <http://www.airmonitoring.utah.gov/dataarchive/2016DailyMaxPM25.pdf>

File Formats

Data Filtering and Processing

Final Variable(s)

Methods

1.

2.

Quality Control

Script Names

1.

Original Data File Names

1.

2.

Processed/Cleaned Data File Names

1.

2.

Utah DEQ Time Series




Figure 15: Utah DEQ time series.

5.11 Utah DEQ Plots

5.12 MODIS AOD

Data Source

- Contact
- Citation/Link
- Data (local)
- Geographic Extent
- Temporal Extent
- Acknowledgment

Brief Description

We will use AOD estimates from the Deep Blue retrieval algorithm for AOD from the MODIS instrument on the NASA Terra and Aqua satellites (MOD04_L2 and MYD04_L2) ([Sayer et al., 2013](#)). The MODIS product is available twice daily at a 10 km spatial resolution for cloud-free scenes and is available longer than our 2008-2014 study period ([NASA LAADS DAAC, 2017a,b](#)).

AOD products use cloud filtering algorithms that often remove pixels in the center of the smoke plumes because they are assumed to be clouds due to high reflectivity ([Kondragunta and Seybold, 2009](#)). Given that these can be in the middle of smoke plumes, often the locations most heavily impacted by smoke have missing data for a key variable, AOD. In our previous work in summer in California when rain clouds are incredibly rare, we could be confident that missing values not along the coast were not clouds. However, for this larger study region and time period, this will be a bigger challenge. We will attempt to isolate smoke plumes from true clouds using satellite imagery and smoke plume polygons from NOAA's Hazard Mapping System Fire Smoke Product ([NOAA OSPO, 2017](#)). We will then estimate missing values within validated smoke plumes, but not within clouds, using radial basis functions as was done in our previous work ([Reid et al., 2015](#)). Radial basis functions are exact interpolation functions that will return observed AOD values where they exist but can interpolate higher values than nearby observations in missing locations, which is needed since the missing values were removed due to their high reflectivity ([Reid et al., 2015](#)).

Notes

File Format

.hdf

Data Filtering and Processing

Final Variable(s)

Methods

1. Step 1: Download the MODIS AOD data sets from both Terra and Aqua sensors:

Using the [NASA EarthData online search tool](#), search for the 'MOD04' (Terra) data set. Set temporal extent by drawing polygon and set spatial extent by adjusting the appropriate filter on the web interface. Select the collection and proceed to download data. For data download options, specify "Stage for Delivery" through the "FTPPull" distribution option. Specify the email address for orders to be sent to. Orders will be sent to your email with instructions on how to connect to the FTP server and pull the ordered data into your local workspace through the command line. Because the amount of data being requested is large, the orders will come through several separate emails. Repeat this step for the 'MYD04'

(Aqua) data set. All of the raw downloaded data from this step will be in .hdf file format.

2. Step 2: Set up file system for data processing:

Create a directory locally named ‘collected_data’. In this directory, make two child directories named "MOD04_terra" and "MYD04_aqua". Follow instructions in email to download data through FTP into the appropriate MODIS directory ('MOD04_terra' or 'MYD04_aqua') depending on whether the order is from the Terra or Aqua sensor.

3. Step 3: Extract lat, long, and aod values from .hdf files and save into .csv files

Run script ‘modis_aod_create_csv_file.py’. This script will take all the .hdf files that you have downloaded and store the lat, long and aod value for non-null pixels from the ‘Deep_Blue_Aerosol_Optical_Satellite’ dataset. A .csv file will be created for each corresponding .hdf file.

4. Step 4: Create .shp file for each .csv file

Run ‘modis_aod_convert_csv_to_shapefile.py’. This script will read in the .csv files and convert them to .shp files using multiprocessing, which speeds up the process.

5. Step 5: Project .shp files to US Albers Equal Area Conic

Run ‘modis_aod_project_to_albers.py’. This script will reproject the .shp files to be US Albers Equal Area Conic (ESRI:102003).

6. Step 6: Combine .shp files for same date and convert to raster with 10km resolution

Run ‘modis_aod_create_daily_averages.py’. This will combine all .shp files from the same date and then produce a raster for each with a 10km resolution. Then, the interpolated grids are clipped to the 11 western states (our study area) with a 100km buffer.

7. Step 7: Extract MODIS AOD value at EPA monitor locations

Using ExtractValuesToPoints tool in ArcGIS.

Quality Control

Script Names

1. modis_aod_create_csv_file.py
2. modis_aod_convert_csv_to_shapefile.py
3. modis_aod_project_to_albers.py
4. modis_aod_create_daily_averages.py

Data File Names

1. western_states_merge.shp

5.13 GASP-West AOD

Data Source

- Contact
- Citation/Link
- Data (local)
- Geographic Extent
- Temporal Extent
- Acknowledgment

Brief Description

We will use AOD estimates from the Geostationary Operational Environmental Satellite West (GOES-West) Aerosol Smoke Product (GASP-West AOD). The GASP product is available at a 4 km resolution at nadir with retrievals every 30 minutes during daylight hours and is available from 2006 onward ([NOAA NCEI, 2017](#)).

AOD products use cloud filtering algorithms that often remove pixels in the center of the smoke plumes because they are assumed to be clouds due to high reflectivity ([Kondragunta and Seybold, 2009](#)). Given that these can be in the middle of smoke plumes, often the locations most heavily impacted by smoke have missing data for a key variable, AOD. In our previous work in summer in California when rain clouds are incredibly rare, we could be confident that missing values not along the coast were not clouds. However, for this larger study region and time period, this will be a bigger challenge. We will attempt to isolate smoke plumes from true clouds using satellite imagery and smoke plume polygons from NOAA's Hazard Mapping System Fire Smoke Product ([NOAA OSPO, 2017](#)). We will then estimate missing values within validated smoke plumes, but not within clouds, using radial basis functions as was done in our previous work ([Reid et al., 2015](#)). Radial basis functions are exact interpolation functions that will return observed AOD values where they exist but can interpolate higher values than nearby observations in missing locations, which is needed since the missing values were removed due to their high reflectivity ([Reid et al., 2015](#)).

Notes

websites: <https://www.ncdc.noaa.gov/data-access/satellite-data/satellite-data-access-datasets>
<https://www.ncdc.noaa.gov/data-access/satellite-data/satellite-data-access-datasets>

Order form for data: <https://www.ncdc.noaa.gov/has/has.dsselect>

<https://www.ncdc.noaa.gov/doclib/index.php?choice=dsi&searchstring=3635&submitted=1&submitted=Search>

File Format

Data Filtering and Processing

Final Variable(s)

Methods

1. Download Data

Navigate to NCEI's [Archive Information Request System \(AIRS\)](#). Scroll down and click on 'Satellite' to expand menu. Click on 'Goes Products' to expand menu. Click on 'Order Data'. Select GOES-West for satellite ID, GASP-AOD-GZ for data type, and appropriate start and end date. Select "Yes" for Submit Batch. Enter email address and submit order. You

will get emails later on with FTP links to your data. Run ‘Generic_FTP_download_to_S3.py’ on an EC2 instance passing in the FTP url as the argument. This will download the data and upload it to S3 (and then delete it off the EC2 instance).

Quality Control

Script Names

1. Generic_FTP_download_to_S3.py

Data File Names

- 1.

5.14 MERRA-2

Data Source

- Contact
- Citation/Link
- Data (local)
- Geographic Extent
- Temporal Extent
- Acknowledgment

Brief Description

Notes

File Formats

Data Filtering and Processing

Final Variable(s)

Methods

1.

2.

Quality Control

Script Names

1.

Original Data File Names

1.

2.

Processed/Cleaned Data File Names

1.

2.

5.15 MAIAC

Data Source

- Contact
- Citation/Link
- Data (local)
- Geographic Extent
- Temporal Extent
- Acknowledgment

Brief Description

Notes

File Format

Data Filtering and Processing

Final Variable(s)

Methods

1.

Quality Control

Script Names

1. Contacted NASA DeepBlue team via email and was given the [FTP](#) site for their research data output. Public data set not yet available. But should be in several months under the name ‘MCD19’.

Data File Names

1. n/a

5.16 MODIS Thermal Anomalies/Fire Daily L3 Global 1km (MCD14DL)

Data Source

- Contact
- Citation/Link
- Data (local)
- Geographic Extent
- Temporal Extent
- Acknowledgment

Brief Description

We will collect data about fire detection locations, size, and fire radiative power from the MODIS Thermal Anomalies/Fire Daily L3 Global 1km (MOD14 and MYD14) ([Giglio et al., 2006](#); [Hawbaker et al., 2017](#)). Using GIS techniques, we will create daily clusters of fire points and use these to calculate: (1) the distance to the nearest fire cluster by day and (2) the sum of Fire Radiative Power (FRP) of the nearest clusters of fires by day as it is likely that smoke levels are higher closer to fires. The MODIS product spans longer than our study period (2008-2014) at daily temporal resolution and has a spatial resolution of 1 km.

Notes

File Format

.shp

Data Filtering and Processing

Final Variable(s)

Methods

1. Navigate to the [NASA EarthData FIRMS Archive Download site](#)
2. Select "Create new Request"
3. In the dropdown for region, select "Custom Region" and draw a bounding box around study area
4. In the dropdown for fire data source, select "MODIS C6"
5. Select dates for study time period
6. In the dropdown for file type, select "Shapefile (.shp)"
7. Enter your email address
8. You will get an email with a download link containing a zipfile with the data

Quality Control

Script Names

1. n/a

Data File Names

1. n/a

5.17 Landsat-derived burned area essential climate variable (BAECV) fire activity data

Data Source

- Contact
- Citation/Link
- Data (local)
- Geographic Extent
- Temporal Extent
- Acknowledgment

Brief Description

We will collect data about fire detection locations, size, and fire radiative power from the Landsat-derived burned area essential climate variable (BAECV) fire activity data, ([LP DAAC, 2017](#)). Using GIS techniques, we will create daily clusters of fire points and use these to calculate: (1) the distance to the nearest fire cluster by day and (2) the sum of Fire Radiative Power (FRP) of the nearest clusters of fires by day as it is likely that smoke levels are higher closer to fires. The BAECV can detect fires larger than 4 km² and provides an estimate of the date of the fire and is available from 1984-2015.

Notes

File Format

.shp

Data Filtering and Processing

Final Variable(s)

Methods

1. BAECV data set already downloaded by EarthLab fire group. Navigate to the ‘earthlab-ls-fire’ S3 bucket, then the v1.1 subdirectory. Here you will find yearly .tar.gz files. Have not spent time decompressing files and exploring data yet but my guess is that within each yearly file, we will find more detailed, daily burn data.

Quality Control

Script Names

1. n/a

Data File Names

1. n/a

5.18 MODIS/Terra and Aqua Burned Area Monthly L3 Global 500 m SIN Grid V006 (MCD64A1)

Data Source

- **Contact**
- **Citation/Link**
- **Data (local)**
- **Geographic Extent**
- **Temporal Extent**
- **Acknowledgment**

Brief Description

We will collect data about fire detection locations, size, and fire radiative power from MODIS/Terra and Aqua Burned Area Monthly L3 Global 500 m SIN Grid V006 (MCD64A1) ([Schroeder et al., 2014](#)). Using GIS techniques, we will create daily clusters of fire points and use these to calculate: (1) the distance to the nearest fire cluster by day and (2) the sum of Fire Radiative Power (FRP) of the nearest clusters of fires by day as it is likely that smoke levels are higher closer to fires.

Notes

File Format

.hdf

Data Filtering and Processing

Final Variable(s)

Methods

1. Run script ‘MODIS_FTP_download.py‘ and pass two arguments: the first is the data set name and the second is the local directory path to save files to (i.e. "MCD64A1" "C:/Users/User/MCD64A1_ Update: ‘MODIS_FTP_download.py‘ is obsolete because NASA LAADS decommissioned their FTP site in favor of HTTPS. So, a new all-purpose script will need to be written to do this download that does HTTPS retrievals instead.

Quality Control

Script Names

1. MODIS_FTP_Download.py

Data File Names

- 1.

5.19 Visible Infrared Imaging Radiometer Suite (VIIRS) (VNP14IMGTDL_NRT)

Data Source

- Contact
- Citation/Link
- Data (local)
- Geographic Extent
- Temporal Extent
- Acknowledgment

Brief Description

We will collect data about fire detection locations, size, and fire radiative power from the Visible Infrared Imaging Radiometer Suite (VIIRS) (VNP14IMGTDL_NRT) ([Schroeder et al., 2014](#)). Using GIS techniques, we will create daily clusters of fire points and use these to calculate: (1) the distance to the nearest fire cluster by day and (2) the sum of Fire Radiative Power (FRP) of the nearest clusters of fires by day as it is likely that smoke levels are higher closer to fires. The MODIS product spans longer than our study period (2008-2014) at daily temporal resolution and has a spatial resolution of 1 km. VIIRS was launched in 2011 and has 12 h temporal resolution with 750 m resolution. The BAECV can detect fires larger than 4 km² and provides an estimate of the date of the fire and is available from 1984-2015.

Notes

File Format

.csv

Data Filtering and Processing

Final Variable(s)

Methods

1. Navigate to the [NASA EarthData FIRMS Archive Download site](#)
2. Select "Create new Request"
3. In the dropdown for region, select "Custom Region" and draw a bounding box around study area
4. In the dropdown for fire data source, select "VIIRS"
5. Select dates for study time period
6. In the dropdown for file type, select "Shapefile (.shp)"
7. Enter your email address
8. You will get an email with a download link containing a zipfile with the data

Quality Control

Script Names

1. n/a

Data File Names

1. fire_archive_V1_2770.csv

5.20 Classified land cover information from the Landsat-derived NLCD 2011

Data Source

- Contact
- Citation/Link
- Data (local)
- Geographic Extent
- Temporal Extent
- Acknowledgment

Brief Description

Classified land cover information from the Landsat-derived NLCD 2011 ([Homer et al., 2017](#)) will be used to calculate estimates of the percentage of urban development (codes 22, 23, and 24), agriculture (codes 81 and 82), and vegetated area other than agricultural land (codes 21, 41, 42, 43, 52, and 71) within buffer radii of 100 m, 250 m, 500 m, and 1000 m around each monitor. The buffer distance that is most highly correlated with PM_{2.5} will be entered into each model. NLCD 2011 has a spatial resolution of 30 m and uses circa 2011 Landsat satellite data.

Notes

File Format

.shp

Data Filtering and Processing

Final Variable(s)

Methods

1. Navigate to the [National Map Viewer](#) and find products for "National Land Cover Database (NLCD)" at the National extent. From the search results, download "NLCD 2011 Land Cover (2011 Edition, amended 2014)". This will download a zipfile with the data.

Quality Control

Script Names

1. n/a

Data File Names

- 1.

5.21 MODIS Snow Cover Daily L3 Global 500m Grid, Version 6 (MOD10A1 and MYD10A1)

Data Source

- **Contact**
- **Citation/Link**
- **Data (local)**
- **Geographic Extent**
- **Temporal Extent**
- **Acknowledgment**

Brief Description

We will use snow cover data from the MODIS Snow Cover Daily L3 Global 500m Grid, Version 6 (MOD10A1 and MYD10A1) ([Hall and Riggs, 2016](#)) because snow coverage is a known contributor to wintertime PM_{2.5} concentrations in mountain valleys ([Whiteman et al., 2014](#)). Daily MOD10A1 and MYD10A1 data are available since 2002 and have 500 m spatial resolution.

Notes

File Format

Data Filtering and Processing

Final Variable(s)

Methods

1. Step 1: Download the MODIS AOD data sets from both Terra and Aqua sensors:

Using the [NASA EarthData online search tool](#), search for the 'MOD10A1' (Terra) data set. Set temporal extent by drawing polygon and set spatial extent by adjusting the appropriate filter on the web interface. Select the collection and proceed to download data. For data download options, specify "Stage for Delivery" through the "FTPPull" distribution option. Specify the email address for orders to be sent to. Orders will be sent to your email with instructions on how to connect to the FTP server and pull the ordered data into your local workspace through the command line. Because the amount of data being requested is large, the orders will come through several separate emails. Repeat this step for the 'MYD10A1' (Aqua) data set. All of the raw downloaded data from this step will be in .hdf file format.

Quality Control

Script Names

- 1.

Data File Names

- 1.

5.22 Elevation

Data Source

- Contact
- Citation/Link
- Data (local)
- Geographic Extent
- Temporal Extent
- Acknowledgment

Brief Description

Elevation can influence PM_{2.5} concentrations; for example, PM_{2.5} can accumulate in mountain valleys during persistent cold air pools (commonly referred to as inversions) during winter ([Whiteman et al., 2014](#)). We will get elevation data from the 3D Elevation Program, which has resolution of 1 arc-second. This resolution is approximately 10 m north/south and varies east/west with latitude ([USGS, 2017](#)).

Notes

File Format

Data Filtering and Processing

Final Variable(s)

Methods

1. Navigate to the [National Map Viewer](#) site and find products for Elevation Products (3DEP), 1 arc-second DEM, IMG file format. Once results are returned, select "Save as Text", which will download a text file containing server links to each NED tile.
2. Download the data using the [download_tiles.py](#) script, which will access the text file that you just downloaded.
3. Extract the elevation values using the [extract_values_to_points.py](#) script.

Quality Control

Script Names

1. `download_tiles.py`
2. `extract_values_to_points.py`

Data File Names

1. n/a

5.23 Meteorological Data

Data Source

North American Mesoscale, Analysis (NAM)

- **Contact**
- **Citation/Link** <https://www.ncdc.noaa.gov/data-access/model-data/model-datasets/north-american-mesonet>
<https://nomads.ncdc.noaa.gov/data/namarl/>
- **Geographic Extent** North America
- **Temporal Extent** Available March, 2004 - present with slight delay
- **Acknowledgment**

Brief Description

We will obtain meteorological data from the North American Mesoscale, Analysis (NAM) because it includes all of the standard meteorological variables, including planetary boundary layer height, which has proved to be an important variable for converting AOD to PM_{2.5} (Liu et al., 2005). We will calculate 24-hour averages from 6-hourly data for temperature, relative humidity, sea level pressure, surface pressure, planetary boundary layer height, dew point temperature, precipitation, snow coverage, and the U and V components of wind speed. NAM has 12 km resolution and is available 2004 onward.

Notes

File Format

Prior to 2018, the files are in *.grb (“grib1”) format, while 2018 data is in *.grb2 (“grib2”) format.

Resources about this file type:

- rNOMADS is an R package for accessing grb* files. It is mostly geared for grib2 files.
<https://cran.r-project.org/web/packages/rNOMADS/rNOMADS.pdf>
- Explanation of what grib files are: http://www.cpc.ncep.noaa.gov/products/wesley/reading_grib.html,
- wgrib program information: <http://www.cpc.ncep.noaa.gov/products/wesley/wgrib.html>

Data Filtering and Processing

1. Use the earthlab/r-reidgroup docker image, which has wgrib and wgrib <http://www.cpc.ncep.noaa.gov/products/wesley/wgrib.html> and wgrib2 <http://www.cpc.ncep.noaa.gov/products/wesley/wgrib2/> installed on it.
2. Run “Define_directories.R” before running each of the steps below. (Need to figure out how to automate this.)
3. Process_NAM_data_step1.R reads in Locations_Dates_of_PM25_Obs_DeDuplicate.csv and outputs Locations_Dates_of_PM25_Obs_DeDuplicate_wNextDay.csv, which includes the next day for each location/day listed in the first file. The purpose of this is so all of the necessary NAM files can be processed. UTC dates can go into the next day for western US time zones. This step uses these input files and R packages and functions:

- (a) Locations_Dates_of_PM25_Obs_DeDuplicate.csv
 - (b) add_next_day_date_loc_function.R.
4. Process_NAM_data_step2_parallel.R downloads each NAM file, extracts relevant data, and deletes the original NAM data. (All of the NAM files together would be about 1.6 Tb.) This file operates in parallel, and will use n-1 cores, where n is the number of cores on the computer. The output is 1 csv with all locations of interest for a given date and time step. The time steps for the NAM are 0Z, 6Z, 12Z, and 18Z. The output files have the format Locations_Dates_of_PM25_Obs_DeDuplicate_YYYY_MM_DD_XXUTC.csv where XX refers to the timestep. Change the study start and stop dates for the dates to be processed. This step uses these input files and R packages and functions:
- (a) Locations_Dates_of_PM25_Obs_DeDuplicate.csv - Data file with dates (local) and locations where you want the NAM data
 - (b) MeteoVariablesNAM.csv - listing of meteorological variables to be extracted from NAM data
 - (c) rNOMADS R package (which calls wgrib and wgrib2) <https://cran.r-project.org/web/packages/rNOMADS/rNOMADS.pdf>
 - (d) parallel R package
 - (e) grb1to2_conversion_prep_function.R - This script downloads the files that will be necessary to run grb1to2.pl, created by the Climate Prediction Center <http://www.cpc.ncep.noaa.gov/products/wesley/grb1to2.html>
 - (f) loop_NAM_run_times.parallel_function.R - this function loops through the time steps on a given day and calls function (listed below) to extract meteo data at locations of interest
 - (g) define_project_bounds_function.R - the bounding box for the study area is defined in this function. The scripts can run faster if the entire NAM domain does not need to be loaded into memory.
 - (h) extract_NAM_data_parallel_function.R - this function extracts the NAM data at points of interest
 - (i) which_type_of_grib_file_function.R - this function determines whether the data for a given time step are grib1 or grib2 format
 - (j) convert_grib1to2_function.R - convert file type from grib1 to grib2, unless it's already a grib2 file. This is essentially a wrapper for grb1to2.pl created by the Climate Prediction Center <http://www.cpc.ncep.noaa.gov/products/wesley/grb1to2.html>
5. Process_NAM_data_step3.R merges all of the files from step 2 into a single file and adds a column at the beginning giving the UTC time stamp.
6. Process_NAM_data_step4.R To Do: deal with time zones and compiling 24-hr summaries
7. To Do: Merge_NAM_times.R will merge the 4 time steps to give a 24-hr summary. Min, max, mean, etc. is set in MeteoVariablesNAM.csv.

Final Variable(s)

See MeteoVariablesNAM.csv

Quality Control

5.24 Dust Storms

Data Source

- Contact
- Citation/Link
- Data (local)
- Geographic Extent
- Temporal Extent
- Acknowledgment

Brief Description

Dust storm records will be included in the machine learning algorithm because they can be a significant indicator of airborne particulate matter from sources other than fires. Dust storm records are available from 1993-2017. The spatial resolution varies, but includes either forecast zone or county ([US National Weather Service, 2017b,c,a](#)).

Notes

File Format

Data Filtering and Processing

Final Variable(s)

Methods

1.

2.

Quality Control

Script Names

1.

Data File Names

1.

6 Data Sources for CAMx Modeling of Source-Attributed Air Quality Modeling

For meteorological inputs, the CAMx modeling will use archived daily 27-km Advanced Research Weather Research and Forecasting (WRF-ARW) grids available via NOAA Real-time Environmental Applications and Display sYstem (READY) servers for the entire study area and time period (Wang et al., 2007; Rolph et al., 2017). For the study years 2008-2012 and 2014, we will use fire emissions datasets prepared by the Western Regional Air Partnership (WRAP) and the National Emissions Inventory (NEI) (US EPA, 2017b) based on aggregated source-tagged fire occurrence data sources, the FCCS (Ottmar et al., 2007), and Consume (Prichard et al., 2009) modeling. For the study year 2013, we will prepare a fire emissions dataset using the same aggregated source-tagged fire occurrence data sources and FCCS/Consume modeling framework in the NASA-funded Wildland Fire Emissions Information System (WFEIS) (MTRI, 2017) developed by Co-I's French and Billmire (French et al., 2014). Fire occurrence datasets include MODIS (MOD14/MYD14 and MCD64A1) and VIIRS (VNP14IMGTDL_NRT) fire data products (Giglio et al., 2006; LP DAAC, 2017; Schroeder et al., 2014). For non-fire emissions during the entire study period, we will use the dataset prepared by WRAP for year 2008.

Look into using spot forecasts to help distinguish between wild and prescribed fires: <http://www.weather.gov/spot/monitor/>

7 CAMx Modeling

8 Compiling Data

8.1 Processing PM2.5 data

These are the scripts that process and compile the PM2.5 data:

1. Script1_Install_Pkgs.R » install packages
2. Define_directories.R » clears all variables and defines directories. Needs to be ran between each of the following scripts. (Want to automate this eventually.) When processing a new batch of data, iterate “processed_data_version” by one letter and create a new subfolder in /home/Processed_Data/ named PM25_data_part_* where * is the new processed_data_version.
3. Process_PM25_data_step1.R » compiles the various PM2.5 data sources into a single data frame. The only eliminations of data are geographic, to remove states that are neither in our study area. Update time frame of study if necessary. The output from this script is a csv file and sink .txt for each PM_{2.5} data source as well as a file with all of the PM2.5 data sources merged together (“PM25_Step1_part_*.csv”).
 - (a) For DRI data, put in flags for voltage data outside the range 11-17 V. (These thresholds are somewhat arbitrary, but it was noticed that when the voltage was outside this range, the PM_{2.5} concentrations were often absurdly high, e.g., greater than 24,000 ug/m³.)
4. Process_PM25_data_step2.R » cleans the data. The following is a list of the quality cuts and changes made to the data:
 - (a) Replace “UNKNOWN” datum in EPA data with “NAD27” per Colleen’s advice.
 - (b) Remove negative and NA PM_{2.5} concentrations. This includes removing all data for a monitor on a given day if any of the hourly observations were negative.
 - (c) For the hourly data, remove monitor-days that do not have at least 18/24 observations.
 - (d) For DRI data, remove data with voltage flags (which includes flags that came with the data and flags that were put in because the battery voltage was outside the range 11-17 V.
 - (e) For DRI data, remove data at or below 0 L/min for flow. Think about whether a minimum value of flow should be set (higher than zero).
 - (f) June 6, 2014 24-hr average PM_{2.5} concentration from monitor “Smoke NCFS E-BAM #1” (Fire_Cache_Smoke_DRI_Smoke_NCFS_E_BAM_N1.csv) is 24,203 ug/m³. There’s nothing apparent wrong with the hourly data, however, this is the only day of data that made it through the other quality checks from this data file. This suggests that this monitor is suspect, and will be removed.
 - (g) Remove data points with lat/lon outside this box: (50,-126) to (25,-101)
 - (h) **To Do** think about making cuts on any unrealistic air temperatures for DRI data
 - (i) **To Do** need to convert missing values that have a -9999 etc to NA value
 - (j) **To Do** merge "24-HR BLK AVG" and "24 HOUR" data together in Sample Duration variable
 - (k) **To Do** figure out why Observation percent has a max value of 200%
 - (l) **To Do** figure out if max AQI value of 546 is reasonable
 - (m) **To Do** Some DRI files looked like they had hour 20:00 data shifted a couple of columns - look into this and fix it.
 - (n) **To Do** Finish filling in Year, month, day information based on date

- (o) **To Do** look over summary() output and plots of every variable and determine if any other cuts are necessary

Process_PM25_data_step3.R » convert all PM2.5 data to the same datum (NAD83) and project coordinates. Take the reprojected location info and put it into the data frame with the daily PM_{2.5} data.

Process_PM25_data_step4.R » combine locations and location/date files for parts b and c. (Part a should be disregarded.) **To Do** write code to take difference between parts bc and d to get a list of just the dates/locations that are new in part d. **To Do** process part d in this file

- 5. Process_PM25_data_step5.R » composite replicate data - in process. **To Do** finish colocated version of code to go with aves version of code. Calls these functions:

- (a) `Combine_true_replicates_R_function.R`
 - (b) `fill_in_aves_coloc_unique_PC_POC_MN_function.R`
 - i. `concatinate_within_column_function.R`
 - (c) `set_data_types_by_column_R_function.R`
- 6. Plot_ML_Input_File.R » create plots, maps, and statistical summary - needs to be changed to take input from De-duplicate code instead of file from Create_ML_Input_File.R
- 7. **to be written** » merge with satellite and other data

8.1.1 Notes about very high data points

June 15, 2012 24-hr average PM_{2.5} concentration from monitor “Smoke #22” (Fire_Cache_Smoke_DRI_Smok is 5,638 ug/m³ - can’t find any reason, so far, to remove this data point, though it’s very odd that the concentrations were low single-digits except for two hours which were extremely high (123,000 and 1000 ug/m³).

88101 and 88502 Time Series




Figure 16: Time series of 88101 and 88502 PM2.5 data.

8.2 Compare 88101 to 88502 PM2.5

9 Machine Learning Methods

setting aside a portion of the PM2.5 data set and then doing 10-fold cross validation on the rest of the data

see <http://www.cvent.com/events/nasa-aist-machine-learning-workshop/event-summary-1f5144a5d1734ca.aspx> and particularly the very end of <https://global.gotomeeting.com/public/recording-player.html?id=owZDmUustOjaW9sJGQ5u9cUG2pBa4D> for list of resources and papers to read.

9.1 ML Scripts

1. ML_PM25_estimation_step0.R » Merge the various predictor variables together with the monitor data or dates/locations of interest
2. ML_PM25_estimation_step0a.R » Plot the training input file
 - (a) predictor variables vs date
 - (b) predictor variables vs PM_{2.5}
3. ML_PM25_estimation_step1.R » ML training algorithms
4. ML_PM25_estimation_step1.R » create data frame of the dates/locations for which we want to predict PM2.5

10 Machine Learning Results

Geometric Centroids of Counties




Figure 17: Geometric Centroids of Counties

10.1 Geometric Centroids of Counties Images




Figure 18: PM2.5 Obs ML Inputs Time Series

10.2 ML Inputs Time Series Images




Figure 19: Date ML Inputs Time Series




Figure 20: Latitude ML Inputs Time Series




Figure 21: Longitude ML Inputs Time Series




Figure 22: A 100 ML Inputs Time Series




Figure 23: C 100 ML Inputs Time Series




Figure 24: Both 100 ML Inputs Time Series




Figure 25: A 250 ML Inputs Time Series




Figure 26: C 250 ML Inputs Time Series




Figure 27: Both 250 ML Inputs Time Series




Figure 28: A 500 ML Inputs Time Series




Figure 29: C 500 ML Inputs Time Series




Figure 30: Both 500 ML Inputs Time Series




Figure 31: A 1000 ML Inputs Time Series




Figure 32: C 1000 ML Inputs Time Series




Figure 33: Both 1000 ML Inputs Time Series




Figure 34: AOD ML Inputs Time Series




Figure 35: MAIAC AOD ML Inputs Time Series




Figure 36: HPBL.surface ML Inputs Time Series




Figure 37: TMP.2.m.above.ground ML Inputs Time Series




Figure 38: RH.2.m.above.ground ML Inputs Time Series




Figure 39: DPT.2.m.above.ground ML Inputs Time Series




Figure 40: APCP.surface ML Inputs Time Series




Figure 41: WEASD.surface ML Inputs Time Series




Figure 42: SNOWC.surface ML Inputs Time Series




Figure 43: UGRD.10.m.above.ground ML Inputs Time Series




Figure 44: VGRD.10.m.above.ground ML Inputs Time Series




Figure 45: PRMSL.mean.sea.level ML Inputs Time Series




Figure 46: PRES.surface ML Inputs Time Series




Figure 47: DZDT.850(mb) ML Inputs Time Series




Figure 48: DZDT.700.mb ML Inputs Time Series




Figure 49: elevation ML Inputs Time Series




Figure 50: NLCD ML Inputs Time Series




Figure 51: PM2.5-Obs 2012-09-18

10.3 ML Inputs Map subset of days Images

PM2.5_Obs 2012-09-19




Figure 52: PM2.5-Obs 2012-09-19

PM2.5_Obs 2012-09-20




Figure 53: PM2.5-Obs 2012-09-20




Figure 54: PM2.5-Obs 2011-05-16




Figure 55: PM2.5-Obs 2011-05-15




Figure 56: PM2.5-Obs 2011-10-05

A_100 2012-09-18




Figure 57: A-100 2012-09-18

A_100 2012-09-19




Figure 58: A-100 2012-09-19

A_100 2012-09-20




Figure 59: A-100 2012-09-20

A_100 2011-05-16




Figure 60: A-100 2011-05-16

A_100 2011-05-15




Figure 61: A-100 2011-05-15

A_100 2011-10-05




Figure 62: A-100 2011-10-05

C_100 2012-09-18




Figure 63: C-100 2012-09-18

C_100 2012-09-19




Figure 64: C-100 2012-09-19

C_100 2012-09-20




Figure 65: C-100 2012-09-20

C_100 2011-05-16




Figure 66: C-100 2011-05-16

C_100 2011-05-15




Figure 67: C-100 2011-05-15

C_100 2011-10-05




Figure 68: C-100 2011-10-05

Both_100 2012-09-18




Figure 69: Both-100 2012-09-18

Both_100 2012-09-19




Figure 70: Both-100 2012-09-19

Both_100 2012-09-20




Figure 71: Both-100 2012-09-20

Both_100 2011-05-16




Figure 72: Both-100 2011-05-16

Both_100 2011-05-15




Figure 73: Both-100 2011-05-15

Both_100 2011-10-05




Figure 74: Both-100 2011-10-05

A_250 2012-09-18




Figure 75: A-250 2012-09-18

A_250 2012-09-19




Figure 76: A-250 2012-09-19

A_250 2012-09-20




Figure 77: A-250 2012-09-20

A_250 2011-05-16




Figure 78: A-250 2011-05-16

A_250 2011-05-15




Figure 79: A-250 2011-05-15

A_250 2011-10-05




Figure 80: A-250 2011-10-05

C_250 2012-09-18




Figure 81: C-250 2012-09-18

C_250 2012-09-19




Figure 82: C-250 2012-09-19

C_250 2012-09-20




Figure 83: C-250 2012-09-20

C_250 2011-05-16




Figure 84: C-250 2011-05-16

C_250 2011-05-15




Figure 85: C-250 2011-05-15

C_250 2011-10-05




Figure 86: C-250 2011-10-05

Both_250 2012-09-18




Figure 87: Both-250 2012-09-18

Both_250 2012-09-19




Figure 88: Both-250 2012-09-19

Both_250 2012-09-20




Figure 89: Both-250 2012-09-20

Both_250 2011-05-16




Figure 90: Both-250 2011-05-16

Both_250 2011-05-15




Figure 91: Both-250 2011-05-15

Both_250 2011-10-05




Figure 92: Both-250 2011-10-05

A_500 2012-09-18




Figure 93: A-500 2012-09-18

A_500 2012-09-19




Figure 94: A-500 2012-09-19

A_500 2012-09-20




Figure 95: A-500 2012-09-20

A_500 2011-05-16




Figure 96: A-500 2011-05-16

A_500 2011-05-15




Figure 97: A-500 2011-05-15

A_500 2011-10-05




Figure 98: A-500 2011-10-05

C_500 2012-09-18




Figure 99: C-500 2012-09-18

C_500 2012-09-19




Figure 100: C-500 2012-09-19

C_500 2012-09-20




Figure 101: C-500 2012-09-20

C_500 2011-05-16




Figure 102: C-500 2011-05-16

C_500 2011-05-15




Figure 103: C-500 2011-05-15

C_500 2011-10-05




Figure 104: C-500 2011-10-05

Both_500 2012-09-18




Figure 105: Both-500 2012-09-18

Both_500 2012-09-19




Figure 106: Both-500 2012-09-19

Both_500 2012-09-20




Figure 107: Both-500 2012-09-20

Both_500 2011-05-16




Figure 108: Both-500 2011-05-16

Both_500 2011-05-15




Figure 109: Both-500 2011-05-15

Both_500 2011-10-05




Figure 110: Both-500 2011-10-05

A_1000 2012-09-18




Figure 111: A-1000 2012-09-18

A_1000 2012-09-19




Figure 112: A-1000 2012-09-19

A_1000 2012-09-20




Figure 113: A-1000 2012-09-20

A_1000 2011-05-16




Figure 114: A-1000 2011-05-16

A_1000 2011-05-15




Figure 115: A-1000 2011-05-15

A_1000 2011-10-05




Figure 116: A-1000 2011-10-05

C_1000 2012-09-18




Figure 117: C-1000 2012-09-18

C_1000 2012-09-19




Figure 118: C-1000 2012-09-19

C_1000 2012-09-20




Figure 119: C-1000 2012-09-20

C_1000 2011-05-16




Figure 120: C-1000 2011-05-16

C_1000 2011-05-15




Figure 121: C-1000 2011-05-15

C_1000 2011-10-05




Figure 122: C-1000 2011-10-05

Both_1000 2012-09-18




Figure 123: Both-1000 2012-09-18




Figure 124: Both-1000 2012-09-19

Both_1000 2012-09-20




Figure 125: Both-1000 2012-09-20

Both_1000 2011-05-16




Figure 126: Both-1000 2011-05-16

Both_1000 2011-05-15




Figure 127: Both-1000 2011-05-15




Figure 128: Both-1000 2011-10-05

AOD 2012-09-18




Figure 129: AOD 2012-09-18

AOD 2012-09-19




Figure 130: AOD 2012-09-19

AOD 2012-09-20




Figure 131: AOD 2012-09-20

AOD 2011-05-16




Figure 132: AOD 2011-05-16

AOD 2011-05-15




Figure 133: AOD 2011-05-15

AOD 2011-10-05




Figure 134: AOD 2011-10-05

MAIAC_AOD 2012-09-18




Figure 135: MAIAC-AOD 2012-09-18

MAIAC_AOD 2012-09-19




Figure 136: MAIAC-AOD 2012-09-19




Figure 137: MAIAC-AOD 2012-09-20




Figure 138: MAIAC-AOD 2011-05-16




Figure 139: MAIAC-AOD 2011-05-15




Figure 140: MAIAC-AOD 2011-10-05

HPBL.surface 2012-09-18




Figure 141: HPBL.surface 2012-09-18

HPBL.surface 2012-09-19




Figure 142: HPBL.surface 2012-09-19

HPBL.surface 2012-09-20




Figure 143: HPBL.surface 2012-09-20

HPBL.surface 2011-05-16




Figure 144: HPBL.surface 2011-05-16

HPBL.surface 2011-05-15




Figure 145: HPBL.surface 2011-05-15

HPBL.surface 2011-10-05




Figure 146: HPBL.surface 2011-10-05

TMP.2.m.above.ground 2012-09-18




Figure 147: TMP.2.m.above.ground 2012-09-18

TMP.2.m.above.ground 2012-09-19




Figure 148: TMP.2.m.above.ground 2012-09-19

TMP.2.m.above.ground 2012-09-20




Figure 149: TMP.2.m.above.ground 2012-09-20

TMP.2.m.above.ground 2011-05-16




Figure 150: TMP.2.m.above.ground 2011-05-16

TMP.2.m.above.ground 2011-05-15




Figure 151: TMP.2.m.above.ground 2011-05-15

TMP.2.m.above.ground 2011-10-05




Figure 152: TMP.2.m.above.ground 2011-10-05

RH.2.m.above.ground 2012-09-18




Figure 153: RH.2.m.above.ground 2012-09-18

RH.2.m.above.ground 2012-09-19




Figure 154: RH.2.m.above.ground 2012-09-19

RH.2.m.above.ground 2012-09-20




Figure 155: RH.2.m.above.ground 2012-09-20

RH.2.m.above.ground 2011-05-16




Figure 156: RH.2.m.above.ground 2011-05-16

RH.2.m.above.ground 2011-05-15




Figure 157: RH.2.m.above.ground 2011-05-15

RH.2.m.above.ground 2011-10-05




Figure 158: RH.2.m.above.ground 2011-10-05

DPT.2.m.above.ground 2012-09-18




Figure 159: DPT.2.m.above.ground 2012-09-18

DPT.2.m.above.ground 2012-09-19




Figure 160: DPT.2.m.above.ground 2012-09-19

DPT.2.m.above.ground 2012-09-20




Figure 161: DPT.2.m.above.ground 2012-09-20

DPT.2.m.above.ground 2011-05-16




Figure 162: DPT.2.m.above.ground 2011-05-16

DPT.2.m.above.ground 2011-05-15




Figure 163: DPT.2.m.above.ground 2011-05-15

DPT.2.m.above.ground 2011-10-05




Figure 164: DPT.2.m.above.ground 2011-10-05

APCP.surface 2012-09-18




Figure 165: APCP.surface 2012-09-18

APCP.surface 2012-09-19




Figure 166: APCP.surface 2012-09-19

APCP.surface 2012-09-20




Figure 167: APCP.surface 2012-09-20

APCP.surface 2011-05-16




Figure 168: APCP.surface 2011-05-16

APCP.surface 2011-05-15




Figure 169: APCP.surface 2011-05-15

APCP.surface 2011-10-05




Figure 170: APCP.surface 2011-10-05

WEASD.surface 2012-09-18




Figure 171: WEASD.surface 2012-09-18

WEASD.surface 2012-09-19




Figure 172: WEASD.surface 2012-09-19

WEASD.surface 2012-09-20




Figure 173: WEASD.surface 2012-09-20

WEASD.surface 2011-05-16




Figure 174: WEASD.surface 2011-05-16

WEASD.surface 2011-05-15




Figure 175: WEASD.surface 2011-05-15

WEASD.surface 2011-10-05




Figure 176: WEASD.surface 2011-10-05

SNOWC.surface 2012-09-18




Figure 177: SNOWC.surface 2012-09-18




Figure 178: SNOWC.surface 2012-09-19




Figure 179: SNOWC.surface 2012-09-20




Figure 180: SNOWC.surface 2011-05-16

SNOWC.surface 2011-05-15




Figure 181: SNOWC.surface 2011-05-15




Figure 182: SNOWC.surface 2011-10-05

UGRD.10.m.above.ground 2012-09-18




Figure 183: UGRD.10.m.above.ground 2012-09-18

UGRD.10.m.above.ground 2012-09-19




Figure 184: UGRD.10.m.above.ground 2012-09-19

UGRD.10.m.above.ground 2012-09-20




Figure 185: UGRD.10.m.above.ground 2012-09-20

UGRD.10.m.above.ground 2011-05-16




Figure 186: UGRD.10.m.above.ground 2011-05-16

UGRD.10.m.above.ground 2011-05-15




Figure 187: UGRD.10.m.above.ground 2011-05-15

UGRD.10.m.above.ground 2011-10-05




Figure 188: UGRD.10.m.above.ground 2011-10-05

VGRD.10.m.above.ground 2012-09-18




Figure 189: VGRD.10.m.above.ground 2012-09-18

VGRD.10.m.above.ground 2012-09-19




Figure 190: VGRD.10.m.above.ground 2012-09-19

VGRD.10.m.above.ground 2012-09-20




Figure 191: VGRD.10.m.above.ground 2012-09-20

VGRD.10.m.above.ground 2011-05-16




Figure 192: VGRD.10.m.above.ground 2011-05-16

VGRD.10.m.above.ground 2011-05-15




Figure 193: VGRD.10.m.above.ground 2011-05-15

VGRD.10.m.above.ground 2011-10-05




Figure 194: VGRD.10.m.above.ground 2011-10-05

PRMSL.mean.sea.level 2012-09-18




Figure 195: PRMSL.mean.sea.level 2012-09-18

PRMSL.mean.sea.level 2012-09-19




Figure 196: PRMSL.mean.sea.level 2012-09-19

PRMSL.mean.sea.level 2012-09-20




Figure 197: PRMSL.mean.sea.level 2012-09-20

PRMSL.mean.sea.level 2011-05-16




Figure 198: PRMSL.mean.sea.level 2011-05-16

PRMSL.mean.sea.level 2011-05-15




Figure 199: PRMSL.mean.sea.level 2011-05-15

PRMSL.mean.sea.level 2011-10-05




Figure 200: PRMSL.mean.sea.level 2011-10-05

PRES.surface 2012-09-18




Figure 201: PRES.surface 2012-09-18

PRES.surface 2012-09-19




Figure 202: PRES.surface 2012-09-19

PRES.surface 2012-09-20




Figure 203: PRES.surface 2012-09-20

PRES.surface 2011-05-16




Figure 204: PRES.surface 2011-05-16

PRES.surface 2011-05-15




Figure 205: PRES.surface 2011-05-15

PRES.surface 2011-10-05




Figure 206: PRES.surface 2011-10-05

DZDT.850.mb 2012-09-18




Figure 207: DZDT.850.mb 2012-09-18

DZDT.850.mb 2012-09-19




Figure 208: DZDT.850.mb 2012-09-19

DZDT.850.mb 2012-09-20




Figure 209: DZDT.850.mb 2012-09-20

DZDT.850.mb 2011-05-16




Figure 210: DZDT.850.mb 2011-05-16

DZDT.850.mb 2011-05-15




Figure 211: DZDT.850.mb 2011-05-15

DZDT.850.mb 2011-10-05




Figure 212: DZDT.850.mb 2011-10-05

DZDT.700.mb 2012-09-18




Figure 213: DZDT.700.mb 2012-09-18

DZDT.700.mb 2012-09-19




Figure 214: DZDT.700.mb 2012-09-19

DZDT.700.mb 2012-09-20




Figure 215: DZDT.700.mb 2012-09-20

DZDT.700.mb 2011-05-16




Figure 216: DZDT.700.mb 2011-05-16

DZDT.700.mb 2011-05-15




Figure 217: DZDT.700.mb 2011-05-15

DZDT.700.mb 2011-10-05




Figure 218: DZDT.700(mb) 2011-10-05




Figure 219: elevation 2012-09-18




Figure 220: elevation 2012-09-19




Figure 221: elevation 2012-09-20




Figure 222: elevation 2011-05-16




Figure 223: elevation 2011-05-15




Figure 224: elevation 2011-10-05

NLCD 2012-09-18




Figure 225: NLCD 2012-09-18

NLCD 2012-09-19




Figure 226: NLCD 2012-09-19

NLCD 2012-09-20




Figure 227: NLCD 2012-09-20

NLCD 2011-05-16




Figure 228: NLCD 2011-05-16

NLCD 2011-05-15




Figure 229: NLCD 2011-05-15

NLCD 2011-10-05




Figure 230: NLCD 2011-10-05




Figure 231: Date ML Inputs Plot against PM2.5

10.4 ML Inputs Plot against PM2.5 Images




Figure 232: Latitude ML Inputs Plot against PM2.5




Figure 233: Longitude ML Inputs Plot against PM2.5




Figure 234: A 100 ML Inputs Plot against PM2.5




Figure 235: C 100 ML Inputs Plot against PM2.5




Figure 236: Both 100 ML Inputs Plot against PM2.5




Figure 237: A 250 ML Inputs Plot against PM2.5




Figure 238: C 250 ML Inputs Plot against PM2.5




Figure 239: Both 250 ML Inputs Plot against PM2.5




Figure 240: A 500 ML Inputs Plot against PM2.5




Figure 241: C 500 ML Inputs Plot against PM2.5




Figure 242: Both 500 ML Inputs Plot against PM2.5




Figure 243: A 1000 ML Inputs Plot against PM2.5




Figure 244: C 1000 ML Inputs Plot against PM2.5




Figure 245: Both 1000 ML Inputs Plot against PM2.5




Figure 246: AOD ML Inputs Plot against PM2.5




Figure 247: MAIAC AOD ML Inputs Plot against PM2.5




Figure 248: HPBL.surface ML Inputs Plot against PM2.5




Figure 249: TMP.2.m.above.ground ML Inputs Plot against PM2.5




Figure 250: RH.2.m.above.ground ML Inputs Plot against PM2.5




Figure 251: DPT.2.m.above.ground ML Inputs Plot against PM2.5




Figure 252: APCP.surface ML Inputs Plot against PM2.5




Figure 253: WEASD.surface ML Inputs Plot against PM2.5




Figure 254: SNOWC.surface ML Inputs Plot against PM2.5




Figure 255: UGRD.10.m.above.ground ML Inputs Plot against PM2.5




Figure 256: VGRD.10.m.above.ground ML Inputs Plot against PM2.5




Figure 257: PRMSL.mean.sea.level ML Inputs Plot against PM2.5




Figure 258: PRES.surface ML Inputs Plot against PM2.5




Figure 259: DZDT.850(mb) ML Inputs Plot against PM2.5




Figure 260: DZDT.700(mb) ML Inputs Plot against PM2.5




Figure 261: elevation ML Inputs Plot against PM2.5




Figure 262: NLCD ML Inputs Plot against PM2.5




Figure 263: PM2.5-Obs Month 1

10.5 ML Inputs Map monthly medians Images




Figure 264: PM2.5-Obs Month 2




Figure 265: PM2.5-Obs Month 3




Figure 266: PM2.5-Obs Month 4




Figure 267: PM2.5-Obs Month 5




Figure 268: PM2.5-Obs Month 6




Figure 269: PM2.5-Obs Month 7




Figure 270: PM2.5-Obs Month 8




Figure 271: PM2.5-Obs Month 9




Figure 272: PM2.5-Obs Month 10




Figure 273: PM2.5-Obs Month 11




Figure 274: PM2.5-Obs Month 12

A_100 Month 1




Figure 275: A-100 Month 1

A_100 Month 2




Figure 276: A-100 Month 2

A_100 Month 3




Figure 277: A-100 Month 3

A_100 Month 4




Figure 278: A-100 Month 4

A_100 Month 5




Figure 279: A-100 Month 5

A_100 Month 6




Figure 280: A-100 Month 6

A_100 Month 7




Figure 281: A-100 Month 7

A_100 Month 8




Figure 282: A-100 Month 8

A_100 Month 9




Figure 283: A-100 Month 9

A_100 Month 10




Figure 284: A-100 Month 10

A_100 Month 11




Figure 285: A-100 Month 11

A_100 Month 12




Figure 286: A-100 Month 12

C_100 Month 1




Figure 287: C-100 Month 1

C_100 Month 2




Figure 288: C-100 Month 2

C_100 Month 3




Figure 289: C-100 Month 3

C_100 Month 4




Figure 290: C-100 Month 4

C_100 Month 5




Figure 291: C-100 Month 5

C_100 Month 6




Figure 292: C-100 Month 6

C_100 Month 7




Figure 293: C-100 Month 7

C_100 Month 8




Figure 294: C-100 Month 8

C_100 Month 9




Figure 295: C-100 Month 9

C_100 Month 10




Figure 296: C-100 Month 10

C_100 Month 11




Figure 297: C-100 Month 11

C_100 Month 12




Figure 298: C-100 Month 12

Both_100 Month 1




Figure 299: Both-100 Month 1

Both_100 Month 2




Figure 300: Both-100 Month 2

Both_100 Month 3




Figure 301: Both-100 Month 3

Both_100 Month 4




Figure 302: Both-100 Month 4

Both_100 Month 5




Figure 303: Both-100 Month 5

Both_100 Month 6




Figure 304: Both-100 Month 6

Both_100 Month 7




Figure 305: Both-100 Month 7

Both_100 Month 8




Figure 306: Both-100 Month 8

Both_100 Month 9




Figure 307: Both-100 Month 9

Both_100 Month 10




Figure 308: Both-100 Month 10

Both_100 Month 11




Figure 309: Both-100 Month 11

Both_100 Month 12




Figure 310: Both-100 Month 12

A_250 Month 1




Figure 311: A-250 Month 1

A_250 Month 2




Figure 312: A-250 Month 2

A_250 Month 3




Figure 313: A-250 Month 3

A_250 Month 4




Figure 314: A-250 Month 4

A_250 Month 5




Figure 315: A-250 Month 5

A_250 Month 6




Figure 316: A-250 Month 6

A_250 Month 7




Figure 317: A-250 Month 7

A_250 Month 8




Figure 318: A-250 Month 8

A_250 Month 9




Figure 319: A-250 Month 9

A_250 Month 10




Figure 320: A-250 Month 10

A_250 Month 11




Figure 321: A-250 Month 11

A_250 Month 12




Figure 322: A-250 Month 12

C_250 Month 1




Figure 323: C-250 Month 1

C_250 Month 2




Figure 324: C-250 Month 2

C_250 Month 3




Figure 325: C-250 Month 3

C_250 Month 4




Figure 326: C-250 Month 4

C_250 Month 5




Figure 327: C-250 Month 5

C_250 Month 6




Figure 328: C-250 Month 6

C_250 Month 7




Figure 329: C-250 Month 7

C_250 Month 8




Figure 330: C-250 Month 8

C_250 Month 9




Figure 331: C-250 Month 9

C_250 Month 10




Figure 332: C-250 Month 10

C_250 Month 11




Figure 333: C-250 Month 11

C_250 Month 12




Figure 334: C-250 Month 12

Both_250 Month 1




Figure 335: Both-250 Month 1

Both_250 Month 2




Figure 336: Both-250 Month 2

Both_250 Month 3




Figure 337: Both-250 Month 3

Both_250 Month 4




Figure 338: Both-250 Month 4

Both_250 Month 5




Figure 339: Both-250 Month 5

Both_250 Month 6




Figure 340: Both-250 Month 6

Both_250 Month 7




Figure 341: Both-250 Month 7

Both_250 Month 8




Figure 342: Both-250 Month 8

Both_250 Month 9




Figure 343: Both-250 Month 9

Both_250 Month 10




Figure 344: Both-250 Month 10

Both_250 Month 11




Figure 345: Both-250 Month 11

Both_250 Month 12




Figure 346: Both-250 Month 12

A_500 Month 1




Figure 347: A-500 Month 1

A_500 Month 2




Figure 348: A-500 Month 2

A_500 Month 3




Figure 349: A-500 Month 3

A_500 Month 4




Figure 350: A-500 Month 4

A_500 Month 5




Figure 351: A-500 Month 5

A_500 Month 6




Figure 352: A-500 Month 6

A_500 Month 7




Figure 353: A-500 Month 7

A_500 Month 8




Figure 354: A-500 Month 8

A_500 Month 9




Figure 355: A-500 Month 9

A_500 Month 10




Figure 356: A-500 Month 10

A_500 Month 11




Figure 357: A-500 Month 11

A_500 Month 12




Figure 358: A-500 Month 12

C_500 Month 1




Figure 359: C-500 Month 1

C_500 Month 2




Figure 360: C-500 Month 2

C_500 Month 3




Figure 361: C-500 Month 3

C_500 Month 4




Figure 362: C-500 Month 4

C_500 Month 5




Figure 363: C-500 Month 5

C_500 Month 6




Figure 364: C-500 Month 6

C_500 Month 7




Figure 365: C-500 Month 7

C_500 Month 8




Figure 366: C-500 Month 8

C_500 Month 9




Figure 367: C-500 Month 9

C_500 Month 10




Figure 368: C-500 Month 10

C_500 Month 11




Figure 369: C-500 Month 11

C_500 Month 12




Figure 370: C-500 Month 12

Both_500 Month 1




Figure 371: Both-500 Month 1

Both_500 Month 2




Figure 372: Both-500 Month 2

Both_500 Month 3




Figure 373: Both-500 Month 3

Both_500 Month 4




Figure 374: Both-500 Month 4

Both_500 Month 5




Figure 375: Both-500 Month 5

Both_500 Month 6




Figure 376: Both-500 Month 6

Both_500 Month 7




Figure 377: Both-500 Month 7

Both_500 Month 8




Figure 378: Both-500 Month 8

Both_500 Month 9




Figure 379: Both-500 Month 9

Both_500 Month 10




Figure 380: Both-500 Month 10

Both_500 Month 11




Figure 381: Both-500 Month 11

Both_500 Month 12




Figure 382: Both-500 Month 12

A_1000 Month 1




Figure 383: A-1000 Month 1

A_1000 Month 2




Figure 384: A-1000 Month 2

A_1000 Month 3




Figure 385: A-1000 Month 3

A_1000 Month 4




Figure 386: A-1000 Month 4

A_1000 Month 5




Figure 387: A-1000 Month 5

A_1000 Month 6




Figure 388: A-1000 Month 6

A_1000 Month 7




Figure 389: A-1000 Month 7

A_1000 Month 8




Figure 390: A-1000 Month 8

A_1000 Month 9




Figure 391: A-1000 Month 9

A_1000 Month 10




Figure 392: A-1000 Month 10

A_1000 Month 11




Figure 393: A-1000 Month 11

A_1000 Month 12




Figure 394: A-1000 Month 12

C_1000 Month 1




Figure 395: C-1000 Month 1

C_1000 Month 2




Figure 396: C-1000 Month 2

C_1000 Month 3




Figure 397: C-1000 Month 3

C_1000 Month 4




Figure 398: C-1000 Month 4

C_1000 Month 5




Figure 399: C-1000 Month 5

C_1000 Month 6




Figure 400: C-1000 Month 6

C_1000 Month 7




Figure 401: C-1000 Month 7

C_1000 Month 8




Figure 402: C-1000 Month 8

C_1000 Month 9




Figure 403: C-1000 Month 9

C_1000 Month 10




Figure 404: C-1000 Month 10

C_1000 Month 11




Figure 405: C-1000 Month 11

C_1000 Month 12




Figure 406: C-1000 Month 12

Both_1000 Month 1




Figure 407: Both-1000 Month 1

Both_1000 Month 2




Figure 408: Both-1000 Month 2

Both_1000 Month 3




Figure 409: Both-1000 Month 3

Both_1000 Month 4




Figure 410: Both-1000 Month 4

Both_1000 Month 5




Figure 411: Both-1000 Month 5

Both_1000 Month 6




Figure 412: Both-1000 Month 6

Both_1000 Month 7




Figure 413: Both-1000 Month 7

Both_1000 Month 8




Figure 414: Both-1000 Month 8

Both_1000 Month 9




Figure 415: Both-1000 Month 9

Both_1000 Month 10




Figure 416: Both-1000 Month 10

Both_1000 Month 11




Figure 417: Both-1000 Month 11

Both_1000 Month 12




Figure 418: Both-1000 Month 12




Figure 419: AOD Month 1

AOD Month 2




Figure 420: AOD Month 2

AOD Month 3




Figure 421: AOD Month 3




Figure 422: AOD Month 4




Figure 423: AOD Month 5

AOD Month 6




Figure 424: AOD Month 6

AOD Month 7




Figure 425: AOD Month 7

AOD Month 8




Figure 426: AOD Month 8




Figure 427: AOD Month 9

AOD Month 10




Figure 428: AOD Month 10

AOD Month 11




Figure 429: AOD Month 11

AOD Month 12




Figure 430: AOD Month 12




Figure 431: MAIAC-AOD Month 1




Figure 432: MAIAC-AOD Month 2




Figure 433: MAIAC-AOD Month 3




Figure 434: MAIAC-AOD Month 4




Figure 435: MAIAC-AOD Month 5




Figure 436: MAIAC-AOD Month 6




Figure 437: MAIAC-AOD Month 7




Figure 438: MAIAC-AOD Month 8




Figure 439: MAIAC-AOD Month 9




Figure 440: MAIAC-AOD Month 10




Figure 441: MAIAC-AOD Month 11




Figure 442: MAIAC-AOD Month 12




Figure 443: HPBL.surface Month 1




Figure 444: HPBL.surface Month 2




Figure 445: HPBL.surface Month 3




Figure 446: HPBL.surface Month 4




Figure 447: HPBL.surface Month 5




Figure 448: HPBL.surface Month 6

HPBL.surface Month 7




Figure 449: HPBL.surface Month 7




Figure 450: HPBL.surface Month 8




Figure 451: HPBL.surface Month 9




Figure 452: HPBL.surface Month 10

HPBL.surface Month 11




Figure 453: HPBL.surface Month 11

HPBL.surface Month 12




Figure 454: HPBL.surface Month 12




Figure 455: TMP.2.m.above.ground Month 1

TMP.2.m.above.ground Month 2




Figure 456: TMP.2.m.above.ground Month 2




Figure 457: TMP.2.m.above.ground Month 3




Figure 458: TMP.2.m.above.ground Month 4




Figure 459: TMP.2.m.above.ground Month 5




Figure 460: TMP.2.m.above.ground Month 6




Figure 461: TMP.2.m.above.ground Month 7




Figure 462: TMP.2.m.above.ground Month 8




Figure 463: TMP.2.m.above.ground Month 9




Figure 464: TMP.2.m.above.ground Month 10

TMP.2.m.above.ground Month 11




Figure 465: TMP.2.m.above.ground Month 11

TMP.2.m.above.ground Month 12




Figure 466: TMP.2.m.above.ground Month 12

RH.2.m.above.ground Month 1




Figure 467: RH.2.m.above.ground Month 1

RH.2.m.above.ground Month 2




Figure 468: RH.2.m.above.ground Month 2

RH.2.m.above.ground Month 3




Figure 469: RH.2.m.above.ground Month 3

RH.2.m.above.ground Month 4




Figure 470: RH.2.m.above.ground Month 4

RH.2.m.above.ground Month 5




Figure 471: RH.2.m.above.ground Month 5




Figure 472: RH.2.m.above.ground Month 6

RH.2.m.above.ground Month 7




Figure 473: RH.2.m.above.ground Month 7




Figure 474: RH.2.m.above.ground Month 8

RH.2.m.above.ground Month 9




Figure 475: RH.2.m.above.ground Month 9

RH.2.m.above.ground Month 10




Figure 476: RH.2.m.above.ground Month 10

RH.2.m.above.ground Month 11




Figure 477: RH.2.m.above.ground Month 11

RH.2.m.above.ground Month 12




Figure 478: RH.2.m.above.ground Month 12

DPT.2.m.above.ground Month 1




Figure 479: DPT.2.m.above.ground Month 1




Figure 480: DPT.2.m.above.ground Month 2

DPT.2.m.above.ground Month 3




Figure 481: DPT.2.m.above.ground Month 3

DPT.2.m.above.ground Month 4




Figure 482: DPT.2.m.above.ground Month 4




Figure 483: DPT.2.m.above.ground Month 5

DPT.2.m.above.ground Month 6




Figure 484: DPT.2.m.above.ground Month 6

DPT.2.m.above.ground Month 7




Figure 485: DPT.2.m.above.ground Month 7

DPT.2.m.above.ground Month 8




Figure 486: DPT.2.m.above.ground Month 8

DPT.2.m.above.ground Month 9




Figure 487: DPT.2.m.above.ground Month 9

DPT.2.m.above.ground Month 10




Figure 488: DPT.2.m.above.ground Month 10

DPT.2.m.above.ground Month 11




Figure 489: DPT.2.m.above.ground Month 11

DPT.2.m.above.ground Month 12




Figure 490: DPT.2.m.above.ground Month 12




Figure 491: APCP.surface Month 1

APCP.surface Month 2




Figure 492: APCP.surface Month 2

APCP.surface Month 3




Figure 493: APCP.surface Month 3




Figure 494: APCP.surface Month 4




Figure 495: APCP.surface Month 5

APCP.surface Month 6




Figure 496: APCP.surface Month 6

APCP.surface Month 7




Figure 497: APCP.surface Month 7

APCP.surface Month 8




Figure 498: APCP.surface Month 8

APCP.surface Month 9




Figure 499: APCP.surface Month 9




Figure 500: APCP.surface Month 10




Figure 501: APCP.surface Month 11

APCP.surface Month 12




Figure 502: APCP.surface Month 12




Figure 503: WEASD.surface Month 1




Figure 504: WEASD.surface Month 2




Figure 505: WEASD.surface Month 3




Figure 506: WEASD.surface Month 4




Figure 507: WEASD.surface Month 5




Figure 508: WEASD.surface Month 6




Figure 509: WEASD.surface Month 7




Figure 510: WEASD.surface Month 8




Figure 511: WEASD.surface Month 9




Figure 512: WEASD.surface Month 10




Figure 513: WEASD.surface Month 11




Figure 514: WEASD.surface Month 12




Figure 515: SNOWC.surface Month 1




Figure 516: SNOWC.surface Month 2




Figure 517: SNOWC.surface Month 3




Figure 518: SNOWC.surface Month 4




Figure 519: SNOWC.surface Month 5




Figure 520: SNOWC.surface Month 6




Figure 521: SNOWC.surface Month 7




Figure 522: SNOWC.surface Month 8




Figure 523: SNOWC.surface Month 9




Figure 524: SNOWC.surface Month 10




Figure 525: SNOWC.surface Month 11




Figure 526: SNOWC.surface Month 12

UGRD.10.m.above.ground Month 1




Figure 527: UGRD.10.m.above.ground Month 1

UGRD.10.m.above.ground Month 2




Figure 528: UGRD.10.m.above.ground Month 2

UGRD.10.m.above.ground Month 3




Figure 529: UGRD.10.m.above.ground Month 3

UGRD.10.m.above.ground Month 4




Figure 530: UGRD.10.m.above.ground Month 4

UGRD.10.m.above.ground Month 5




Figure 531: UGRD.10.m.above.ground Month 5

UGRD.10.m.above.ground Month 6




Figure 532: UGRD.10.m.above.ground Month 6

UGRD.10.m.above.ground Month 7




Figure 533: UGRD.10.m.above.ground Month 7

UGRD.10.m.above.ground Month 8




Figure 534: UGRD.10.m.above.ground Month 8

UGRD.10.m.above.ground Month 9




Figure 535: UGRD.10.m.above.ground Month 9

UGRD.10.m.above.ground Month 10




Figure 536: UGRD.10.m.above.ground Month 10

UGRD.10.m.above.ground Month 11




Figure 537: UGRD.10.m.above.ground Month 11

UGRD.10.m.above.ground Month 12




Figure 538: UGRD.10.m.above.ground Month 12

VGRD.10.m.above.ground Month 1




Figure 539: VGRD.10.m.above.ground Month 1

VGRD.10.m.above.ground Month 2




Figure 540: VGRD.10.m.above.ground Month 2

VGRD.10.m.above.ground Month 3




Figure 541: VGRD.10.m.above.ground Month 3

VGRD.10.m.above.ground Month 4




Figure 542: VGRD.10.m.above.ground Month 4

VGRD.10.m.above.ground Month 5




Figure 543: VGRD.10.m.above.ground Month 5




Figure 544: VGRD.10.m.above.ground Month 6

VGRD.10.m.above.ground Month 7




Figure 545: VGRD.10.m.above.ground Month 7




Figure 546: VGRD.10.m.above.ground Month 8

VGRD.10.m.above.ground Month 9




Figure 547: VGRD.10.m.above.ground Month 9

VGRD.10.m.above.ground Month 10




Figure 548: VGRD.10.m.above.ground Month 10

VGRD.10.m.above.ground Month 11




Figure 549: VGRD.10.m.above.ground Month 11

VGRD.10.m.above.ground Month 12




Figure 550: VGRD.10.m.above.ground Month 12




Figure 551: PRMSL.mean.sea.level Month 1

PRMSL.mean.sea.level Month 2




Figure 552: PRMSL.mean.sea.level Month 2




Figure 553: PRMSL.mean.sea.level Month 3




Figure 554: PRMSL.mean.sea.level Month 4




Figure 555: PRMSL.mean.sea.level Month 5




Figure 556: PRMSL.mean.sea.level Month 6




Figure 557: PRMSL.mean.sea.level Month 7




Figure 558: PRMSL.mean.sea.level Month 8




Figure 559: PRMSL.mean.sea.level Month 9




Figure 560: PRMSL.mean.sea.level Month 10




Figure 561: PRMSL.mean.sea.level Month 11




Figure 562: PRMSL.mean.sea.level Month 12




Figure 563: PRES.surface Month 1




Figure 564: PRES.surface Month 2




Figure 565: PRES.surface Month 3




Figure 566: PRES.surface Month 4




Figure 567: PRES.surface Month 5




Figure 568: PRES.surface Month 6




Figure 569: PRES.surface Month 7




Figure 570: PRES.surface Month 8




Figure 571: PRES.surface Month 9




Figure 572: PRES.surface Month 10




Figure 573: PRES.surface Month 11




Figure 574: PRES.surface Month 12

DZDT.850.mb Month 1




Figure 575: DZDT.850.mb Month 1

DZDT.850(mb) Month 2




Figure 576: DZDT.850(mb) Month 2

DZDT.850.mb Month 3




Figure 577: DZDT.850.mb Month 3

DZDT.850.mb Month 4




Figure 578: DZDT.850.mb Month 4

DZDT.850(mb) Month 5




Figure 579: DZDT.850(mb) Month 5

DZDT.850.mb Month 6




Figure 580: DZDT.850.mb Month 6

DZDT.850(mb) Month 7




Figure 581: DZDT.850(mb) Month 7

DZDT.850.mb Month 8




Figure 582: DZDT.850.mb Month 8

DZDT.850(mb) Month 9




Figure 583: DZDT.850(mb) Month 9

DZDT.850(mb Month 10)




Figure 584: DZDT.850(mb Month 10)

DZDT.850(mb) Month 11




Figure 585: DZDT.850(mb) Month 11

DZDT.850(mb) Month 12




Figure 586: DZDT.850(mb) Month 12

DZDT.700(mb) Month 1




Figure 587: DZDT.700(mb) Month 1

DZDT.700(mb) Month 2




Figure 588: DZDT.700(mb) Month 2

DZDT.700(mb) Month 3




Figure 589: DZDT.700(mb) Month 3

DZDT.700(mb) Month 4




Figure 590: DZDT.700(mb) Month 4

DZDT.700(mb) Month 5




Figure 591: DZDT.700(mb) Month 5

DZDT.700(mb) Month 6




Figure 592: DZDT.700(mb) Month 6

DZDT.700(mb) Month 7




Figure 593: DZDT.700(mb) Month 7

DZDT.700(mb) Month 8




Figure 594: DZDT.700(mb) Month 8




Figure 595: DZDT.700(mb) Month 9




Figure 596: DZDT.700.mb Month 10

DZDT.700(mb) Month 11




Figure 597: DZDT.700(mb) Month 11

DZDT.700(mb) Month 12




Figure 598: DZDT.700(mb) Month 12




Figure 599: elevation Month 1




Figure 600: elevation Month 2




Figure 601: elevation Month 3




Figure 602: elevation Month 4




Figure 603: elevation Month 5




Figure 604: elevation Month 6




Figure 605: elevation Month 7




Figure 606: elevation Month 8




Figure 607: elevation Month 9




Figure 608: elevation Month 10




Figure 609: elevation Month 11




Figure 610: elevation Month 12

NLCD Month 1




Figure 611: NLCD Month 1

NLCD Month 2




Figure 612: NLCD Month 2

NLCD Month 3




Figure 613: NLCD Month 3

NLCD Month 4




Figure 614: NLCD Month 4

NLCD Month 5




Figure 615: NLCD Month 5

NLCD Month 6




Figure 616: NLCD Month 6

NLCD Month 7




Figure 617: NLCD Month 7

NLCD Month 8




Figure 618: NLCD Month 8




Figure 619: NLCD Month 9

NLCD Month 10




Figure 620: NLCD Month 10

NLCD Month 11




Figure 621: NLCD Month 11

NLCD Month 12




Figure 622: NLCD Month 12

References

- Abatzoglou, J. T. and Williams, A. P. (2016). Impact of anthropogenic climate change on wildfire across western US forests. *Proceedings of the National Academy of Sciences*, 113(42):11770–11775.
- Anyenda, E. O., Higashi, T., Kambayashi, Y., Thao, N. T. T., Michigami, Y., Fujimura, M., Hara, J., Tsujiguchi, H., Kitaoka, M., Asakura, H., Hori, D., Yamada, Y., Hayashi, K., Hayakawa, K., and Nakamura, H. (2016). Exposure to daily ambient particulate polycyclic aromatic hydrocarbons and cough occurrence in adult chronic cough patients: A longitudinal study. *Atmospheric Environment*, 140(Supplement C):34 – 41.
- Brokamp, C., Jandarov, R., Rao, M. B., LeMasters, G., and Ryan, P. (2017). Exposure assessment models for elemental components of particulate matter in an urban environment: A comparison of regression and random forest approaches. *Atmospheric Environment*, 151:1–11.
- French, N. H. F., McKenzie, D., Erickson, T., Koziol, B., Billmire, M., Endsley, K. A., Scheinerman, N. K. Y., Jenkins, L., Miller, M. E., Ottmar, R., and Prichard, S. (2014). Modeling Regional-Scale Wildland Fire Emissions with the Wildland Fire Emissions Information System. *Earth Interactions*, 18(16):1–26.
- Giglio, L., Csiszar, I., and Justice, C. O. (2006). Global distribution and seasonality of active fires as observed with the Terra and Aqua Moderate Resolution Imaging Spectroradiometer (MODIS) sensors. *Journal of Geophysical Research: Biogeosciences*, 111(G2). G02016; <https://modis.gsfc.nasa.gov/data/dataproducts/mod14.php>.
- Hall, D. K. and Riggs, G. A. (2016). MODIS/Aqua Snow Cover Daily L3 Global 500m Grid, Version 6. NASA National Snow and Ice Data Center Distributed Active Archive Center. <http://dx.doi.org/10.5067/MODIS/MYD10A1.006>.
- Hawbaker, T. J., Vanderhoof, M. K., Beal, Y.-J., Takacs, J. D., Schmidt, G. L., Falgout, J. T., Williams, B., Fairaux, N. M., Caldwell, M. K., Picotte, J. J., Howard, S. M., Stitt, S., and Dwyer, J. L. (2017). Mapping burned areas using dense time-series of Landsat data. *Remote Sensing of Environment*, 198(Supplement C):504 – 522.
- Homer, C., Dewitz, J., Yang, L., Jin, S., Danielson, P., Xian, G., Coulston, J., Herold, N., Wickham, J., and Megown, K. (2017). Completion of the 2011 National Land Cover Database for the Conterminous United States – Representing a Decade of Land Cover Change Information. *Photogrammetric Engineering & Remote Sensing*, 81(5):345 – 354. <https://www.mrlc.gov/nlcd2011.php>.
- J., F. E., T., A. J., K., B. J., T., F. J., and A., B. B. (2016). Quantifying the human influence on fire ignition across the western usa. *Ecological Applications*, 26(8):2390–2401.
- Kollanus, V., Tiittanen, P., Niemi, J. V., and Lanki, T. (2016). Effects of long-range transported air pollution from vegetation fires on daily mortality and hospital admissions in the Helsinki metropolitan area, Finland. *Environ Res*, 151:351–358.

- Kondragunta, S. and Seybold, M. (2009). Revisions to GOES Aerosol and Smoke Product (GASP) Algorithm. <http://www.ssd.noaa.gov/PS/FIRE/GASP/gasp.html>.
- Larsen, A. E., Reich, B. J., Ruminski, M., and Rappold, A. G. (2017). Impacts of fire smoke plumes on regional air quality, 2006-2013. *Journal of Exposure Science & Environmental Epidemiology*.
- Liu, J. C., Wilson, A., Mickley, L. J., Dominici, F., Ebisu, K., Wang, Y., Sulprizio, M. P., Peng, R. D., Yue, X., Anderson, G. B., and Bell, M. L. (2016). Wildfire-specific Fine Particulate Matter and Risk of Hospital Admissions in Urban and Rural Counties. *Epidemiology*, 28:77–85.
- Liu, Y., Sarnat, J. A., Kilaru, V., Jacob, D. J., and Koutrakis, P. (2005). Estimating ground-level PM_{2.5} in the eastern United States using satellite remote sensing. *Environ Sci Technol*, 39(9):3269–78.
- LP DAAC (2017, accessed November 12, 2017). MCD64A1: MODIS/Terra and Aqua Burned Area Monthly L3 Global 500 m SIN Grid V006. https://lpdaac.usgs.gov/dataset_discovery/modis/modis_products_table/mcd64a1_v006.
- MTRI (2017 accessed November 7, 2017). *Wildland Fire Emissions Information System*. <http://wfeis.mtri.org/>.
- NASA LAADS DAAC (2017, accessed November 2, 2017a). MOD04_L2 - MODIS/Terra Aerosol 5-Min L2 Swath 10km. https://ladsweb.modaps.eosdis.nasa.gov/api/v1/productPage/product=MOD04_L2.
- NASA LAADS DAAC (2017, accessed November 2, 2017b). MYD04_L2 - MODIS/Aqua Aerosol 5-Min L2 Swath 10km. https://ladsweb.modaps.eosdis.nasa.gov/api/v1/productPage/product=MYD04_L2.
- NOAA NCEI (2017, accessed November 2, 2017). *Satellite Data Access by Datasets*. <https://www.ncdc.noaa.gov/data-access/satellite-data/satellite-data-access-datasets>.
- NOAA OSPO (2017, accessed November 3, 2017). *Hazard Mapping System Fire and Smoke Product*. <http://www.ospo.noaa.gov/Products/land/hms.html>.
- Ottmar, R. D., Sandberg, D. V., Riccardi, C. L., and Prichard, S. J. (2007). An overview of the Fuel Characteristic Classification System — Quantifying, classifying, and creating fuelbeds for resource planning. *Canadian Journal of Forest Research*, 37(12):2383–2393.
- Prichard, S. J., Ottmar, R. D., and Anderson, G. A. (2009). Consume 3.0 user's guide. *USDA Forest Service Pacific Wildland Fire Sciences Laboratory Rep.*, page 239.
- Reid, C. E., Jerrett, M., Petersen, M. L., Pfister, G. G., Morefield, P. E., Tager, I. B., Raffuse, S. M., and Balmes, J. R. (2015). Spatiotemporal prediction of fine particulate matter during the 2008 northern California wildfires using machine learning. *Environ Sci Technol*, 49(6):3887–96.
- Rolph, G., Stein, A., and Stunder, B. (2017). Real-time Environmental Applications and Display sYstem: READY. *Environmental Modelling & Software*, 95(Supplement C):210 – 228.

- Salimi, F., Henderson, S. B., Morgan, G. G., Jalaludin, B., and Johnston, F. H. (2016). Ambient particulate matter, landscape fire smoke, and emergency ambulance dispatches in Sydney, Australia. *Environ Int.*
- Sampson, P. D., Richards, M., Szpiro, A. A., Bergen, S., Sheppard, L., Larson, T. V., and Kaufman, J. D. (2013). A regionalized national universal kriging model using partial least squares regression for estimating annual pm2.5 concentrations in epidemiology. *Atmospheric Environment*, 75:383 – 392.
- Sayer, A. M., Hsu, N. C., Bettenhausen, C., and Jeong, M.-J. (2013). Validation and uncertainty estimates for MODIS Collection 6 “Deep Blue” aerosol data. *Journal of Geophysical Research: Atmospheres*, 118(14):7864–7872.
- Schroeder, W., Oliva, P., Giglio, L., and Csiszar, I. A. (2014). The New VIIRS 375m active fire detection data product: Algorithm description and initial assessment. *Remote Sensing of Environment*, 143(Supplement C):85 – 96.
- Torvela, T., Tissari, J., Sippula, O., Kaivosoja, T., Leskinen, J., Virén, A., Lähde, A., and Jokiniemi, J. (2014). Effect of wood combustion conditions on the morphology of freshly emitted fine particles. *Atmospheric Environment*, 87(Supplement C):65 – 76.
- US EPA (2017, accessed November 2, 2017a). *AQS Memos - Technical Note on Reporting PM2.5 Continuous Monitoring and Speciation Data to the Air Quality System (AQS)*. <https://www.epa.gov/aqs/aqs-memos-technical-note-reporting-pm25-continuous-monitoring- and-speciation-data-air-quality>.
- US EPA (2017, accessed November 2, 2017c). *Outdoor Air Quality Data Download Daily Data*. <https://www.epa.gov/outdoor-air-quality-data/download-daily-data>.
- US EPA (2017, accessed November 2, 2017d). *Parameters*. <https://aqs.epa.gov/aqsweb/documents/codetables/parameters.html>.
- US EPA (2017, accessed November 2, 2017e). *PM 2.5 - Visibility (IMPROVE)*. <https://www3.epa.gov/ttnamti1/visdata.html>.
- US EPA (2017, accessed November 2, 2017f). *Sampling Methods for All Parameters*. https://aqs.epa.gov/aqsweb/documents/codetables/methods_all.html.
- US EPA (2017, accessed October 23, 2017b). *National Emissions Inventory (NEI)*. <https://www.epa.gov/air-emissions-inventories/national-emissions-inventory-nei>.
- US National Weather Service (2016, accessed November 2, 2017a). *National Weather Service Instruction 10-1605*. <https://www.ncdc.noaa.gov/stormevents/pd01016005curr.pdf>.
- US National Weather Service (2017, accessed November 2, 2017b). *Storm Events Database*. <https://www.ncdc.noaa.gov/stormevents/>.
- US National Weather Service (2017, accessed November 2, 2017c). *Storm Events Database: Database Details*. <https://www.ncdc.noaa.gov/stormevents/details.jsp>.

USGS (2017, accessed November 6, 2017). *About 3DEP Products and Services*. https://nationalmap.gov/3DEP/3dep_prodserv.html.

Wang, W., Barker, D., Bray, J., Bruyere, C., Duda, M., Dudhia, J., Gill, D., and Michalakes, J. (2007). User's Guide for Advanced Research WRF (ARW) Modeling System Version 3. *Mesoscale and Microscale Meteorology Division–National Center for Atmospheric Research (MMM-NCAR)*.

Westerling, A. L. (2016a). Correction to ‘increasing western us forest wildfire activity: sensitivity to changes in the timing of spring’. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, 371(1707).

Westerling, A. L. (2016b). Increasing western US forest wildfire activity: sensitivity to changes in the timing of spring. *Philos Trans R Soc Lond B Biol Sci*, 371(1696). bibtex: westerling_increasing_2016.

Whiteman, C. D., Hoch, S. W., Horel, J. D., and Charland, A. (2014). Relationship between particulate air pollution and meteorological variables in Utah’s Salt Lake Valley. *Atmospheric Environment*, 94(Supplement C):742 – 753.