SpectralBridge: An Auditable Software Workflow for Reflectance Comparisons Across NEON, Drones, and Landsat

Ty Tuff[1,*], Aashish Mukund[1], Nayani T. Ilangakoon[2], Erick Verleye[2]
Megan E. Cattau[4], Janushi Shastri[2,3], Pavan Appanna[2,3], Tyler L. McIntosh[2]
Matthew E. Bitters[2], Carol A. Wessman[5], Jennifer K. Balch[1,6], and Cibele Amaral[1]

Revised 12 October 2026; source snapshots identified in text

[1]Earth Lab and the Environmental Data Science Innovation & Impact Lab (ESIIL), Cooperative Institute for Research in Environmental Sciences (CIRES), University of Colorado Boulder, Boulder, Colorado, USA
[2]Earth Lab, Cooperative Institute for Research in Environmental Sciences (CIRES), University of Colorado Boulder, Boulder, Colorado, USA
[3]Department of Computer Science, University of Colorado Boulder, Boulder, Colorado, USA
[4]Human-Environment Systems, School of the Environment, Boise State University, Boise, Idaho, USA
[5]Department of Ecology and Evolutionary Biology, University of Colorado Boulder, Boulder, Colorado, USA
[6]Department of Geography, University of Colorado Boulder, Boulder, Colorado, USA

[*]Corresponding author: ty.tuff@colorado.edu

ORCID iDs: Ty Tuff, 0000-0001-5249-5197; Nayani T. Ilangakoon, 0000-0002-6607-0605; Erick Verleye, 0009-0009-8854-0127; Megan E. Cattau, 0000-0003-2164-3809; Tyler L. McIntosh, 0000-0002-6383-1755; Carol A. Wessman, 0000-0002-7325-0153; Jennifer K. Balch, 0000-0002-3983-7970; Cibele Amaral, 0000-0001-7597-2427

1 Summary

Reflectance describes how a surface reflects light at different wavelengths. SpectralBridge is Python research software that helps teams prepare and check reflectance products from NEON airborne sensors, drones, and Landsat. It retains named intermediate rasters, extracted values, quality checks, and provenance so users can inspect how each product was made. Separate routes preserve sensor-specific operations, while collection tools summarize selected flightlines and sites. The package organizes established corrections rather than proposing a new algorithm. Tests and team research examples demonstrate software contracts and limitations, not agreement among sensors.

2 Statement of need

Band names invite direct comparison, but spectral response, footprint, viewing direction, and illumination help define the reflectance recorded by each instrument. A drone may isolate sunlit foliage; NEON records a dense spectrum; Landsat mixes canopy, gaps, soil, and shadow across broader pixels (Kampe et al., 2010; National Ecological Observatory Network, 2026; Woodcock

and Strahler, 1987). Matching labels therefore do not make measurements equivalent, and these differences can change an ecological interpretation (Nicodemus et al., 1977; Schaepman-Strub et al., 2006; Teillet et al., 1997; Roy et al., 2016; Huang et al., 2002).

The intended users are remote-sensing researchers and developers who need to construct sensor-oriented products, inspect intermediate states, verify outputs, or summarize flightlines. They remain responsible for temporal matching, co-registration, and common spatial support. Independent validation also requires characterized reference measurements and uncertainty analysis; process completion alone cannot establish it (Origo et al., 2020).

SpectralBridge preserves the sequence from source through configured transformations to extracted values, helping users locate where an unexpected pattern first appears. It does not diagnose discrepancies or choose a valid spatial/temporal design. Figure 1 illustrates the separate roles of spatial support and spectral response.




Figure 1: Two independent sources of measurement mismatch. (A) The same illustrated vegetation, soil, and shadow are integrated over different pixel footprints; grids are schematic and not to scale. (B) Read from top to bottom: the upper curve is the input reflectance spectrum; the lower curves show how strongly each target band weights each wavelength. They are response functions, not output spectra. Applying a response curve to the input produces one weighted reflectance value, shown schematically by the matching colored output label and defined in Equation 1. The Gaussian profiles use packaged Landsat 8 OLI and MicaSense centers and full widths at half maximum (FWHM), not official measured response curves (Earth Lab, 2026k); the spectrum is illustrative.

3 State of the field

HyTools is SpectralBridge's closest methodological relative and part of its NEON lineage. HLS and Sen2Like harmonize satellite products; FORCE processes satellite archives; Snakemake and Nextflow orchestrate general pipelines (Chlus et al., 2025; Claverie et al., 2018; Ju et al., 2025;

Saunier et al., 2022; Frantz, 2019; Köster and Rahmann, 2012; Di Tommaso et al., 2017). In the systems compared here, none combines the separate NEON and drone routes with collection-level artifact and QA contracts (Earth Lab, 2026x; Earth Lab, 2026j). SpectralBridge's contribution is this sensor-aware integration, restart checks, and grouped summaries—not a new correction method or harmonization product.

4 Software design

The workflow addresses geometry normalization and target-band weighting, while preserving geospatial products for downstream spatial reconciliation. It does not create a common spatial support; nominally matching bands may also require bandpass adjustment (Chander et al., 2013; Roy et al., 2016; Earth Lab, 2026z; Earth Lab, 2026m).

NEON provides the dense spectrum from which configured MicaSense- and Landsat-like products are constructed. Independently acquired drone data follow a separate preparation path (National Ecological Observatory Network, 2026; Earth Lab, 2026z; Earth Lab, 2026h). Thus, Drone → NEON → Landsat describes a comparison design, not a physical conversion chain. A named target product reflects configured weighting and any recorded adjustment, while retaining NEON's original spatial support.

Geometry and terrain

For NEON, the workflow estimates bandwise SCS+C topographic terms, then applies RossThick and LiDenseR BRDF terms using available solar and view geometry (Queally et al., 2022; Earth Lab, 2026r; Earth Lab, 2026ab). These are established methods, with code lineage to HyTools acknowledged in the implementation (Ross, 1981; Roujean et al., 1992; Li and Strahler, 1992; Soenen et al., 2005; Wanner et al., 1995; Schaaf et al., 2002; Chlus et al., 2025). Invalid geometry ratios receive neutral factors, which QA should report. Both corrections share one output cube, so their separate effects cannot be isolated from that artifact.

Spectral response

Different instruments weight different parts of the spectrum, so matching band names do not guarantee matching measurements (Chander et al., 2013; Roy et al., 2016). Given source reflectance samples ρ_i and target-band weights $s_{b,i}$ sampled on the same source-band grid, SpectralBridge computes

$$\hat{\rho}_b = \frac{\sum_i \rho_i s_{b,i}}{\sum_i s_{b,i}}. \tag{1}$$

The implementation normalizes supplied response weights by their discrete sum; it does not perform wavelength-interval quadrature. Weights must therefore match the source grid and intended spectral support. Actual Landsat weighting requires measured relative spectral responses, not band centers and widths alone (Barsi et al., 2014). A deterministic fixture checks this arithmetic to an absolute tolerance of 2×10^{-7} (Earth Lab, 2026s; Earth Lab, 2026ae).

Source atmospheric correction is inherited from NEON; some configured Landsat-like products also receive a separate empirical brightness adjustment (National Ecological Observatory Network,

2026; Earth Lab, 2026l). These configured representations are not proof of physical agreement with another sensor (Claverie, 2023; Ju et al., 2025).

Spatial support

Matching band spaces does not make a crown-scale spectrum and a Landsat pixel the same spatial mixture (Teillet et al., 1997; Woodcock and Strahler, 1987; Huang et al., 2002). SpectralBridge preserves georeferenced rasters and polygon extractions but leaves grid alignment, overlap, and aggregation to the user, because these choices determine which observations contribute (Dungan et al., 2002; Earth Lab, 2026m; Earth Lab, 2026h).

Architecture and data flow

Figure 2 aligns the two observation routes and the shared collection-analysis path. NEON and drone processing share most stages; their main differences are at the edges: drone processing adds local-image staging and may use an affine mapping, while NEON's dense spectrum supports target-band convolution.

Each route writes named artifacts for distinct measurement states, linking the source to corrected and target-band products (Earth Lab, 2026e; Earth Lab, 2026m).




Figure 2: The NEON and drone routes share most stages and retain similar rasters, tables, and QA/provenance. Drone processing adds local input staging and may apply a configured affine mapping; NEON processing convolves its dense spectrum with target-band responses. The shared collection path streams selected, aligned products into grouped fits and diagnostics. The diagram shows intended workflow structure, not one campaign's results (Earth Lab, 2026j; Earth Lab, 2026m; Earth Lab, 2026h; Earth Lab, 2026x; Earth Lab, 2026a).

The architecture evolved from a HyTools-based workflow; HyTools is no longer a runtime dependency, but lineage remains in NEON access, ENVI writing, and correction components (Chlus et al., 2025; Earth Lab, 2025a; Earth Lab, 2025b; Earth Lab, 2026i; Earth Lab, 2026t; Earth Lab, 2026r).

The NEON route applies configured geometry corrections and spectral weighting to its source cube. The drone route retains native bands and may apply reviewed affine coefficients, without runtime coefficient fitting or spectral convolution (Earth Lab, 2026e; Earth Lab, 2026m; Earth Lab, 2026h; Earth Lab, 2026c). Collection analysis streams aligned ENVI windows into pooled, flightline-, and site-level fits. It accumulates compact statistics but still reads source rasters; no performance advantage is claimed without a matched benchmark (Earth Lab, 2026a; Earth Lab, 2026aa; Earth Lab, 2026b).

Named files serve as stage interfaces, not disposable cache. This design lets users inspect intermediate states, check stage-specific QA, and reuse valid products after interruption; it costs storage and I/O, and does not serialize every realized parameter. Restart therefore supports recovery, not

guaranteed exact replay. Completion is request-specific: every configured output must exist and be readable (Earth Lab, 2026j; Earth Lab, 2026d; Earth Lab, 2026ad; Earth Lab, 2026u).




Figure 3: Four questions govern a multi-stage scientific workflow. Normal process termination does not establish production of the requested scientific state; artifact existence does not establish QA; and QA does not establish agreement with independent observations. Remote-sensing validation frameworks likewise treat validation as staged evidence rather than a binary declaration (Morisette et al., 2006; Origo et al., 2020). The examples in the Research impact statement illustrate these separate checks (SpectralBridge project team, 2026; Earth Lab, 2026p).

The package is OS-independent, requires Python 3.10 or newer, and is distributed under the GNU General Public License version 3 (GPLv3); installation and usage are documented in its README (Earth Lab, 2026v; Earth Lab, 2026x). The evidence spans version 2.2.0 at `f7e7ddd` for the historical NEON route and `c86da6a` for later bulk work and 2.3.0rc1 preparation, but supplied run records do not link every result to an exact revision (Earth Lab, 2026y; Earth Lab, 2026g). The public release page listed `v1.0.0` at commit `9d1de28` as latest on 12 October 2026; the cited Zenodo DOI is a pre-rename archive, not a record for these results (Earth Lab, 2026w; Earth Lab, 2026f). The submitted source, evidence, and permanent archive citation must identify a consistent version.

Reuse potential

Reuse is most direct for NEON products and drone observations that match the supported input schemas. Users can configure target-band responses or reviewed affine registries, while preserving exact wavelength definitions and coefficient provenance; adding spatial aggregation or a new sensor remains a study-specific extension. The ENVI, Parquet, and JSON products can then be inspected or consumed by downstream analysis tools (Earth Lab, 2026m; Earth Lab, 2026h; Earth Lab, 2026x).

5 Research impact statement

The examples below come from the authors' ongoing research workflow for comparing NEON, drone, and Landsat reflectance; they are developer-team applications, not evidence of external adoption. Deterministic fixtures cover convolution, correction, extraction, QA, restart, and collection-analysis contracts (Kanewala and Bieman, 2014; Earth Lab, 2026ac; Earth Lab, 2026n). In the R10C

NEON example, the declared route produced 18 readable Parquet outputs, while QA retained warnings, including 68 poor-quality wavelength bands that were reported but not masked. This demonstrates artifact completion and warning preservation, not correction efficacy or agreement with Landsat (Earth Lab, 2026p; Earth Lab, 2026q).

A separate 43-flight drone attempt exposes the value of explicit completion checks: orchestration labeled all flights processed, but required ENVI and Parquet outputs were absent (SpectralBridge project team, 2026). A supplied bulk analysis summarizes 122 flightlines, but its executed software revision and run manifest were not provided (*SpectralBridge bulk evidence bundle* 2026; Earth Lab, 2026o). Together, these cases show how the workflow can expose incomplete production, preserve warnings, and summarize collections while making the limits of the evidence visible. They do not establish independent cross-sensor accuracy, a performance advantage, improved ecological inference, or adoption beyond the developer team.

6 AI usage disclosure

OpenAI Codex assisted with source and documentation review, proposing software edits, investigating the drone defect, manuscript editing, LaTeX and citation checks, exported-table calculations, and figure preparation; a partial prompt log records some work (Earth Lab, 2026g). Human team members reviewed proposed software changes, citations, and calculations. AI output was not treated as scientific evidence; the authors retain responsibility for the software, analyses, and manuscript. [Authors: confirm the model/version and identify any other AI tools and versions used.]

7 Acknowledgements

Funding and other acknowledgements were not specified in the materials reviewed for this draft. The authors should identify financial support and any funder role, or state that no financial support was received, and complete any other acknowledgements before submission.

References

Barsi, Julia A., Kenton Lee, Geir Kvaran, Brian L. Markham, and Jeffrey A. Pedelty (2014). "The Spectral Response of the Landsat-8 Operational Land Imager". In: *Remote Sensing* 6.10, pp. 10232–10251. DOI: 10.3390/rs61010232.

Chander, Gyanesh, Nischal Mishra, Dennis L. Helder, David B. Aaron, Amit Angal, Taeyoung Choi, Xiaoxiong Xiong, and David R. Doelling (2013). "Applications of spectral band adjustment factors (SBAF) for cross-calibration". In: *IEEE Transactions on Geoscience and Remote Sensing* 51.3, pp. 1267–1281. DOI: 10.1109/TGRS.2012.2228007.

Chlus, Adam, Zhiwei Ye, Ting Zheng, Natalie Queally, Evan Greenberg, and Philip A. Townsend (2025). *HyTools: Hyperspectral image processing library*. Version 1.6.1. Software release. DOI: 10.5281/zenodo.16900026.

Claverie, Martin (2023). "Evaluation of surface reflectance bandpass adjustment techniques". In: *ISPRS Journal of Photogrammetry and Remote Sensing* 198, pp. 210–222. DOI: 10.1016/j.isprsjprs.2023.03.011.

Claverie, Martin, Junchang Ju, Jeffrey G. Masek, Jennifer L. Dungan, Eric F. Vermote, Jean-Claude Roger, Sergii V. Skakun, and Christopher Justice (2018). "The Harmonized Landsat and Sentinel-2 surface reflectance data set". In: *Remote Sensing of Environment* 219, pp. 145–161. DOI: 10.1016/j.rse.2018.09.002.

Di Tommaso, Paolo, Maria Chatzou, Evan W. Floden, Pablo Prieto Barja, Emilio Palumbo, and Cedric Notredame (2017). "Nextflow enables reproducible computational workflows". In: *Nature Biotechnology* 35.4, pp. 316–319. DOI: 10.1038/nbt.3820.

Dungan, Jennifer L., James N. Perry, Mark R. T. Dale, Pierre Legendre, Sheila Citron-Pousty, Marie-Josée Fortin, Anna Jakomulska, Maria Miriti, and Michael S. Rosenberg (2002). "A balanced view of scale in spatial statistical analysis". In: *Ecography* 25.5, pp. 626–640. DOI: 10.1034/j.1600-0587.2002.250510.x.

Earth Lab (2025a). *SpectralBridge history: HyTools compatibility and dependency work, 8–23 October 2025*. https://github.com/earthlab/spectralbridge/commits/main/?since=2025-10-08&until=2025-10-23. Includes commits 89178b6, 92c2bc7, 447d237, e3d3a41, 53e8206, 8deb979, a6656ea, and 4642eba; inspected 4 September 2026.

— (2025b). *SpectralBridge history: transition to internal NEON processing components, 27–28 October 2025*. https://github.com/earthlab/spectralbridge/commits/main/?since=2025-10-27&until=2025-10-28. Includes commits fc370c4, b5969f5, fa0235a, e73b739, 5677078, 6942cdb, e88473d, 9aa0159, 328922b, a241cb1, 38c42dc, and 1c6a55f; inspected 4 September 2026.

— (2026a). *Build a bulk cross-run analysis*. Source snapshot inspected 9 September 2026. URL: https://github.com/earthlab/spectralbridge/blob/c86da6a642a8a0bef37108df31af1fed1a882bd2/docs/vignettes/bulk-analysis.md.

— (2026b). *Bulk pipeline implementation: September snapshot*. Source snapshot inspected 9 September 2026. URL: https://github.com/earthlab/spectralbridge/blob/c86da6a642a8a0bef37108df31af1fed1a882bd2/src/spectralbridge/pipelines/bulk.py.

— (2026c). *Bulk population-analysis pipeline source*. https://github.com/earthlab/spectralbridge/blob/f7e7ddd2e01ee5f7cd74688eb479fd12fbc0b705/src/spectralbridge/pipelines/bulk.py. Accessed 4 September 2026.

— (2026d). *Carry On My Wayward Son: resume a run*. https://github.com/earthlab/spectralbridge/blob/f7e7ddd2e01ee5f7cd74688eb479fd12fbc0b705/docs/vignettes/carry-on-wayward-son.md.

— (2026e). *Core NEON pipeline source*. https://github.com/earthlab/spectralbridge/blob/f7e7ddd2e01ee5f7cd74688eb479fd12fbc0b705/src/spectralbridge/pipelines/pipeline.py. Accessed 4 September 2026.

— (2026f). *DOI and Zenodo status*. https://github.com/earthlab/spectralbridge/blob/f7e7ddd2e01ee5f7cd74688eb479fd12fbc0b705/docs/dev/doi-zenodo.md. Reviewed 3 June 2026.

— (2026g). *Finalize SpectralBridge 2.3.0rc1 preparation*. Commit inspected 9 September 2026; preparation is distinct from publication. URL: https://github.com/earthlab/spectralbridge/commit/c86da6a642a8a0bef37108df31af1fed1a882bd2.

— (2026h). *Module vignette 6: process drone imagery*. https://github.com/earthlab/spectralbridge/blob/f7e7ddd2e01ee5f7cd74688eb479fd12fbc0b705/docs/vignettes/drone-processing.md.

— (2026i). *NeonCube and NEON ancillary-data implementation*. https://github.com/earthlab/spectralbridge/blob/f7e7ddd2e01ee5f7cd74688eb479fd12fbc0b705/src/spectralbridge/neon_cube.py. Source identifies portions adapted and simplified from HyTools under GPLv3; accessed 4 September 2026.

— (2026j). *Package architecture*. https://github.com/earthlab/spectralbridge/blob/f7e7ddd2e01ee5f7cd74688eb479fd12fbc0b705/docs/dev/architecture.md.
— (2026k). *Packaged Landsat and MicaSense band definitions*. https://github.com/earthlab/spectralbridge/blob/f7e7ddd2e01ee5f7cd74688eb479fd12fbc0b705/src/spectralbridge/data/landsat_band_parameters.json. Accessed 4 September 2026.
— (2026l). *Packaged Landsat brightness adjustments derived from Table Mountain HLS medians*. https://github.com/earthlab/spectralbridge/tree/f7e7ddd2e01ee5f7cd74688eb479fd12fbc0b705/src/spectralbridge/data/brightness. Immutable coefficient-directory snapshot inspected 4 September 2026.
— (2026m). *Pipeline outputs and file structure*. https://github.com/earthlab/spectralbridge/blob/f7e7ddd2e01ee5f7cd74688eb479fd12fbc0b705/docs/pipeline/outputs.md.
— (2026n). *Production validation record*. https://github.com/earthlab/spectralbridge/blob/f7e7ddd2e01ee5f7cd74688eb479fd12fbc0b705/docs/dev/production-validation-record.md. Accessed 4 September 2026.
— (2026o). *Production validation record: September snapshot*. Operator-reported production evidence; inspected 9 September 2026. URL: https://github.com/earthlab/spectralbridge/blob/c86da6a642a8a0bef37108df31af1fed1a882bd2/docs/dev/production-validation-record.md.
— (2026p). *R10C combined QA record*. https://github.com/earthlab/spectralbridge/blob/f7e7ddd2e01ee5f7cd74688eb479fd12fbc0b705/docs/validation/artifacts/r10c-l002-20210915/qa/combined/combined_qa.json. Archived at commit f7e7ddd; report provenance records generating commit d88e8d0947718d1e33b534080c224b703987e536.
— (2026q). *Real flightline QA walkthrough: R10C, D10, L002, 2021-09-15*. https://github.com/earthlab/spectralbridge/blob/f7e7ddd2e01ee5f7cd74688eb479fd12fbc0b705/docs/validation/real-data-example.md.
— (2026r). *SCS+C and kernel-driven BRDF correction implementation*. https://github.com/earthlab/spectralbridge/blob/f7e7ddd2e01ee5f7cd74688eb479fd12fbc0b705/src/spectralbridge/corrections.py. Adapted from HyTools; accessed 4 September 2026.
— (2026s). *Spectral resampling source*. https://github.com/earthlab/spectralbridge/blob/f7e7ddd2e01ee5f7cd74688eb479fd12fbc0b705/src/spectralbridge/resample.py. Accessed 4 September 2026.
— (2026t). *SpectralBridge ENVI header and writer implementation*. https://github.com/earthlab/spectralbridge/blob/f7e7ddd2e01ee5f7cd74688eb479fd12fbc0b705/src/spectralbridge/envi_writer.py. Source identifies portions adapted from HyTools under GPLv3; accessed 4 September 2026.
— (2026u). *SpectralBridge outputs and file structure*. Documentation inspected 25 September 2026; defines the public per-flightline artifact contract, success signals, restart checks, and QA reuse fingerprints. URL: https://earthlab.github.io/spectralbridge/pipeline/outputs/.
— (2026v). *SpectralBridge package metadata*. https://github.com/earthlab/spectralbridge/blob/f7e7ddd2e01ee5f7cd74688eb479fd12fbc0b705/pyproject.toml. Accessed 4 September 2026.
— (2026w). *SpectralBridge public GitHub releases*. Inspected 12 October 2026; v1.0.0 was listed as latest; no public 2.3.0rc1 GitHub release was listed. URL: https://github.com/earthlab/spectralbridge/releases.
— (2026x). *SpectralBridge repository README: September release-candidate snapshot*. Pinned source inspected 25 September 2026; documents the affine drone-translation registry, optional Landsat QA, compact bulk analysis, spectral-library reporting, and exact-artifact

smoke scope. URL: https://github.com/earthlab/spectralbridge/blob/c86da6a642a8a0bef37108df31af1fed1a882bd2/README.md.
— (2026y). *SpectralBridge source repository, commit f7e7ddd2e01ee5f7cd74688eb479fd12fbc0b705*. https://github.com/earthlab/spectralbridge/tree/f7e7ddd2e01ee5f7cd74688eb479fd12fbc0b705. Accessed 4 September 2026.
— (2026z). *SpectralBridge: Start Here*. https://github.com/earthlab/spectralbridge/blob/f7e7ddd2e01ee5f7cd74688eb479fd12fbc0b705/START_HERE.md.
— (2026aa). *Streaming bulk statistics implementation*. Source snapshot inspected 9 September 2026. URL: https://github.com/earthlab/spectralbridge/blob/c86da6a642a8a0bef37108df31af1fed1a882bd2/src/spectralbridge/bulk/streaming.py.
— (2026ab). *Topographic and BRDF implementation*. https://github.com/earthlab/spectralbridge/blob/f7e7ddd2e01ee5f7cd74688eb479fd12fbc0b705/src/spectralbridge/brdf_topo.py. See also corrections.py; accessed 4 September 2026.
— (2026ac). *Validation evidence index*. https://github.com/earthlab/spectralbridge/blob/f7e7ddd2e01ee5f7cd74688eb479fd12fbc0b705/docs/validation/index.md. Updated 4 September 2026.
— (2026ad). *Validation—save and restart behavior*. https://github.com/earthlab/spectralbridge/blob/f7e7ddd2e01ee5f7cd74688eb479fd12fbc0b705/docs/validation/save_restart.md.
— (2026ae). *Validation—sensor convolution*. https://github.com/earthlab/spectralbridge/blob/f7e7ddd2e01ee5f7cd74688eb479fd12fbc0b705/docs/validation/sensor_convolution.md.
Frantz, David (2019). "FORCE—Landsat + Sentinel-2 analysis ready data and beyond". In: *Remote Sensing* 11.9, p. 1124. DOI: 10.3390/rs11091124.
Huang, Chengquan, John R. G. Townshend, Shunlin Liang, Satish N. V. Kalluri, and Ruth S. DeFries (2002). "Impact of sensor's point spread function on land cover characterization: Assessment and deconvolution". In: *Remote Sensing of Environment* 80.2, pp. 203–212. DOI: 10.1016/S0034-4257(01)00298-X.
Ju, Junchang, Qiang Zhou, Brian Freitag, David P. Roy, Hankui Zhang, Madhu Sridhar, John Mandel, Saeed Arab, Gail L. Schmidt, Christopher J. Crawford, Ferran Gascon, Peter A. Strobl, Jeffrey G. Masek, and Christopher S. R. Neigh (2025). "The Harmonized Landsat and Sentinel-2 version 2.0 surface reflectance dataset". In: *Remote Sensing of Environment* 324, p. 114723. DOI: 10.1016/j.rse.2025.114723.
Kampe, Thomas U., Brian R. Johnson, Michele Kuester, and Michael Keller (2010). "NEON: The first continental-scale ecological observatory with airborne remote sensing of vegetation canopy biochemistry and structure". In: *Journal of Applied Remote Sensing* 4.1, p. 043510. DOI: 10.1117/1.3361375.
Kanewala, Upulee and James M. Bieman (2014). "Testing scientific software: A systematic literature review". In: *Information and Software Technology* 56.10, pp. 1219–1232. DOI: 10.1016/j.infsof.2014.05.006.
Köster, Johannes and Sven Rahmann (2012). "Snakemake—a scalable bioinformatics workflow engine". In: *Bioinformatics* 28.19, pp. 2520–2522. DOI: 10.1093/bioinformatics/bts480.
Li, Xiaowen and Alan H. Strahler (1992). "Geometric-optical bidirectional reflectance modeling of the discrete crown vegetation canopy: Effect of crown shape and mutual shadowing". In: *IEEE Transactions on Geoscience and Remote Sensing* 30.2, pp. 276–292. DOI: 10.1109/36.134078.
Morisette, Jeffrey T. et al. (2006). "Validation of global moderate-resolution LAI products: A framework proposed within the CEOS land product validation subgroup". In: *IEEE Transactions on Geoscience and Remote Sensing* 44.7, pp. 1804–1817. DOI: 10.1109/TGRS.2006.872529.

National Ecological Observatory Network (2026). *Spectrometer orthorectified surface directional reflectance—flightline, RELEASE-2026*. Data product DP1.30006.001. Accessed 4 September 2026. DOI: 10.48443/jhbd-n155.

Nicodemus, Fred E., Joseph C. Richmond, Jack J. Hsia, Irving W. Ginsberg, and Thomas Limperis (1977). *Geometrical Considerations and Nomenclature for Reflectance*. NBS Monograph 160. National Bureau of Standards. DOI: 10.6028/NBS.MONO.160.

Origo, Niall, Javier Gorroño, James Ryder, Joanne Nightingale, and Agnieszka Bialek (2020). "Fiducial Reference Measurements for validation of Sentinel-2 and Proba-V surface reflectance products". In: *Remote Sensing of Environment* 241, p. 111690. DOI: 10.1016/j.rse.2020.111690.

Queally, Natalie, Zhiwei Ye, Ting Zheng, Adam Chlus, Fabian Schneider, Ryan P. Pavlick, and Philip A. Townsend (2022). "FlexBRDF: A flexible BRDF correction for grouped processing of airborne imaging spectroscopy flightlines". In: *Journal of Geophysical Research: Biogeosciences* 127.1, e2021JG006622. DOI: 10.1029/2021JG006622.

Ross, Juhan (1981). *The Radiation Regime and Architecture of Plant Stands*. Vol. 3. Tasks for Vegetation Science. The Hague: Dr W. Junk Publishers. DOI: 10.1007/978-94-009-8647-3.

Roujean, Jean-Louis, Marc Leroy, and Pierre-Yves Deschamps (1992). "A bidirectional reflectance model of the Earth's surface for the correction of remote sensing data". In: *Journal of Geophysical Research: Atmospheres* 97.D18, pp. 20455–20468. DOI: 10.1029/92JD01411.

Roy, David P., Valerii Kovalskyy, Hankui K. Zhang, Eric F. Vermote, Lin Yan, S. S. Kumar, and Alexander Egorov (2016). "Characterization of Landsat-7 to Landsat-8 reflective wavelength and normalized difference vegetation index continuity". In: *Remote Sensing of Environment* 185, pp. 57–70. DOI: 10.1016/j.rse.2015.12.024.

Saunier, Sophie, Bringfried Pflug, Ignacio M. Lobos, Belen Franch, Jerome Louis, Raquel De Los Reyes, Virginie Debaecker, Elisabetta G. Cadau, Valentina Boccia, Ferran Gascon, and Sultan Kocaman (2022). "Sen2Like: Paving the way towards harmonization and fusion of optical data". In: *Remote Sensing* 14.16, p. 3855. DOI: 10.3390/rs14163855.

Schaaf, Crystal B., Feng Gao, Alan H. Strahler, Wolfgang Lucht, Xiaowen Li, Trevor Tsang, Nicholas C. Strugnell, Xiaoyang Zhang, Yufang Jin, Jan-Peter Muller, Philip Lewis, Michael Barnsley, Paul Hobson, Mathias Disney, Gareth Roberts, Michael Dunderdale, Christopher Doll, Robert P. D'Entremont, Baoxin Hu, Shunlin Liang, Jeffrey L. Privette, and David Roy (2002). "First operational BRDF, albedo nadir reflectance products from MODIS". In: *Remote Sensing of Environment* 83.1–2, pp. 135–148. DOI: 10.1016/S0034-4257(02)00091-3.

Schaepman-Strub, Gabriela, Michael E. Schaepman, Thomas H. Painter, Stefan Dangel, and John V. Martonchik (2006). "Reflectance quantities in optical remote sensing—definitions and case studies". In: *Remote Sensing of Environment* 103.1, pp. 27–42. DOI: 10.1016/j.rse.2006.03.002.

Soenen, Scott A., Derek R. Peddle, and Craig A. Coburn (2005). "SCS+C: A modified sun-canopy-sensor topographic correction in forested terrain". In: *IEEE Transactions on Geoscience and Remote Sensing* 43.9, pp. 2148–2159. DOI: 10.1109/TGRS.2005.852480.

SpectralBridge bulk evidence bundle (2026). *SpectralBridge bulk results visual report and numerical exports*. Author-supplied local evidence bundle. Report and CSV exports reviewed 9 September 2026. Primary filenames, analysis identifier, and SHA-256 checksums are recorded in the supplement. No permanent public identifier was supplied.

SpectralBridge project team (2026). *2023–2024 drone campaign production inventory*. Author-supplied production inventory. Reviewed 18 September 2026. Records 43 staged working HDF5 observations, approximately 7.15 GB of output, QA and JSON artifacts, absent ENVI and Parquet products, and orchestration status of 43 processed and zero failed. Exact production

revision, manifest, checksums, resolved configuration, root cause, and corrective revision were not supplied.

Teillet, Philippe M., Karl Staenz, and Darrel J. Williams (1997). "Effects of spectral, spatial, and radiometric characteristics on remote sensing vegetation indices of forested regions". In: *Remote Sensing of Environment* 61.1, pp. 139–149. DOI: 10.1016/S0034-4257(96)00248-9.

Wanner, Wolfgang, Xiaowen Li, and Alan H. Strahler (1995). "On the derivation of kernels for kernel-driven models of bidirectional reflectance". In: *Journal of Geophysical Research: Atmospheres* 100.D10, pp. 21077–21089. DOI: 10.1029/95JD02371.

Woodcock, Curtis E. and Alan H. Strahler (1987). "The factor of scale in remote sensing". In: *Remote Sensing of Environment* 21.3, pp. 311–332. DOI: 10.1016/0034-4257(87)90015-0.