


Parallel Algorithm for Closest Pair Problem

Ge Wu | July 24, 2013

INSTITUTE FOR THEORETICAL INFORMATICS



Closest Pair Problem

- Given n **different unordered** points $P = \{p_1, p_2, \dots, p_n\}$ in **unit square**


$$p_i = (x_i, y_i) \in (0, 1) \times (0, 1) \subset \mathbb{R}^2$$

- Find a pair of points with closest euclidean distance between them
- Find any pair if there's a tie.

- $O(n \log n)$ lower bound in comparison tree model
- **Bentley and Shamos 1976**
 $O(n \log n)$ algorithm using divide and conquer
- **Rabin 1976**
 $O(n)$ randomized algorithm with $O(1)$ floor function
- **Fortune and Hopcroft 1978**
 $O(n \log \log n)$ deterministic algorithm with $O(1)$ floor function
- **Khuller and Matias 1995**
Another $O(n)$ randomized algorithm


Algorithm

- Sample ($n^{\frac{2}{3}}$ Points from P)
- Partition
- Compute




Algorithm

- Sample ($n^{\frac{2}{3}}$ Points from P)
- Partition
- Compute




Algorithm

- Sample ($n^{\frac{2}{3}}$ Points from P)
- Partition
- Compute




Algorithm

- Sample ($n^{\frac{2}{3}}$ Points from P)
- Partition
- Compute




Algorithm

- Sample ($n^{\frac{2}{3}}$ Points from P)
- Partition
- Compute




Algorithm

- Sample ($n^{\frac{2}{3}}$ Points from P)
- Partition
- Compute




Algorithm

- Sample ($n^{\frac{2}{3}}$ Points from P)
- Partition
- Compute




Algorithm

- Sample ($n^{\frac{2}{3}}$ Points from P)
- Partition
- Compute




Algorithm

- Sample ($n^{\frac{2}{3}}$ Points from P)
- Partition
- Compute




Algorithm

- Sample ($n^{\frac{2}{3}}$ Points from P)
- Partition
- Compute




Algorithm

- Sample ($n^{\frac{2}{3}}$ Points from P)
- Partition
- Compute




Algorithm

- Sample ($n^{\frac{2}{3}}$ Points from P)
- Partition
- Compute



Algorithm

- Sample ($n^{\frac{2}{3}}$ Points from P)
- Partition
- Compute





- Compute all pair of distances: $O((n^{\frac{2}{3}})^2) = O(n^{\frac{4}{3}})$
- Divide & Conquer: $O(n^{\frac{2}{3}} \log n^{\frac{2}{3}}) = O(n^{\frac{2}{3}} \log n) \subset O(n)$
More samples possible: $O(\frac{n}{\log n} \log \frac{n}{\log n}) \subset O(n)$
- Other approach?

- Map points to the cells
- Divide the coordinates by cell length and **truncate** them to integer
- Partition $G = \{G_1, G_2, \dots, G_k\}$ with:

$$\bigcup_{i=1}^k G_i = P, G_i \cap G_j = \emptyset, k \leq n$$

- $O(n)$ with hashing

Partition



$$H(cell_{x_idx}, cell_{y_idx}) = (cell_{x_idx} + 2 \cdot cell_{y_idx}) \bmod 7$$

Theorem

Let G_i with $1 \leq i \leq k$ be the ***i-th*** non-empty cell and c, d some positive constants, then

$$\text{Prob} \left(\sum_{i=1}^k |G_i|^2 \leq c \cdot n \right) \geq 1 - \frac{1}{2^{n^d}}$$

Runtime for computing all pairs distances

$$O\left(\sum_{i=1}^k |G_i| \cdot |G_i \cup \text{neighbor}(G_i)|\right) = O\left(\sum_{i=1}^k |G_i|^2\right) = O(n)$$

Recursively calculate the closest distance on samples $S = \{s_1, \dots, s_{n^{\frac{2}{3}}}\}$

- Sample another $n^{\frac{4}{9}}$ points S' from S
- Get closest distance on S' by calculating all pair distances: $O(n^{\frac{8}{9}})$
- Partition and compute the closest distance on S

```
0 Cell set  $H = \emptyset$ , Result  $rst = \infty$ 

1 Choose set  $S$  of  $n^{\frac{2}{3}}$  samples  $O(n^{\frac{2}{3}})$ 
    $\delta = \min_{x,y \in S, x \neq y} (dist(x, y))$   $O(n^{\frac{8}{9}})$ 
2 for  $i = 1$  to  $n$  do  $O(n)$ 
    $c = cell(p_i)$ 
   if not  $H.findCell(c)$  then
      $H.addCell(\{c, idx = \{c.x, c.y\}\})$ 
      $H.findCell(c).addPoint(p_i)$ 
3 foreach  $c_1$  in  $H$  do  $O(n)$ 
   foreach  $c_2$  in  $\{c_1\} \cup neighbor(c_1)$  do
     foreach  $(p_i, p_j)$  in  $c_1 \times c_2$ 
        $rst = \min(rst, dist(p_i, p_j))$ 
```

Runtime: $O(n)$ with high probability, Worst: case $O(n^2)$

0 Cell set $H = \emptyset$, Result $rst = \infty$ **## H synchronized Hashmap**

1 Choose set S of $n^{\frac{2}{3}}$ samples **## allocate p PEs**

$\delta = \min_{x,y \in S, x \neq y} (\text{dist}(x, y))$ **## recurse once**

2 **for** $i = 1$ **to** n **do** **## allocate p PEs**

$c = \text{cell}(p_i)$

if not $H.\text{findCell}(c)$ **then**

$H.\text{addCell}(\{c, \text{idx} = \{c.x, c.y\}\})$ **## maximal n times**

$H.\text{findCell}(c).\text{addPoint}(p_i)$

3 **foreach** c_1 **in** H **do**

foreach c_2 **in** $\{c_1\} \cup \text{neighbor}(c_1)$ **do**

foreach (p_i, p_j) **in** $c_1 \times c_2$ **## allocate** $\lfloor p \cdot \frac{\# \text{Pairs}}{|c_1| |c_2|} \rfloor$ **PEs**

$rst = \min(rst, \text{dist}(p_i, p_j))$

Total runtime: $O(\frac{n}{p} + \log p)$ **## factor $\log p$ for collecting distances**

$$O(\frac{n^{\frac{2}{3}}}{p})$$

$$O(\frac{n^{\frac{8}{9}}}{p} + \log p)$$

$$O(\frac{n}{p})$$

$$O(\frac{n}{p} + \log p)$$

Theorem

Let G_i with $1 \leq i \leq k$ be the ***i-th*** non-empty cell and c, d some positive constants, then

$$\text{Prob} \left(\sum_{i=1}^k |G_i|^2 \leq c \cdot n \right) \geq 1 - \frac{1}{2^{n^d}}$$

Sampling Lemma

Let $G = \{G_1, G_2, \dots, G_k\}$ be a partition of set P , for which $N(D) \geq n$, where

$$N(G) = \sum_{i=1}^k \frac{|G_i| \cdot (|G_i| - 1)}{2}$$

If $n^{\frac{2}{3}}$ pairwise different elements are drawn at random from P then the probability, that two elements will be chosen from the same G_i , is at least $1 - 2^{-n^c}$ for some positive constant c

Proof see

- **Rabin 1976**
- **Dietzfelbinger et al. 1997**

Estimate the probability using Chebyshev's inequality

Lemma 2

Let G_δ be a grid with gap δ and $G_{2\delta}$ another grid with gap 2δ , which is obtained by ignoring every second line of G_δ , then


$$N(G_{2\delta}) \leq 4N(G_\delta) + \frac{3}{2}n$$

Let $k = \sum_{i=1}^4 k_i$

$f(x) = x(x-1)$ convex $\Rightarrow f(\frac{1}{4}k) \leq \frac{1}{4} \sum_{i=1}^4 f(k_i)$

$$\begin{aligned} \frac{1}{2}k(k-1) &= 8 \cdot \frac{1}{4}k(\frac{1}{4}k-1) + \frac{3}{2}k \\ &\leq 8 \cdot \frac{1}{4} \sum_{i=1}^4 k_i * (k_i - 1) + \frac{3}{2}k \end{aligned}$$

$$\Rightarrow N(G_{2\delta}) \leq 8 \cdot \frac{1}{2}N(G_\delta) + \frac{3}{2}n$$




Lemma 3

For any grid G_δ , $G_{\delta'}$ with $\delta' \leq \delta$ the following applies

$$N(G_{\delta'}) \leq 16N(G_\delta) + 6n$$

$$\begin{aligned} N(G_{\delta'}) &\leq \sum_{i=1}^4 N(G_{2\delta}^i) \\ &\leq \sum_{i=1}^4 (4N(G_\delta) + \frac{3}{2}n) \\ &= 16N(G_\delta) + 6n \end{aligned}$$




Lemma 3

For any grid G_δ , $G_{\delta'}$ with $\delta' \leq \delta$ the following applies

$$N(G_{\delta'}) \leq 16N(G_\delta) + 6n$$

$$\begin{aligned}
 N(G_{\delta'}) &\leq \sum_{i=1}^4 N(G_{2\delta}^i) \\
 &\leq \sum_{i=1}^4 (4N(G_\delta) + \frac{3}{2}n) \\
 &= 16N(G_\delta) + 6n
 \end{aligned}$$




Lemma 3

For any grid G_δ , $G_{\delta'}$ with $\delta' \leq \delta$ the following applies

$$N(G_{\delta'}) \leq 16N(G_\delta) + 6n$$

$$\begin{aligned} N(G_{\delta'}) &\leq \sum_{i=1}^4 N(G_{2\delta}^i) \\ &\leq \sum_{i=1}^4 (4N(G_\delta) + \frac{3}{2}n) \\ &= 16N(G_\delta) + 6n \end{aligned}$$




Lemma 3

For any grid G_δ , $G_{\delta'}$ with $\delta' \leq \delta$ the following applies

$$N(G_{\delta'}) \leq 16N(G_\delta) + 6n$$

$$\begin{aligned} N(G_{\delta'}) &\leq \sum_{i=1}^4 N(G_{2\delta}^i) \\ &\leq \sum_{i=1}^4 (4N(G_\delta) + \frac{3}{2}n) \\ &= 16N(G_\delta) + 6n \end{aligned}$$




Lemma 3

For any grid G_δ , $G_{\delta'}$ with $\delta' \leq \delta$ the following applies

$$N(G_{\delta'}) \leq 16N(G_\delta) + 6n$$

$$\begin{aligned} N(G_{\delta'}) &\leq \sum_{i=1}^4 N(G_{2\delta}^i) \\ &\leq \sum_{i=1}^4 (4N(G_\delta) + \frac{3}{2}n) \\ &= 16N(G_\delta) + 6n \end{aligned}$$



- Two of $n^{\frac{2}{3}}$ samples will be very likely in the same cell, if $N(G) \geq n$
- $N(G_{2\delta}) \leq 4N(G_\delta) + \frac{3}{2}n$
- $N(G_{\delta'}) \leq 16N(G_\delta) + 6n$ if $\delta' \leq \delta$

Proof.

Let δ^* be the grid gap, which makes $n \leq N(G_{\delta^*}) < 5.5n$. (It exists!)
If $n^{\frac{2}{3}}$ samples are randomly chosen, two of them will be in one cell with high probability. Let δ be the distance between them with $\delta < 2\delta^*$, then

$$\begin{aligned} N(G_\delta) &\leq 16N(G_{2\delta^*}) + 6n \\ &\leq 16(4N(G_{\delta^*}) + \frac{3}{2}n) + 6n \in O(n) \end{aligned}$$



- [BS76] Jon Louis Bentley and Michael Ian Shamos.
“Divide-and-conquer in multidimensional space”. In:
Proceedings of the eighth annual ACM symposium on Theory of computing. STOC '76. Hershey, Pennsylvania, USA: ACM, 1976, pp. 220–230. DOI: 10.1145/800113.803652. URL: <http://doi.acm.org/10.1145/800113.803652>.
- [CG89] Benny Chor and Oded Goldreich. “On the power of two-point based sampling”. In: *Journal of Complexity* 5.1 (1989), pp. 96–106.
- [Die+97] Martin Dietzfelbinger et al. “A reliable randomized algorithm for the closest-pair problem”. In: *Journal of Algorithms* 25.1 (1997), pp. 19–51.

- [FH78] Steven Fortune and John E Hopcroft. *A note on Rabin's nearest-neighbor algorithm*. Tech. rep. Cornell University, 1978.
- [KM95] Samir Khuller and Yossi Matias. “A simple randomized sieve algorithm for the closest-pair problem”. In: *Information and Computation* 118.1 (1995), pp. 34–37.
- [KT06] J. Kleinberg and E. Tardos. “Algorithm Design”. In: Pearson Education, 2006. Chap. 13 Randomized Algorithms.
- [Lip09] Richard J. Lipton. *Rabin Flips a Coin*. 2009. URL: <http://rjlipton.wordpress.com/2009/03/01/rabin-flips-a-coin/> (visited on 07/15/2013).
- [Rab76] Michael Oser Rabin. “Probabilistic algorithms”. In: *Algorithms and Complexity: New Directions and Recent Results*. Ed. by Joseph Frederick Traub. Academic Press, 1976, pp. 21–39.

- [Weba] URL: http://www.cse.ust.hk/tcsc/comp670r/Class_1_Notes.ppt/.
- [Webb] URL: http://en.wikipedia.org/wiki/Closest_pair_of_points_problem/.