ὼ ḡ╥

p0jX K_jX

I p l—Q l—。

[illegible] 5

[illegible] 5
1. [illegible] 5
2. [illegible] 5
3. [illegible] 6
4. [illegible] 6
5. [illegible] 6

Chapter 1. [illegible] Yule-Simpson [illegible] 7
1. [illegible] 7
2. [illegible] 7
3. [illegible] 7
4. [illegible] 7
5. Yule-Simpson [illegible] 8
6. [illegible] Yule-Simpson [illegible] 8
7. LaLonde [illegible] 9
8. [illegible] 9
9. [illegible] 9
10. 9

Chapter 2. [illegible] 11
1. [illegible] 11
2. [illegible] 11
3. [illegible] Science Table[illegible] 11
4. [illegible] 12
5. [illegible] 12
6. [illegible] 13
7. [illegible] Yule-Simpson [illegible] 13
8. [illegible] 13
9. SUTVA[illegible] 13
10. [illegible] 14
11. [illegible] 15
12. [illegible] 15
13. [illegible] 15
14. 15

Chapter 3. [illegible] 17
1. [illegible] 17
2. Fisher [illegible] 17

3. [illegible] 17
4. Sharp Null [illegible] FRT [illegible] 18
5. [illegible] p 18
6. Monte Carlo [illegible] 19
7. [illegible] 19
8. FRT [illegible] t [illegible] 26
9. LaLonde [illegible] 26
10. [illegible] 26
11. Sharp Null [illegible] 27
12. [illegible] 27
13. 27

Chapter 4. Neymanian [illegible] 29
1. [illegible] 29
2. [illegible] 29
3. [illegible] 30
4. Neyman (1923) [illegible] 30
5. [illegible] 31
6. Neyman (1923) [illegible] 31
7. [illegible] 32
8. Neymanian [illegible] OLS [illegible] 32
9. [illegible] 33
10. [illegible] 33
11. 34

Chapter 5. [illegible] 35
1. [illegible] 35
2. SRE [illegible] 35
3. [illegible] 36
4. SRE [illegible] Fisher [illegible] 36
5. Neymanian [illegible] 38
6. SRE CRE [illegible] 39
7. CRE [illegible] 40
8. [illegible] 40
9. [illegible] 40
10. 41

Chapter 6. [illegible] 43
1. [illegible] 43
2. [illegible] 43
3. ReM [illegible] 44
4. [illegible] 45
5. [illegible] 48
6. [illegible] 48
7. [illegible] 49
8. [illegible] 49
9. [illegible] 50
10. 50

Chapter 7. 53
1. 53
2. MPE 53
3. Fisher 53
4. Neymanian 56
5. 57
6. Darwin FRT 58
7. Workshop 59
8. 59
9. 59

Chapter 8. Fisher Studentized 61
1. Chapter 3 FRT Chapter 8 61
2. Studentized 61
3. Studentized 62
4. FRT 63
5. 64
6. 64
7. 65
8. 66
9. 66

Chapter 9. 69
1. 69
2. CRE 69
3. 71
4. SRE 71
5. 71
6. 72

Chapter 10. 73
1. 73
2. 73
3. 74
4. 74
5. 75
6. 76
7. 77
8. 77
9. 77

Chapter 11. 79
1. 79
2. 79
3. 80
4. 81
5. 82
6. 82

7. 83
8. 83
9. 83

Chapter 12. 85
1. 85
2. DR 85
3. 86
4. DR 86
5. 87
6. 87
7. 88
8. 88
9. 88
10. 89
11. 89
12. IPW DR 89
13. DR 90
14. 90
15. 90

Chapter 13. 93
1. 93
2. 93
3. 94
4. K 95
5. SRE 95
6. 95
7. 96
8. 97

Chapter 14. 99
1. IPW 99
2. Hajek WLS 99
3. Lin (2013) 100
4. WLS 101
5. 102
6. 102
7. 102

Chapter 15. 105
1. 105
2. 105
3. 105
4. 106
5. ATT 107
6. 107
7. LaLonde 107

8. [illegible] 108
9. 108

Chapter 16. [illegible] 109
1. [illegible] 109
2. [illegible] 109
3. [illegible] 110
4. [illegible] 111
5. 114
6. [illegible] 115

Chapter 17. E-Value[illegible] 117
1. [illegible] 117
2. Cornfield [illegible] 117
3. E-value 119
4. [illegible] 119
5. [illegible] 120
6. [illegible] 121
7. [illegible] 122
8. 122
9. [illegible] 122

Chapter 18. [illegible] 123
1. [illegible] 123
2. Manski [illegible] 123
3. [illegible] 124
4. R [illegible] 126
5. [illegible] 127
6. 128
7. [illegible] 129

Chapter 19. Rosenbaum [illegible] p [illegible] 131
1. [illegible] 131
2. [illegible] 131
3. [illegible] p 132
4. [illegible] 133
5. [illegible] 133
6. [illegible] 134

Chapter 20. [illegible] 135
1. [illegible] 135
2. [illegible] 135
3. [illegible] 136
4. [illegible] 136
5. [illegible] 137
[illegible] 138

Chapter 21. [illegible] 141
1. [illegible] 141

2. [illegible] 141
3. [illegible] 142
4. [illegible] 142

[illegible]

[illegible] Causal Inference [illegible]
[illegible]
[illegible]

[illegible]

[illegible]
[illegible]
[illegible]
Rubin [illegible] 1974 [illegible] Potential Outcomes Framework [illegible]
[illegible]
[illegible]
[illegible] Peng Ding [illegible] A First Course in Causal Inference [illegible]

- [illegible]
- Fisher [illegible] Neyman [illegible]
- [illegible]
- [illegible]
- [illegible]

[illegible]

[illegible]

(1) [illegible] **K**Peng Ding [illegible]
(2) [illegible] **K**`https://arxiv.org/abs/2305.18793`
(3) [illegible] **K**2023
(4) [illegible] **K**490
(5) [illegible] **K**
   - [illegible] Fisher [illegible] Neyman [illegible]
   - [illegible] CRE [illegible]
   - [illegible]
   - [illegible]

(6) [illegible] **K**This book provides an introduction to causal inference for students and researchers interested in estimating treatment effects from randomized experiments and observational studies. It unifies the potential outcomes framework with a focus on randomization-based inference, covering topics from basic concepts like the Yule-Simpson paradox to advanced methods including propensity score matching, instrumental variables, and sensitivity analysis. The book emphasizes both finite-sample exact inference (Fisher's approach) and asymptotic inference (Neyman's approach), providing readers with a comprehensive toolkit for causal analysis.

(7) [illegible] Causal Inference§ Potential Outcomes§ Randomization Test§ Neyman Inference§ Observational Studies§ Propensity Score

[illegible]

- [illegible] *Causal Inference for Statistics, Social, and Biomedical Sciences.* Cambridge University Press. [illegible]
- [illegible] "Estimating Causal Effects of Treatments in Randomized and Non-Randomized Studies." *Journal of Educational Psychology.* [illegible]
- [illegible] "The Central Role of the Propensity Score in Observational Studies for Causal Effects." *Biometrika.* [illegible]
- [illegible] *Mastering 'Metrics.* Princeton University Press. [illegible]

[illegible]

[illegible]

- [illegible]
- [illegible]
- [illegible]

[illegible]

- Markov [illegible] Chebyshev [illegible]
- Hoeffding [illegible] Chernoff [illegible]
- CLT [illegible]

[illegible]

(1) [illegible] 1-2 [illegible]
(2) [illegible] 3-4 [illegible] Fisher [illegible] Neyman [illegible]
(3) [illegible] 5 [illegible]
(4) [illegible] 6-8 [illegible]
(5) [illegible] 9 [illegible]
(6) [illegible] 10-18 [illegible]

[illegible]

[illegible]

- [illegible] [illegible] Fisher [illegible] Neyman [illegible]
- [illegible] [illegible]
- [illegible] [illegible]
- [illegible] [illegible]

CHAPTER 1

[illegible]

[illegible]

[illegible]

[illegible]

- ” [illegible] ”
- ” [illegible] ”
- ” [illegible] ”

[illegible] correlation [illegible]

[illegible]

[illegible]

[illegible] 1.1. 一家公司想要知道某种新药是否有效。他们给一组患者服用新药，另一组服用安慰剂。结果显示服药组的康复率更高。这能否说明药物有效？

[illegible]

- [illegible]
- [illegible]
- [illegible]

[illegible] causal inference [illegible]

[illegible]

[illegible]

- [illegible]
- [illegible]
- [illegible]

[illegible] 相关性可以是因果的，也可以是非因果的。

[illegible]

[illegible] Z [illegible] Y [illegible]

	$Y = 1$	$Y = 0$
$Z = 1$	p_{11}	p_{10}
$Z = 0$	p_{01}	p_{00}

[illegible]

$$\mathrm{RD} = \mathbb{P}(Y = 1 \mid Z = 1) - \mathbb{P}(Y = 1 \mid Z = 0) = \frac{p_{11}}{p_{11} + p_{10}} - \frac{p_{01}}{p_{01} + p_{00}}$$

[illegible]

$$\mathrm{RR} = \frac{\mathbb{P}(Y = 1 \mid Z = 1)}{\mathbb{P}(Y = 1 \mid Z = 0)} = \frac{p_{11}}{p_{11} + p_{10}} \Big/ \frac{p_{01}}{p_{01} + p_{00}}$$

[illegible]

$$\mathrm{OR} = \frac{\mathbb{P}(Y = 1 \mid Z = 1)/\mathbb{P}(Y = 0 \mid Z = 1)}{\mathbb{P}(Y = 1 \mid Z = 0)/\mathbb{P}(Y = 0 \mid Z = 0)} = \frac{p_{11}p_{00}}{p_{10}p_{01}}$$

[illegible] $\perp\!\!\!\perp$ [illegible]

[illegible] (1) *以下命题等价：$Z \perp\!\!\!\perp Y$、$RD = 0$、$RR = 1$、$OR = 1$。*

(2) *若所有 $p_{zy} > 0$，则 $RD > 0$ 等价于 $RR > 1$，也等价于 $OR > 1$。*

(3) *若 $\mathbb{P}(Y = 1 \mid Z = 1)$ 和 $\mathbb{P}(Y = 1 \mid Z = 0)$ 都很小，则 $OR \approx RR$。*

[illegible]

[illegible]

[illegible] 1.2 ([illegible] Yule-Simpson [illegible]). 考虑一个大学录取数据：

院系	男生录取率	女生录取率
A	*62% (341/550)*	*82% (137/167)*
B	*63% (28/44)*	*68% (24/35)*
	45% (369/816)	***30% (161/525)***

有趣的现象：

- 在 *A* 院系，男女生录取率分别是 *62%* 和 *82%*
- 在 *B* 院系，男女生录取率分别是 *63%* 和 *68%*
- 但总体来看，男生的录取率反而更高！

这就是 *Yule-Simpson* 悖论：[illegible]。

[illegible] 如果我们不控制混杂因素（*confounding factors*），就可能得出完全错误的因果结论

[illegible]

[illegible]

$$\begin{pmatrix} X \\ Y \\ Z \end{pmatrix} \sim N\left(\begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 1 & \rho_{XY} & \rho_{XZ} \\ \rho_{XY} & 1 & \rho_{YZ} \\ \rho_{XZ} & \rho_{YZ} & 1 \end{pmatrix}\right)$$

Y [illegible] Z [illegible] $\rho_{YZ|X}$ [illegible] $(Y, Z) \mid X$ [illegible]

[illegible]

$$\rho_{YZ|X} = \frac{\rho_{YZ} - \rho_{YX}\rho_{ZX}}{\sqrt{1 - \rho_{YX}^2}\sqrt{1 - \rho_{ZX}^2}}$$

[illegible] $\rho_{YZ} > 0$ [illegible] $\rho_{YZ|X}$ [illegible] Yule-Simpson [illegible]

[illegible]

LaLonde [illegible] National Supported Work Demonstration [illegible] $2^{10} = 1024$ [illegible] treatment [illegible]

[illegible]

Bertrand and Mullainathan (2004) [illegible]

[illegible] ” [illegible] ” [illegible]

[illegible]

[illegible]

(1) [illegible] $\neq$ [illegible]
(2) [illegible] RD、RR、OR [illegible]
(3) 《 [illegible]
(4) [illegible]
(5) [illegible]

[illegible]

[illegible]

[illegible] 1.1. { {独立性命题的证明 证明 *Proposition* { {中的命题 *(1)* 和 *(2)*。

[illegible] 1.2. { {*Yule-Simpson* 悖论的更多例子

(1) 给出一个 $2 \times 2 \times 2$ 列联表的数值例子，其中出现 *Yule-Simpson* 悖论。
(2) 找一个现实生活中的 *Yule-Simpson* 悖论例子。

[illegible] 1.3. { {相关与偏相关 考虑三维正态随机向量 $(X, Y, Z)^\top \sim N(\mathbf{0}, \Sigma)$。
证明偏相关系数公式（见 *Proposition* { }{。
给出一个 $\rho_{YZ} > 0$ 但 $\rho_{YZ|X} < 0$ 的数值例子。
：这是正态随机向量情形下的 *Yule-Simpson* 悖论。

[illegible] 1.4. { {规格搜索 重做 *Hainmueller (2012)* 的数据分析。使用结局 ***re78*** 和治疗 ***treat***。数据集还包含 *10* 个协变量，共有 $2^{10} = 1024$ 种可能的协变量子集组合。运行所有可能的线性回归子集，并报告 *treatment* 系数的分布：有多少是正显著、负显著、不显著的？

[illegible] 1.5. { {种族歧视的进一步分析 重做 *Bertrand and Mullainathan (2004)* 的简历数据分析。分别对男性和女性进行分析，讨论亚组分析的结果与总体结论的差异。

ΰ́ 1.6. { {推荐阅读 ┤ḓ : { 是关于 *Berkeley* 录取数据悖论的原始论文。{ 从因果推断角度重新审视了该研究。

α̈ɥ e : { 是统计因果推断领域的经典综述文章，它推广了 ”*Rubin* 因果模型 ” 这一名称。在 *UC Berkeley*，我们称之为 ”*Neyman* 模型 ”——原因显而易见。

CHAPTER 2

[illegible]

[illegible]

[illegible] Yule-Simpson [illegible]

[illegible]

[illegible]

[illegible] Neyman [illegible] 1923 [illegible] Rubin [illegible] 1970 [illegible] counterfactual [illegible]

[illegible]

[illegible] i [illegible]

- $T_i = 1$ [illegible]
- $T_i = 0$ [illegible]

[illegible]

- $Y_i(1)$ [illegible]
- $Y_i(0)$ [illegible]

[illegible] $Y_i(1)$ [illegible] $Y_i(0)$ [illegible]

2.1 (2.1 [illegible]). [illegible] (*counterfactual*) 指的是未被观测到的潜在结果。具体而言：

- 对于治疗组个体 i（$T_i = 1$），我们观察到 $Y_i(1)$，而 $Y_i(0)$ 是反事实
- 对于对照组个体 i（$T_i = 0$），我们观察到 $Y_i(0)$，而 $Y_i(1)$ 是反事实

正因为潜在结果框架讨论的是“如果接受相反治疗会怎样”，所以它也被称为[illegible] (*counterfactual framework*)。

2.2. [illegible] “[illegible]” [illegible]**：**因为它们描述的是“如果... 会怎样”的情形，这种结果是“潜在的”而非“现实的”。这正是 *counterfactual*（反事实）思维的本质。

[illegible]

[illegible] $n \times 2$ [illegible]

i	$Y_i(1)$	$Y_i(0)$
1	$Y_1(1)$	$Y_1(0)$
2	$Y_2(1)$	$Y_2(0)$
$\vdots$	$\vdots$	$\vdots$
n	$Y_n(1)$	$Y_n(0)$

[illegible]

[illegible]

[illegible] i [illegible]

$$\tau_i = Y_i(1) - Y_i(0)$$

[illegible] “[illegible]” [illegible] 1923 [illegible] Jerzy Neyman [illegible]

对于每个个体，我们只能观察到一种潜在结果——实际发生的那种。另一种潜在结果是潜在的、永不可见的。

[illegible]

2.3. [illegible]：因为科学上感兴趣的大多数因果问题，都涉及将现实与反事实进行比较。我们可以问“这种药对这个患者有效吗？”——但这个问题要求我们同时知道患者吃药和不吃药的结果，而这是不可能的。

[illegible]

[illegible]

$$\tau = \mathbb{E}[Y_i(1) - Y_i(0)] = \bar{Y}(1) - \bar{Y}(0) \tag{2.1}$$

[illegible][1]

$$\tau = \frac{1}{n}\sum_{i=1}^{n}\{Y_i(1) - Y_i(0)\} = \frac{1}{n}\sum_{i=1}^{n} Y_i(1) - \frac{1}{n}\sum_{i=1}^{n} Y_i(0) \tag{2.2}$$

[illegible]

$$\tau = \bar{Y}(1) - \bar{Y}(0) = \frac{1}{n}\sum_{i=1}^{n}\tau_i \tag{2.3}$$

---

[1] [illegible] **??**

[illegible]

[illegible] X_i [illegible]

- [illegible] $X_i = 1$ [illegible] $X_i = 0$ [illegible]
- [illegible] / [illegible]
- [illegible] / [illegible]

[illegible][2]

$$\tau_x = \frac{\sum_{i=1}^{n} \mathbb{I}(X_i = x)\{Y_i(1) - Y_i(0)\}}{\sum_{i=1}^{n} \mathbb{I}(X_i = x)}, \quad (x = 0, 1) \tag{2.4}$$

[illegible]

[illegible][3]

$$\tau = \pi_1\tau_1 + \pi_0\tau_0 \tag{2.5}$$

[illegible] $\pi_x = \frac{1}{n}\sum_{i=1}^{n} \mathbb{I}(X_i = x)$ [illegible] $X_i = x$ [illegible] $\tau_1 > 0$ [illegible] $\tau_0 > 0$ [illegible] $\tau > 0$ [illegible]

[illegible]

[illegible]

[illegible] T_i [illegible] i [illegible] Y_i [illegible]

$$\begin{aligned}
Y_i &= \begin{cases} Y_i(1), & \text{if } T_i = 1 \\ Y_i(0), & \text{if } T_i = 0 \end{cases} && (1)\\
&= T_i Y_i(1) + (1 - T_i) Y_i(0) && (2)\\
&= Y_i(0) + T_i\{Y_i(1) - Y_i(0)\} && (3)\\
&= Y_i(0) + T_i\tau_i && (4)
\end{aligned}$$

[illegible] $\tau_i = Y_i(1) - Y_i(0)$ [illegible]

[illegible]

[illegible]

[illegible] 2.1 ([illegible]). 单元 i 的潜在结果不依赖于其他单元的治疗分配。这有时被称为[illegible]假设。

[illegible] 2.2 ( ). 治疗水平没有其他版本。等价地，我们要求治疗水平是良好定义的，至少对关注的结局是如此。这有时被称为 假设。

[illegible] 2.3 (SUTVA). 假设 [illegible]和 [illegible]同时成立。

---

[2] [illegible] ??

[3] [illegible] ??

¹I¹ alho ÂN×1$ ÂT†í-Wi Ä €Í çl,ü' *¹.
²T $Y_i(1)$ „ $Y_i(0)$ ü,^ r2HH@/¥}iMñ2ì¥´||%¿ ð¦¥›¹ Í ¥
8™T œñ•Oüö)ïb

´Q alho ¥è0

íÄ L! MQ@BMi2`72`2M+2 ´bQ¥mf™KTiBQM

UR.V™hÐfì¤Õ BM72+iBQmb /Bb2 b2b M/'%þh++µBM™ŸiBQM
H Bñ¦^Ž¤Õfì,Ç•Yð1-¥Ñh„B Î•YðÛ¶¦$Ë™¥
ÀqbèÂ ÂTá¥]Y¤Õ @Ëfì áÑ@Ë¥Àq9ö†®ii'P
á1-Àµ¤Õb ƒÕfƒ/ †í i ¥. ²T $Y_i(1)$ „ $Y_i(0)$ ûG ¿ ð
¦¥Í s¥b

Uk©VŽr‹L M2irQ`F 2tT2`BK2MŒi©bŽL Ï ÂTBñ¦lž
6 +2#Q<QáF i`2 iK2ðM¥i´¶AžƒH<áisž•Yª V?9ö ™
¾ÁÖii'P´¶'¦Àµ°¤lž<ábƒü^îÌ¥ n r‹ bTBHHQp2`
2z2+ib

UjVÛ uFL Ï¥M#¹v M2B;?#Q`BM; THQib BM ` M/QKBx2/ #HQ+
K2Mibj<L Ï M#¹v¥TþV?Mº•YiiÂTM#¹vP¨ g
V?öYVr›1i•Y'¹v¥Á b

BÁŸL! +QMbBbi2M+v bb´mQK¥f¯î™BQM

UR¯V†Í +QKTQbBi2 i`2 iKÍ2M&iµb¯ Fî†sH M]¥ óÍ
ôS&V? ‹,]¥L=ÄçbèÂ
ùî õ bKQFBkM´;¥•YH óõ ôƒñÍ µ Õ™TiiPõ
Ë˜a õÔqa Æ' © ƒtûV?•Á,]¥. ²Tb
ùîvÐ•â +QHH2;2 2/mlÆ¥BYQHM óvÐ ôƒñÍ "
vÐË˜ N38fkRèRYpýbna <aÓÝ©)© ƒtµsö/î.
²T¥,BÁb

Uk4V r‹ /Qb ;2 2z2+iDbÐùîÏ M]¥0þ ëV? ‹,]4
74 °¤•Y rbèÂ ó›¨-³Ä ôT¹Í Œ,]4 ¥-³Ä ö
Á3›†,]¥. ²Tb

UjVL@Ms BKTH2K2Mi iBQM p'`FBÍ iB QMM] ,]L@€V
?„ÆµsbèÂ •ØÍ ùîÏ ,]Í =¥„ì„/?V?•ÁM]¥
ó•ØÍ ôÁ3,]rTb

' alho ,îëH . ²T•O³1©žb C}yTw ÓD Â \ ž ù
îÂ…)Øi Ä Í ñ' "¥f™b

ÿ kXX9alho ¥åN×1Ÿ , _m#BM U|RN¸3hyoV 'ë¹. ²T•O¥ $
L! Àµ alho yTw ¥"ÐÄ^,V?¥b

RyXQYL±»¥†

9¸Föz ¹I¹,?e†¹Ù óL=?3¥²T^I¹ ô$
sÃ^ yTr‹'É ^BñQYL +QmMi2`7 ÀiQm bH'áÌÙ óƒÕ0^
Žµr ôH áÌL= ^ Ù

óÂTƒñÑ€ù 0 ²Tö8"$ŒCL^ ðV?Àµù0 XXXô

ƒÕ±»ZTV?ž¦$¤ `.À}ÐÄ Œñ^áÌ ,yT¥·BøÈZTb.
²T•O¥N´ ¿ ñ|ƒÕ}Ðó³ Ä¹b ¥"ÐÔýb

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

$$\delta_1 = \text{median}\{Y_i(1)\}_{i=1}^n - \text{median}\{Y_i(0)\}_{i=1}^n \tag{2.6}$$

[illegible]

$$\delta_2 = \text{median}\{Y_i(1) - Y_i(0)\}_{i=1}^n \tag{2.7}$$

[illegible][4]

[illegible]

[illegible]

(1) [illegible] $Y_i(1)$ [illegible] $Y_i(0)$
(2) [illegible] $n \times 2$ [illegible] $\{Y_i(1), Y_i(0)\}_{i=1}^n$
(3) [illegible] $\tau_i = Y_i(1) - Y_i(0)$
(4) [illegible] $\tau = \bar{Y}(1) - \bar{Y}(0) = n^{-1}\sum_{i=1}^n \tau_i$
(5) [illegible]
(6) [illegible] $Y_i = T_iY_i(1) + (1 - T_i)Y_i(0)$
(7) [illegible] + [illegible]
(8) [illegible]

[illegible]

[illegible]

[illegible] 2.1. [illegible] 完美的医生 假设潜在结果由以下模型生成：

$$Y_i(0) \sim N(0,1), \quad \tau_i = -0.5 + Y_i(0), \quad Y_i(1) = Y_i(0) + \tau_i$$

治疗分配为 $T_i = \mathbb{I}(\tau_i \geq 0)$，观测结果为 $Y_i = T_iY_i(1) + (1 - T_i)Y_i(0)$。

计算条件均值差 $\mathbb{E}(Y_i \mid T_i = 1) - \mathbb{E}(Y_i \mid T_i = 0)$。

[illegible]：需要使用截断正态分布的均值公式。当 $X \sim N(\mu, \sigma^2)$ 时，

$$\mathbb{E}(X \mid a < X < b) = \mu - \sigma\frac{\phi\left(\frac{b-\mu}{\sigma}\right) - \phi\left(\frac{a-\mu}{\sigma}\right)}{\Phi\left(\frac{b-\mu}{\sigma}\right) - \Phi\left(\frac{a-\mu}{\sigma}\right)}$$

---

[4] [illegible] ??

其中 ϕ 和 Φ 分别是标准正态分布的密度函数和分布函数。

ΰ 2.2. { {非线性因果参数 证明平均因果效应是线性因果参数的例子，即 $\bar{Y}(1) - \bar{Y}(0) = n^{-1}\sum_{i=1}^{n}\{Y_i(1) - Y_i(0)\}$。

其他参数可能不是线性的。例如，定义中位数治疗效应为：

$$\delta_1 = median\{Y_i(1)\} - median\{Y_i(0)\}$$

这通常不同于个体治疗效应的中位数：

$$\delta_2 = median\{Y_i(1) - Y_i(0)\}$$

(1) 给出数值例子分别满足 $\delta_1 = \delta_2$、$\delta_1 > \delta_2$ 和 $\delta_1 < \delta_2$。
(2) 哪个参数更有意义？为什么？请用例子来证明你的结论。

ΰ 2.3. { {平均效应与个体效应 给出一个数值例子，其中 $\tau = n^{-1}\sum_{i=1}^{n}\{Y_i(1) - Y_i(0)\} > 0$，但有不到一半的单元满足 $Y_i(1) > Y_i(0)$。即平均因果效应为正，但治疗对不到一半的单元有益。

ΰ 2.4. { {推荐阅读 ⊣ḓ : { 是统计因果推断领域的经典综述文章，它推广了 *“Rubin* 因果模型*”* 这一名称。在 *UC Berkeley*，我们称之为 *“Neyman* 模型*”*——原因显而易见。

CHAPTER 3

[illegible]

[illegible]

[illegible]

$$\tau = \mathbb{E}[Y_i(1) - Y_i(0)] = \bar{Y}(1) - \bar{Y}(0) \tag{5}$$

[illegible] $Y_i(1)$ [illegible] $Y_i(0)$ [illegible][1] [illegible]

[illegible] X [illegible] Y [illegible] X [illegible] Y [illegible]

[illegible]

Ronald Fisher 1935 [illegible][3] Design of Experiments [illegible]

(1) [illegible] treatment [illegible] control

(2) [illegible]

[illegible] treatment [illegible] control [illegible]

Fisher [illegible][2,3] [illegible]

[illegible]

[illegible] n [illegible] n_1 [illegible] n_0 [illegible] $n = n_1 + n_0$

3.1 ([illegible] Complete Randomized Experiment, CRE). 固定 n_1 和 n_0 (其中 $n = n_1 + n_0$)。完全随机实验的治疗 *assignment* 机制为:

$$\mathbb{P}(\mathbf{Z} = \mathbf{z}) = \frac{1}{\binom{n}{n_1}}, \tag{6}$$

其中 $\mathbf{z} = (z_1, \ldots, z_n)$ 满足 $\sum_{i=1}^{n} z_i = n_1$ 和 $\sum_{i=1}^{n}(1 - z_i) = n_0$。

---

[1] [illegible] **??**

[2] [illegible] Fisher Randomization Test [illegible] **??**

[3] [illegible] Fisher [illegible] **??**

[illegible] $\mathbf{Y}(1)$ [illegible] $\mathbf{Y}(0)$ [illegible]

$$\mathbb{P}\{\mathbf{Z}=\mathbf{z} \mid \mathbf{Y}(1),\mathbf{Y}(0)\} = \frac{1}{\binom{n}{n_1}} \tag{7}$$

CRE $\mathbf{Z} \perp\!\!\!\perp \{\mathbf{Y}(1),\mathbf{Y}(0)\}$ [illegible] CRE [illegible] $\mathbf{Z}$ [illegible] n_1 [illegible] 1 [illegible] n_0 [illegible] 0 [illegible]

[illegible]

[illegible]

$$H_{0F}: Y_i(1) = Y_i(0) \quad [illegible] \quad i = 1,\dots,n. \tag{8}$$

[illegible] $\mathbf{Y}(1) = \mathbf{Y}(0) = \mathbf{Y} = (Y_1,\dots,Y_n)$ [illegible][4] [illegible] H_{0F} [illegible] FRT [illegible]

$$T = T(\mathbf{Z},\mathbf{Y}), \tag{9}$$

[illegible] $\mathbf{Y}$ [illegible] H_{0F} [illegible] T [illegible] $\mathbf{Z}$ [illegible] $\mathbf{Z}$ [illegible] T [illegible] p [illegible] CRE [illegible] $\mathbf{Z}$ [illegible] $\{\mathbf{z}^1,\dots,\mathbf{z}^M\}$ [illegible] $M = \binom{n}{n_1}$ [illegible] $\mathbf{z}^m$ [illegible] n_1 [illegible] 1 [illegible] n_0 [illegible] 0 [illegible][5]

[illegible]

[illegible] T [illegible]

$$p_{\text{frt}} = M^{-1}\sum_{m=1}^{M} \mathbb{I}\{T(\mathbf{z}^m,\mathbf{Y}) \ge T(\mathbf{Z},\mathbf{Y})\}. \tag{10}$$

[illegible] Fisher [illegible] p [illegible] p_{frt} [illegible] H_{0F} [illegible] finite-sample exact[illegible][6]

$$\mathbb{P}(p_{\text{frt}} \le u) \le u \quad \text{for all} \quad 0 \le u \le 1. \tag{11}$$

3.1. [illegible] ” [illegible] ” [illegible]：这意味着 p 值的分布（虽然在离散情况下是阶梯函数）满足上式，因此 p 值在 $[0,1]$ 上均匀分布（对于连续情况）。

---

[4] [illegible] H_{0F} [illegible] $\mathbf{Y}$ [illegible] ??

[5] [illegible] $\{\mathbf{z}^1,\dots,\mathbf{z}^M\}$ [illegible] ??

[6] [illegible] ??

[illegible]

[illegible] M [illegible] $n = 100, n_1 = 50$ [illegible] $M > 10^{29}$ [illegible] Monte Carlo [illegible] p_{frt} [illegible] $\tilde{p}_{\text{frt}}$ [illegible] [7][8]

$$\hat{p}_{\text{frt}} = R^{-1} \sum_{r=1}^{R} \mathbb{I}\{T(\mathbf{z}^r, \mathbf{Y}) \geq T(\mathbf{Z}, \mathbf{Y})\}, \tag{12}$$

[illegible] $\mathbf{z}^r$ [illegible] $\mathbf{Z}$ [illegible] R [illegible] p_{frt} [illegible] $\mathbf{Z}$ [illegible] FRT [illegible] CRE [illegible]

$$\tilde{p}_{\text{frt}} = \frac{1 + \sum_{r=1}^{R} \mathbb{I}\{T(\mathbf{z}^r, \mathbf{Y}) \geq T(\mathbf{Z}, \mathbf{Y})\}}{1 + R}. \tag{13}$$

[illegible] R [illegible] p

[illegible]

FRT [illegible] p [illegible] H_{0F} [illegible]

[illegible]

$$\hat{\tau} = \hat{\bar{Y}}(1) - \hat{\bar{Y}}(0), \tag{14}$$

[illegible]

$$\hat{\bar{Y}}(1) = n_1^{-1} \sum_{Z_i=1} Y_i = n_1^{-1} \sum_{i=1}^{n} Z_i Y_i \tag{15}$$

[illegible] treatment [illegible]

$$\hat{\bar{Y}}(0) = n_0^{-1} \sum_{Z_i=0} Y_i = n_0^{-1} \sum_{i=1}^{n} (1 - Z_i) Y_i \tag{16}$$

[illegible] control [illegible] H_{0F} [illegible] $\hat{\tau}$ [illegible] 0 [illegible]

$$\text{var}(\hat{\tau}) = \frac{n}{n_1 n_0} s^2, \tag{17}$$

[illegible] $s^2 = (n-1)^{-1} \sum_{i=1}^{n} (Y_i - \bar{Y})^2$ [illegible] $\bar{Y} = n^{-1} \sum_{i=1}^{n} Y_i$ [9] [illegible] Finite Population Central Limit Theorem [illegible]

$$\frac{\hat{\tau}}{\sqrt{\frac{n}{n_1 n_0} s^2}} \rightarrow \mathcal{N}(0, 1) \tag{18}$$

in distribution.

---

[7] [illegible] Monte Carlo [illegible] **??**

[8] [illegible] $\mathbf{Z}$ [illegible] **??**

[9] [illegible] **??**

[illegible] H_{0F} [illegible] p [illegible] t [illegible]
p
[illegible] **: k o “ :** [illegible][10] [illegible]

$$(n-1)s^2 = \sum_{Z_i=1}\{Y_i - \hat{\bar{Y}}(1)\}^2 + \sum_{Z_i=0}\{Y_i - \hat{\bar{Y}}(0)\}^2 + \frac{n_1 n_0}{n}\hat{\tau}^2. \tag{19}$$

[illegible] ANOVA [illegible] $Y_i - \bar{Y}$ [illegible] $(Y_i - \hat{\bar{Y}}(Z_i))$ [illegible] $(\hat{\bar{Y}}(Z_i) - \bar{Y})$ [illegible] treatment [illegible] control [illegible] $\frac{n_1 n_0}{n}\hat{\tau}^2$ [11]

3.2. [illegible] ***FRT*** [illegible] ***t*** [illegible] **:** 传统 *t* 检验需要正态性假设。但 *FRT* 框架表明，只要随机化分配正确执行，*t* 检验程序在有限总体 + 随机化情境下可以使用——渐近理论只依赖于有限总体 *CLT*，而不依赖于数据的原始分布。

[illegible]

$$t = \frac{\hat{\bar{Y}}(1) - \hat{\bar{Y}}(0)}{\sqrt{\frac{\hat{S}^2(1)}{n_1} + \frac{\hat{S}^2(0)}{n_0}}}, \tag{20}$$

[illegible]

$$\hat{S}^2(1) = (n_1 - 1)^{-1}\sum_{Z_i=1}[Y_i - \hat{\bar{Y}}(1)]^2, \quad \hat{S}^2(0) = (n_0 - 1)^{-1}\sum_{Z_i=0}[Y_i - \hat{\bar{Y}}(0)]^2 \tag{21}$$

[illegible] treatment [illegible] control [illegible] H_{0F} [illegible] CLT [illegible] t [illegible] $t \to \mathcal{N}(0,1)$ [illegible] `t.test(var.equal = FALSE)` [illegible] p [illegible]

3.3. [illegible] **:** 当 *treatment* 组和 *control* 组的方差差异较大时，差值均值的检验可能失效。学生化通过分别估计两组的方差，能够更好地处理异方差情况。

[illegible] Chi-Square Test [illegible]

[illegible]

Insight [illegible] t [illegible]

[illegible] Karl Pearson [illegible] 1900 [illegible]

---

[10] [illegible] ??

[11] ANOVA [illegible] ??

7.3.1. 拟合优度检验. [illegible] k [illegible] $O_1, O_2, \ldots, O_k$ [illegible] n [illegible] H_0 [illegible] i [illegible] $E_i = np_{i0}$ [illegible] p_{i0} [illegible] H_0 [illegible]

[illegible]

$$\chi^2 = \sum_{i=1}^{k} \frac{(O_i - E_i)^2}{E_i}. \tag{22}$$

[illegible] $(O_i - E_i)^2/E_i$ [illegible] E_i [illegible] 100 [illegible] 100 [illegible]

[illegible] H_0 [illegible] n [illegible]

$$\chi^2 \xrightarrow{d} \chi^2_{k-1}, \tag{23}$$

[illegible] $k-1$ [illegible] $\sum O_i = \sum E_i = n$

Insight [illegible] $k-1$ [illegible] k [illegible] n [illegible] k [illegible] $k-1$ [illegible]

[illegible] 3.1 ([illegible]). [illegible]：投掷一枚骰子 *120* 次，想检验它是否均匀（各面概率均为 1/6）。

[illegible]：

面	*1*	*2*	*3*	*4*	*5*	*6*
观测频数 O_i	*15*	*22*	*18*	*25*	*17*	*23*
期望频数 E_i	*20*	*20*	*20*	*20*	*20*	*20*

[illegible]：

$$\chi^2 = \frac{(15-20)^2}{20} + \frac{(22-20)^2}{20} + \frac{(18-20)^2}{20} + \frac{(25-20)^2}{20} + \frac{(17-20)^2}{20} + \frac{(23-20)^2}{20} = 4.3$$

自由度 $k-1=5$，查表得 $\chi^2_{0.05,5} = 11.07$。

由于 $4.3 < 11.07$，在 $\alpha = 0.05$ 水平下[illegible] 均匀性假设。

R [illegible]：

```
> obs <- c(15, 22, 18, 25, 17, 23)
> chisq.test(obs, p = rep(1/6, 6))

	Chi-squared test for given probabilities

X-squared = 4.3, df = 5, p-value = 0.507
```

p 值为 0.507，远大于 0.05，与手工计算一致。

□

Insight [illegible] $E_i = 20$ [illegible] < 5 [illegible] exact test[illegible]

[illegible] 3.2 ([illegible]). [illegible]：孟德尔杂交黄色圆粒豌豆与绿色皱粒豌豆，预期后代比例为：

$$\text{黄圆} : \text{黄皱} : \text{绿圆} : \text{绿皱} = 9 : 3 : 3 : 1$$

[illegible] (共 *556* 株)：

$$O = (315, 101, 108, 32), \quad E = (556 \times \tfrac{9}{16}, \dots) = (312.75, 104.25, 104.25, 34.75)$$

[illegible]：

$$\chi^2 = \frac{(315-312.75)^2}{312.75} + \frac{(101-104.25)^2}{104.25} + \frac{(108-104.25)^2}{104.25} + \frac{(32-34.75)^2}{34.75} \approx 0.47$$

自由度 $k-1=3$，查表得 $\chi^2_{0.05,3}=7.81$。由于 $0.47<7.81$，[illegible] 孟德尔比例。

R [illegible]：

```
> obs <- c(315, 101, 108, 32)
> prob <- c(9/16, 3/16, 3/16, 1/16)
> chisq.test(obs, p = prob)

    Chi-squared test for given probabilities

X-squared = 0.47, df = 3, p-value = 0.925
```

p 值为 0.925——数据与孟德尔比例高度吻合。

□

7.3.2. 独立性检验. [illegible]

[illegible] $r \times c$ [illegible] X [illegible] Y [illegible] O_{ij} [illegible] (i,j) [illegible] $O_{i\cdot} = \sum_j O_{ij}$ [illegible] i [illegible] $O_{\cdot j} = \sum_i O_{ij}$ [illegible] j [illegible] n

[illegible] $H_0 : X \perp Y$ [illegible]

$$E_{ij} = \frac{O_{i\cdot} \cdot O_{\cdot j}}{n}. \tag{24}$$

[illegible]

$$\chi^2 = \sum_{i=1}^{r} \sum_{j=1}^{c} \frac{(O_{ij} - E_{ij})^2}{E_{ij}}. \tag{25}$$

[illegible] H_0 [illegible] $(r-1)(c-1)$

Insight [illegible] 2×2 [illegible] 1 [illegible] p [illegible]

[illegible] 2×2 [illegible] R [illegible] `chisq.test` [illegible] Yates [illegible]

$$\chi^2_{\text{Yates}} = \sum \frac{(|O-E| - 0.5)^2}{E}.$$

[illegible] 3.3 (2×2 [illegible]). 考虑二元结局的随机实验，数据汇总为：

	成功 $(Y=1)$	失败 $(Y=0)$	合计
治疗 $(Z=1)$	$n_{11}=30$	$n_{10}=20$	$n_1=50$
对照 $(Z=0)$	$n_{01}=20$	$n_{00}=30$	$n_0=50$

1 :

$$E_{11}=\frac{50\times 50}{100}=25,\quad E_{10}=\frac{50\times 50}{100}=25,\quad E_{01}=\frac{50\times 50}{100}=25,\quad E_{00}=25$$

2 χ^2:

$$\chi^2=\frac{(30-25)^2}{25}+\frac{(20-25)^2}{25}+\frac{(20-25)^2}{25}+\frac{(30-25)^2}{25}=4$$

自由度 $(2-1)(2-1)=1$，临界值 $\chi^2_{0.05,1}=3.84$。由于 $4>3.84$， ——治疗与结局存在关联。

R :

```
> tb <- matrix(c(30, 20, 20, 30), nrow = 2, byrow = TRUE)
> chisq.test(tb)

    Pearson's Chi-squared test with Yates' continuity correction

X-squared = 4, df = 1, p-value = 0.0455
```

注意 *Yates* 修正后的 χ^2 从 4 降到 3.24，p 值 $=0.072$——这说明连续性修正在小样本时影响显著。

□

3.4 (3×2). : 研究吸烟状态与肺癌诊断的关联，数据如下：

	患肺癌	未患肺癌	合计
重度吸烟	*60*	*40*	*100*
轻度吸烟	*20*	*30*	*50*
不吸烟	*10*	*40*	*50*

1 :

$$E_{11}=\frac{100\times 90}{200}=45,\quad E_{12}=\frac{100\times 110}{200}=55$$
$$E_{21}=\frac{50\times 90}{200}=22.5,\quad E_{22}=\frac{50\times 110}{200}=27.5$$
$$E_{31}=\frac{50\times 90}{200}=22.5,\quad E_{32}=\frac{50\times 110}{200}=27.5$$

2 χ^2:

$$\chi^2=\frac{(60-45)^2}{45}+\frac{(40-55)^2}{55}+\frac{(20-22.5)^2}{22.5}+\frac{(30-27.5)^2}{27.5}+\frac{(10-22.5)^2}{22.5}+\frac{(40-27.5)^2}{27.5}\approx 20.4$$

自由度 $(3-1)(2-1)=2$，查表 $\chi^2_{0.05,2}=5.99$。由于 $20.4>5.99$， ——吸烟与肺癌存在显著关联。

R :

```
> tb <- matrix(c(60, 40, 20, 30, 10, 40), nrow = 3, byrow = TRUE)
> rownames(tb) <- c("重度吸烟", "轻度吸烟", "不吸烟")
> colnames(tb) <- c("患肺癌", "未患肺癌")
> chisq.test(tb)

    Pearson's Chi-squared test
```

```
X-squared = 20.4, df = 2, p-value = 3.7e-05
```

p 值极小，结论稳健。

□

[illegible] 3.5 (4×3 [illegible]). [illegible]：调查 *500* 人的教育程度与收入水平：

	低收入	中等收入	高收入	合计
初中及以下	*80*	*50*	*20*	*150*
高中	*60*	*80*	*30*	*170*
本科	*30*	*70*	*40*	*140*
研究生	*10*	*30*	*100*	*140*
合计	*180*	*230*	*190*	*500*

R [illegible]：

```
> edu <- c(80, 50, 20, 60, 80, 30, 30, 70, 40, 10, 30, 100)
> tb <- matrix(edu, nrow = 4, byrow = TRUE)
> rownames(tb) <- c("初中及以下", "高中", "本科", "研究生")
> colnames(tb) <- c("低收入", "中等收入", "高收入")
> chisq.test(tb)

    Pearson's Chi-squared test

X-squared = 186.7, df = 6, p-value < 2.2e-16
```

p 值极小，拒绝独立性——教育程度与收入水平存在强关联。

[illegible]：对于 $r \times c$ 表，还可以计算 *Cramér* V 来衡量关联强度：

$$V = \sqrt{\frac{\chi^2}{n \times \min(r-1, c-1)}}. \tag{26}$$

对于本例，$V = \sqrt{186.7/(500 \times 2)} \approx 0.43$，表示中等偏强的关联。

□

7.3.3. 二元结局下的卡方检验：与 *Fisher* 精确检验的对比. [illegible]

[illegible] $Z \perp Y$ [illegible]

U a。[illegible] **Q** [illegible]

	Chi-square test	Fisher exact test
[illegible]	[illegible]	[illegible]
p	[illegible]	[illegible]
[illegible]	[illegible] > 5	[illegible] < 5

Insight [illegible] Fisher [illegible] p [illegible]
[illegible] Chi-square [illegible]

[illegible] Bertrand & Mullainathan, 2004[illegible]
[illegible]
[illegible]

```
resume = read.csv("resume.csv")
Alltable = table(resume$race, resume$call)
Alltable
```

```
#          0    1
# black 2278  157
# white 2200  235
```

[illegible] 2×2 [illegible]

		[illegible] $(Y=1)$	[illegible] $(Y=0)$	[illegible]
[illegible]	$(Z=1)$	157	2278	2435
[illegible]	$(Z=0)$	235	2200	2435

[illegible]

$$E_{11} = \frac{2435 \times 392}{4870} \approx 195.7, \quad E_{12} = \frac{2435 \times 4478}{4870} \approx 2239.3$$

$$E_{21} = \frac{2435 \times 392}{4870} \approx 196.3, \quad E_{22} = \frac{2435 \times 4478}{4870} \approx 2238.7$$

[illegible] 5[illegible]

```
> chisq.test(Alltable)

	Pearson's Chi-squared test with Yates' continuity correction

X-squared = 17.6, df = 1, p-value = 2.7e-05
```

p [illegible]

[illegible] [illegible] $H_0 : \tau = 0$[illegible] $\widehat{\mathrm{RD}} = \hat{p}_1 - \hat{p}_0$ [illegible]

$$\mathrm{var}(\widehat{\mathrm{RD}}) \approx \frac{\hat{p}_1(1-\hat{p}_1)}{n_1} + \frac{\hat{p}_0(1-\hat{p}_0)}{n_0}$$

[illegible] χ^2 [illegible] $(\widehat{\mathrm{RD}})^2 / \mathrm{var}(\widehat{\mathrm{RD}})$ [illegible]

[illegible]

(1) [illegible] < 5[illegible] 2×2 [illegible] Fisher [illegible]

(2) [illegible]

(3) [illegible] `chisq.test(tb, simulate.p.value = TRUE)`

Insight [illegible] ” [illegible] ” [illegible] > 5 [illegible] < 5[illegible] p [illegible] Fisher [illegible]

[illegible] [illegible]

[illegible] Wilcoxon [illegible]

$$W = \sum_{i=1}^{n} Z_i R_i,$$

[illegible] R_i [illegible] Y_i [illegible][12][illegible] H_{0F} [illegible] **Z**[illegible] R_i [illegible]

$$\mathbb{E}(W) = \frac{n_1(n+1)}{2}, \quad \mathrm{var}(W) = \frac{n_1 n_0 (n+1)}{12}.$$

[illegible]

---

[12] [illegible] ??

- [illegible]
- [illegible]
- [illegible]

[illegible] KS [illegible]

[illegible]

$$D = \max_y |\hat{F}_1(y) - \hat{F}_0(y)|,$$

[illegible] $\hat{F}_1(y) = n_1^{-1} \sum_{i=1}^n Z_i \mathbb{I}(Y_i \le y)$ [illegible]

[illegible]

[illegible]

[illegible]

[illegible] t [illegible] FRT [illegible] t [illegible] + [illegible] CLT [illegible]

[illegible]

1986 [illegible] Robert LaLonde [illegible] National Supported Work Demonstration [illegible]

[illegible] R [illegible] FRT [illegible]

```
library(Matching)
data(lalonde)
z <- lalonde$treat
y <- lalonde$re78
```

[illegible]

[illegible]	[illegible]	[illegible]
$\hat{\tau}$	2.84	[illegible] t [illegible]
t	2.67	[illegible]
W	27402.5	Wilcoxon [illegible]
D	0.132	KS [illegible]

[illegible] Monte Carlo [illegible] Monte Carlo [illegible] $R = 10^4$ [illegible]

```
> exact.pv = c(mean(Tauhat >= tauhat),
+              mean(Student >= student),
+              mean(Wilcox >= W),
+              mean(Ks >= D))
> round(exact.pv, 3)
[1] 0.002 0.002 0.006 0.040
```

[illegible] p [illegible] 0.05 [illegible]

[illegible]

FRT [illegible]

[illegible] James Lind (1716-1794) [illegible] 12 [illegible] 6 [illegible]

[illegible] 2×2 [illegible] H_{0F} [illegible]

$$p_{\text{frt}} = \frac{1}{\binom{12}{2}} = \frac{1}{66} = 0.015.$$

[illegible] [illegible] 8 [illegible] 4 [illegible] 4 [illegible] $X = 4$ [illegible] p [illegible]

$$p_{\text{frt}} = \frac{1}{70} = 0.014.$$

[illegible][13] [illegible] H_{0F} [illegible] X [illegible] Hypergeometric$(N = 8, K = 4, n = 4)$ [illegible] "[illegible]" [illegible] $\mathbb{P}(X = 4) = 1/70$ [illegible] p $p_{\text{frt}} = 1/70 \approx 0.014$ [illegible] Fisher [illegible]

(1) [illegible] p [illegible]
(2) [illegible] FRT [illegible]

[illegible]

FRT [illegible] H_{0F} [illegible] sharp null [illegible]

$$H_0(\boldsymbol{\tau}) : Y_i(1) - Y_i(0) = \tau_i \quad \text{[illegible]} \quad i. \tag{27}$$

[illegible] $\boldsymbol{\tau}$ [illegible] $(1 - \alpha)$ [illegible]

[illegible]

[illegible]

(1) [illegible] $\mathbb{P}(\mathbf{Z} = \mathbf{z}) = 1/\binom{n}{n_1}$
(2) [illegible] + [illegible]
(3) [illegible] $H_{0F}: Y_i(1) = Y_i(0)$ [illegible] i
(4) [illegible] $p_{\text{frt}} = M^{-1} \sum_{m=1}^{M} \mathbb{I}\{T(\mathbf{z}^m, \mathbf{Y}) \geq T(\mathbf{Z}, \mathbf{Y})\}$
(5) [illegible] $\hat{\tau}$ [illegible] t [illegible] Wilcoxon W [illegible] KS D
(6) [illegible] FRT [illegible] t [illegible]
(7) [illegible] Lind [illegible] Fisher [illegible]

[illegible] Neymanian [illegible]

[illegible]

[illegible] 3.1. *3.1 —p_{frt} 的精确性 证明在 sharp null 下，p_{frt} 满足：*

$$\mathbb{P}(p_{frt} \leq u) \leq u \quad \textit{for all} \quad 0 \leq u \leq 1.$$

---

[13] [illegible] **??**

3.2. *3.2 —Monte Carlo* 误差 设 p_{frt} 是固定值，$\hat{p}_{frt}$ 是其 *Monte Carlo* 估计（见 { }。证明：

$$\mathbb{E}_{mc}(\hat{p}_{frt}) = p_{frt}, \quad \mathrm{var}_{mc}(\hat{p}_{frt}) \le \frac{1}{4R},$$

其中下标”*mc*”表示由于 *Monte Carlo* 的随机性。

3.3. *3.3* —有限样本有效的 *Monte Carlo* 近似 证明修正的 *Monte Carlo* 近似

$$\tilde{p}_{frt} = \frac{1 + \sum_{r=1}^{R} \mathbb{I}\{T(\mathbf{z}^r, \mathbf{Y}) \ge T(\mathbf{Z}, \mathbf{Y})\}}{1+R}$$

在任意有限的 R 下都是有限样本有效的 p 值。

：可以利用以下两个基本概率结果：

1. (1) 对于两个二项分布 $X_1 \sim Binomial(R, p_1)$ 和 $X_2 \sim Binomial(R, p_2)$，若 $p_1 \ge p_2$，则 $\mathbb{P}(X_1 \le x) \le \mathbb{P}(X_2 \le x)$。
2. (2) 若 $p \sim Uniform(0, 1)$ 且 $X \mid p \sim Binomial(R, p)$，则 X 在 $\{0, 1, \ldots, R\}$ 上均匀分布。

3.4. *3.4* —*Fisher* 精确检验 考虑二元结局的 *CRE*，数据汇总为 2×2 列联表：

	$Y=1$	$Y=0$	合计
$Z=1$	n_{11}	n_{10}	n_1 着
$Z=0$	n_{01}	n_{00}	n_0

在 H_{0F} 下，证明任何检验统计量 $T(n_{11}, n_{10}, n_{01}, n_{00})$ 是 n_{11} 的函数，且 n_{11} 的精确分布是超几何分布。请说明超几何分布的参数。

3.5. *3.5* —女士品茶详细计算 回忆女士品茶实验（见 { }。计算 $\mathbb{P}(X = k)$，其中 X 是女士正确识别 k 杯先加牛奶的茶的个数，$k = 0, 1, 2, 3, 4$。

3.6. *3.6* —协变量调整的 *FRT* 使用 *LaLonde* 数据集（见 { }，添加协变量进行 *FRT* 分析。假设所有潜在结果和协变量都是固定数值。

有两种策略：

1. (1) 运行结局对协变量的线性回归，得到残差（即”伪结局”）。基于残差定义四种检验统计量，进行 *FRT* 并报告 p 值。
2. (2) 定义检验统计量为结局对治疗和协变量线性回归中治疗系数的 *FRT*。

解释为什么上述两种策略的五个 p 值都是有限样本精确的。

3.7. *3.7* —*FRT* 与广义线性模型 使用与 { }相同的数据集，但将结局改为 *re78 > 0* 的二元指示变量。运行结局对治疗和协变量的 *logistic* 回归。治疗系数是否显著？*FRT* p 值是多少？

3.8. *3.8* —代数细节 验证 { 。{

3.9. *3.9* —推荐阅读 { 是一篇倡导在复杂实验中同时报告 p 值及其相应随机化分布的论文。

CHAPTER 4

[illegible]

[illegible]

[illegible] Fisher's randomization test, FRT [illegible] sharp null [illegible] p [illegible]

[illegible] FRT [illegible] null [illegible] FRT [illegible]

[illegible] Fisher [illegible] p [illegible] " [illegible] " [illegible] " [illegible] " [illegible] Fisher [illegible] distribution-free [illegible]

Jerzy Neyman [illegible] 1923 [illegible] p [illegible] Fisherian [illegible] Neymanian [illegible]

4.1. [illegible] : *FRT* 适用于小样本精确推断和假设检验，*Neymanian* 适用于大样本渐近推断和置信区间构建。两者互为补充。

[illegible] 1

Neymanian [illegible] Neyman [illegible] Fisher [illegible]

[illegible] n [illegible] n_1 [illegible] $n_0 = n - n_1$ [illegible] $i = 1, \ldots, n$ [illegible]

- $Y_i(1)$ [illegible] i [illegible]
- $Y_i(0)$ [illegible] i [illegible]
- $\tau_i = Y_i(1) - Y_i(0)$ [illegible] i [illegible]

[illegible] " [illegible] " [illegible]

$$\bar{Y}(1) = \frac{1}{n}\sum_{i=1}^{n} Y_i(1), \quad \bar{Y}(0) = \frac{1}{n}\sum_{i=1}^{n} Y_i(0)$$

[illegible] $n - 1$ [illegible]

$$S^2(1) = \frac{1}{n-1}\sum_{i=1}^{n}\{Y_i(1) - \bar{Y}(1)\}^2, \quad S^2(0) = \frac{1}{n-1}\sum_{i=1}^{n}\{Y_i(0) - \bar{Y}(0)\}^2$$

[illegible]

$$S(1,0) = \frac{1}{n-1}\sum_{i=1}^{n}\{Y_i(1)-\bar{Y}(1)\}\{Y_i(0)-\bar{Y}(0)\}$$

[illegible]

$$\tau = \frac{1}{n}\sum_{i=1}^{n}\tau_i = \bar{Y}(1)-\bar{Y}(0)$$

[illegible]

$$S^2(\tau) = \frac{1}{n-1}\sum_{i=1}^{n}(\tau_i-\tau)^2$$

[illegible]

4.2. [illegible]：$Y(1)$ 和 $Y(0)$ 之间可能有相关性。若治疗组和对照组的个体是匹配的（正相关），估计的方差会减小；若是负相关，方差会增大。

[illegible]

[illegible] Neymanian [illegible]

[illegible] (Lemma 4.1 [illegible])

$$2S(1,0) = S^2(1)+S^2(0)-S^2(\tau)$$

其中 $\tau_i = Y_i(1)-Y_i(0)$ 是个体因果效应。

[illegible] $S^2(\tau)=0$ [illegible] $Y_i(1)$ [illegible] $Y_i(0)$ [illegible] $S(1,0)$ [illegible] $[S^2(1)+S^2(0)]/2$ [illegible] $S^2(\tau)>0$ [illegible] $2S(1,0) < S^2(1)+S^2(0)$[1] [illegible]

[illegible]

[1] [illegible]

$$\hat{\bar{Y}}(1) = \frac{1}{n_1}\sum_{i=1}^{n}Z_iY_i, \quad \hat{\bar{Y}}(0) = \frac{1}{n_0}\sum_{i=1}^{n}(1-Z_i)Y_i$$

[illegible] $Z_i \in \{0,1\}$ [illegible]

$$\hat{S}^2(1) = \frac{1}{n_1-1}\sum_{i=1}^{n}Z_i\{Y_i-\hat{\bar{Y}}(1)\}^2, \quad \hat{S}^2(0) = \frac{1}{n_0-1}\sum_{i=1}^{n}(1-Z_i)\{Y_i-\hat{\bar{Y}}(0)\}^2$$

[illegible] $S(1,0)$ [illegible] $S^2(\tau)$ [illegible] $Y_i(1)$ [illegible] $Y_i(0)$ [illegible] Neyman [illegible]

[illegible] (Neyman (1923) [illegible]) 在完全随机化实验（*CRE*）下，令 $\hat{\tau} = \hat{\bar{Y}}(1)-\hat{\bar{Y}}(0)$。则：

---

[1] [illegible] **??**

(1) [illegible]：$\hat{\tau}$ 对 τ 是无偏的：

$$\mathbb{E}(\hat{\tau}) = \tau$$

(2) [illegible]：$\hat{\tau}$ 的方差为

$$\operatorname{var}(\hat{\tau}) = \frac{S^2(1)}{n_1} + \frac{S^2(0)}{n_0} - \frac{S^2(\tau)}{n} \tag{28}$$

$$= \frac{n_0}{n_1 n}S^2(1) + \frac{n_1}{n_0 n}S^2(0) + \frac{2}{n}S(1,0) \tag{29}$$

(3) [illegible]：方差估计量

$$\hat{V} = \frac{\hat{S}^2(1)}{n_1} + \frac{\hat{S}^2(0)}{n_0}$$

对 $\operatorname{var}(\hat{\tau})$ 是保守的，即

$$\mathbb{E}(\hat{V}) - \operatorname{var}(\hat{\tau}) = \frac{S^2(\tau)}{n} \geq 0$$

等号当且仅当 $\tau_i \equiv \tau$（所有单元个体效应恒定）时成立。

[illegible] $\{Y_i(1), Y_i(0)\}$ [illegible] Z_i [illegible] n_1 [illegible] 1 [illegible] n_0 [illegible] 0 [illegible]

4.3. [illegible]：直觉上：若个体因果效应存在变异（$S^2(\tau) > 0$），治疗组和对照组之间的差异不仅来自随机抽样误差，还来自因果效应本身的真实变异。这种”真实变异”不应被算作估计误差。

Ww[illegible]

[illegible] $S^2(\tau)$

[illegible] $\hat{\tau}$ [illegible]

[illegible] $\{Y_i(1)\}$ [illegible]

- [illegible] $S(1,0) > 0$ [illegible] $\hat{\tau}$
- [illegible] $S(1,0) < 0$ [illegible] $\hat{\tau}$

[illegible] $\hat{\tau}$ [illegible]

[illegible] $S^2(\tau) = 0$ [illegible] $\operatorname{var}(\hat{\tau}) = S^2(1)/n_1 + S^2(0)/n_0$ [illegible] Neymanian [illegible]

[illegible]

[illegible][2]

[illegible][3]

[illegible][4]

---

[2] [illegible] ??

[3] [illegible] ??

[4] [illegible] ??

[illegible] $\hat{V}$ [illegible] HC2 [illegible] $\hat{V}$[6] [illegible] LaLonde [illegible]

```
> tauhat = mean(y[z==1]) - mean(y[z==0])
> vhat   = var(y[z==1])/n1 + var(y[z==0])/n0
> sehat  = sqrt(vhat)
> tauhat
[1] 1794.343
> sehat
[1] 670.9967
```

OLS [illegible]

```
> olsfit = lm(y ~ z)
> summary(olsfit)$coef[2, 1:2]
  Estimate Std. Error
 1794.3431   632.8536
```

[illegible] EHW [illegible] HC2 [illegible]

```
> sqrt(hccm(olsfit, type = "hc2")[2, 2])
[1] 670.9967
```

[illegible]

[illegible] CLT [illegible]

[illegible] Cauchy [illegible]

[illegible]

[illegible] Neymanian [illegible]

(1) [illegible] hat [illegible] $\{\bar{Y}(1), \bar{Y}(0), S^2(1), S^2(0), S(1,0), S^2(\tau)\}$

(2) [illegible] $2S(1,0) = S^2(1) + S^2(0) - S^2(\tau)$

(3) [illegible]

  - $\mathbb{E}(\hat{\tau}) = \tau$ [illegible]
  - $\mathrm{var}(\hat{\tau}) = \frac{S^2(1)}{n_1} + \frac{S^2(0)}{n_0} - \frac{S^2(\tau)}{n}$
  - $\mathbb{E}(\hat{V}) = \mathrm{var}(\hat{\tau}) + \frac{S^2(\tau)}{n}$ [illegible]

(4) [illegible] $\hat{V} = \frac{\hat{S}^2(1)}{n_1} + \frac{\hat{S}^2(0)}{n_0}$

(5) [illegible] $\hat{\tau} \pm z_{1-\alpha/2}\sqrt{\hat{V}}$

(6) [illegible] $\frac{\hat{\tau}-\tau}{\sqrt{\mathrm{var}(\hat{\tau})}} \xrightarrow{d} N(0,1)$

(7) [illegible] $\hat{\beta} = \hat{\tau}$ [illegible] EHW [illegible]

---

[6] [illegible] ??

q q w

ǘ 4.1. { {引理的证明 证明 *Lemma* { :{ $2S(1,0) = S^2(1) + S^2(0) - S^2(\tau)$。

ǘ 4.2. { {*Neyman (1923)* 定理的另一种证明 在 *CRE* 下，计算：

$$\operatorname{var}\{\hat{\bar{Y}}(1)\}, \quad \operatorname{var}\{\hat{\bar{Y}}(0)\}, \quad \operatorname{cov}\{\hat{\bar{Y}}(1), \hat{\bar{Y}}(0)\}$$

并利用这些公式推导 $\operatorname{var}(\hat{\tau})$。

ǘ 4.3. { {*Neymanian* 推断与 *OLS* 证明 { {(*OLS* 系数等于差值均值)。证明 *EHW* 稳健方差估计量的 *HC2* 变体恰好等于 $\hat{V}$。
þ‖ȯh { ˳{

ǘ 4.4. { {治疗效应的异质性 证明 $S^2(\tau) = 0$ 蕴含 $S^2(1) = S^2(0)$。给出反例说明 $S^2(1) = S^2(0)$ 但 $S^2(\tau) \neq 0$。
证明 $S^2(1) < S^2(0)$ 蕴含 $S(Y(0), \tau) < 0$。给出反例说明 $S^2(1) > S^2(0)$ 但 $S(Y(0), \tau) < 0$。

ǘ 4.5. { {方差公式的改进上界 证明方差公式的改进上界：

$$\operatorname{var}(\hat{\tau}) \leq \frac{1}{n}\left\{\sqrt{\frac{n_0}{n_1}}S(1) + \sqrt{\frac{n_1}{n_0}}S(0)\right\}^2$$

并讨论等号成立的条件。

ǘ 4.6. { {*Neyman (1923)* 的向量版本 将 *Neyman (1923)* 的结果推广到多结局情形。考虑向量结局 $\mathbf{V} \in \mathbb{R}^K$ 的平均因果效应：

$$\tau_{\mathbf{V}} = \frac{1}{n}\sum_{i=1}^{n}\{\mathbf{V}_i(1) - \mathbf{V}_i(0)\}$$

证明 $\hat{\tau}_{\mathbf{V}}$ 对 $\tau_{\mathbf{V}}$ 是无偏的，并求出其协方差矩阵。

ǘ 4.7. { {完全随机化实验中的推断 回忆在 { {中定义的完全随机化实验。在这种设定下，推导平均因果效应的方差公式，并比较其与 *CRE* 中方差的差异。

ǘ 4.8. { {推荐阅读 ┤ḓ : { 重新审视了因果推断中的一些经典问题。{ 是关于因果推断的权威综述。
ä̤ɇ e : *Neyman 1923* 年的原始论文 { 奠定了有限总体推断的基础。当代统计学家将其视为因果推断范式的里程碑。

CHAPTER 5

[illegible]

[illegible]

"Block what you can and randomize what you cannot." — ([illegible] page 103)

[illegible] George Box [illegible] " [illegible]

[illegible] CRE [illegible] chance [illegible]

[illegible] covariate imbalance [illegible] treatment [illegible]

[illegible]

[illegible] n [illegible] $X_i \in \{1, \dots, K\}$ [illegible]

$$n_{[k]} = \#\{i : X_i = k\}, \quad \pi_{[k]} = n_{[k]}/n$$

[illegible] k [illegible] $k = 1, \dots, K$ [illegible] CRE [illegible] n_1 [illegible] n_0 [illegible] $n_{[k]1}$ [illegible] $n_{[k]0}$ [illegible] CRE [illegible][1]

$$\mathbb{E}\left(\frac{n_{[k]1}}{n_1} - \frac{n_{[k]0}}{n_0}\right) = 0, \quad \frac{n_{[k]1}}{n_1} - \frac{n_{[k]0}}{n_0} \neq 0 \quad .$$

[illegible]

[illegible] 5.1 ([illegible] (SRE)). 固定 $n_{[k]1}$ 或 $n_{[k]0}$（$k = 1, \dots, K$），在离散的协变量 X 的每一层内独立进行 *CRE*。

[illegible] blocks [illegible] block randomization [illegible]

$$e_{[k]} = \frac{n_{[k]1}}{n_{[k]}}.$$

[illegible] SRE CRE [illegible]

(1) SRE [illegible] CRE [illegible]

---

[1] [illegible] ??

(2) SRE $e_{[k]}$ [illegible] CRE [illegible]

[illegible]

[illegible] i [illegible] $Y_i(1)$ [illegible] $Y_i(0)$ [illegible] $\tau_i = Y_i(1) - Y_i(0)$ [illegible] k [illegible]

$$\tau_{[k]} = n_{[k]}^{-1} \sum_{X_i=k} \tau_i.$$

[illegible]

$$\tau = n^{-1} \sum_{i=1}^{n} \tau_i = \sum_{k=1}^{K} \pi_{[k]} \tau_{[k]}.$$

[illegible] τ [illegible]

$$\hat{\tau}_{\mathrm{SRE}} = \sum_{k=1}^{K} \pi_{[k]} \hat{\tau}_{[k]},$$

[illegible]

$$\hat{\tau}_{[k]} = n_{[k]1}^{-1} \sum_{i=1}^{n} \mathbb{I}(X_i = k, Z_i = 1) Y_i - n_{[k]0}^{-1} \sum_{i=1}^{n} \mathbb{I}(X_i = k, Z_i = 0) Y_i$$

[illegible] k [illegible] $\tau_{[k]}$ [illegible] k [illegible] CRE [illegible] τ [illegible]

[illegible]

CRE [illegible] SRE [illegible] Fisher [illegible] FRT [illegible] Sharp null [illegible]

$$H_{0F} : Y_i(1) = Y_i(0) \quad i = 1, \ldots, n.$$

FRT [illegible] $T = T(\mathbf{Z}, \mathbf{Y}, \mathbf{X})$ [illegible] $\mathbf{Z}, \mathbf{Y}, \mathbf{X}$ [illegible] SRE [illegible] H_{0F} [illegible] T [illegible] $\mathbf{Z}$ SRE [illegible]

[illegible] FRT [illegible]

(1) [illegible] X [illegible]

(2) [illegible] SRE [illegible]

[illegible]

4.1.1. 分层差值均值估计量. [illegible] FRT [illegible] $\hat{\tau}_{\mathrm{SRE}}$ [illegible]

[illegible] 5.1 ([illegible]). 在 *FRT* 中，我们可以使用分层估计量作为检验统计量：

$$T = \hat{\tau}_{\mathrm{S}} = \sum_{k=1}^{K} \pi_{[k]} \hat{\tau}_{[k]},$$

其中 $\hat{\tau}_{[k]}$ 由 [illegible] 定义。这个统计量直接衡量了跨层的平均因果效应。

4.1.2. 学生化分层估计量. [illegible] FRT [illegible]

$$t_{\mathrm{SRE}} = \frac{\hat{\tau}_{\mathrm{SRE}}}{\sqrt{\hat{V}_{\mathrm{SRE}}}},$$

[illegible]

$$\hat{V}_{\mathrm{SRE}} = \sum_{k=1}^{K} \pi_{[k]}^2 \left(\frac{\hat{S}_{[k]}^2(1)}{n_{[k]1}} + \frac{\hat{S}_{[k]}^2(0)}{n_{[k]0}} \right).$$

[illegible] $\hat{S}_{[k]}^2(1)$ [illegible] $\hat{S}_{[k]}^2(0)$ [illegible] k [illegible][2]

[illegible] 5.2 ([illegible]). 对于 [illegible]中的学生化统计量 t_{SRE}，我们需要计算其方差估计量 $\hat{V}_{SRE}$（由 [illegible]给出）。在 *FRT* 中，我们通过模拟所有可行的 *treatment* 分配来计算 t_{SRE} 的精确分布。

4.1.3. 合并 *Wilcoxon* 秩和统计量. [illegible] k [illegible] Wilcoxon [illegible] $W_{[k]}$[illegible]

$$W_{\mathrm{SRE}} = \sum_{k=1}^{K} c_{[k]} W_{[k]}.$$

[illegible] $c_{[k]} = 1/(n_{[k]1} n_{[k]0})$[illegible] $c_{[k]} = 1/(n_{[k]} + 1)$

[illegible] 5.3 (Van Elteren [illegible]). 在第 k 层内，我们计算 *Wilcoxon* 秩和统计量 $W_{[k]}$（回忆第 [illegible]章的 *Wilcoxon* 秩和检验），然后使用 [illegible]合并各层统计量。*Van Elteren (1960)* 的两种加权方法在不同场景下各有优势。

4.1.4. *Hodges–Lehmann* 对齐秩统计量. Van Elteren (1960) [illegible]

$$\tilde{Y}_i = Y_i - \bar{Y}_{[k]}, \quad [illegible] \quad X_i = k,$$

[illegible] $\bar{Y}_{[k]} = n_{[k]}^{-1} \sum_{X_i = k} Y_i$ [illegible] k [illegible] $(\tilde{Y}_1, \ldots, \tilde{Y}_n)$ [illegible] $(\tilde{R}_1, \ldots, \tilde{R}_n)$[illegible]

$$\tilde{W} = \sum_{i=1}^{n} Z_i \tilde{R}_i.$$

[illegible] 5.4 (Hodges–Lehmann [illegible]). [illegible]的方法首先通过 [illegible]对齐结果，这消除了层间差异的影响。然后对池化结果求秩，并使用 [illegible]作为检验统计量。这种方法在多个小层情况下特别有效。

---

[2] [illegible] ?? [illegible] ?? [illegible] ??

d X * _ 1Ï ¥ ª s ª

µ Ö Û x M X ¥ * _ 1 Ï » k ª = ¤ s Í „ v ¥ † í ” ^ Û ¥ b
a _ 1Ï ƒ t ” 3 ^ % ç ¥ b Œ Â T á Ì ó ç $M = \{n_{[k]1}; n_{[k]0}\}_{k=1}^{K}$ ¥ H q / É › H q w
* ¹ * _ 1 ü M î a _ 1b
” Ð Â T M¥ î µ s û , ¹ , * ¹ [3]

$$S_{*\_1}(w = x_j \mid M) = \frac{S_{*\_1}(w = x_j M)}{S_{*\_1}(M)} = \prod_{k=1}^{K} \frac{1}{\frac{n_{[k]}}{n_{[k]1}}} :$$

' w ó ç MHq/V * _ 1 ¤ ž ¥ H q s ƒ Ð V a _ 1¤ ž ¥ s ƒ M] b
y N ó ç M¥ H q / á Ì V [^ s a _ 1B " s { µ Ö Û x M X ¥ * _ 1b
6 _ hM ¹ H q 6 _ h L 2 v K M B **M** ¹ ª s ª T Q b i @ b i ` i B } + i B Q M

$$\hat{S}_a = \sum_{k=1}^{K} {}_{[k]} \hat{}_{[k]} :$$

™ T Ð $\hat{}_{a\_1}$ M] b
s ª ! 9 ¨ P ¨ X ª s ª s ¨ P ¨ X b ñ Ì ^ P ¨ X ¥ } Z E b

3 XL I Ï ¥ Ù 5

Â … Ê 4 X Ÿ y a _ 1$ Ø , X ‹ ¾ ? ç © . ² T b t f ƒ / L € ô
5 • ù î © ç © x M µ @ ¥ ß © M * ¹ X ¥ Ê 4 V [° ¤ ’ ç b ð f
ƒ / ƒ Õ ß © M , b L € Q 7 ô Z L Ÿ Ê 4 X i i è Â X V [^ ù î u
× ¥ · U M Ð 3 M) b
K ¥ Ê 4 ^ B ñ M 1 Ù 5 b Ø , Â T î µ ª û @ v 9 F K ö 4 Ú 9 r q b
– 7 v ¥ K . À V ? † ® 9 r q b E ù î Ï á Ì 4 ³ ž 9 F K ¥ H = l
m ¾ h b Ü K = 5 Y È @ [¤ r q 9 m b
» õ Ã x M 8 ¹ ÷ $ Â T á Ì µ 5 • ù î V [y ë ó ç ƒ t x M H . ² T
$Y(0)$ ¥ ˜ – ª Ê 4 X ¹ ç © $\hat{Y}(0)$ ¥ Ö Û Ä ñ ' b B î ¹ Â T á Ì À µ ƒ "
¥ 5 • ù î € , X S + ! ¥ Ö Û Ä á Ì V [P ¨ B Õ ÷ Y ¨ ¥ ¦ { ë ¹ × Û Ä
` 2 ` M / Q K B x bB Q M

N X c l ²

' c º s ª Ð ª s ª Û L ö 1 = ¸ Â /

- U RaV _ 1 ¥ î * _ 1 V ? Á 3 , ù ¬ ¥ Í s ¥ Y V s ª V [ö î E ç + Q p ` B i 2 B K # H bM + 2
- U kaV _ 1 ç l \\ Ö Û x M ¥ ò ª = ÿ ë É › * _ 1 % ç ò ª ¥ Í s ¥ ” b
- U j Vr q 9 m ’ x M ? ç © . ² T H a _ 11 * _ 1 ÷ µ r y ¹ s ª h " ª W M s n \\ b
- U 96V _ h „ L 2 v K M B wM û V [w < ž a _ 1 s Y \\ „ \\ Ï) , b
- U 8ªVs ª * _ 1 Ï ó ç s ª ” H q ª s Z E Ð a _ 1M] i i ƒ ^ P ¨ x M ¥ } Z E b

---

[3] £ ü n ¸ c \\

10. 习题

练习 5.1. *CRE 中的协变量平衡* 证明 ??：在 *CRE* 下，

$$\mathbb{E}\left(\frac{n_{[k]1}}{n_1}-\frac{n_{[k]0}}{n_0}\right)=0.$$

练习 5.2. *常数层别倾向得分的结果* 证明 ??：当所有层的层别倾向得分相等 $e_{[k]}=e$（对所有 k）时，

$$\hat{\tau}=\hat{\tau}_{SRE}.$$

练习 5.3. *常数个体因果效应的后果* 假设个体因果效应是常数 $\tau_i=\tau$（对所有 $i=1,\ldots,n$）。考虑以下 τ 的加权估计量类：

$$\hat{\tau}_w=\sum_{k=1}^{K}w_{[k]}\hat{\tau}_{[k]},$$

其中权重 $w_{[k]}$ 对所有 k 非负。

(1) 找出使 $\hat{\tau}_w$ 对 τ 无偏的 $w_{[k]}$ 条件。
(2) 在所有无偏估计量中，找出使 $\hat{\tau}_w$ 方差最小的权重。

练习 5.4. *比较 CRE 和 SRE* 证明 ??：当层样本量较大时，$var_{CRE}(\hat{\tau})$ 与 $var_{SRE}(\hat{\tau}_{SRE})$ 之差约为

$$\sum_{k=1}^{K}\frac{\pi_{[k]}}{n}\left\{\sqrt{\frac{n_0}{n_1}}\{\bar{Y}_{[k]}(1)-\bar{Y}(1)\}+\sqrt{\frac{n_1}{n_0}}\{\bar{Y}_{[k]}(0)-\bar{Y}(0)\}\right\}^2\geq 0.$$

练习 5.5. *从 CRE 到 SRE* 证明 ??：数学上，如果 $\mathbf{n}$ 的所有分量都不为零，那么

$$\Pr_{CRE}(\mathbf{Z}=\mathbf{z}\mid\mathbf{n})=\frac{1}{\prod_{k=1}^{K}\binom{n_{[k]}}{n_{[k]1}}}.$$

即，$\mathbf{Z}$ 在给定 $\mathbf{n}$ 条件下从 *CRE* 得到的条件分布与从 *SRE* 得到的分布相同。

练习 5.6. *更多 FRT 统计量* 在 ?? 中，我们介绍了多种检验统计量。选择另一种统计量（如 *Wilcoxon* 秩和统计量或 *Hodges–Lehmann* 对齐秩统计量），重新进行 *FRT* 分析。

练习 5.7. *Imbens and Rubin (2015) 中的 SRE 的 FRT* *Imbens and Rubin (2015)* 讨论了 *1985-1986* 年在田纳西州进行的 *SRE*（*Student/Teacher Achievement Ratio* 实验）。*kindergarten* 数据如下：

层对应于学校，分析单元是教师或班级。*treatment* 等于 *1* 表示小班（每教师 *13-17* 名学生），等于 *0* 表示普通班（*22-25* 名学生）。*outcome* 是标准化的平均数学成绩。

使用 *FRT* 重新分析 *Project STAR* 数据。使用 $\hat{\tau}_{SRE}$、W_{SRE} 和 $\tilde{W}$ 作为 *FRT* 的检验统计量。比较 p 值。

注：本书使用 Z 表示治疗指示变量，但 *Imbens and Rubin (2015)* 使用 W。

练习 5.8. *多中心试验* *Gould (1998, Table 1)* 报告了以下来自多中心试验的数据：

这是一个以中心为层的 *SRE*。该试验旨在研究非那雄胺（用于治疗良性前列腺增生的药物）的疗效和耐受性。在每个中心的 *29* 个中心内，患者被随机分配到三个组：对照组、非那雄胺 *1mg* 和非那雄胺 *5mg*。

由于没有报告个体水平的数据，我们无法实施 *FRT*。然而，*Neymanian* 推断只需要汇总统计。分别报告非那雄胺 *1mg* 和非那雄胺 *5mg* 与对照组比较的点估计量和方差估计。

练习 5.9. 数据分析重新分析 重新分析第 ?? 章 ?? 节中使用的 *LaLonde* 数据。同时进行 *Fisherian* 和 *Neymanian* 推断。

Mahalanobis 距离的一个优美性质是它对协变量的非奇异线性变换保持不变。

引理 6.1 (引理 6.1 —Mahalanobis 距离的不变性). *式* **??** *中的 M 在变换 $X_i \mapsto b_0 + BX_i$（其中 $b_0 \in \mathbb{R}^K$，$B \in \mathbb{R}^{K\times K}$ 可逆）下保持不变。*

为什么这个性质重要？因为它意味着在使用 Mahalanobis 距离时，我们不必担心协变量的量纲问题——无论是用米还是厘米，用摄氏度还是华氏度，M 的数值都不会改变。

有限总体中心极限定理（Li and Ding, 2017）保证：当 n 足够大时，在 CRE 下 M 近似服从 χ^2_K 分布，渐近期望为 K，方差为 $2K$。因此，在 CRE 下 M 的实现值很可能很大。

重随机化的核心思想是：**拒绝那些协变量不平衡的治疗分配**。

2.3. ReM 的正式定义.

定义 6.2 (重随机化（ReM）). *从 CRE 中抽取治疗向量* $\mathbf{Z}$*，当且仅当* $M \le a$ *时接受，其中* $a > 0$ *是预定阈值。*

阈值 a 的选择类似于 SRE 中层数的选择——这是一个在实践中非平凡的问题。当 $a = \infty$ 时，ReM 退化为普通 CRE；当 $a = 0$ 时，只有极少数可行的治疗分配，实验几乎没有随机性。折中方案是选择一个小但不极小的 a，例如取 χ^2_K 分布的上分位数（如 0.001 分位数）。

几何直观：ReM 通过约束协变量均值之差 $\hat{\tau}_X$ 的大小，在设计阶段”塑造”了实验的几何结构。Morgan 和 Rubin (2012) 的研究表明，这种设计具有优雅的数学理论和良好的几何性质。

3. ReM 下的统计推断

3.1. Fisher 随机化检验在 ReM 下的推广. 一个关键问题是：如何在 ReM 下分析数据？Bruhn and McKenzie (2009) 和 Morgan and Rubin (2012) 论证，只要我们在约束 $M \le a$ 下模拟治疗向量 $\mathbf{Z}$，就仍然可以使用 FRT。这在 sharp null 假设下总是给出有限样本精确 p 值。

在非 null 假设下，推导 ReM 的有限样本性质是一个具有挑战性的问题。Li et al. (2018b) 在正则条件（Condition 6.1）下推导了 $\hat{\tau}$ 的渐近分布。

条件 6.1 (条件 6.1). *当* $n \to \infty$ *时：*

(1) n_1/n *和* n_0/n *具有正极限；*
(2) $\{X_i, Y_i(1), Y_i(0), \tau_i\}$ *的有限总体协方差具有有限极限；*
(3) $\max_{1\le i\le n}\{Y_i(1)-\bar{Y}(1)\}^2/n \to 0$, $\max_{1\le i\le n}\{Y_i(0)-\bar{Y}(0)\}^2/n \to 0$, *且* $\max_{1\le i\le n} X_i^\top X_i/n \to 0$。

3.2. ReM 的渐近分布：几何解释. 下面给出 ReM 的主要渐近结果。为符号简洁，记

$$L_{K,a} \sim D_1 \mid \mathbf{D}^\top \mathbf{D} \le a,$$

其中 $\mathbf{D} = (D_1, \ldots, D_K)$ 服从 K 维标准正态分布；ε 服从一元标准正态分布；$L_{K,a} \perp \varepsilon$。

定理 6.3 (定理 6.1 —ReM 下的渐近分布). *在 ReM（$M \le a$）和条件* **??** *下，*

$$\hat{\tau} - \tau \dot{\sim} \sqrt{var(\hat{\tau})}\left\{\sqrt{R^2}L_{K,a} + \sqrt{1-R^2}\varepsilon\right\},$$

其中

$$var(\hat{\tau}) = \frac{S^2(1)}{n_1} + \frac{S^2(0)}{n_0} - \frac{S^2(\tau)}{n}$$

是 *Neyman (1923)* 的方差公式（见第 [illegible]章），且

$$R^2 = corr^2(\hat{\tau}, \hat{\tau}_X)$$

是 $\hat{\tau}$ 与 $\hat{\tau}_X$ 在 *CRE* 下的平方多重相关系数。

[illegible] Li et al. (2018b) [illegible] $\hat{\tau}$ [illegible] $\hat{\tau}_X$ [illegible] $\hat{\tau}_X$ [illegible] $\cos^2\theta = R^2$ [illegible] θ [illegible] $\hat{\tau}$ [illegible] $\hat{\tau}_X$ [illegible] ReM [illegible] $L_{K,a}$ [illegible] ReM [illegible]

[illegible]

- [illegible] $a = \infty$ [illegible] CRE [illegible] $\hat{\tau} - \tau \dot{\sim} \sqrt{\text{var}(\hat{\tau})}\varepsilon$ [illegible]
- [illegible] $a \to 0$ [illegible] $\hat{\tau} - \tau \dot{\sim} \sqrt{\text{var}(\hat{\tau})(1-R^2)}\varepsilon$

[illegible] a [illegible] ReM [illegible] R^2 [illegible] R^2 [illegible][2]

[illegible] 6.1 [illegible] R^2 [illegible]）在 *CRE* 下，

$$R^2 = corr^2(\hat{\tau}, \hat{\tau}_X) = \frac{n_1^{-1}S^2(1 \mid X) + n_0^{-1}S^2(0 \mid X) - n^{-1}S^2(\tau \mid X)}{n_1^{-1}S^2(1) + n_0^{-1}S^2(0) - n^{-1}S^2(\tau)},$$

其中 $\{S^2(1), S^2(0), S^2(\tau)\}$ 是 $\{Y_i(1), Y_i(0), \tau_i\}$ 的有限总体方差，$\{S^2(1 \mid x), S^2(0 \mid x), S^2(\tau \mid x)\}$ 是它们在 $(1, X_i)$ 上线性投影的相应有限总体方差。在常因果效应假设（$\tau_i = \tau$）下，R^2 退化为 $S^2(0 \mid X)/S^2(0)$，即 $Y_i(0)$ 与 X_i 之间的有限总体平方多重相关系数。

[illegible] R^2 [illegible] ReM [illegible]

[illegible] ReM [illegible] Neyman (1923) [illegible] overcover τ [illegible] Li et al. (2018b) [illegible]$_1$ [illegible]

[illegible]

[illegible]$_2$ [illegible]

[illegible] [illegible] **X** [illegible] H_{0F} [illegible] $T = T(\mathbf{Z}, \mathbf{Y}, \mathbf{X})$ [illegible] p [illegible] FRT [illegible]$_1$ [illegible]

Zhao and Ding (2021a) [illegible]

[illegible] 6.5 ([illegible] FRT [illegible]). 基于拟合统计模型后的残差构造检验统计量。将 Y_i 对 X_i 回归得到残差 $\hat{\varepsilon}_i$，然后将 $\hat{\varepsilon}_i$ 视为伪结果来构造检验统计量。

[illegible] 6.6 ([illegible] FRT [illegible]). 使用回归系数作为检验统计量。将 Y_i 对 (Z_i, X_i) 回归，取 Z_i 的系数作为检验统计量。

[illegible] Y_i [illegible] X_i [illegible] Z_i [illegible] H_{0F} [illegible] H_{0F} [illegible] Y_i [illegible] (Z_i, X_i) [illegible] Z_i [illegible] Z_i [illegible] H_{0F} [illegible] ”[illegible]” [illegible] logistic [illegible]$_2$ [illegible] H_{0F} [illegible]

---

[2] [illegible] ??

[illegible]

[illegible] Li et al. (2018b) [illegible] ReM [illegible] Lin (2013) [illegible] a [illegible] ReM [illegible] $\hat{\tau}$ [illegible] CRE [illegible] $\hat{\tau}_{\mathrm{L}}$ [illegible] ReM [illegible] Lin (2013) [illegible] a [illegible] ReM [illegible]

[illegible] C_i [illegible] K [illegible] $K-1$ [illegible]

$$X_i = \left(\mathbb{I}(C_i = 1) - \pi_{[1]}, \ldots, \mathbb{I}(C_i = K-1) - \pi_{[K-1]}\right),$$

[illegible] $\pi_{[k]}$ [illegible] $C_i = k$ [illegible] Lin (2013) [illegible]

[illegible] 6.3 [illegible]）基于 X_i 的 $\hat{\tau}_{\mathrm{L}}$ 在数值上与基于 C_i 的后分层估计量 $\hat{\tau}_{\mathrm{PS}}$（见第 [illegible] 章）相同。

[illegible]

[illegible] X [illegible] Y [illegible] X [illegible] Y [illegible] X [illegible] X [illegible] $\beta_1 = \beta_0 = 1$ [illegible]

$$\hat{\tau}(1,1) = n_1^{-1}\sum_{i=1}^{n} Z_i(Y_i - X_i) - n_0^{-1}\sum_{i=1}^{n}(1 - Z_i)(Y_i - X_i).$$

[illegible] $g_i = Y_i - X_i$ [illegible] $\hat{\tau}(1,1)$ [illegible] $\beta_1 = \beta_0 = 1$ [illegible] Lin (2013) [illegible] β [illegible] $\hat{\tau}(1,1)$ [illegible] Lin (2013) [illegible] Lin (2013) [illegible] $\hat{\tau}(1,1)$ [illegible]

[illegible]

[illegible] C [illegible] X [illegible] Lin (2013) [illegible] $\hat{\tau}_{\mathrm{L},[k]}$ [illegible]

$$\hat{\tau}_{\mathrm{L,S}} = \sum_{k=1}^{K} \pi_{[k]}\hat{\tau}_{\mathrm{L},[k]}.$$

[illegible]

$$\hat{V}_{\mathrm{L,S}} = \sum_{k=1}^{K} \pi_{[k]}^2 \hat{V}_{\mathrm{EHW},[k]},$$

[illegible] $\hat{V}_{\mathrm{EHW},[k]}$ [illegible] k [illegible] OLS [illegible] EHW [illegible]

[illegible]

Li and Ding (2020) [illegible] Lin (2013) [illegible] EHW [illegible]

[illegible] Neyman (1923) [illegible] Li and Ding (2020) [illegible]

TABLE 1. [illegible]

	[illegible]	
[illegible]	$\hat{\tau}$ (Neyman, 1923)	$\hat{\tau}_{\mathrm{L}}$ (Lin, 2013)
CRE	[illegible]	[illegible]
ReM	$\hat{\tau}$ (Li et al., 2018b)	$\hat{\tau}_{\mathrm{L}}$ (Li and Ding, 2020)

[illegible]

- [illegible] 1[illegible] CRE [illegible] $\hat{\tau}_{\mathrm{L}}$ [illegible] $\hat{\tau}$ [illegible]
- [illegible] 2[illegible] ReM [illegible] ReM [illegible] $\hat{\tau}$ [illegible] CRE [illegible] $\hat{\tau}$ [illegible]
- [illegible] 3 [illegible] 4[illegible] ReM CRE [illegible]

R [illegible]

Angrist et al. (2009) [illegible] 8 [illegible] GPA [illegible] GPA

[illegible] 6.1 ([illegible]). 我们使用 *Angrist et al. (2009)* 数据的一个子集，评估四种设计与分析策略的表现。

代码实现（省略数据读取和缺失值填补）：

```
# 未调整估计量
fit_unadj = lm(y ~ z)
ace_unadj = coef(fit_unadj)[2]
se_unadj = sqrt(hccm(fit_unadj, type = "hc2"))[2, 2]

# 回归调整
x = scale(x)
fit_adj = lm(y ~ z*x)
ace_adj = coef(fit_adj)[2]
se_adj = sqrt(hccm(fit_adj, type = "hc2"))[2, 2]
```

模拟结果（*2000* 次蒙特卡洛复制，*ReM* 阈值 $a = 0.05$）：

估计量	估计值	标准误	t 统计量	p 值
Neyman	*0.054*	*0.076*	*0.719*	*0.472*
Lin	*0.075*	*0.072*	*1.036*	*0.300*

调整后的估计量有更小的标准误，尽管它给出了与未调整估计量相同的（不显著）结论。

[illegible] ReM [illegible] 0.1 [illegible] 0.25 [illegible]

[illegible] ReM [illegible]

[illegible]

[illegible] ReM[illegible] Mahalanobis [illegible] **Z** [illegible] **X** [illegible] $X_i = (x_{i1}, \dots, x_{iK})^\top$ [illegible] $|\hat{\tau}_{x_k}/\sqrt{n/(n_1 n_0) S^2_{x_k}}| \le a$ [illegible] $k = 1, \dots, K$ [illegible] **Z** [illegible] Zhao and Ding (2021b) [illegible]

[illegible] Lin (2013) [illegible] logistic [illegible] logistic [illegible] logistic [illegible] Lin (2013) [illegible] CRE [illegible] Guo and Basse (2023) [illegible] Lin (2013) [illegible]

[illegible] Bloniarz et al. (2016) [illegible] LASSO [illegible] OLS [illegible] Lei and Ding (2021) [illegible] Lin (2013) [illegible] Wager et al. (2016) [illegible] ReM [illegible]

[illegible]

[illegible] 6.1. *FRT under ReM* 描述在 *ReM* 下的 *FRT*。

[illegible] 6.2. *Mahalanobis* 距离的不变性 证明引理 { 。{

[illegible] 6.3. 重随机化下差值均值估计量的无偏性 假设从 *CRE* 中抽取 $\mathbf{Z} = (Z_1, \dots, Z_n)$，当且仅当 $\phi(\mathbf{Z}, \mathbf{X}) = 1$ 时接受，其中 ϕ 是预定的平衡准则。证明：如果 $n_1 = n_0$ 且 $\phi(\mathbf{Z}, \mathbf{X}) = \phi(\mathbf{1}_n - \mathbf{Z}, \mathbf{X})$，则 $\hat{\tau}$ 对 τ 无偏。验证当 $n_1 = n_0$ 时，使用 *Mahalanobis* 距离的重随机化满足此条件。给出当这两个条件不满足时 $\hat{\tau}$ 对 τ 有偏的反例。

[illegible] 6.4. R^2 的等价形式 证明命题 { 。{

[illegible] 6.5. 协变量调整 *FRT* 在协变量调整 *FRT* 中，分别在伪结果策略和模型输出策略下推导 p 值的计算方法。

[illegible] 6.6. *Lin* 估计量的偏 使用蒙特卡洛模拟研究 *Lin* 估计量在小样本下的有限样本偏。将样本量从 $n = 20$ 变化到 $n = 500$，重复 *1000* 次，报告平均偏和均方误差。

[illegible] 6.7. *Lin* 估计量的 *OLS* 表示 证明命题 { {(这是纯线性代数事实)。

[illegible] 6.8. 预测估计量的等价性 证明 $\hat{\tau}_{pred} = \hat{\tau}_{\mathrm{L}}$。

[illegible] 6.9. 投影估计量的等价性 证明 $\hat{\tau}_{proj} = \hat{\tau}_{\mathrm{L}}$。

[illegible] 6.10. 协变量不平衡调整形式 证明 $\hat{\tau}(\beta_1, \beta_0) = \hat{\tau} - \gamma^\top \hat{\tau}_X$ 和 $\hat{\tau}_{\mathrm{L}} = \hat{\tau} - \hat{\gamma}^\top \hat{\tau}_X$。

ǘ 6.11. 差分估计量的方差 推导 $\hat{\tau}(1,1)$ 的保守方差估计量 $\hat{V}(1,1)$。

ǘ 6.12. 回归调整与后分层的等价性 证明命题 { 。{

ǘ 6.13. 协变量不平衡与效率 模拟研究当协变量与结果的相关性从 0.1 变化到 0.9 时，回归调整的效率增益如何变化。

ǘ 6.14. 高维协变量的 *LASSO* 调整 使用 *LASSO* 代替 *OLS* 进行回归调整，比较两者在协变量稀疏时的表现。

ǘ 6.15. 缺失数据的填补 在 { {的模拟中，评估不同缺失值填补策略对估计量的影响。

*> Sh1_ d

¥ Û L

RX„ý Vsªž¥ ¥ Á p

» \\c¥sªÛ L a_1Ï áÌ|†í¿xM sª Äª=ÿëÉ› ›†Û L b’ÄªºµBñÍ †í„Bñ v†íH áÌ¤ž a_1¥K ™Tii ¥ Û L J i+?2/@S B`b 1tT2`BK2MiH J S A1 Ä B ª9ë¹B ñ T Bb

JS1A»º^ a_1¥Bñ+è Œñšµÿ+ 9„w ¦{ OÐ4©ŸùîÏ ¥Ä¥ZEáMM1 | » \\cW%),byN '€| ÿëîcb

,•Ù5 ’xM ? ç©. ²TH |M»¥†í¥ ª Û s¥Í ?Ž ÉB„4Ú 9Ú $

kX JS1¥çlÐ:|

InBñ c 2n ñ†í¥L bÂTáÌšµ? ç©²T¥xM V[ô xM ¥M»Ÿ|†í¥ b ¿†S xM áÌV[ô ¾xM †í†½ –ª¥ M#¥†íb ¿ »xM V[çl†í-W¥î Ö –ªP¨ ¢ØE K ªd=sÄ¥ØEÉ›¥ b[R]

: (i;j) VU» i Ï¥» j ñ†í Ï $i = 1;\dots;n$ $j = 1;2$b†í (i;j) ¥. ² T¹ $Y_{ij}(1)$ ¤sÍ „ $Y_{ij}(0)$ ¤s vb ÄB = Û s¥Bñ†í¤sÍ 6Bñ¤s vbçl

UjjV
$$Z_i = \begin{cases} 1; & \text{Ï»Bñ†í¤sÍ}; \\ 0; & \text{Ï»=ñ†í¤sÍ}: \end{cases}$$

çl dX P¥ Ŵ L UJS1XáVÌµ $(Z_i)_{i=1}^n$ AA·"2`MQ(1/2)H:HB

' Í ·UM $(Z_1;\dots;Z_n)$ ÿë]sƒ ›V•”¹ 1/2 ¥umæsƒb

» i =¥4©²T¹

$$Y_{i1} = Z_i Y_{i1}(1) + (1 \quad Z_i) Y_{i1}(0); \quad Y_{i2} = Z_i Y_{i2}(0) + (1 \quad Z_i) Y_{i2}(1):$$

yN 4©” ¹ $(Z_i; Y_{i1}; Y_{i2})_{i=1}^n$ b

jX 6Bb?Û2`Ä_

Ð-•L B" áÌSÔV[P¨ 6Bb?Û2`Ä_ 6_hŸ_ UÌ,L!

Uj9V
$$H_{0F} : Y_{ij}(1) = Y_{ij}(0) \quad \text{îµ} \quad i = 1;\dots;n \ \text{„} \ j = 1;2:$$

6_hÏ áÌ³1V \\ E $(Z_1;\dots;Z_n)$ ¥sƒb[/^ ¿¥ =Í FÐ vF²T-µ¥+ÕÜÅ_ d9 b

---

[R] ¥ ØE¥W%=¸n \b

3.1. 配对内差异. 定义配对 i 内的**配对差异**:

$$\hat{\tau}_i = \text{治疗组结果} - \text{对照组结果} = (2Z_i - 1)(Y_{i1} - Y_{i2}) = S_i(Y_{i1} - Y_{i2}), \tag{35}$$

其中 $S_i = 2Z_i - 1$ 是独立同分布的随机符号，均值为 0，方差为 1。由于 $\hat{\tau}_i = 0$ 的配对对随机化分布没有贡献，我们在 FRT 讨论中剔除这些配对。

配对差异的均值为

$$\hat{\tau} = n^{-1} \sum_{i=1}^{n} \hat{\tau}_i. \tag{36}$$

3.2. 配对 t 统计量.

例 7.1 (配对 t 统计量). 在 H_{0F} 下，$\hat{\tau}_i$ 是固定的，而 S_i 是随机的。因此

$$\mathbb{E}(\hat{\tau}) = 0, \quad var(\hat{\tau}) = n^{-2} \sum_{i=1}^{n} var(S_i)(Y_{i1} - Y_{i2})^2 = n^{-2} \sum_{i=1}^{n} \hat{\tau}_i^2.$$

基于独立随机变量之和的中心极限定理，我们有正态近似：[2]

$$\frac{\hat{\tau}}{\sqrt{n^{-2}\sum_{i=1}^{n} \hat{\tau}_i^2}} \to \mathrm{N}(0,1) \tag{37}$$

依分布收敛。许多标准教科书建议在 MPE 中使用以下配对 t 统计量:

$$t_{pair} = \frac{\hat{\tau}}{\sqrt{\{n(n-1)\}^{-1}\sum_{i=1}^{n}(\hat{\tau}_i - \hat{\tau})^2}}. \tag{38}$$

在 H_{0F} 下，当 n 较大且 $\hat{\tau}$ 较小时，t_{pair} 与 **??** 几乎相同。

在经典统计学中，使用 t_{pair} 的动机基于不同框架。当 $\hat{\tau}_i \overset{IID}{\sim} \mathrm{N}(0,\sigma^2)$ 时，可以证明 $t_{pair} \sim t(n-1)$，即 t_{pair} 的精确分布是自由度为 $n-1$ 的 t 分布，在大 n 下接近 $\mathrm{N}(0,1)$。R 函数 `t.test` 可以实现此检验。**??** 的讨论给出了经典配对 t 检验的另一种证明，无需假设数据的正态性。

3.3. Wilcoxon 符号秩统计量.

例 7.2 (Wilcoxon 符号秩统计量). 基于 $(|\hat{\tau}_1|, \ldots, |\hat{\tau}_n|)$ 的秩 $(R_1, \ldots, R_n)$，我们可以定义检验统计量

$$W = \sum_{i=1}^{n} \mathbb{I}(\hat{\tau}_i > 0) R_i. \tag{39}$$

在 H_{0F} 下，$|\hat{\tau}_i|$ 是固定的，因此 R_i 也是固定的。由此可得

$$\mathbb{E}(W) = \frac{1}{2}\sum_{i=1}^{n} R_i = \frac{1}{2}\sum_{i=1}^{n} i = \frac{n(n+1)}{4},$$

$$var(W) = \frac{1}{4}\sum_{i=1}^{n} R_i^2 = \frac{1}{4}\sum_{i=1}^{n} i^2 = \frac{n(n+1)(2n+1)}{24}.$$

---

[2] 推导见附录 **??**

独立随机变量之和的中心极限定理保证以下正态近似：[3]

$$\frac{W - n(n+1)/4}{\sqrt{n(n+1)(2n+1)/24}} \to \mathrm{N}(0,1) \tag{40}$$

依分布收敛。R 函数 *wilcox.test* 可以实现精确和渐近检验。

∼wXw dpgjpZp p ǎ 7、jwa l p”

ǎ 7.3 (Kolmogorov–Smirnov Ω). 在 H_{0F} 下，绝对值 $(|\hat\tau_1|, \ldots, |\hat\tau_n|)$ 是固定的，但符号是随机的。因此 $(\hat\tau_1, \ldots, \hat\tau_n)$ 和 $-(\hat\tau_1, \ldots, \hat\tau_n)$ 应具有相同分布。

设

$$\hat F(t) = n^{-1} \sum_{i=1}^{n} \mathbb{I}(\hat\tau_i \le t)$$

是 $(\hat\tau_1, \ldots, \hat\tau_n)$ 的经验分布函数，

$$1 - \hat F(-t-) = n^{-1} \sum_{i=1}^{n} \mathbb{I}(-\hat\tau_i \le t)$$

是 $-(\hat\tau_1, \ldots, \hat\tau_n)$ 的经验分布函数，其中 $\hat F(-t-)$ 是 $\hat F(\cdot)$ 在 $-t$ 处的左极限。*Kolmogorov–Smirnov* 型统计量定义为

$$D = \max_t |\hat F(t) + \hat F(-t-) - 1|. \tag{41}$$

{ 提出了这个检验统计量并推导了其精确和渐近分布。标准软件包中未实现此统计量，但我们可以基于 *FRT* 模拟其精确分布并计算 p 值。

∼wWw w

ǎ 7.4 (Åῷ Ω). 符号统计量仅使用配对内差异的符号：

$$\Delta = \sum_{i=1}^{n} \mathbb{I}(\hat\tau_i > 0). \tag{42}$$

在 H_{0F} 下，

$$\mathbb{I}(\hat\tau_i > 0) \overset{IID}{\sim} Bernoulli(1/2),$$

因此 $\Delta \sim Binomial(n, 1/2)$。这给出了一个基于二项分布的精确检验，$R$ 函数 *binom.test* 可以实现。利用中心极限定理，我们还可以基于以下正态近似进行检验：[4]

$$\frac{\Delta - n/2}{\sqrt{n/4}} \to \mathrm{N}(0,1) \tag{43}$$

依分布收敛。

---

[3] þ‖Òh ??

[4] þ‖Òh ??

[illegible] MPE [illegible] 1[5] [illegible] 2×2 [illegible]

	[illegible] =1	[illegible] =0
[illegible] =1	m_{11}	m_{10}
[illegible] =0	m_{01}	m_{00}

[illegible] 7.5 (McNemar [illegible]). 在 H_{0F} 下，一致对的数量 m_{11} 和 m_{00} 是固定的，$m_{10}+m_{01}$ 也是固定的。因此唯一的随机分量是 m_{10}，其分布为

$$m_{10} \sim Binomial(m_{10}+m_{01}, 1/2).$$

这给出了一个基于二项分布的精确检验。*R* 函数 *mcnemar.test* 给出基于二项分布正态近似的渐近检验：

$$\frac{m_{10}-(m_{10}+m_{01})/2}{\sqrt{(m_{10}+m_{01})/4}} = \frac{m_{10}-m_{01}}{\sqrt{m_{10}+m_{01}}} \rightarrow \mathrm{N}(0,1) \tag{44}$$

依分布收敛。精确 *FRT* 和渐近检验都不依赖于 m_{11} 或 m_{00}，只有不一致对的数量在这些检验中起作用。

[illegible]

[illegible] i [illegible]

$$\tau_i = \frac{1}{2}\{Y_{i1}(1)+Y_{i2}(1)-Y_{i1}(0)-Y_{i2}(0)\}, \tag{45}$$

[illegible]

$$\tau = n^{-1}\sum_{i=1}^{n}\tau_i = (2n)^{-1}\sum_{i=1}^{n}\sum_{j=1}^{2}\{Y_{ij}(1)-Y_{ij}(0)\}. \tag{46}$$

[illegible] $\hat{\tau}_i$ [illegible] τ_i [illegible] $\hat{\tau}$ [illegible] τ [illegible][5] [illegible] MPE [illegible] SRE [illegible] $\hat{\tau}$ [illegible] i [illegible] $\hat{\tau}_i$ [illegible] MPE [illegible] $\hat{\tau}$ [illegible] MPE [illegible] $\hat{\tau}_i$ [illegible] μ [illegible] σ^2 [illegible] $\hat{\tau} = n^{-1}\sum_{i=1}^{n}\hat{\tau}_i$ [illegible] σ^2/n σ^2 [illegible] $(n-1)^{-1}\sum_{i=1}^{n}(\hat{\tau}_i-\hat{\tau})^2$ [illegible] $\mathrm{var}(\hat{\tau})$ [illegible]

$$\hat{V} = \{n(n-1)\}^{-1}\sum_{i=1}^{n}(\hat{\tau}_i-\hat{\tau})^2. \tag{47}$$

[illegible] MPE [illegible] $\hat{V}$ [illegible] $\hat{\tau}_i$ [illegible] $\hat{\tau}$ [illegible] MPE [illegible]

[illegible] **(**[illegible]**)** 在 *MPE* 下，由 [illegible] 定义的 $\hat{V}$ 是 $\hat{\tau}$ 真实方差的一个保守估计：

$$\mathbb{E}(\hat{V}) - var(\hat{\tau}) = \{n(n-1)\}^{-1}\sum_{i=1}^{n}(\tau_i-\tau)^2 \geq 0. \tag{48}$$

---

[5] [illegible] **??**

当 τ_i 在各配对间恒定时，$\mathbb{E}(\hat{V}) = var(\hat{\tau})$。

[illegible] MPE [illegible] $\hat{V}$ [illegible] $\hat{V}$ [illegible] $\hat{\tau}_i$ [illegible] $\mathrm{var}(\hat{\tau})$ [illegible] $\hat{\tau}_i$ [illegible][6]

[illegible] Neymanian [illegible]

$$\frac{\hat{\tau} - \tau}{\sqrt{\mathrm{var}(\hat{\tau})}} \to \mathrm{N}(0,1) \tag{49}$$

$n \to \infty$ [illegible] Wald [illegible]

$$\hat{\tau} \pm z_{1-\alpha/2}\sqrt{\hat{V}} \tag{50}$$

[illegible] $1-\alpha$ [illegible] τ

[illegible] $\hat{\tau}$ [illegible] $\hat{V}$ [illegible] OLS [illegible]

[illegible] **w**(OLS [illegible])**w** $\hat{\tau}$ 和 $\hat{V}$ 等于将向量 $(\hat{\tau}_1, \ldots, \hat{\tau}_n)^\top$ 仅对截距进行 *OLS* 拟合的系数和方差估计量。

$\hat{\tau}$ [illegible] $\hat{V}$ [illegible] OLS [illegible] R [illegible] `lm(tau_hat ~ 1)` [illegible][7]

Ww[illegible]

[illegible] $X_{i1} \neq X_{i2}$ [illegible][8]

[illegible] (i,j) [illegible] X_{ij} [illegible] MPE [illegible]

Ww q w [illegible] **w** FRT [illegible] OLS [illegible] $\hat{\varepsilon}_{ij}$ [illegible] $\hat{\varepsilon}_{ij}$ [illegible] MPE [illegible]

Ww [illegible] **w** **w** [illegible] τ [illegible] $\hat{\tau}_{X,i}$ [illegible] $\hat{\tau}_X$ [illegible]

$$\mathbb{E}(\hat{\tau}_{X,i}) = 0, \quad \mathbb{E}(\hat{\tau}_X) = 0,$$

[illegible]

$$\mathrm{cov}(\hat{\tau}_X) = n^{-2}\sum_{i=1}^{n} \hat{\tau}_{X,i}\hat{\tau}_{X,i}^\top.$$

[illegible] MPE [illegible] $\hat{\tau}_{X,i}$ [illegible] $\mathrm{cov}(\hat{\tau}_X)$ [illegible] $(Z_1, \ldots, Z_n)$ [illegible] $\hat{\tau}_X$ [illegible]

[illegible] γ [illegible]

$$\hat{\boldsymbol{\tau}}(\gamma) = \hat{\boldsymbol{\tau}} - \boldsymbol{\gamma}^\top \hat{\boldsymbol{\tau}}_X. \tag{51}$$

[illegible] γ [illegible] γ [illegible] $\hat{\tau}(\gamma)$ [illegible]

---

[6] [illegible] **??**

[7] [illegible] **??**

[8] [illegible] **??**

[illegible] $cov(\hat{\tau}_X, \hat{\tau})$ 的一个无偏估计为

$$\hat{\theta} = \{n(n-1)\}^{-1} \sum_{i=1}^{n} (\hat{\tau}_{X,i} - \hat{\tau}_X)(\hat{\tau}_i - \hat{\tau}). \tag{52}$$

[illegible]

$$\hat{\gamma} = \left(n^{-2} \sum_{i=1}^{n} \hat{\tau}_{X,i} \hat{\tau}_{X,i}^{\top} \right)^{-1} \hat{\theta} \approx \left(\sum_{i=1}^{n} (\hat{\tau}_{X,i} - \hat{\tau}_X)(\hat{\tau}_{X,i} - \hat{\tau}_X)^{\top} \right)^{-1} \sum_{i=1}^{n} (\hat{\tau}_{X,i} - \hat{\tau}_X)(\hat{\tau}_i - \hat{\tau}),$$

[illegible] $\hat{\tau}_i$ [illegible] $\hat{\tau}_{X,i}$ [illegible] OLS [illegible] $\hat{\tau}_{X,i}$ [illegible]

$$\hat{\tau}_{\text{adj}} = \hat{\tau}(\hat{\gamma}) = \hat{\tau} - \hat{\gamma}^{\top} \hat{\tau}_X. \tag{53}$$

OLS [illegible] $\hat{\tau}_{\text{adj}}$ [illegible] $\hat{\tau}_i$ [illegible] 1 [illegible] $\hat{\tau}_{X,i}$ [illegible] OLS [illegible] $\hat{\tau}_{\text{adj}}$ [illegible]

$$\hat{V}_{\text{adj}} = \hat{V} + \hat{\gamma}^{\top} \text{cov}(\hat{\tau}_X) \hat{\gamma} - 2\hat{\gamma}^{\top} \hat{\theta} = \hat{V} - \hat{\theta}^{\top} \text{cov}(\hat{\tau}_X)^{-1} \hat{\theta}.$$

[illegible] OLS [illegible] 在 *MPE* 下，协变量调整估计量 $\hat{\tau}_{adj}$ 和相关方差估计量 $\hat{V}_{adj}$ 可以方便地近似为将 $\hat{\tau}_i$ 的向量对 1 的向量和 $\hat{\tau}_{X,i}$ 的矩阵进行 *OLS* 拟合的截距和相关方差估计量。

[illegible] EHW [illegible] Eicker–Huber–White [illegible] MPE [illegible] CRE [illegible] Lin (2013) [illegible]

[illegible]

[illegible] 15 [illegible] R [illegible] HistData [illegible] cross [illegible] self [illegible] diff [illegible] MPE [illegible] $2^{15} = 32768$ [illegible] $\hat{\tau}$ [illegible] p [illegible]

```
> difference = ZeaMays$diff
> n.pairs = length(difference)
> abs.diff = abs(difference)
> t_obs = mean(difference)
> t_ran = sapply(1:2^15,
+ function(x) {
+ sum(MP_enumerate(x, 15)*abs.diff)
+ })/n.pairs
> pvalue = mean(t_ran >= t_obs)
> pvalue
[1] 0.02633667
```

[illegible] p [illegible] 0.026[illegible] 5% [illegible] H_{0F} [illegible] $\hat{\tau}$ [illegible]

Figure 1. Darwin [illegible] $\hat{\tau}$ [illegible]

[illegible]

[illegible] Workshop [illegible] 10 [illegible] 8 [illegible]

[illegible]	[illegible]			[illegible]		
	[illegible] 1	[illegible] 2	[illegible]	[illegible] 1	[illegible] 2	[illegible]
1	24	25	-1	51	56	-5
2	42	40	2	64	65	-1
3	52	52	0	75	72	3
4	71	70	1	79	77	2
5	29	27	2	54	49	5
6	50	50	0	75	78	-3
7	69	69	0	91	92	-1
8	36	34	2	64	55	9

1 Neymanian [illegible][9]

[illegible]

[illegible] MPE [illegible]

(1) [illegible] MPE [illegible] SRE [illegible]

(2) [illegible] $Z_i \overset{\text{IID}}{\sim} \text{Bernoulli}(1/2)$

(3) [illegible] t [illegible] Wilcoxon [illegible] Kolmogorov–Smirnov [illegible]

(4) [illegible] $\hat{V}$ [illegible] OLS [illegible]

(5) [illegible] $\hat{\tau}_{\text{adj}}$ [illegible] $\hat{V}_{\text{adj}}$ [illegible] OLS [illegible]

[illegible]

[illegible] 7.1. *MPE* 的方差 证明在 *MPE* 下，$\hat{\tau}$ 的方差为

$$var(\hat{\tau}) = n^{-2} \sum_{i=1}^{n} var(\hat{\tau}_i).$$

[illegible] 7.2. 配对 t 统计量的渐近分布 证明 [illegible] 在 H_{0F} 下，

$$\frac{\hat{\tau}}{\sqrt{n^{-2} \sum_{i=1}^{n} \hat{\tau}_i^2}} \to \text{N}(0, 1)$$

依分布收敛。

[illegible] 7.3. *Wilcoxon* 符号秩统计量的矩 证明 [illegible] $\mathbb{E}(W)$ 和 $var(W)$ 的公式。

[illegible] 7.4. *McNemar* 统计量的精确分布 证明在 H_{0F} 下，$m_{10} \sim Binomial(m_{10}+m_{01}, 1/2)$。

---

[9] [illegible] **??**

ů 7.5. 方差估计量的偏 证明 { :{ 在 *MPE* 下，

$$\mathbb{E}(\hat{V}) - var(\hat{\tau}) = \{n(n-1)\}^{-1} \sum_{i=1}^{n} (\tau_i - \tau)^2 \geq 0.$$

ů 7.6. *OLS* 解释 证明 { :{ $\hat{\tau}$ 和 $\hat{V}$ 等于将 $(\hat{\tau}_1, \ldots, \hat{\tau}_n)^\top$ 仅对截距进行 *OLS* 拟合的系数和方差估计量。

ů 7.7. 协方差估计量 证明 { :{ $cov(\hat{\tau}_X, \hat{\tau})$ 的无偏估计为

$$\hat{\theta} = \{n(n-1)\}^{-1} \sum_{i=1}^{n} (\hat{\tau}_{X,i} - \hat{\tau}_X)(\hat{\tau}_i - \hat{\tau}).$$

ů 7.8. 调整估计量的 *OLS* 解释 证明 { :{ 协变量调整估计量 $\hat{\tau}_{adj}$ 可以近似等于将 $\hat{\tau}_i$ 对 1 和 $\hat{\tau}_{X,i}$（带截距）进行 *OLS* 拟合的截距。

ů 7.9. *Darwin* 数据的 *Neymanian* 分析 使用 *Neymanian* 方法重新分析 *Darwin* 数据。报告点估计、方差估计和 *95%* 置信区间。

ů 7.10. 儿童电视 *Workshop* 实验的 *FRT* 分析 使用 *FRT* 重新分析儿童电视 *Workshop* 实验数据，使用配对 t 统计量作为检验统计量。

CHAPTER 8

[illegible]

[illegible]

[illegible] Fisher [illegible] FRT [illegible] sharp null [illegible] $H_{0F}: Y_i(1) = Y_i(0)$ [illegible] FRT [illegible] p [illegible]

$$\mathbb{P}(p_{\text{frt}} \le u) \le u \quad \text{for all} \quad 0 \le u \le 1.$$

[illegible] FRT [illegible] H_{0F} [illegible] $H_{0N}: \tau = 0$ [illegible] $\hat{\tau}$ [illegible] FRT [illegible] H_{0N} [illegible] anti-conservative [illegible] dual guarantees [illegible] $t = \hat{\tau}/\sqrt{\hat{V}}$ [illegible] H_{0F} [illegible] H_{0N} [illegible] Stratified Randomized Experiment, SRE [illegible]

[illegible]

[illegible]

[illegible] CRE [illegible] n_1 [illegible] n_0 [illegible] $n = n_1 + n_0$ [illegible]

(1) [illegible]

$$H_{0F}: Y_i(1) = Y_i(0) \qquad i = 1, \ldots, n.$$

(2) [illegible]

$$H_{0N}: \tau = 0 \iff \bar{Y}(1) = \bar{Y}(0).$$

H_{0F} [illegible] FRT [illegible] p [illegible] H_{0N} [illegible] $\hat{\tau}$ [illegible] H_{0F} [illegible] studentized [illegible]

$$t = \frac{\hat{\tau}}{\sqrt{\hat{V}}},$$

[illegible] $\hat{\tau}$ [illegible] $\hat{V}$ [illegible][1]

---

[1] [illegible] ?? [illegible] ??

[illegible]

[illegible] Ding and Dasgupta (2017) [illegible] 在 *CRE* 中，使用 *studentized* 统计量 t 的 *FRT* 具有以下双重保证：

(1) H_{0F} ：p_{frt} 是有限样本精确的；
(2) H_{0N} ：p_{frt} 是渐近保守的（*asymptotically conservative*）。

[illegible] $\hat{\tau}$ [illegible] t [illegible] H_{0N} [illegible]

H_{0N} [illegible] $\hat{\tau}$ [illegible]

$$\hat{\tau} \doteq \mathrm{N}\left(0, \frac{S^2(1)}{n_1} + \frac{S^2(0)}{n_0} - \frac{S^2(\tau)}{n}\right).$$

[illegible] FRT pretends [illegible] Science Table [illegible] H_{0F} [illegible] $(Y_i, Y_i)_{i=1}^n$ [illegible] $\hat{\tau}$ [illegible]

$$\hat{\tau}^{\pi} \doteq \mathrm{N}\left(0, \frac{s^2}{n_1} + \frac{s^2}{n_0}\right),$$

[illegible] s^2 [illegible] FRT [illegible] H_{0N} [illegible] p

[illegible] studentized [illegible] $t = \hat{\tau}/\sqrt{\hat{V}}$ [illegible] H_{0N} [illegible]

$$t \doteq \mathrm{N}(0, C^2),$$

[illegible] $C^2 \leq 1$ [illegible] $Y_i(1) - Y_i(0) = \tau$ [illegible] constant treatment effect [illegible]

[illegible] FRT [illegible]

$$t^{\pi} \doteq \mathrm{N}(0, 1).$$

[illegible] C^2 FRT [illegible] 1 [illegible] p [illegible]

[illegible] studentization [illegible] FRT [illegible]

8.1. [illegible] 证明了这个双重保证是 *studentized* 统计量独有的性质。其他统计量（如原始 $\hat{\tau}$）的 *FRT* 在 H_{0N} 下可能不再控制第一类错误率。

[illegible]

[illegible] SRE [illegible] SRE [illegible] K [illegible] strata [illegible] k [illegible] $\tau_{[k]}$ [illegible] $\tau = \sum_{k=1}^K \pi_{[k]} \tau_{[k]}$ [illegible]

$$\hat{\tau}_{\mathrm{SRE}} = \sum_{k=1}^{K} \pi_{[k]} \hat{\tau}_{[k]},$$

$$\hat{V}_{\mathrm{SRE}} = \sum_{k=1}^{K} \pi_{[k]}^2 \left(\frac{\hat{S}^2_{[k]}(1)}{n_{[k]1}} + \frac{\hat{S}^2_{[k]}(0)}{n_{[k]0}}\right).$$

[illegible] SRE [illegible] studentized [illegible]

$$t_{\mathrm{SRE}} = \frac{\hat{\tau}_{\mathrm{SRE}}}{\sqrt{\hat{V}_{\mathrm{SRE}}}}.$$

[illegible] **(SRE FRT [illegible])** 在 *SRE* 中，使用 *studentized* 统计量 t_{SRE} 的 *FRT* 具有以下双重保证：

(1) H_{0F} ：p_{frt} 是有限样本精确的；
(2) H_{0N} ：p_{frt} 是渐近保守的。

SRE FRT [illegible] CRE [illegible] X [illegible]

[illegible]

[illegible] [illegible]

[illegible] studentized Lin [illegible]

$$t_{\mathrm{L}} = \frac{\hat{\tau}_{\mathrm{L}}}{\sqrt{\hat{V}_{\mathrm{L}}}},$$

[illegible] $\hat{\tau}_{\mathrm{L}}$ [illegible] $\hat{V}_{\mathrm{L}}$ [illegible] OLS [illegible] t_{L} [illegible] Y_i on $(1, Z_i, X_i, Z_iX_i)$ [illegible] Z_i [illegible] studentized [illegible]

[illegible] **(Zhao and Ding (2021a) [illegible])** 在 *CRE* 中，使用 *studentized Lin* 统计量 t_{L} 的 *FRT* 具有以下多重保证：

(1) H_{0F} ：p_{frt} 是有限样本精确的；
(2) H_{0N} ：p_{frt} 是渐近保守的；
(3) [illegible] H_{0N} [illegible] ：p_{frt} 渐近地比使用 t 的 *FRT* 更高效（功效更高）；
(4) [illegible]。

[illegible] $\hat{\tau}_{\mathrm{L}}$ [illegible] FRT with t_{L} [illegible] t_{L} [illegible] FRT [illegible] FRT [illegible] H_{0N} [illegible] t_{L}

[illegible] SRE [illegible]

$$t_{\mathrm{L,SRE}} = \frac{\hat{\tau}_{\mathrm{L,SRE}}}{\sqrt{\hat{V}_{\mathrm{L,SRE}}}}.$$

[illegible] Lin [illegible]

8X Eùî

\ „ \ YV E £ bim/2MidB9x2¥/µK"'ŸÉbáÌ Nº BñÅ˜!
Âb

InBñ c $n = 100$ ñ†í¥µK98 Í F $n_1 = 20$ vF $n_0 = 80$ b Äñ
†í i 3îxM X_i lMB(7;1) . ²T¹

$$Y_i(1) \quad L(X_i^3; 1); \quad Y_i(0) \quad L(\ X_i^3; 0{:}5^2):$$

$Y_i(1)$ „ $Y_i(0)$ É›Ï•Ä[' =0 '¡@ H_{0N} b
E1 RkÕ_ d9

URLV2vK M URN¥kØVÕ PGaLC Í F"”a ¿ÜÅS ¥ t d9 a
¿ 1>qS ¥ t d9
UkÀV²T¦{ \ n5 Y_i $(1; X_i)$ —µÄ –ª‹¨ •ØÕd9
UjV6Bb?2` UR¥NkP8GVaLC n» \\c
U9GVBM Uky¥RPjVGaLC n» \\c KÔØÕ bim/2MidB9x2b/

²T 3XXR H_{0N} / ºµ¹Õ bim/2MidB9x2/ GB¥MØÕ„ 1>q¥BÕ ?
û l»BËp q OL=p qÊ® ob ÅÖÕd9 ¥L=p q(Ú¿
l£Ü yN,a†¨¿_ Ð,L!b

ƒñ E²TÐ \\¥Ø,ç©›†BÁb

eXLlyp

¿ •Ø,„ E²T áÌó [/Llyp

eXRX *¥1yp X

URÂVTº1• H_{0F} '^Ži ©…)Ør‹ P¨©…_ d9 ¥ 6_hûó
µK"'Ú' T´bV[Ê4ÿr Ú¥d9 Â ^b
UkÂVT1• H_{0N} 'Ü(yTr‹^Ž¹, A¶P¨ bim/2MiBdx92/ w
iP¨ t t_Gb
UjVÂTµxM V¨ P¨ t_G YÈ1 t ÷Úrb

eXkX a¥1yp X

URíVxM P¨ t_{a_1}
UkµVxM P¨ t_{G,a_1}b

eXjXÛ Ä _2J ¥+Yªü X ¿P¨ J ? H MQ@ÕEÉb×Û Ä¥L
_2J \ ?CP¨ðS t d9 ¥ 6_h, µ \\Ï¥ × £bŒP¨ t_G ¥ 6_h
– ûƒt £bƒjA xM Øœ„ bim/2MiBx iB_Q2MJ Ï¥×1Ÿ b

eX9ÄX¥ L JS1 ¥yp X ¿Ä¥ L JS1 ypÜ›¿ a_1¥yp
íxM P¨ t d9 PGaE† $^\wedge_i$ Q1Mª¥© [
µxM P¨ t d9 PGaE† $^\wedge_i$ Q1M„ $^\wedge_{X;i}$ ª¥© [b

eX8,X², XP¨ bim/2MiBdx92/¥ 6_h ^÷½†¥Ê4b 'v"'žÿÍ
» 'H ¿ LQ`KÍH»„ ¿ 6_h¥ T´+¬M]b'žÿÍ», 'H 6_h
À ? £ b? `T MLm/H HT´^µr¥bƒ^'€¥wiSEb

[illegible]

[illegible] " [illegible] " vs [illegible] " [illegible] " vs [illegible]$_1$ [illegible]

[illegible]

- **I** ~ **P** [illegible] t [illegible] t_{L} [illegible] FRT[illegible]
- [illegible] ~ **P** [illegible] t_{SRE} [illegible] $t_{\mathrm{L,SRE}}$ [illegible] FRT

[illegible] soccer vs control [illegible] physician vs control [illegible]

Table 1. [illegible] soccer vs control

	est	s.e.	p_{Normal}	p_{frt}
class 1 (N)	0.051	0.502	0.919	0.924
class 1 (L)	0.050	0.489	0.919	0.929
class 2 (N)	−0.158	0.451	0.726	0.722
class 2 (L)	−0.176	0.452	0.698	0.700
class 3 (N)	0.005	0.403	0.990	0.989
class 3 (L)	−0.096	0.385	0.803	0.806
class 4 (N)	−0.492	0.447	0.271	0.288
class 4 (L)	−0.511	0.447	0.253	0.283
class 5 (N)	0.390	0.369	0.291	0.314
class 5 (L)	0.443	0.318	0.164	0.186
all (N)	−0.051	0.204	0.802	0.800
all (L)	−0.074	0.200	0.712	0.712

N[illegible] Neyman (1923) [illegible] L[illegible] Lin (2013) [illegible]

Table 2. [illegible] physician vs control

	est	s.e.	p_{Normal}	p_{frt}
class 1 (N)	0.567	0.426	0.183	0.192
class 1 (L)	0.588	0.418	0.160	0.174
class 2 (N)	0.193	0.438	0.659	0.666
class 2 (L)	0.265	0.409	0.517	0.523
class 3 (N)	1.305	0.494	0.008	0.012
class 3 (L)	1.501	0.462	0.001	0.003
class 4 (N)	−0.273	0.413	0.508	0.515
class 4 (L)	−0.313	0.417	0.454	0.462
class 5 (N)	−0.050	0.379	0.895	0.912
class 5 (L)	−0.067	0.279	0.811	0.816
all (N)	0.406	0.202	0.045	0.047
all (L)	0.463	0.190	0.015	0.017

N[illegible] Neyman (1923) [illegible] L[illegible] Lin (2013) [illegible]

[illegible]

(1) [illegible] t_{L} [illegible] t [illegible]

(2) [illegible] FRT p [illegible] p [illegible]

(3) [illegible] soccer [illegible] physician [illegible] $p_{\mathrm{frt}} \approx 0.017$ [illegible] class 3 [illegible] $p_{\mathrm{frt}} \approx 0.003$ [illegible]

(4) [illegible] class 2 [illegible] class 4 [illegible]

[illegible] studentized [illegible] N(0, 1) [illegible]

Figure 1. [illegible] N(0, 1) [illegible] N(0, 1) [illegible]

[illegible]

[illegible] Fisher [illegible]

(1) [illegible] studentized [illegible] $t = \hat{\tau}/\sqrt{\hat{V}}$ [illegible] FRT [illegible] H_{0F} [illegible] H_{0N} [illegible]

(2) [illegible] $\hat{\tau}$ [illegible] FRT [illegible] H_{0N} [illegible] FRT [illegible] Studentization [illegible]

(3) [illegible] studentized Lin [illegible] t_{L} [illegible] FRT [illegible] H_{0F} [illegible] H_{0N} [illegible]

(4) [illegible] SRE [illegible] t_{SRE} [illegible] SRE [illegible] $t_{\mathrm{L,SRE}}$ [illegible]

(5) [illegible] studentized [illegible] FRT [illegible] p [illegible] sharp null [illegible]

[illegible]

[illegible] 8.1. *8.1 —Angrist and Lavy (2009) 数据重新分析 这是第 7 章 Problem 7.8 的 Fisher counterpart。报告使用 studentized 统计量的 FRT 所得到的* p_{frt}。

[illegible] 8.2. *8.2 —Zhao and Ding (2021a) Figure 1 的复制 Zhao and Ding (2021a) 使用模拟来评估各种检验统计量的 FRT 的有限样本性质。基于他们的 Figure 1，他们推荐使用* $t_{\mathrm{L},SRE}$ *来分析 SRE。复制他们的 Figure 1。*

[illegible] 8.3. *8.3 —Studentization 的直觉 解释为什么 studentization 能够修复* $\hat{\tau}$ *在* H_{0N} *下的反保守问题。从方差结构的角度阐述你的理由。*

[illegible] 8.4. *8.4 —推导* t *在* H_{0N} *下的渐近分布 在 CRE 中，推导 studentized 统计量* $t = \hat{\tau}/\sqrt{\hat{V}}$ *在* H_{0N} *下的渐近分布。证明其渐近方差不超过 1。*

[illegible] 8.5. *8.5 —Lin 估计量的性质 描述 Lin (2013) 协变量调整估计量* $\hat{\tau}_{\mathrm{L}}$ *的构造。证明它在线性结果模型正确设定时是高效的。*

Ǘ 8.6. *8.6* —推荐阅读 { 和 { 在 *permutation tests* 中使用了 *studentized* 统计量。在设计-*based* 框架下，{ 和 { 给出了 *multi-armed experiments* 的推荐，{ 将推荐推广到协变量调整情形。阅读这些文献并撰写一页摘要。

[illegible] $\hat{\tau}$ [illegible] $\mathrm{var}(\hat{\tau} \mid \mathbf{Z})$ [illegible] X [illegible]

[illegible]

[illegible] ANCOVA [illegible]

$$Y(1) = \gamma_1 + \beta_1^T X + \varepsilon(1), \qquad E\{\varepsilon(1)\} = 0,$$

$$Y(0) = \gamma_0 + \beta_0^T X + \varepsilon(0), \qquad E\{\varepsilon(0)\} = 0.$$

[illegible]

$$\tau = E\{Y(1) - Y(0)\} = (\gamma_1 - \gamma_0) + (\beta_1 - \beta_0)^T E(X).$$

[illegible] $E(X)$ [illegible] $\varepsilon(1)$ [illegible] $\varepsilon(0)$ [illegible] τ [illegible]

[illegible] $\hat{\gamma}_1, \hat{\beta}_1, \hat{\gamma}_0, \hat{\beta}_0$ [illegible] OLS [illegible] τ [illegible]

$$\hat{\tau}_{\mathrm{adj}} = \hat{\gamma}_1 - \hat{\gamma}_0 + (\hat{\beta}_1 - \hat{\beta}_0)^T \bar{X}.$$

[illegible] $\bar{X} = 0$ [illegible]

$$\hat{\tau}_L = \hat{\gamma}_1 - \hat{\gamma}_0.$$

[illegible] OLS [illegible]

$$Y = \alpha_0 + \alpha_Z Z + \alpha_X^T X + \alpha_{ZX}^T (Z \cdot X) + \varepsilon,$$

[illegible] $\hat{\tau}_L = \hat{\alpha}_Z$[2] [illegible] $\hat{\tau}_L$ [illegible] τ [illegible]

[illegible] OLS [illegible] $\hat{\tau}_L$ [illegible] OLS [illegible] EHW [illegible] $\bar{X}$ [illegible] OLS [illegible][3] [illegible] EHW [illegible]

$$\hat{V}_L = \hat{V}_{\mathrm{EHW}} + \frac{1}{n} (\hat{\beta}_1 - \hat{\beta}_0)^T S_X^2 (\hat{\beta}_1 - \hat{\beta}_0),$$

[illegible] - $\hat{V}_{\mathrm{EHW}}$ [illegible] OLS [illegible] EHW [illegible] - S_X^2 [illegible] X [illegible] - $(\hat{\beta}_1 - \hat{\beta}_0)$ [illegible] $Z \cdot X$ [illegible] 95% [illegible][4]

---

[2] [illegible] OLS [illegible] ??

[3] [illegible] ??

[4] [illegible] ?? [illegible]

~[illegible]

[illegible] $n = 500$[illegible] 2000 [illegible] 0 [5] [illegible]

(1) **f a** [illegible] -0.0001[illegible]
(2) **P]** [illegible] EHW [illegible] 0.139[illegible] 0.151[illegible] 95% [illegible] 92.7%[illegible]
(3) [illegible] 0.153[illegible] 95% [illegible] 95.3%

[illegible]

Xw 、[illegible]**P**

[illegible] SRE[illegible] $X_i \in \{1, \dots, K\}$ [illegible]

[illegible] 9.2 (SRE [illegible]). 在超级总体框架下，分层随机实验满足：

$$Z \perp \{Y(1), Y(0)\} \mid X.$$

[illegible] $X = k$ [illegible]

$$\tau_{[k]} = E\{Y(1) - Y(0) \mid X = k\} = E(Y \mid Z = 1, X = k) - E(Y \mid Z = 0, X = k).$$

[illegible]

$$\tau = \sum_{k=1}^{K} \mathrm{pr}(X = k)\tau_{[k]}.$$

SRE [illegible] CRE[illegible] CRE [illegible] Lin (2013) [illegible][6] [illegible] 2 [illegible] $\hat{V}_S$ [illegible] $\mathrm{var}(\hat{\tau}_S)$ [illegible]

1

Ww[illegible]

[illegible]

(1) [illegible] $Z \perp \{Y(1), Y(0), X\}$ [illegible] τ [illegible]
(2) [illegible] X [illegible]
(3) **f a l t** [illegible]**q ~ u** [illegible] OLS [illegible]
(4) [illegible] EHW [illegible]
(5) 、 ~ **P**[illegible]

---

[5] R [illegible] **??**

[6] [illegible] **??**

[illegible]

[illegible]

[illegible] 9.1. *CRE* 的协变量平衡 在超级总体框架下，证明 [illegible] 平均因果效应可以写为 $E(Y \mid Z=1) - E(Y \mid Z=0)$。

[illegible] 9.2. *OLS* 分解与 *pooled regression* 基于 [illegible] 和 [illegible] 证明观测结果的 *OLS* 分解为：

$$Y = \alpha_0 + \alpha_Z Z + \alpha_X^T X + \alpha_{ZX}^T (Z \cdot X) + \varepsilon,$$

其中

$$\alpha_0 = \gamma_0, \quad \alpha_Z = \gamma_1 - \gamma_0, \quad \alpha_X = \beta_0, \quad \alpha_{ZX} = \beta_1 - \beta_0.$$

[illegible] 9.3. *Lin (2013)* 估计量的性质 证明在正则条件下，*Lin (2013)* 估计量 $\hat{\tau}_L$ 是 τ 的半参数有效估计量。

[illegible] 9.4. 方差估计量的比较 在超级总体框架下，模拟 $\{X_i, Z_i, Y_i(1), Y_i(0)\}_{i=1}^n$，其中 $\beta_1 \neq \beta_0$。比较以下四种方差估计量的表现：

(1) 标准的 *EHW* 稳健方差估计量；
(2) 带校正项 [illegible] 的 *EHW* 估计量；
(3) *Bootstrap* 方差估计量（协变量在开始时中心化，但不每次重新中心化）；
(4) *Bootstrap* 方差估计量（协变量在每次 *bootstrap* 样本中重新中心化）。

[illegible] 9.5. *SRE* 中的协变量调整 在 *SRE* 的超级总体框架下，推导 τ 的协变量调整估计量及其方差估计量。

[illegible] 9.6. 协变量调整与后分层的关系 证明：当各层的层别倾向得分 $e_{[k]} = n_{[k]1}/n_{[k]}$ 相等时，协变量调整与后分层给出相同的点估计量。

[illegible] 9.7. 文献阅读 阅读 [illegible] 的原文，总结其核心贡献和主要结论。

[illegible] 9.8. 回归调整的几何解释 从几何角度解释协变量调整：为什么减去 $(\hat{\beta}_1 - \hat{\beta}_0)^T \bar{X}$ 可以提高估计精度？

CHAPTER 10

CRE

Part III

earnings

LaLonde

NHANES 2005-2006

homocysteine

female

age3 ed3 BMI bmi3 pov2

BMI

NHANES 2007-2008

BMI

[illegible]

[illegible] i [illegible] X_i [illegible] $Z_i \in \{0, 1\}$ [illegible] Y_i [illegible] $Y_i(1)$ [illegible] $Y_i(0)$ [illegible]

$$\{X_i, Z_i, Y_i(1), Y_i(0)\}_{i=1}^{n} \overset{\text{iid}}{\sim} \{X, Z, Y(1), Y(0)\}.$$

[illegible]

- [illegible]

$$\tau = \mathbb{E}\{Y(1) - Y(0)\}.$$

- [illegible]

$$\tau_{\mathrm{T}} = \mathbb{E}\{Y(1) - Y(0) \mid Z = 1\}.$$

- [illegible]

$$\tau_{\mathrm{C}} = \mathbb{E}\{Y(1) - Y(0) \mid Z = 0\}.$$

[illegible]

$$\tau_{\mathrm{T}} = \mathbb{E}(Y \mid Z = 1) - \mathbb{E}\{Y(0) \mid Z = 1\}.$$

[illegible] $\mathbb{E}(Y \mid Z = 1)$ [illegible] $\mathbb{E}\{Y(0) \mid Z = 1\}$ [illegible]

[illegible] τ_{C} [illegible]

$$\tau_{\mathrm{C}} = \mathbb{E}\{Y(1) \mid Z = 0\} - \mathbb{E}(Y \mid Z = 0),$$

[illegible] $\mathbb{E}\{Y(1) \mid Z = 0\}$ [illegible]

$$\tau_{\mathrm{PF}} = \mathbb{E}(Y \mid Z = 1) - \mathbb{E}(Y \mid Z = 0).$$

[illegible]

$$\tau_{\mathrm{PF}} - \tau_{\mathrm{T}} = \mathbb{E}\{Y(0) \mid Z = 1\} - \mathbb{E}\{Y(0) \mid Z = 0\},$$

$$\tau_{\mathrm{PF}} - \tau_{\mathrm{C}} = \mathbb{E}\{Y(1) \mid Z = 1\} - \mathbb{E}\{Y(1) \mid Z = 0\}.$$

[illegible] τ_{PF} [illegible] CRE [illegible] $Z \perp \{Y(1), Y(0)\}$ [illegible]

[illegible]

[illegible]

1 [illegible][8]

$$\hat{\tau} = n^{-1}\sum_{i=1}^{n}\left\{\frac{e^{\hat{\beta}_0+\hat{\beta}_z+\hat{\beta}_x X_i}}{1+e^{\hat{\beta}_0+\hat{\beta}_z+\hat{\beta}_x X_i}} - \frac{e^{\hat{\beta}_0+\hat{\beta}_x X_i}}{1+e^{\hat{\beta}_0+\hat{\beta}_x X_i}}\right\}.$$

[illegible] Z [illegible]

[illegible] ” [illegible]

” 5 ” [illegible] ”

[illegible]

[illegible] Part III [illegible] ”Four Pillars” [illegible]

(1) [illegible]

(2) [illegible] $Y(z) \perp Z \mid X$ [illegible] X [illegible]

(3) [illegible]

(4) [illegible] $\mathbb{E}(Y \mid Z, X)$ [illegible]

(5) [illegible]

[illegible]

[illegible]

(1) [illegible] $\tau = \Pr(Z=1)\tau_{\mathrm{T}} + \Pr(Z=0)\tau_{\mathrm{C}}$

(2) [illegible]

$$Y(1) = g_1(X, V_1), \quad Y(0) = g_0(X, V_0), \quad Z = \mathbb{I}\{g(X, V) \geq 0\},$$

[illegible] $(V_1, V_0) \perp V$ [illegible] V [illegible] U [illegible]

(3) [illegible] Y [illegible]

$$\mathbb{E}(Y \mid Z, X) = \frac{e^{\beta_0+\beta_z Z+\beta_x X+\beta_{xz} XZ}}{1+e^{\beta_0+\beta_z Z+\beta_x X+\beta_{xz} XZ}}.$$

[illegible] τ [illegible]

(4) [illegible]

[illegible]

- [illegible]
- [illegible]
- [illegible]
- [illegible] g-formula [illegible]
- [illegible]

---

[8] [illegible] **??**

CHAPTER 11

[illegible]

[illegible]

[illegible] randomization [illegible] treatment [illegible]

[illegible] X [illegible]

[illegible]

[illegible] $\Pr(Z = 1 \mid X)$ [illegible] X [illegible]

[illegible] 11.1 ([illegible] (Propensity Score)). 倾向得分定义为给定协变量 X 时接受治疗的概率：

$$e(X) = \Pr(Z = 1 \mid X).$$

[illegible] $[0, 1]$ [illegible] $e(X_i) = 0.8$ [illegible] X_i [illegible] 80% [illegible]

[illegible] $e(X) = \Pr(Z = 1)$ [illegible] randomization [illegible] X [illegible]

[illegible] X [illegible] $e(X)$ [illegible] X [illegible]

[illegible] (Rosenbaum & Rubin, 1983) 在强可忽略性（*strong ignorability*）假设下，如果

$$\{Y(1), Y(0)\} \perp\!\!\!\perp Z \mid X,$$

则

$$\{Y(1), Y(0)\} \perp\!\!\!\perp Z \mid e(X).$$

[illegible] $e(X)$ [illegible] X [illegible]

¹I¹ƒñ²T×1$ ðSxM X V[^©i»”¥ M-alÆa•âa¯<ll Œ`_¤s e(X) ^Bñ [0;1] uW¥S bYVØœƒñB»M áÌ©N¿Øœ îµðSxM iiƒü^ _Qb2M#„ m_Km#BěM¹ ô†»ý ô¥clb[R] ÐsªÛ L ¥ó" » \\ cÏ áÌ), sªÛ L a_1bÂTáÌ ¿`_¤s e(X) ¥|´ †íÉ›sª *¹ÄBª=¥4©Ÿ” Ð a_1 µM]¥ ² iiy¹óçM]¥`_¤s Í s¥Ð. ²THqÿëb

jX`_¤ssª

jXRX©ÖÛ`_¤s XL!`_¤s e(X) X© OÇ| K ñ,]¥´ $f e_1;:::;e_K g$b ® \\ ÄBª e_k = Í s¥Ð. ²THqÿëbyN áÌµ K ñÿë¥›†Û L *_1 iiƒž^» \\ c),¥ a_1² b
98Ü(yTr‹V[YVsª 9 É› 9

$$\hat{}_{Sa} = \sum_{k=1}^{K} {}_{[k]}\hat{}_{[k]};$$

Ï $_{[k]}$ ^» k ª¥981è $\hat{}_{[k]}$ ^¾ª=¥µ´(´b ^{k} LlÏ¥Ù5 K ¥Ê4³1ƒ™bÂT K þl sªª¥HqÿëŸV?,îë ÂTK þv Äª=¥"' ,@ ¸ ªV?º cÍ F vF¥†íbô \ ¥yp K =5 ¸ Æß/? h"v†sÊµiiƒñ ô* ¥”3 8ô| Q C 'c¥),Ïb

jXkÃ©õÃ`_¤s X L=ffƒÏ `_¤sYÈ^Ã©¥ áÌ³1YVd9 ˜É› 9bKÈn¥ZE^E†Bñ HQ;BbBB†

$$\mathrm{HQ} f_{;} e(X i) g = {}_0 + {}_1X_1 + \quad + {}_pX_p:$$

9¤ž¥`_¤s ê(X) V[| r n ñ,]¥´ ’Äñ†í¥xM û,›† M]Hb¹ ‹¨sªZE áÌ³1 ê(X) É›ÖÛÄbBÕÈ¨¦{^¿ sÊ” É›sª[j]

è RRXR UL>”L1¥a`_¤ssª VXln \\ Ï¥ L> L1a” báÌP ¨ HQ;Bb˜iB9+`_¤s –ª¿sÊ”|"'s¹ K ªb \\ ZU ,] K ´ K =5;10;30 / Í F„ vF`_¤ssƒ¥°Zmb

> 6B;m`2 RX L1a” "Ï 9`_¤s¥°Zm âä¹ vF ç ä¹Í Fb

’`_¤sÃ©OõÃH `_¤ssª 9 ¥µrŸG ¿[/ ñHq
URžV!ç ˜ S(Z =1 j X) ž’!ç P¤ ê(X) ? } ‡L¥`_¤sb
Uk¥Vß Hq Pp2`H T +QM/B0i<Be(X M) < 1 îµ†íîëiiƒ’ Äñ[9] †íûµ¤sÍ „ v¥V?Ÿb

---

[R] w•n¸c \\
[k] W%w•n¸c \\
[j] W%),n¸c \\
[9] ×ßHq¥W%),n¸c \\

稳健性观察：倾向得分分层估计量只需要 $\hat{e}(X)$ 的排序正确，而不需要其精确值——这使得它相对于其他方法更加稳健。这一性质在许多数值例子中得到验证，但其严格量化理论仍在研究中。

4. 逆概率加权估计量

4.1. Horvitz–Thompson 估计量. 基于倾向得分，我们可以构造平均因果效应的估计量。一个直接的想法是使用**逆概率加权（Inverse Probability Weighting, IPW）**：

$$\hat{\tau}_{\mathrm{HT}} = \frac{1}{n}\sum_{i=1}^{n}\left\{\frac{Z_iY_i}{e(X_i)} - \frac{(1-Z_i)Y_i}{1-e(X_i)}\right\}.$$

?? 被称为 **Horvitz–Thompson（HT）估计量**。? 在抽样调查中首次提出这一估计量，? 将其引入因果推断领域。

为什么这个估计量有效？直观上，对于治疗组的单元，我们用 $1/e(X_i)$ 作为权重进行加权——倾向得分越小的单元（即越”不像”会接受治疗的单元），其权重越大。这相当于对这些单元进行”过度代表”，以消除由选择性偏差造成的治疗组和对照组之间的系统性差异。[5]

稳定性问题：当某些单元的倾向得分接近 0 或 1 时，权重 $1/e(X_i)$ 或 $1/(1-e(X_i))$ 会变得非常大，导致估计量不稳定。HT 估计量在有限样本中可能表现极差——某些估计值可能远离其他所有方法的结果。[6]

4.2. 标准化逆概率加权估计量. 为了解决 HT 估计量的不稳定性问题，一个自然的修正是对权重进行**标准化**：

$$\hat{\tau}_{\mathrm{IPW}} = \frac{1}{\hat{\mathbf{1}}_T}\sum_{i=1}^{n}\frac{Z_iY_i}{e(X_i)} - \frac{1}{\hat{\mathbf{1}}_C}\sum_{i=1}^{n}\frac{(1-Z_i)Y_i}{1-e(X_i)},$$

其中

$$\hat{\mathbf{1}}_T = \sum_{i=1}^{n}\frac{Z_i}{e(X_i)}, \quad \hat{\mathbf{1}}_C = \sum_{i=1}^{n}\frac{1-Z_i}{1-e(X_i)}.$$

直观理解：标准化权重的思想是将权重之和调整为 1——即 $\hat{\mathbf{1}}_T$ 和 $\hat{\mathbf{1}}_C$ 分别代表治疗组和对照组的”有效样本量”。标准化后的估计量对常数平移不敏感，而原始 HT 估计量则不具备这一性质。[7]

例 11.2 (NHANES 数据的逆概率加权估计). *延续* **??**，*我们对 NHANES 数据应用不同截断水平的逆概率加权估计量。*

TABLE 1. 不同截断水平下的 IPW 估计量

截断水平	点估计	标准误
$(0,1)$（无截断）	-52.3	18.2
$(0.01, 0.99)$	-8.7	4.3
$(0.05, 0.95)$	-6.2	3.8
$(0.10, 0.90)$	-5.1	3.5

---

[5] HT 估计量的期望推导见附录 ??

[6] HT 估计量的方差分析见附录 ??

[7] 标准化估计量的性质证明见附录 ??

[illegible] HT [illegible] IPW [illegible]

[illegible]

[illegible] 倾向得分 $e(X)$ 满足以下性质：

(1) 条件独立性：$X \perp\!\!\!\perp Z \mid e(X)$；

(2) 加权平衡：对于任何协变量函数 $g(X)$，有

$$\mathbb{E}\left\{\frac{Zg(X)}{e(X)}\right\} = \mathbb{E}\{g(X)\}.$$

[illegible] X [illegible] Z [illegible][8] [illegible] Z [illegible] X [illegible]

[illegible] $\hat{e}(X)$ [illegible] $\hat{e}(X)$ [illegible]

$$\mathrm{Bias}_{[k]} = \frac{1}{n_{[k]}} \sum_{i:\hat{e}\ (X_i)=e_k} Z_i g(X_i) - \frac{1}{n_{[k]}} \sum_{i:\hat{e}\ (X_i)=e_k} (1 - Z_i) g(X_i) \approx 0.$$

[illegible]

[illegible]

Table 2. [illegible] vs. [illegible]

	[illegible]	[illegible]
[illegible]	$\Pr(Z = 1 \mid X)$	[illegible] $\mathbb{E}(Y \mid Z, X)$
[illegible]		[illegible]
[illegible]		[illegible]
[illegible]		[illegible]

---

[8] [illegible] ??

[illegible]

[illegible] $0 < e(X) < 1$ [illegible]

[illegible] $e(X_i) = 1$ [illegible] i [illegible] $Y_i(0)$ [illegible]

[illegible]

(1) [illegible]

(2) [illegible] 0 ⁵ 1 [illegible]

[illegible] 0 ⁵ 1 [illegible]

[illegible]

[illegible]

(1) [illegible] $e(X) = \Pr(Z = 1 \mid X)$ [illegible]

(2) [illegible]

(3) [illegible]

(4) [illegible] Horvitz–Thompson [illegible]

(5) [illegible]

(6) [illegible] $0 < e(X) < 1$ [illegible]

[illegible]

[illegible] 11.1. 倾向得分的降维性质 证明 [illegible] 在强可忽略性假设下，如果 $\{Y(1), Y(0)\} \perp\!\!\!\perp Z \mid X$，则 $\{Y(1), Y(0)\} \perp\!\!\!\perp Z \mid e(X)$。

提示：利用概率论中的推论——如果 $A \perp\!\!\!\perp B \mid C$，且 D 是 C 的函数，则 $A \perp\!\!\!\perp B \mid D$。

[illegible] 11.2. *HT* 估计量的期望 证明 *HT* 估计量 [illegible] 是总体平均因果效应 τ 的无偏估计。

提示：分别计算 $\mathbb{E}\left(\frac{ZY(1)}{e(X)}\right)$ 和 $\mathbb{E}\left(\frac{(1-Z)Y(0)}{1-e(X)}\right)$，利用 $e(X) = \Pr(Z = 1 \mid X)$。

[illegible] 11.3. 标准化权重的性质 验证 [illegible] 中标准化权重的性质：$\hat{\mathbf{1}}_T$ 和 $\hat{\mathbf{1}}_C$ 在使用真实倾向得分时，其期望均为 n。

注：这也是 [illegible] 中使用特殊记号 $\hat{1}_T$ 和 $\hat{1}_C$ 的原因——这两个量的期望都是 *1*。

[illegible] 11.4. 倾向得分的平衡性质证明 证明 [illegible] 的第二个结论：对于任意协变量函数 $g(X)$，

$$\mathbb{E}\left\{\frac{Zg(X)}{e(X)}\right\} = \mathbb{E}\{g(X)\}.$$

提示：使用 *tower property* 和条件概率的定义。

ǘ 11.5. 协变量平衡的数值检验 使用 { {中的 *NHANES* 数据，检验不同分层数量 K 下的协变量平衡。报告治疗组和对照组在各层内关键协变量（如年龄、*BMI*）均值之差的绝对值。

ǘ 11.6. 截断水平的敏感性分析 对 { {中的结果进行敏感性分析：比较不同截断水平对点估计和标准误的影响，讨论截断水平选择的实际准则。

ǘ 11.7. *IPW* 与结果回归的比较 使用第 { {章中的模拟数据或真实数据，分别应用 *IPW* 估计量和结果回归估计量，比较两者的点估计和置信区间。

ǘ 11.8. 极端反事实的影响 { 讨论了极端反事实问题。构造一个例子，其中某些单元的倾向得分接近 *1*，讨论这种情况下因果推断面临的挑战。

CHAPTER 12

[illegible]

[illegible]

[illegible] outcome regression [illegible]

$$\tau = E\{\mu_1(X)\} - E\{\mu_0(X)\},$$

[illegible] $\mu_1(X) = E(Y \mid Z = 1, X)$ [illegible] $\mu_0(X) = E(Y \mid Z = 0, X)$ [illegible] X [illegible] $\mu_1(X)$ [illegible] $\mu_0(X)$ [illegible] τ [illegible] IPW [illegible]

$$\tau = E\left\{\frac{ZY}{e(X)}\right\} - E\left\{\frac{(1-Z)Y}{1-e(X)}\right\},$$

[illegible] $e(X) = \Pr(Z = 1 \mid X)$ [illegible] $e(X)$ [illegible] James Robins [illegible] doubly robust estimator [illegible]

[illegible]

[illegible] working models [illegible]

- [illegible] $\mu_1(X, \beta_1)$ [illegible] $\mu_0(X, \beta_0)$ [illegible] $E(Y(1) \mid X)$ [illegible] $E(Y(0) \mid X)$ [illegible] $\mu_1(X, \beta_1) = \mu_1(X)$
- [illegible] $e(X, \alpha)$ [illegible] $\Pr(Z = 1 \mid X)$ [illegible] $e(X, \alpha) = e(X)$

[illegible] ” [illegible] ” $_1$ [illegible]

$$\tilde{\mu}_1^{\text{dr}} = E\left[\frac{Z\{Y - \mu_1(X, \beta_1)\}}{e(X, \alpha)} + \mu_1(X, \beta_1)\right], \tag{60}$$

$$\tilde{\mu}_0^{\text{dr}} = E\left[\frac{(1-Z)\{Y - \mu_0(X, \beta_0)\}}{1 - e(X, \alpha)} + \mu_0(X, \beta_0)\right]. \tag{61}$$

[illegible]

$$\tilde{\mu}_1^{\mathrm{dr}} = E\left[\frac{ZY}{e(X,\alpha)} - \frac{Z - e(X,\alpha)}{e(X,\alpha)}\mu_1(X,\beta_1)\right], \tag{62}$$

$$\tilde{\mu}_0^{\mathrm{dr}} = E\left[\frac{(1-Z)Y}{1-e(X,\alpha)} - \frac{e(X,\alpha) - Z}{1-e(X,\alpha)}\mu_0(X,\beta_0)\right]. \tag{63}$$

[illegible]

- [illegible]
- [illegible] IPW [illegible]

[illegible] augmented inverse propensity score weighting, AIPW[illegible]

[illegible]

[illegible]

[illegible] [illegible]假设可忽略性 $Z \perp \{Y(1), Y(0)\} \mid X$ 和 *overlap* $0 < e(X) < 1$。

(1) 若 $e(X,\alpha) = e(X)$ 或 $\mu_1(X,\beta_1) = \mu_1(X)$ 之一成立，则 $\tilde{\mu}_1^{\mathrm{dr}} = E\{Y(1)\}$。
(2) 若 $e(X,\alpha) = e(X)$ 或 $\mu_0(X,\beta_0) = \mu_0(X)$ 之一成立，则 $\tilde{\mu}_0^{\mathrm{dr}} = E\{Y(0)\}$。
(3) 若 $e(X,\alpha) = e(X)$ 或 $\{\mu_1(X,\beta_1) = \mu_1(X), \mu_0(X,\beta_0) = \mu_0(X)\}$ 之一成立，则 $\tilde{\mu}_1^{\mathrm{dr}} - \tilde{\mu}_0^{\mathrm{dr}} = \tau$。

[illegible] $\tilde{\mu}_1^{\mathrm{dr}} - \tilde{\mu}_0^{\mathrm{dr}}$ [illegible] τ [illegible][1]

[illegible]

[illegible] analog [illegible]

[illegible] 12.2 ([illegible]). 基于数据 $(X_i, Z_i, Y_i)_{i=1}^n$，按以下步骤构造 τ 的双重稳健估计量：

(1) 拟合倾向得分：得到 $e(X_i, \hat{\alpha})$；
(2) 拟合结果均值：得到 $\mu_1(X_i, \hat{\beta}_1)$ 和 $\mu_0(X_i, \hat{\beta}_0)$；
(3) 构造双重稳健估计量 $\hat{\tau}^{\mathrm{dr}} = \hat{\mu}_1^{\mathrm{dr}} - \hat{\mu}_0^{\mathrm{dr}}$，其中

$$\hat{\mu}_1^{\mathrm{dr}} = \frac{1}{n}\sum_{i=1}^{n}\left[\frac{Z_i\{Y_i - \mu_1(X_i,\hat{\beta}_1)\}}{e(X_i,\hat{\alpha})} + \mu_1(X_i,\hat{\beta}_1)\right],$$

$$\hat{\mu}_0^{\mathrm{dr}} = \frac{1}{n}\sum_{i=1}^{n}\left[\frac{(1-Z_i)\{Y_i - \mu_0(X_i,\hat{\beta}_0)\}}{1-e(X_i,\hat{\alpha})} + \mu_0(X_i,\hat{\beta}_0)\right].$$

[illegible][2]

$$\hat{\tau}^{\mathrm{dr}} = \hat{\tau}^{\mathrm{reg}} + \frac{1}{n}\sum_{i=1}^{n}\frac{Z_i\{Y_i - \mu_1(X_i,\hat{\beta}_1)\}}{e(X_i,\hat{\alpha})} - \frac{1}{n}\sum_{i=1}^{n}\frac{(1-Z_i)\{Y_i - \mu_0(X_i,\hat{\beta}_0)\}}{1-e(X_i,\hat{\alpha})}.$$

---

[1] [illegible] **??** [illegible] **??**
[2] [illegible] **??**

[illegible] $\hat{\tau}^{\text{reg}}$ [illegible][3]

$$\hat{\tau}^{\text{dr}} = \hat{\tau}^{\text{ipw}} - \frac{1}{n}\sum_{i=1}^{n}\frac{Z_i - e(X_i,\hat{\alpha})}{e(X_i,\hat{\alpha})}\mu_1(X_i,\hat{\beta}_1) + \frac{1}{n}\sum_{i=1}^{n}\frac{e(X_i,\hat{\alpha}) - Z_i}{1 - e(X_i,\hat{\alpha})}\mu_0(X_i,\hat{\beta}_0).$$

[illegible] bootstrap [illegible] $\hat{\tau}^{\text{dr}}$ [illegible]

[illegible]

[illegible] IPW [illegible]

[illegible] [illegible] **[illegible]** IPW [illegible] $\mu_1 = E\left\{\frac{ZY}{e(X)}\right\}$ [illegible] Y [illegible]

[illegible] $\mu_1(X,\beta_1)$ [illegible][4]

$$\mu_1 = E\{Y(1)\} = E\{Y(1) - \mu_1(X,\beta_1)\} + E\{\mu_1(X,\beta_1)\} = E\left\{\frac{Z\{Y - \mu_1(X,\beta_1)\}}{e(X)} + \mu_1(X,\beta_1)\right\}.$$

[illegible]

[illegible] [illegible] $\tilde{\mu}_1 = E\{\mu_1(X,\beta_1)\}$ [illegible] μ_1 [illegible] $E\{\mu_1(X,\beta_1) - Y(1)\}$ [illegible] IPW [illegible][5]

$$B = E\left\{\frac{Z\{\mu_1(X,\beta_1) - Y\}}{e(X)}\right\},$$

[illegible] $\tilde{\mu}_1 - B$ [illegible]

[illegible]

[illegible][6]

(1) [illegible]
(2) [illegible]
(3) [illegible]
(4) [illegible]

[illegible] 1-3 [illegible] DR [illegible] 4 [illegible] DR [illegible][7] ” [illegible] ” [illegible]

---

[3] [illegible] ??

[4] [illegible] ??

[5] [illegible] ??

[6] [illegible] ??

[7] [illegible] ??

7. 进一步讨论：为何脆弱？

回顾 ?? 的证明，双重稳健性的关键在于以下乘积结构：[8]

$$\tilde{\mu}_1^{\mathrm{dr}} - E\{Y(1)\} = E\left[\frac{e(X) - e(X,\alpha)}{e(X,\alpha)} \times \{\mu_1(X) - \mu_1(X,\beta_1)\}\right].$$

只要 $e(X) = e(X,\alpha)$ 或 $\mu_1(X) = \mu_1(X,\beta_1)$ 之一成立，上式就为零。然而，当两个模型都错误时，两个误差的乘积可能放大偏差——这就是”双重脆弱性”的数学本质。模拟研究证实了这一点。

尽管 ? 基于模拟研究批评了双重稳健估计量，但自 ? 的开创性工作以来，双重稳健估计量一直是因果推断的标准策略。近年来，它在理论统计和计量经济学文献中以更响亮的名字”双机器学习”（double machine learning）重新兴起。? 的基本思想是用更灵活的机器学习工具替代传统的参数工作模型。

8. 本章小结

本章介绍了双重稳健估计量，主要内容如下：

(1) **核心思想**：结合结果回归和 IPW，只要其中一个模型正确就能保证一致性。
(2) **公式结构（??–??）**：增强逆概率加权估计量（AIPW）。
(3) **双重稳健性（??）**：只要倾向得分模型或结果模型之一正确，$\tilde{\mu}_1^{\mathrm{dr}} - \tilde{\mu}_0^{\mathrm{dr}} = \tau$。
(4) **两种直觉视角**：从 IPW 降方差（??）和从结果回归降偏倚（??）。
(5) **双重脆弱性**：当两个模型都错误时，两个误差的乘积可能放大偏差。

9. 习题

练习 12.1. *验证双重稳健性 考虑协变量离散的情况* $X \in \{1, \ldots, K\}$*。证明在这种情况下，分层估计量、结果回归估计量、Horvitz-Thompson 估计量、Hajek 估计量和双重稳健估计量在数值上是完全相同的。(参见 ?, Theorem 12.1)*

练习 12.2. *DR 估计量的另一种形式 基于 ??，给出* $\mu_1 = E\{Y(1)\}$ *的双重稳健估计量的另一种形式：*

$$\tilde{\mu}_1^{\mathrm{dr2}} = \frac{E\left[\frac{Z\{Y - \mu_1(X,\beta_1)\}}{e(X,\alpha)}\right]}{E\left[\frac{Z}{e(X,\alpha)}\right]} + E\{\mu_1(X,\beta_1)\}.$$

证明若 $e(X,\alpha) = e(X)$ *或* $\mu_1(X,\beta_1) = \mu_1(X)$ *之一成立，则* $\tilde{\mu}_1^{\mathrm{dr2}} = \mu_1$*。给出估计* μ_0 *的类似公式，并给出基于这些公式的* τ *的双重稳健估计量的样本 analog。(参见 ?, 式 (12.7))*

练习 12.3. *DR 估计量的方差 证明 ?? 和 ?? 的等价性。(参见 ?, 式 (12.9))*

---

[8] 详细推导见附录 ??

[illegible]

[illegible] ignorability [illegible] overlap [illegible] $\tilde{\mu}_1^{\mathrm{dr}} = E\{Y(1)\}$ [illegible] [9]

$$
\begin{aligned}
\tilde{\mu}_1^{\mathrm{dr}} - E\{Y(1)\} &= E\left[\frac{Z\{Y(1)-\mu_1(X,\beta_1)\}}{e(X,\alpha)} - \{Y(1)-\mu_1(X,\beta_1)\}\right] \\
&= E\left[\frac{Z-e(X,\alpha)}{e(X,\alpha)}\{Y(1)-\mu_1(X,\beta_1)\}\right] \\
&= E\left(E\left[\frac{Z-e(X,\alpha)}{e(X,\alpha)}\{Y(1)-\mu_1(X,\beta_1)\} \mid X\right]\right) \quad \text{(tower property)} \\
&= E\left[E\left\{\frac{Z-e(X,\alpha)}{e(X,\alpha)} \mid X\right\} \times E\left\{Y(1)-\mu_1(X,\beta_1) \mid X\right\}\right] \quad \text{(ignorability)} \\
&= E\left[\frac{e(X)-e(X,\alpha)}{e(X,\alpha)} \times \{\mu_1(X)-\mu_1(X,\beta_1)\}\right].
\end{aligned}
$$

[illegible] $e(X) = e(X,\alpha)$ [illegible] $\mu_1(X) = \mu_1(X,\beta_1)$ [illegible] □

[illegible]

[illegible]

$$
\hat{\tau}^{\mathrm{dr}} = \hat{\mu}_1^{\mathrm{dr}} - \hat{\mu}_0^{\mathrm{dr}} = \frac{1}{n}\sum_{i=1}^{n}\left[\mu_1(X_i,\hat{\beta}_1) - \mu_0(X_i,\hat{\beta}_0)\right] + \frac{1}{n}\sum_{i=1}^{n}\left[\frac{Z_i\{Y_i-\mu_1(X_i,\hat{\beta}_1)\}}{e(X_i,\hat{\alpha})} - \frac{(1-Z_i)\{Y_i-\mu_0(X_i,\hat{\beta}_0)\}}{1-e(X_i,\hat{\alpha})}\right]
$$

[illegible] $\hat{\tau}^{\mathrm{reg}}$ [illegible]

[illegible]

$$
\frac{Z\{Y-\mu_1(X,\beta_1)\}}{e(X,\alpha)} = \frac{ZY}{e(X,\alpha)} - \frac{Z}{e(X,\alpha)}\mu_1(X,\beta_1),
$$

[illegible]

$$
\frac{(1-Z)\{Y-\mu_0(X,\beta_0)\}}{1-e(X,\alpha)} = \frac{(1-Z)Y}{1-e(X,\alpha)} - \frac{1-Z}{1-e(X,\alpha)}\mu_0(X,\beta_0),
$$

[illegible]

[illegible] **M** [illegible]

[illegible]

$$
\mu_1 = E\{Y(1)\} = E\{Y(1)-\mu_1(X,\beta_1)\} + E\{\mu_1(X,\beta_1)\}
$$

[illegible] IPW [illegible] $Y(1)-\mu_1(X,\beta_1)$ [illegible]

$$
E\{Y(1)-\mu_1(X,\beta_1)\} = E\left\{\frac{Z\{Y-\mu_1(X,\beta_1)\}}{e(X)}\right\},
$$

[illegible]

$$
\mu_1 = E\left\{\frac{Z\{Y-\mu_1(X,\beta_1)\}}{e(X)}\right\} + E\{\mu_1(X,\beta_1)\}.
$$

---

[9] [illegible] (**?**, Theorem 12.1)

[illegible]

[illegible] $\tilde{\mu}_1 = E\{\mu_1(X, \beta_1)\}$ [illegible]

$$\text{bias} = E\{\mu_1(X, \beta_1) - Y(1)\}.$$

[illegible] IPW [illegible]

$$B = E\left\{\frac{Z\{\mu_1(X, \beta_1) - Y\}}{e(X)}\right\}.$$

[illegible] $\tilde{\mu}_1 - B$ [illegible]

[illegible]

[illegible]

$$\tilde{\mu}_1^{\mathrm{dr}} - E\{Y(1)\} = E\left[\frac{e(X) - e(X, \alpha)}{e(X, \alpha)} \times \{\mu_1(X) - \mu_1(X, \beta_1)\}\right].$$

[illegible] $e(X) - e(X, \alpha)$ [illegible] $\mu_1(X) - \mu_1(X, \beta_1)$ [illegible]

[illegible]

[illegible] $n = 500$ [illegible] 500 [illegible]

[illegible]

```
x = matrix(rnorm(n*2), n, 2)
x1 = cbind(1, x)
beta.z = c(0, 1, 1)
pscore = 1/(1 + exp(- as.vector(x1 %*% beta.z)))
z = rbinom(n, 1, pscore)
beta.y1 = c(1, 2, 1)
beta.y0 = c(1, 2, 1)
y1 = rnorm(n, x1 %*% beta.y1)
y0 = rnorm(n, x1 %*% beta.y0)
y = z*y1 + (1-z)*y0
```

[illegible]

[illegible]

[illegible]

[illegible] 4 [illegible]

[illegible]

Table 1. [illegible] $n = 500$[illegible] 500 [illegible]

		reg	HT	Hajek	DR
[illegible]	1[illegible]				
[illegible]		0.00	0.02	0.03	0.01
[illegible]		0.11	0.28	0.26	0.13
[illegible]		0.10	0.25	0.23	0.12
[illegible]	2[illegible]				
[illegible]		0.00	−0.76	−0.75	−0.01
[illegible]		0.12	0.59	0.47	0.18
[illegible]		0.13	0.50	0.38	0.18
[illegible]	3[illegible]				
[illegible]		−0.05	0.00	−0.01	0.00
[illegible]		0.11	0.15	0.14	0.14
[illegible]		0.11	0.14	0.13	0.14
[illegible]	4[illegible]				
[illegible]		−0.08	0.11	−0.07	0.16
[illegible]		0.13	0.32	0.20	0.41
[illegible]		0.13	0.25	0.16	0.26

CHAPTER 13

[1]

##

"Block what you can and randomize what you cannot." — (page 103)

SRE

SRE X

treatment control

$e(X) = \Pr(Z = 1)$

$e(X) = \Pr(Z = 1 \mid X)$ X X

X X

X curse of dimensionality

SRE

confounding

$e_{[k]} = n_{[k]1}/n_{[k]}$

13.1 ( ). 倾向得分定义为给定协变量条件下治疗分配的条件概率：

$$e(X) = \Pr(Z = 1 \mid X).$$

$[0, 1]$

X ignorability X

( )如果 $Z \perp \{Y(1), Y(0)\} \mid X$，则有

$$Z \perp \{Y(1), Y(0)\} \mid e(X).$$

ignorability X

$e(X)$

X X $e(X)$

[1]

---

[1] **??**

$\Pr\{Z=1 \mid Y(1),Y(0),e(X)\} = \Pr\{Z=1 \mid e(X)\}$ tower property ignorability

$$\Pr\{Z=1 \mid Y(1),Y(0),e(X)\} = E\{Z \mid Y(1),Y(0),e(X)\} = E\big[E\{Z \mid Y(1),Y(0),X\} \mid Y(1),Y(0),e(X)\big].$$

ignorability $E\{Z \mid Y(1),Y(0),X\} = E\{Z \mid X\} = e(X)$

$$\Pr\{Z=1 \mid Y(1),Y(0),e(X)\} = E\{e(X) \mid Y(1),Y(0),e(X)\} = e(X).$$

$\Pr\{Z=1 \mid e(X)\} = E\{Z \mid e(X)\} = E\{e(X) \mid e(X)\} = e(X)$

Rosenbaum and Rubin 1983 *Biometrika*

Biometrika

ignorability X $e(X)$

$e(X)$ K $\{e_1,\ldots,e_K\}$ K n

$$Z \perp \{Y(1),Y(0)\} \mid e(X)=e_k \quad (k=1,\ldots,K).$$

k K CRE SRE

$$\tau_{[k]} = E(Y \mid Z=1, e(X)=e_k) - E(Y \mid Z=0, e(X)=e_k),$$

$$\tau = \sum_{k=1}^{K} \pi_{[k]}\tau_{[k]},$$

$\pi_{[k]} = \Pr\{e(X)=e_k\}$ k

$e(X) = \Pr(Z=1 \mid X)$ X logistic $\hat{e}(X)$ n

$$\hat{e}'(X_i) = e_k, \quad \hat{e}(X_i) \quad (k-1)/K \quad k/K \quad .$$

$$Z \perp \{Y(1),Y(0)\} \mid \hat{e}'(X)=e_k \quad (k=1,\ldots,K).$$

[illegible] ignorability [illegible] $\hat{e}'(X)$ [illegible] { {[illegible] SRE [illegible]

[illegible]

[illegible] K [illegible]

[illegible] K [illegible]

- [illegible] K [illegible] $Z \perp \{Y(1), Y(0)\} \mid \hat{e}'(X)$ [illegible]
- [illegible] K [illegible]

Rosenbaum and Rubin[illegible] 1983, 1984[illegible] Cochran[illegible] 1968[illegible] $K = 5$ [illegible] $K = 5$ [illegible]

[illegible] $K = 5$ [illegible] { [illegible] K [illegible] { [illegible] K [illegible]

[illegible]

[illegible] { {[illegible] SRE[illegible] SRE [illegible]

(1) [illegible] SRE [illegible] X [illegible] CRE[illegible] $e(X)$ [illegible] ” [illegible] ” [illegible]

(2) [illegible] SRE [illegible] $e_{[k]} = n_{[k]1}/n_{[k]}$ [illegible] e_k [illegible]

(3) [illegible]

[illegible] SRE [illegible] Neyman [illegible] Fisher [illegible]

[illegible]

[illegible] $\hat{e}'(X)$ [illegible] SRE [illegible]

[illegible] **B p p** [illegible]

[illegible]

[illegible]

[illegible] NHANES [illegible]

[illegible] Schoolmeal [illegible] BMI [illegible]

[illegible] $Z_i = 1$ [illegible] $Z_i = 0$ [illegible]

[illegible] BMI [illegible]

[illegible] logistic [illegible]

$$\hat{e}(X_i) = \mathrm{logit}^{-1}(\hat{\alpha}_0 + \hat{\alpha}_1 X_{i1} + \cdots + \hat{\alpha}_p X_{ip}).$$

[illegible] $K = 5, 10, 30$ [illegible]

FIGURE 1. NHANES [illegible]

[illegible] K [illegible] $K \in \{5, 10, 20, 50, 80\}$ [illegible][2]

TABLE 1. [illegible] K [illegible]

	$K = 5$	$K = 10$	$K = 20$	$K = 50$	$K = 80$
[illegible]	-0.116	-0.178	-0.200	-0.265	$-$
[illegible]	0.283	0.282	0.279	0.272	NA

[illegible]

(1) [illegible] K [illegible] 5 [illegible] 50 [illegible] -0.2[illegible]

(2) [illegible] $K = 80$ [illegible] NA [illegible]

(3) [illegible] BMI [illegible]

[illegible]

TABLE 2. [illegible]

	naive	Fisher	Lin
[illegible]	0.534	0.061	-0.017
[illegible]	0.225	0.227	0.226

[illegible]

- [illegible] 0.534 [illegible]
- [illegible] OLS [illegible] 0.061 [illegible]
- [illegible] OLS [illegible] -0.017 [illegible]
- [illegible] -0.17 [illegible] -0.27 [illegible] Lin [illegible] K [illegible]

---

[2] [illegible] ??

[illegible] SRE [illegible]

[illegible]

[illegible]

(1) [illegible] ignorability [illegible] $e(X)$ [illegible] X [illegible]

(2) [illegible] K [illegible] ” [illegible] ” [illegible]

(3) [illegible] SRE [illegible]

(4) K [illegible] $K = 5$ [illegible]

(5) [illegible]

[illegible] doubly robust [illegible]

CHAPTER 14

[illegible]

[illegible]

[illegible] IPW[illegible] Hajek [illegible]

$$\hat{\tau}^{\text{hajek}} = \frac{\sum_{i=1}^{n} \frac{Z_i Y_i}{\hat{e}(X_i)}}{\sum_{i=1}^{n} \frac{Z_i}{\hat{e}(X_i)}} - \frac{\sum_{i=1}^{n} \frac{(1-Z_i)Y_i}{1-\hat{e}(X_i)}}{\sum_{i=1}^{n} \frac{1-Z_i}{1-\hat{e}(X_i)}} \tag{64}$$

[illegible] Imbens, 2004[illegible] { [illegible] WLS[illegible] $\hat{\tau}^{\text{hajek}}$ [illegible] WLS [illegible] Z_i [illegible]

$$(\hat{\alpha}, \hat{\beta}) = \arg\min_{\alpha,\beta} \sum_{i=1}^{n} w_i (Y_i - \alpha - \beta Z_i)^2$$

[illegible]

$$w_i = \frac{Z_i}{\hat{e}(X_i)} + \frac{1-Z_i}{1-\hat{e}(X_i)} = \begin{cases} \frac{1}{\hat{e}(X_i)} & \text{if } Z_i = 1; \\ \frac{1}{1-\hat{e}(X_i)} & \text{if } Z_i = 0. \end{cases} \tag{65}$$

[illegible] IPW [illegible]

[illegible]

[illegible]

[illegible] 14.1 [illegible] Hajek [illegible] WLS)[illegible]在 [illegible]的记号下，$\hat{\tau}^{hajek}$ 严格等于上述 *WLS* 回归中 Z_i 的系数 $\hat{\beta}$。

[illegible] CRE[illegible] $e(X_i) \equiv p$ [illegible] w_i [illegible] WLS [illegible] OLS [illegible] CRE OLS [illegible] τ [illegible] $1/\hat{e}(X_i)$ [illegible] $1/\{1-\hat{e}(X_i)\}$ [illegible] " [illegible] " [illegible] Hajek [illegible] OLS [illegible] treatment [illegible] { [illegible] WLS [illegible] $\hat{\tau}^{\text{hajek}}$ [illegible] WLS [illegible] $\hat{\tau}^{\text{hajek}}$ [illegible] Bootstrap [illegible][1]

[illegible]

---

[1] [illegible] **??**

∼ [illegible] f a l t ‖ [illegible] q ∼ u

CRE [illegible] Lin (2013) [illegible] OLS [illegible] treatment [illegible]

$$Y_i \sim (1, Z_i, X_i, Z_i X_i)$$

[illegible] $\bar{X} = 0$ [illegible] OLS [illegible] WLS [illegible] { [illegible]
1 $\hat{\tau}$ [illegible] WLS [illegible]

$$Y_i \sim (1, Z_i, X_i, Z_i X_i)$$

Z_i [illegible] Hirano and Imbens (2001) [illegible]

∼ [illegible] [illegible]

$$\mathbb{E}(Y \mid Z = 1, X) = \beta_{10} + \beta_{1x}^{\top} X, \quad \mathbb{E}(Y \mid Z = 0, X) = \beta_{00} + \beta_{0x}^{\top} X$$

[illegible]

(1) [illegible] WLS [illegible] OLS [illegible] τ [illegible] Z [illegible] τ

(2) [illegible] WLS Z [illegible] τ [illegible]

[illegible] N p ‘ B g Q p B ‘ 。 — l Q 。 。

[illegible] Robins et al. (2007) [illegible] M. Joffe [illegible]

∼ [illegible] $\hat{e}(X_i, \hat{\alpha})$ [illegible] $(\mu_1(X_i, \hat{\beta}_1), \mu_0(X_i, \hat{\beta}_0))$
1 WLS [illegible]

$$\hat{\tau}_{\text{wls}}^{\text{reg}} = \frac{1}{n} \sum_{i=1}^{n} \mu_1(X_i, \hat{\beta}_1) - \frac{1}{n} \sum_{i=1}^{n} \mu_0(X_i, \hat{\beta}_0)$$

[illegible]

$$\hat{\tau}_{\text{wls}}^{\text{dr}} = \hat{\tau}_{\text{wls}}^{\text{reg}} + \frac{1}{n} \sum_{i=1}^{n} \frac{Z_i\{Y_i - \mu_1(X_i, \hat{\beta}_1)\}}{\hat{e}(X_i, \hat{\alpha})} - \frac{1}{n} \sum_{i=1}^{n} \frac{(1 - Z_i)\{Y_i - \mu_0(X_i, \hat{\beta}_0)\}}{1 - \hat{e}(X_i, \hat{\alpha})}$$

[illegible] WLS [illegible] Z_i [illegible]

[illegible] q X w [illegible] 14.2 [illegible]) 若 $\bar{X} = 0$ 且

$$(\mu_1(X_i, \hat{\beta}_1), \mu_0(X_i, \hat{\beta}_0)) = (\hat{\beta}_{10} + \hat{\beta}_{1x}^{\top} X_i,\ \hat{\beta}_{00} + \hat{\beta}_{0x}^{\top} X_i)$$

基于权重 { 对 $Y_i \sim (1, Z_i, X_i, Z_i X_i)$ 的 WLS 拟合，则

$$\hat{\tau}_{wls}^{dr} = \hat{\tau}_{wls}^{reg} = \hat{\beta}_{10} - \hat{\beta}_{00}$$

即 Z_i 在 WLS 回归中的系数。

[illegible] $\hat{s}$[2]
WLS [illegible]

$$\sum_{i=1}^{n} \frac{Z_i(Y_i - \hat{\beta}_{10} - \hat{\beta}_{1x}^{\top} X_i)}{\hat{e}(X_i)} = 0, \quad \sum_{i=1}^{n} \frac{(1 - Z_i)(Y_i - \hat{\beta}_{00} - \hat{\beta}_{0x}^{\top} X_i)}{1 - \hat{e}(X_i)} = 0.$$

[illegible] $\hat{\tau}^{\mathrm{dr}}_{\mathrm{wls}} - \hat{\tau}^{\mathrm{reg}}_{\mathrm{wls}} = 0$ [illegible] $\bar{X} =$ [illegible] $\hat{\beta}_{10} - \hat{\beta}_{00}$

[illegible] Freedman and Berk (2008)[1] [illegible]

[illegible] WLS [illegible] WLS [illegible] OLS [illegible]

[illegible] WLS [illegible] OLS [illegible] Freedman and Berk (2008) [illegible] WLS [illegible] $\hat{s}$ WLS [illegible] Bootstrap [illegible]

14.1. [illegible] ***WLS*** [illegible]：当（*1*）倾向得分模型可信且（*2*）误差方差可能与倾向得分相关（异方差）时，*WLS* 是更好的选择。当结局模型可信时，简单的 *OLS* 可能更稳定。

[illegible]

[illegible] Average Causal Effect on the Treated, τ_T [illegible]

[illegible] Hajek [illegible]

$$\hat{\tau}_T^{\mathrm{hajek}} = \hat{\bar{Y}}(1) - \frac{\sum_{i=1}^{n} \hat{o}(X_i)(1 - Z_i) Y_i}{\sum_{i=1}^{n} \hat{o}(X_i)(1 - Z_i)}$$

[illegible] $\hat{o}(X_i) = \hat{e}(X_i)/\{1 - \hat{e}(X_i)\}$

[illegible] 14.2 [illegible] Hajek [illegible] WLS) $\hat{\tau}_T^{hajek}$ 严格等于如下 *WLS* 回归中 Z_i 的系数：

$$(\hat{\alpha}, \hat{\beta}) = \arg\min_{\alpha,\beta} \sum_{i=1}^{n} w_{T_i}(Y_i - \alpha - \beta Z_i)^2$$

其中权重为

$$w_{T_i} = Z_i + (1 - Z_i)\hat{o}(X_i) = \begin{cases} 1 & \text{if } Z_i = 1; \\ \hat{o}(X_i) & \text{if } Z_i = 0. \end{cases} \tag{66}$$

[illegible] $\hat{\bar{X}}(1) =$ [illegible] τ_T [illegible] WLS [illegible]

$$Y_i \sim (1, Z_i, X_i, Z_i X_i)$$

[illegible] Z_i [illegible]

---

[2] [illegible] ??

[illegible] 14.3 [illegible]）若 $\hat{\hat{X}}(1)=0$ 且 $\mu_0(X_i,\hat{\beta}_0)=\hat{\beta}_{00}+\hat{\beta}_{0x}^{\top}X_i$ 基于权重 [illegible] 对 $Y_i \sim (1, Z_i, X_i, Z_iX_i)$ 的 *WLS* 拟合，则

$$\hat{\tau}^{dr}_{T,wls}=\hat{\tau}^{reg}_{T,wls}=\hat{\beta}_{10}-\hat{\beta}_{00}$$

即 Z_i 在 *WLS* 回归中的系数。

[illegible][3]

[illegible]

[illegible] CRE [illegible]

Table 1. [illegible] CRE [illegible]

	CRE	[illegible]
[illegible]	$Y_i \sim (1, Z_i)$	$Y_i \sim (1, Z_i)$ [illegible] w_i [illegible]
[illegible]	$Y_i \sim (1, Z_i, X_i, Z_iX_i)$	$Y_i \sim (1, Z_i, X_i, Z_iX_i)$ [illegible] w_i [illegible]

[illegible] w_i [illegible] CRE [illegible] OLS [illegible] OLS [illegible] WLS [illegible]

[illegible]

[illegible] WLS [illegible]

(1) [illegible] Hajek [illegible] WLS [illegible] treatment [illegible]
(2) [illegible] WLS [illegible] X_i [illegible] Z_iX_i [illegible] Lin, 2013[illegible]
(3) [illegible] WLS [illegible] Robins et al., 2007[illegible]
(4) [illegible] WLS [illegible] OLS [illegible]
(5) [illegible] τ_T [illegible]

WLS [illegible] IPW [illegible]

[illegible]

[illegible] 14.1. [illegible]命题 *14.1* 和 *14.2* 的证明 证明 [illegible]和 [illegible]
[illegible]：这些是附录 *B.3* 中一元 *WLS* 问题的特例。利用 *WLS* 的正规方程即可证明。

[illegible] 14.2. [illegible]预测估计量与双重稳健估计量 另一种结局回归估计量是预测估计量

$$\hat{\tau}^{pred}=\hat{\mu}_1^{pred}-\hat{\mu}_0^{pred}$$

其中

$$\hat{\mu}_1^{pred}=\frac{1}{n}\sum_{i=1}^{n}\{Z_iY_i+(1-Z_i)\mu_1(X_i,\hat{\beta}_1)\},\quad \hat{\mu}_0^{pred}=\frac{1}{n}\sum_{i=1}^{n}\{Z_i\mu_0(X_i,\hat{\beta}_1)+(1-Z_i)Y_i\}.$$

证明：当 $(\mu_1(X_i,\hat{\beta}_1),\mu_0(X_i,\hat{\beta}_1))=(\hat{\beta}_{10}+\hat{\beta}_{1x}^{\top}X_i,\hat{\beta}_{00}+\hat{\beta}_{0x}^{\top}X_i)$ 来自基于权重

$$w_i=Z_i/\hat{o}(X_i)+(1-Z_i)\hat{o}(X_i)$$

---

[3] [illegible] ??

对 $Y_i \sim (1, X_i)$ 的 *WLS* 拟合时，双重稳健估计量等于 $\hat{\tau}^{pred}$。
[illegible] ：*Cao et al. (2009)* 和 *Vermeulen and Vansteelandt (2015)* 从其他理论角度 *motivation* 了上述权重。

[illegible] 14.3. { {*WLS* 的效率性质 讨论在何种情形下 *WLS* 估计量比 *OLS* 估计量更有效。在 *Freedman and Berk (2008)* 的模拟设置中，为什么 *OLS* 表现更好？这与本章讨论的异方差情形有何关联？

[illegible] 14.4. { {推荐阅读 [illegible] ：{ 是关于因果推断中加权方法的权威综述。{ 讨论了协变量调整在随机化实验中的应用。

[illegible] ：*WLS* 与 *IPW* 的等价性最早由 *Imbens (2004)* 明确指出。双重稳健性的概念在 *Robins et al. (2007)* 中得到了系统阐述。

CHAPTER 15

[illegible]

[illegible]

“Matching is a way of making observational studies look like randomized experiments.” — ([illegible])

[illegible] **j ; — J ^ a l Z Q — a j ; — p** [illegible] Rubin [illegible]

[illegible]

[illegible] n_1 [illegible] n_0 [illegible] $i = 1, \ldots, n_1$ [illegible] $m(i)$ [illegible] $X_i = X_{m(i)}$ [illegible] matched-pairs experiment, MPE [illegible]

$$\Pr\{i \text{ [illegible]}, m(i) \text{ [illegible]}\} = \frac{e(X_i)\{1 - e(X_{m(i)})\}}{\sum_j \{e(X_j) + e(X_{m(j)})\}}.$$

$X_i = X_{m(i)}$ [illegible] $e(X_i) = e(X_{m(i)})$ [illegible] $e(X_i)/\{e(X_i) + e(X_i)\} = 1/2$ [illegible]

[illegible] $X_i = X_{m(i)}$ [illegible] FRT [illegible] $|\hat{e}(X_i) - \hat{e}(X_{m(i)})|$ [illegible] $|\mathrm{logit}\{\hat{e}(X_i)\} - \mathrm{logit}\{\hat{e}(X_{m(i)})\}|$ [illegible]

[illegible]

[illegible] Abadie–Imbens (AI) [illegible][1]

$$\{X_i, Z_i, Y_i(1), Y_i(0)\}_{i=1}^{n} \overset{\text{IID}}{\sim} \{X, Z, Y(1), Y(0)\}.$$

---

[1] [illegible] **??**

[illegible] P [illegible] $1-M$ [illegible] i [illegible] $Y_i(1)$ [illegible] Y_i[illegible]

$$\hat{Y}_i(1) = Y_i.$$

[illegible] $Y_i(0)$ [illegible]

$$\hat{Y}_i(0) = M^{-1}\sum_{j=1}^{M} Y_{m_j(i)},$$

[illegible] $m_j(i)$ [illegible] j [illegible]

$$\hat{\tau}^{\mathrm{m}} = n^{-1}\sum_{i=1}^{n} Z_i\{\hat{Y}_i(1)-\hat{Y}_i(0)\} + n^{-1}\sum_{i=1}^{n}(1-Z_i)\{\hat{Y}_i(0)-\hat{Y}_i(1)\}.$$

[illegible] $J(i)$ [illegible] i [illegible] $K_i = |J(i)\cap\{j : Z_j = 1\}|$ [illegible]

[illegible] ([illegible]) 在 M 固定的情况下，当 $n\to\infty$ 时，

$$\sqrt{n}(\hat{\tau}^{\mathrm{m}}-\tau) \Rightarrow \mathcal{N}\{0, V^{\mathrm{m}}(\tau)\},$$

其中偏差来自不精确匹配。[2]

[illegible] bootstrap [illegible] bootstrap[illegible] $\hat{V}^{\mathrm{mbc}}$ [illegible]

$$\hat{\tau}^{\mathrm{mbc}} = \hat{\tau}^{\mathrm{m}} + n^{-1}\sum_{i=1}^{n}\hat{\psi}_i,$$

[illegible] $\hat{\psi}_i$ [illegible][3] [illegible] $\hat{\tau}^{\mathrm{mbc}}$ [illegible] $\hat{\psi}_i$ [illegible] bootstrap [illegible] bootstrap [illegible] $\hat{V}^{\mathrm{mbc}}$[illegible]

[illegible]

[illegible] 1 [illegible]

[illegible] ([illegible]ACE [illegible]) 偏差校正匹配估计量满足

$$\hat{\tau}^{\mathrm{mbc}} = \hat{\tau}^{\mathrm{DR}},$$

其中 $\hat{\tau}^{\mathrm{DR}}$ 是第 [illegible]章定义的双稳健估计量。[4]

---

[2] [illegible] ??

[3] [illegible] ??

[4] [illegible] ??

[illegible] $1+K_i/M$ [illegible] $1/\hat{e}(X_i)$ [illegible] $e(X_i)$ [illegible] $1/\hat{e}(X_i)$ [illegible] K_i [illegible] $1+K_i/M$ [illegible] M [illegible] $1+K_i/M$ [illegible] $1/e(X_i)$ [illegible] M [illegible] M [illegible]

[illegible]

[illegible]

$$\tau_{\mathrm{T}} = \mathbb{E}\{Y_i(1) - Y_i(0) \mid Z_i = 1\}.$$

ATT [illegible]

$$\hat{\tau}_{\mathrm{T}}^{\mathrm{m}} = n_1^{-1} \sum_{i:Z_i=1} \{\hat{Y}_i(1) - \hat{Y}_i(0)\},$$

[illegible] $\hat{Y}_i(1) = Y_i$ [illegible] $\hat{Y}_i(0) = M^{-1}\sum_{j=1}^{M} Y_{m_j(i)}$ [illegible]

$$\hat{\tau}_{\mathrm{T}}^{\mathrm{mbc}} = \hat{\tau}_{\mathrm{T}}^{\mathrm{m}} + n_1^{-1} \sum_{i:Z_i=1} \hat{\psi}_i^{\mathrm{T}}.$$

[illegible] (ATT [illegible]) *ATT* 的偏差校正匹配估计量满足

$$\hat{\tau}_{\mathrm{T}}^{\mathrm{mbc}} = \hat{\tau}_{\mathrm{T}}^{\mathrm{DR}},$$

其中 $\hat{\tau}_{\mathrm{T}}^{\mathrm{DR}}$ 是 *ATT* 的双稳健估计量。[5]

[illegible] K_i/M [illegible]

[illegible]

[illegible] M [illegible] M [illegible]

[illegible]

[illegible] OLS [illegible]

[illegible]

[illegible] ATT [illegible] \$1794 [illegible] \$623 [illegible][6]

[illegible] Matching [illegible] 1-1 [illegible] ATT [illegible] \$1658 [illegible] \$1794 [illegible]

---

[5] [illegible] ??

[6] [illegible] ?? ?? [illegible]

[illegible]

[illegible]

(1) [illegible]

(2) [illegible]

(3) [illegible]

(4) [illegible]

(5) [illegible] ATT [illegible]

(6) [illegible]

mw

[illegible] 15.1. 匹配与倾向得分 证明：在精确匹配下，倾向得分对所有配对单元相同，即 $e(X_i) = e(X_{m(i)})$。这如何支持”精确匹配模拟随机实验”的论断？

[illegible] 15.2. 一对一与一对多匹配的偏差比较 考虑 $1-1$ 匹配和 $1-M$ 匹配。当 M 增大时，匹配估计量的偏差如何变化？为什么？

[illegible] 15.3. 匹配估计量的偏差校正项 从 { {出发，推导偏差校正项 $\hat{\psi}_i$ 的具体形式。提示：参考 { 的推导。

[illegible] 15.4. *ATT* 匹配估计量的性质 证明 { :{ *ATT* 的偏差校正匹配估计量等于 *ATT* 的双稳健估计量。

[illegible] 15.5. 协变量平衡 对 *LaLonde* 数据进行匹配分析，检验匹配前后协变量的平衡性。使用不同的 M 值（一对一、一对二、一对四）比较结果。

[illegible] 15.6. 匹配与回归的比较 使用第 { {章 { {节的数据，分别用匹配估计量和结果回归估计量估计 *ATT*。比较两种方法的结果和效率。

[illegible] 15.7. 倾向得分匹配的理论性质 { 提供了基于倾向得分匹配的理论。阅读该论文，总结其主要结论及其与 { 建议的关联。

[illegible] 15.8. 高维协变量的挑战 当协变量维度很高时，直接匹配面临维数灾难。讨论三种可能的解决策略：降维、惩罚回归、机器学习方法。

CHAPTER 16

“Although we cannot see the wind, we can see its effects on the leaves.” —

$$Z \perp \{Y(1), Y(0)\} \mid X. \tag{16.1}$$

X Z

George Box “所有模型都是错的，但有些有用。”

X

U

Pearl (1995)

$$X \to Z \to Y$$

X Z Y X Z Z Y

$$\begin{cases} X \sim F_X(x), \\ Z = f_Z(X, \varepsilon_Z), \\ Y(z) = f_Y(X, z, \varepsilon_Y(z)), \end{cases} \tag{16.2}$$

$\varepsilon_Z \perp \varepsilon_Y(z)$ $Z \perp Y(z) \mid X$

U

$$U \to Z,\ U \to Y,\ X \to Z,\ X \to Y$$

[illegible]

$$\begin{cases} X \sim F_X(x), \\ U \sim F_U(u), \\ Z = f_Z(X, U, \varepsilon_Z), \\ Y(z) = f_Y(X, U, z, \varepsilon_Y(z)), \end{cases} \tag{16.3}$$

[illegible] $\varepsilon_Z \perp \varepsilon_Y(z)$ [illegible] $Z \perp Y(z) \mid (X, U)$ [illegible] $Z \not\perp Y(z) \mid X$ [illegible] (X, U) [illegible] X [illegible] U [illegible]

[illegible] Z [illegible] Y [illegible] backdoor path [illegible]

[illegible]

[illegible]

$$Z \perp Y(1) \mid X, \quad Z \perp Y(0) \mid X. \tag{16.4}$$

[illegible]

$$\Pr\{Y(1) \mid Z = 1, X\} = \Pr\{Y(1) \mid Z = 0, X\},$$

$$\Pr\{Y(0) \mid Z = 1, X\} = \Pr\{Y(0) \mid Z = 0, X\}. \tag{16.5}$$

[illegible] X [illegible] $\Pr\{Y(1) \mid Z = 0, X\}$ [illegible] $\Pr\{Y(0) \mid Z = 1, X\}$ [illegible]

[illegible]

[illegible]

[illegible]

[illegible] $Z \perp Y(z) \mid X$ [illegible] $Z \perp Y^{\mathrm{n}}(z) \mid X$ [illegible] Y^{n} [illegible] Y [illegible] Z [illegible] Y^{n} [illegible] Y^{n} [illegible]

$$\tau(Z \to Y^{\mathrm{n}}) = \mathbb{E}\{Y^{\mathrm{n}}(1) - Y^{\mathrm{n}}(0)\} \tag{16.6}$$

[illegible] $\tau(Z \to Y^{\mathrm{n}}) = 0$ [illegible]

[illegible] Z [illegible] Y^{n} [illegible]

[illegible] 16.1 (Cornfield [illegible] (1959)[illegible]). *Cornfield* 等 *(1959)* 研究了吸烟对肺癌的因果效应。他们在分析中控制了众多重要的背景变量，但仍可能存在某些未测混杂因素使观察到的效应产生偏倚。

为了加强因果关系的证据，他们同时报告了吸烟对车祸的影响——基于生物学知识，吸烟对车祸应该没有因果效应。结果显示关联接近于零。因此，即使无法完全排除未测混杂，这个基于阴性结果的补充分析使吸烟对肺癌因果效应的证据更加有力。

[illegible] 16.2 (Jackson [illegible] (2006)[illegible]). *Jackson* 等 *(2006)* 对观测性研究的结果持怀疑态度。这些研究显示，在调整了测量的协变量后，流感疫苗接种显著降低了流感季节的肺炎/流感住院率和全因死亡率。

他们的补充分析考察了流感季节开始之前的结局指标。疫苗接种通常在秋季开始，但流感传播在冬季之前是极少的。基于生物学，疫苗效应在流感季节应该最为显著。但 *Jackson* 等发现，疫苗效应在流感季节之前反而更大——这暗示观察到的效应可能源于未测混杂。

[illegible] Z [illegible] U [illegible] Y[illegible] Z [illegible] U [illegible] Y [illegible]

Lipsitch [illegible] (2010) [illegible] Rosenbaum (1989) [illegible]

[illegible] Z^{n} [illegible] Z [illegible] Z^{n} [illegible] Y [illegible]

$$\tau(Z^{\mathrm{n}} \to Y) = \mathbb{E}\{Y(1^{\mathrm{n}}) - Y(0^{\mathrm{n}})\} \quad (16.7)$$

[illegible] 16.3 (Sanderson [illegible] (2017)[illegible]). *Sanderson* 等 *(2017)* 给出了利用阴性暴露的多个例子。他们建议比较母体暴露与后代结局的关联，以及父亲暴露与同样结局的关联。

在母体吸烟与后代结局的研究中：母体吸烟可能受到未测混杂（如社会经济地位）的影响，而父亲吸烟不太可能受到同样的混杂影响（因为父亲不经历怀孕）。因此，如果观察到的母体吸烟效应远大于父亲吸烟效应，这支持了因果解释；如果两者效应相似，则暗示可能存在混杂。

[illegible] Z [illegible] Z^{n} [illegible] Y [illegible]

[illegible]

[illegible]

[illegible]

Rosenbaum (2002b) [illegible] “没有理由避免调整描述研究对象治疗前特征的变量。”[illegible] Rubin (2007) [illegible] “通常，条件性越强的假设，越容易接受。”[illegible]

Pearl [illegible]

---

[1] [illegible]

[illegible] M [illegible]

$$U_1 \to X \leftarrow U_2,\ U_1 \to Z,\ U_2 \to Y$$

[illegible] M [illegible] M [illegible]

$$\begin{cases} U_1 \perp U_2, \\ X = f_X(U_1, U_2, \varepsilon_X), \\ Z = f_Z(U_1, \varepsilon_Z), \\ Y = Y(z) = f_Y(U_2, \varepsilon_Y), \end{cases} \tag{16.8}$$

[illegible] $(\varepsilon_X, \varepsilon_Z, \varepsilon_Y)$ [illegible] X [illegible] U_1 [illegible] U_2 [illegible] Z [illegible] Y [illegible] Z [illegible] Y [illegible] Z [illegible] Y [illegible]

[illegible] $Z \perp Y$ [illegible] Z [illegible] Y [illegible] X [illegible]

$$\tau_{\mathrm{PF}} = \mathbb{E}(Y \mid Z = 1) - \mathbb{E}(Y \mid Z = 0) = 0. \tag{16.9}$$

[illegible] X [illegible] X [illegible] U_1 [illegible] U_2 [illegible] $U_1 \not\perp U_2 \mid X$ [illegible] $Z \not\perp Y \mid X$ [illegible]

$$\int \{\mathbb{E}(Y \mid Z = 1, X = x) - \mathbb{E}(Y \mid Z = 0, X = x)\} f(x)\, dx \neq 0. \tag{16.10}$$

[illegible] X [illegible] U_1 [illegible] U_2 [illegible] U_1 [illegible] Z [illegible] U_2 [illegible] Y [illegible] X [illegible] X [illegible] Z [illegible] Y [illegible] X [illegible] U_1 [illegible] U_2 [illegible] Z [illegible] Y [illegible]

[illegible]

$$\begin{cases} X = aU_1 + bU_2 + \varepsilon_X, \\ Z = cU_1 + \varepsilon_Z, \\ Y = dU_2 + \varepsilon_Y, \end{cases} \tag{16.11}$$

[illegible] $(U_1, U_2, \varepsilon_X, \varepsilon_Z, \varepsilon_Y) \overset{\mathrm{IID}}{\sim} \mathrm{N}(0,1)$ [illegible]

$$\mathrm{cov}(Z, Y) = \mathrm{cov}(cU_1 + \varepsilon_Z, dU_2 + \varepsilon_Y) = 0, \tag{16.12}$$

[illegible] X [illegible] Z [illegible] Y [illegible]

$$\rho_{ZY|X} \propto -abcd, \tag{16.13}$$

[illegible] $abcd$ [illegible]

[illegible] 16.4 (M-[illegible]). 考虑样本量 $n = 10^6$ 的大规模蒙特卡洛模拟：

```
U1 = rnorm(n)
U2 = rnorm(n)
X = U1 + U2 + rnorm(n)
Y = U2 + rnorm(n)
Z = U1 + rnorm(n)

# 连续治疗
```

```
summary(lm(Y ~ Z))$coef[2, 1]  # [1] 0.001
summary(lm(Y ~ Z + X))$coef[2, 1]  # [1] -0.20

# 二值治疗
Z = (Z >= 0)
summary(lm(Y ~ Z))$coef[2, 1]  # [1] 0.002
summary(lm(Y ~ Z + X))$coef[2, 1]  # [1] -0.42
```

未调整的简单估计量接近真实值 (*0*)，但调整 X 后的估计量产生了显著偏倚 (约 -0.2 到 -0.4)。这就是 M-偏倚。

M-[illegible] U_1 [illegible] U_2 [illegible] X [illegible] U_1 [illegible] U_2 [illegible]

[illegible] M-[illegible]

$$Z \leftarrow X \rightarrow U \rightarrow Y$$

[illegible] X [illegible] Z [illegible] U [illegible] U [illegible] Z [illegible] Y [illegible]

$$\begin{cases} Z = aX + bU + \varepsilon_Z, \\ Y(z) = \tau z + cU + \varepsilon_Y, \end{cases} \tag{16.14}$$

[illegible] $(U, X, \varepsilon_Z, \varepsilon_Y) \overset{\text{IID}}{\sim} \mathrm{N}(0,1)$ [illegible] $X \perp U$ [illegible] X [illegible] Z [illegible] Y [illegible] X [illegible] Z [illegible] Y [illegible] U [illegible] Z [illegible] X [illegible] U [illegible] X [illegible] Z [illegible] U [illegible] U [illegible]

[illegible] OLS [illegible]

$$\tau_{\text{unadj}} = \tau + \frac{c\ \mathrm{cov}(Z, U)}{\mathrm{var}(Z)} = \tau + \frac{cb}{a^2 + b^2 + 1}, \tag{16.15}$$

[illegible] $bc/(a^2 + b^2 + 1)$ [illegible] OLS [illegible] Y [illegible] (Z, X) [illegible]

$$\tau_{\text{adj}} = \tau + \frac{bc}{b^2 + 1}. \tag{16.16}$$

[illegible] $a \neq 0$ [illegible] $a^2 + b^2 + 1 > b^2 + 1$ [illegible]

$$\left|\frac{bc}{a^2 + b^2 + 1}\right| < \left|\frac{bc}{b^2 + 1}\right|. \tag{16.17}$$

[illegible] X [illegible] Z [illegible] a [illegible]

[illegible] 16.5 (Z-[illegible]). 考虑蒙特卡洛模拟:

```
n = 10^6
X = rnorm(n)
U = rnorm(n)
Z = X + U + rnorm(n)
Y = U + rnorm(n)

# Z ~ X + U
```

```
summary(lm(Y ~ Z))$coef[2, 1]  # [1] 0.333
summary(lm(Y ~ Z + X))$coef[2, 1]  # [1] 0.501

# X-Z 关联更强时
Z = 2*X + U + rnorm(n)
summary(lm(Y ~ Z))$coef[2, 1]  # [1] 0.167
summary(lm(Y ~ Z + X))$coef[2, 1]  # [1] 0.501

# X-Z 关联更强时
Z = 10*X + U + rnorm(n)
summary(lm(Y ~ Z))$coef[2, 1]  # [1] 0.01
summary(lm(Y ~ Z + X))$coef[2, 1]  # [1] 0.50
```

可以看到：随着 X 与 Z 的关联增强，未调整估计量越来越接近真实值 $\tau = 0$，而调整后的估计量始终为 0.5——因为调整反而引入了更大的偏倚。

Z-[illegible] Problem [illegible] Pearl (2010a, 2011) [illegible] Ding [illegible] (2017b) [illegible]

[illegible] Pearl [illegible] Z [illegible] X [illegible] X [illegible] Z [illegible] Z-[illegible] [illegible]

[illegible] [illegible]

$$X_Y \to Y,\, X_Y \to Z,\, X \to Z,\, X \to Y,\, X_I \to Z,\, X_I \to Y$$

[illegible]

(1) X [illegible] X [illegible] X

(2) X_R [illegible]

(3) X_Z [illegible] Z-[illegible] X_Z [illegible]

(4) X_Y [illegible]

(5) X_I [illegible]

[illegible] X [illegible] X_Y [illegible]

[illegible]

[illegible] 16.1. 验证正文中的公式 [illegible]和 [illegible]

TABLE 1. [illegible]

[illegible]	Z [illegible]	Y [illegible]	
X [illegible]	[illegible]	[illegible]	[illegible]
X_R [illegible]	[illegible]	[illegible]	[illegible]
X_Z [illegible]	[illegible]	[illegible]	[illegible]
X_Y [illegible]	[illegible]	[illegible]	[illegible]
X_I [illegible]	[illegible]	[illegible]	[illegible]

[illegible] 16.2. 考虑正文中的 M-偏倚正态线性模型。证明给定 X 时 Z 和 Y 的偏相关系数公式 [illegible]

[illegible] 16.3. 找出一个你自己的研究领域中的阴性结果或阴性暴露的例子。讨论为什么它符合阴性控制的条件，以及它如何帮助评估不可混淆性假设。

[illegible] 16.4. [illegible] 和 *econometricians* 提出了以下 ***Cochran*** [illegible] （也称为遗漏变量偏倚公式）。设 $(y_i, x_{1i}, x_{2i})_{i=1}^n$ 为 *IID* 样本，其中 y_i 是标量，x_{1i} 维数为 K，x_{2i} 维数为 L。考虑 *OLS* 分解：

$$\begin{aligned} y_i &= \beta_1^{\mathrm{T}} x_{1i} + \beta_2^{\mathrm{T}} x_{2i} + \varepsilon_i \quad \text{（长回归）}, \\ y_i &= \gamma^{\mathrm{T}} x_{1i} + e_i \quad \text{（短回归）}, \\ x_{2i} &= \delta^{\mathrm{T}} x_{1i} + v_i. \end{aligned}$$

证明：$\gamma = \beta_1 + \delta\beta_2$。解释为什么 $\delta\beta_2$ 被称为遗漏变量偏倚。

[illegible] 16.5. 设计一个模拟研究，演示以下情形：

(1) 调整混杂变量 X 可以减少偏倚；
(2) 调整与未测混杂相关的变量会引入 M-偏倚；
(3) 调整工具变量 X_Z 在存在未测混杂时会放大偏倚。

对于每种情形，使用大样本量（$n = 10^5$）计算未调整和调整后的估计量。

[illegible]

[illegible] Pearl (2000) [illegible]
VanderWeele [illegible] Shpitser (2011) [illegible]
Rosenbaum (2002b) [illegible]
[illegible]
Ding [illegible] (2017b) [illegible] Z-[illegible]

CHAPTER 17

[illegible]

[illegible]

"It is possible that the entire relationship between cigarette smoking and lung cancer is due to some common cause." — ([illegible])

Part III [illegible] outcome regression [illegible] [illegible] [illegible] [illegible]

$$Z \perp \{Y(1), Y(0)\} \mid X. \tag{67}$$

[illegible] X [illegible] Z [illegible]

[illegible] VanderWeele and Ding, 2017 [illegible] [1]

[illegible]

[illegible]

$$Z \perp \{Y(1), Y(0)\} \mid (X, U), \tag{68}$$

[illegible] U [illegible] (X, U) [illegible] Z [illegible] Y [illegible] U [illegible] U [illegible] Y [illegible] Ding and VanderWeele, 2016 [illegible] Y [illegible]

[illegible] $X = x$ [illegible]

$$\mathrm{RR}^{\text{true}}_{ZY|x} = \frac{\Pr\{Y(1) = 1 \mid X = x\}}{\Pr\{Y(0) = 1 \mid X = x\}}. \tag{69}$$

---

[1] Ding and VanderWeele (2016) [illegible] Rosenbaum [illegible]

$$\mathrm{RR}^{\mathrm{obs}}_{ZY|x} = \frac{\Pr(Y=1 \mid Z=1, X=x)}{\Pr(Y=1 \mid Z=0, X=x)}. \tag{70}$$

e U U U U X

$$Z \leftarrow U \rightarrow Y,$$

U Z Y common cause $Z \perp Y \mid (X, U)$ $Z \not\perp Y \mid X$ U Ding and VanderWeele (2016) U

$$\mathrm{RR}_{ZU|x} = \frac{\Pr(U=1 \mid Z=1, X=x)}{\Pr(U=1 \mid Z=0, X=x)}, \tag{71}$$

$$\mathrm{RR}_{UY|x} = \frac{\Pr(Y=1 \mid U=1, X=x)}{\Pr(Y=1 \mid U=0, X=x)}, \tag{72}$$

E-value

Cornfield-type 在假设 下，假设

$$RR^{\mathrm{obs}}_{ZY|x} > 1, \quad RR_{ZU|x} > 1, \quad RR_{UY|x} > 1. \tag{73}$$

则有

$$RR^{\mathrm{obs}}_{ZY|x} \le \frac{RR_{ZU|x} \cdot RR_{UY|x}}{RR_{ZU|x} + RR_{UY|x} - 1}. \tag{74}$$

$Z \leftarrow U \rightarrow Y$ $\mathrm{RR}_{ZU|x} \cdot \mathrm{RR}_{UY|x}/(\mathrm{RR}_{ZU|x} + \mathrm{RR}_{UY|x} - 1)$ $\mathrm{RR}^{\mathrm{obs}}_{ZY|x}$ Gastwirth et al., 1998

$$\min(\mathrm{RR}_{ZU|x}, \mathrm{RR}_{UY|x}) \ge \mathrm{RR}^{\mathrm{obs}}_{ZY|x}. \tag{75}$$

Cornfield et al. (1959)

$f_{1,x} = \Pr(U=1 \mid Z=1, X=x)$ $f_{0,x} = \Pr(U=1 \mid Z=0, X=x)$ $\mathrm{RR}^{\mathrm{obs}}_{ZY|x}$ $Z \perp Y \mid (X, U)$ ( ) [1]

---

[1] ??

[illegible]

[illegible] (4)[illegible] $w = \max(\mathrm{RR}_{ZU|x}, \mathrm{RR}_{UY|x})$[illegible]

$$\mathrm{RR}^{\mathrm{obs}}_{ZY|x} \le \frac{w^2}{2w-1}. \tag{76}$$

[illegible] ≤ 1[illegible]

$$w \ge \mathrm{RR}^{\mathrm{obs}}_{ZY|x} + \sqrt{\mathrm{RR}^{\mathrm{obs}}_{ZY|x}(\mathrm{RR}^{\mathrm{obs}}_{ZY|x} - 1)}. \tag{77}$$

[illegible] E-value [illegible]

[illegible] 17.2 (E-value). 给定观测条件风险比 $RR^{\mathrm{obs}}_{ZY|x}$，定义 ***E-value*** 为

$$\text{E-value} = RR^{\mathrm{obs}}_{ZY|x} + \sqrt{RR^{\mathrm{obs}}_{ZY|x}(RR^{\mathrm{obs}}_{ZY|x} - 1)}. \tag{78}$$

[illegible] E-value [illegible] $\max(\mathrm{RR}_{ZU|x}, \mathrm{RR}_{UY|x})$ [illegible] R[illegible] $R <$ E-value [illegible]

[illegible] p [illegible] Fisher p [illegible] p [illegible] p [illegible] E-value [illegible]

[illegible] β[illegible])定义 $\beta(w_1, w_2) = w_1 w_2/(w_1 + w_2 - 1)$（$w_1 > 1, w_2 > 1$）。则：

(1) $\beta(w_1, w_2)$ 对称于 w_1 和 w_2；
(2) $\beta(w_1, w_2)$ 对 w_1 和 w_2 均单调递增；
(3) $\beta(w_1, w_2) \le w_1$ 且 $\beta(w_1, w_2) \le w_2$；
(4) $\beta(w_1, w_2) \le w^2/(2w-1)$，其中 $w = \max(w_1, w_2)$。

[illegible] (3) [illegible] Cornfield [illegible][2] [illegible] (4) [illegible] Cornfield [illegible] E-value [illegible]

[illegible]

[illegible] 17.1 (Hammond and Horn, 1958). 研究美国人群中吸烟与肺癌的关系。不调整协变量的数据可列为下表：

	肺癌	无肺癌
吸烟者	*397*	*78,557*
非吸烟者	*51*	*108,778*

基于该数据，风险比估计为 $\widehat{RR} = 10.73$，*95%* 置信区间为 $[8.02, 14.36]$。

- 解释点估计的 *E-value*：

$$\text{E-value} = 10.73 + \sqrt{10.73 \times 9.73} \approx 20.95.$$

- 解释下限的 *E-value*：

$$\text{E-value}_{\mathrm{CI}} = 8.02 + \sqrt{8.02 \times 7.02} \approx 15.52.$$

---

[2] [illegible] **??**

[illegible]：*要完全解释掉观察到的风险比（甚至其置信下限），未测混杂与治疗的关联以及与结果的关联，都必须达到至少 15.52。这意味着未测混杂必须是一个*[illegible]*的风险因子——它与吸烟的关联程度要接近吸烟本身与肺癌的关联程度（10 倍以上）。在领域知识上，这种强度的未知混杂似乎不太合理，因此该研究提供了较强的因果证据。*

[illegible] Hammond and Horn [illegible] $(\mathrm{RR}_{ZU|x}, \mathrm{RR}_{UY|x})$ [illegible]

[illegible]

[illegible] E-value [illegible] Bradford Hill, 1965 [illegible] Bradford Hill [illegible]

[illegible]

[illegible] E-value [illegible]

[illegible] Bradford Hill [illegible] temporality [illegible] biological plausibility [illegible] consistency [illegible] dose-response [illegible] E-value [illegible]

[illegible] 17.4 (Bradford Hill [illegible]). *([illegible])* 提出的九条因果判定准则：

(1) 强度（*strength*）；
(2) 一致性（*consistency*）；
(3) 特异性（*specificity*）；
(4) 时间顺序（*temporality*）；
(5) 生物学梯度（*biological gradient*）；
(6) 生物学合理性（*plausibility*）；
(7) 一致性（*coherence*）；
(8) 实验证据（*experiment*）；
(9) 类比（*analogy*）。

E-value [illegible] (1) [illegible] E-value [illegible]

[illegible]

$$\Pr(Y_i = 1 \mid Z_i, X_i) = \frac{e^{\beta_0+\beta_1 Z_i+\beta_2\, X_i}}{1 + e^{\beta_0+\beta_1 Z_i+\beta_2\, X_i}}. \tag{79}$$

[illegible] β_1 [illegible] X [illegible] common odds ratio [illegible]

[illegible] $\Pr(Y = 1 \mid Z, X)$ [illegible] β_1 [illegible] $\hat{\beta}_1$ [illegible] E-value[illegible]

$$\widehat{\text{E-value}} = \exp(\hat{\beta}_1) + \sqrt{\exp(\hat{\beta}_1)(\exp(\hat{\beta}_1) - 1)}. \tag{80}$$

[illegible] E-value [illegible]

[illegible] 17.2 (NCHS 2003 [illegible]). 使用美国国家卫生统计中心 *2003* 年出生证明数据，研究高龄母亲（≥ 35 岁）对早产的影响。调整社会人口学协变量后，逻辑斯蒂回归得到的条件风险比为 1.31，其 *95%* 置信区间为 $[1.29, 1.33]$。

对应的 *E-value* 约为 1.94（点估计）和 1.91（置信下限）。这意味着要解释掉观察到的关联，未测混杂必须同时与治疗（高龄）和结果（早产）有至少 *1.9* 倍的关联——这个强度在流行病学上看起来是较大的，但远不如吸烟-肺癌例子中的 *20.95*。

[illegible] [illegible] Ding and VanderWeele (2016) [illegible]

[illegible] **[illegible]** [illegible] 修改 $RR_{UY|x}$ 的定义为：

$$RR_{UY|x} = \max_{z=0,1} \frac{\Pr(Y=1 \mid Z=z, U=1, X=x)}{\Pr(Y=1 \mid Z=z, U=0, X=x)}. \tag{81}$$

则在假设 [illegible] 下，

$$RR^{\text{true}}_{ZY|x} \geq \frac{RR^{\text{obs}}_{ZY|x}}{\frac{RR_{ZU\ x}\cdot RR_{UY\ x}}{RR_{ZU\ x}+RR_{UY\ x}-1}}. \tag{82}$$

[illegible] $\mathrm{RR}^{\text{true}}_{ZY|x}$ [illegible] Cornfield [illegible]

[illegible]

E-value [illegible] Ioannidis et al., 2019 [illegible]

[illegible] [illegible] E-value [illegible] $2 \cdot \mathrm{RR}^{\text{obs}}_{ZY|x}$ [illegible] E-value [illegible]

[illegible] E-value [illegible] E-value [illegible] 10.73 [illegible] E-value 20.95 [illegible]

[illegible] E-value [illegible] $\max(\mathrm{RR}_{ZU|x}, \mathrm{RR}_{UY|x})$ [illegible] E-value [illegible] E-value [illegible] X [illegible] E-value [illegible] X [illegible] X_j [illegible] Z-X_j [illegible] X_j-Y [illegible]

[illegible] [illegible] Ding and VanderWeele (2016) [illegible] HR [illegible] E-value [illegible] outcome regression [illegible] IPW [illegible]

[illegible]

[illegible] E-value [illegible]

(1) [illegible]

(2) [illegible] $\mathrm{RR}^{\mathrm{obs}}_{ZY|x}$ [illegible] $\mathrm{RR}^{\mathrm{obs}}_{ZY|x}$ [illegible]

(3) [illegible] $\text{E-value} = \mathrm{RR}^{\mathrm{obs}}_{ZY|x} + \sqrt{\mathrm{RR}^{\mathrm{obs}}_{ZY|x}(\mathrm{RR}^{\mathrm{obs}}_{ZY|x} - 1)}$ [illegible]

(4) E-value [illegible]

(5) E-value [illegible] Bradford Hill [illegible]

[illegible]

[illegible]

17.1 (17.1). 证明函数 $f(x) = (ax+1)/(bx+1)$ 在 $a > b$ 时关于 x 单调递增，在 $a < b$ 时单调递减。

17.2 (17.2). 证明在假设 $Z \perp Y \mid (X, U)$ 下，如果 $RR_{ZU|x} > 1$ 但 $RR_{UY|x} < 1$，则 $RR^{\mathrm{obs}}_{ZY|x} < 1$。这说明 [illegible] 中的第三个条件在某种意义上是冗余的。

17.3 (17.3). 证明引理 [illegible]

17.4 (17.4). （*Schlesselman 1978* 公式）在二值治疗 Z、结果 Y 和未测混杂 U 下，假设层内（$U = 0$ 和 $U = 1$）治疗对结果的风险比相同，$RR_{ZY|U=0} = RR_{ZY|U=1}$，且混杂对结果的风险比在两组治疗下也相同（记为 γ）。证明：

$$\frac{RR^{\mathrm{obs}}_{ZY}}{RR^{\mathrm{true}}_{ZY}} = \frac{1 + (\gamma - 1)\Pr(U = 1 \mid Z = 1)}{1 + (\gamma - 1)\Pr(U = 1 \mid Z = 0)}. \tag{83}$$

并讨论此公式与定理 [illegible] 的关系。

17.5 (17.5). 使用与例 [illegible] 相同的数据集，计算结果为子痫前期（*preeclampsia*）时的 *E-value*。

17.6 (17.6). （风险差的 *Cornfield* 不等式）考虑二值 Z, Y, U。在潜在可忽略性 $Z \perp Y \mid U$ 下，证明风险差的 *Cornfield* 不等式：

$$RD^{\mathrm{obs}}_{ZY} = RD_{ZU} \times RD_{UY}, \tag{84}$$

并由此推出：

$$\min(RD_{ZU}, RD_{UY}) \geq RD^{\mathrm{obs}}_{ZY}, \quad \max(RD_{ZU}, RD_{UY}) \geq \sqrt{RD^{\mathrm{obs}}_{ZY}}. \tag{85}$$

[illegible]

- [illegible] Cornfield [illegible] E-value [illegible]
- [illegible] E-value [illegible]
- [illegible] Bradford Hill [illegible]
- [illegible] E-value [illegible]

CHAPTER 18

[illegible]

[illegible]

Part III [illegible] outcome regression[illegible] X [illegible] Z [illegible]

$$Z \perp \{Y(1), Y(0)\} \mid X.$$

[illegible] $\tau = \mathbb{E}\{Y(1) - Y(0)\}$ [illegible]

[illegible] Part III [illegible]

[illegible] Manski (1990) [illegible]

[illegible] E-value [illegible] outcome regression [illegible] IPW [illegible]

[illegible]

[illegible]

[illegible] 18.1 ([illegible]). 如果观测数据分布与参数 θ 的多个值相容，则称参数 θ 是[illegible]的。

[illegible] θ [illegible]

[illegible] τ [illegible] τ [illegible]

[illegible] outcomes [illegible]

$$\underline{y} \le Y_i(1), Y_i(0) \le \overline{y}.$$

[illegible] $[\underline{y}, \overline{y}] = [0, 1]$ [illegible] $E\{Y(1)\}$ [illegible] $E\{Y(0)\}$ [illegible] $E\{Y(1)\}$ [illegible]

$$E\{Y(1)\} = E\{Y(1) \mid Z = 1\} \Pr(Z = 1) + E\{Y(1) \mid Z = 0\} \Pr(Z = 0).$$

$Y(1) \in [\underline{y}, \overline{y}]$ [illegible]

$$E\{Y(1) \mid Z = 0\} \in [\underline{y}, \overline{y}].$$

[illegible] $E\{Y(1)\}$ [illegible]

$$E\{Y \mid Z = 1\} \Pr(Z = 1) + \underline{y} \Pr(Z = 0),$$

[illegible]

$$E\{Y \mid Z = 1\} \Pr(Z = 1) + \overline{y} \Pr(Z = 0).$$

[1]

[illegible] $E\{Y(0)\}$ [illegible]

$$\underline{y} \Pr(Z = 1) + E\{Y \mid Z = 0\} \Pr(Z = 0),$$

[illegible]

$$\overline{y} \Pr(Z = 1) + E\{Y \mid Z = 0\} \Pr(Z = 0).$$

[illegible] $\tau = E\{Y(1)\} - E\{Y(0)\}$ [illegible]

$$E\{Y \mid Z = 1\} \Pr(Z = 1) + \underline{y} \Pr(Z = 0) - \overline{y} \Pr(Z = 1) - E\{Y \mid Z = 0\} \Pr(Z = 0),$$

[illegible]

$$E\{Y \mid Z = 1\} \Pr(Z = 1) + \overline{y} \Pr(Z = 0) - \underline{y} \Pr(Z = 1) - E\{Y \mid Z = 0\} \Pr(Z = 0).$$

[illegible] $\overline{y} - \underline{y}$ [illegible] $[\underline{y} - \overline{y}, \overline{y} - \underline{y}]$ [illegible] $2(\overline{y} - \underline{y})$ [illegible] 1

[illegible] Cochran (1953) [illegible] Manski (1990) [illegible] Manski (2003) [illegible]

[illegible]

[illegible] Manski [illegible]

[illegible]

$$\varepsilon_1(X) = \frac{E\{Y(1) \mid Z = 1, X\}}{E\{Y(1) \mid Z = 0, X\}}, \quad \varepsilon_0(X) = \frac{E\{Y(0) \mid Z = 1, X\}}{E\{Y(0) \mid Z = 0, X\}}. \tag{86}$$

[illegible] $\varepsilon_1(X)$ [illegible] X [illegible] $\varepsilon_1(X) = 1$ [illegible] $Y(1)$ [illegible] $Y(1)$ [illegible] X [illegible] $\varepsilon_1(X) = \varepsilon_1$ [illegible] $\varepsilon_0(X) = \varepsilon_0$ [illegible]

[illegible] $\mu_1(X) = E(Y \mid Z = 1, X)$ [illegible] $\mu_0(X) = E(Y \mid Z = 0, X)$ [illegible]

---

[1] [illegible] **??**

~**w** [illegible]**w** **w** [illegible]

[illegible]

[illegible] **q R w**([illegible] I[illegible]$_1$ outcome regression)**w**在已知 $\varepsilon_1(X)$ 和 $\varepsilon_0(X)$ 的条件下，我们有

$$E\{Y(1) \mid Z=0\} = E\left\{\frac{\mu_1(X)}{\varepsilon_1(X)} \mid Z=0\right\},$$
$$E\{Y(0) \mid Z=1\} = E\left\{\mu_0(X)\varepsilon_0(X) \mid Z=1\right\}.$$

因此，

$$\tau = E\left[ZY + (1-Z)\frac{\mu_1(X)}{\varepsilon_1(X)}\right] - E\left[Z\mu_0(X)\varepsilon_0(X) + (1-Z)Y\right] \tag{87}$$
$$= E\left[Z\mu_1(X) + (1-Z)\frac{\mu_1(X)}{\varepsilon_1(X)}\right] - E\left[Z\mu_0(X)\varepsilon_0(X) + (1-Z)\mu_0(X)\right]. \tag{88}$$

[illegible] { [illegible] $\varepsilon_1(X) = \varepsilon_0(X) = 1$ [illegible] {[2] [illegible] outcome regression [illegible]

[1] { [illegible]

r Q N a [illegible]**—a** [illegible]

$$\hat{\tau}^{\text{pred}} = \left\{n^{-1}\sum_{i=1}^{n} Z_iY_i + n^{-1}\sum_{i=1}^{n}(1-Z_i)\frac{\hat{\mu}_1(X_i)}{\varepsilon_1(X_i)}\right\} \tag{89}$$
$$-\left\{n^{-1}\sum_{i=1}^{n} Z_i\hat{\mu}_0(X_i)\varepsilon_0(X_i) + n^{-1}\sum_{i=1}^{n}(1-Z_i)Y_i\right\}. \tag{90}$$

(91)

r p c Q [illegible]**—a** [illegible]

$$\hat{\tau}^{\text{proj}} = \left\{n^{-1}\sum_{i=1}^{n} Z_i\hat{\mu}_1(X_i) + n^{-1}\sum_{i=1}^{n}(1-Z_i)\frac{\hat{\mu}_1(X_i)}{\varepsilon_1(X_i)}\right\} \tag{92}$$
$$-\left\{n^{-1}\sum_{i=1}^{n} Z_i\hat{\mu}_0(X_i)\varepsilon_0(X_i) + n^{-1}\sum_{i=1}^{n}(1-Z_i)\hat{\mu}_0(X_i)\right\}. \tag{93}$$

(94)

[illegible] “predictive” [illegible] “projective” [illegible] Firth and Bennett, 1998; Ding and Li, 2018[illegible]

[illegible]

[illegible] **q R w**([illegible] II[illegible]$_1$ IPW)**w**在已知 $\varepsilon_1(X)$ 和 $\varepsilon_0(X)$ 的条件下，

$$E\{Y(1)\} = E\left\{w_1(X)\frac{Z}{e(X)}Y\right\}, \quad E\{Y(0)\} = E\left\{w_0(X)\frac{1-Z}{1-e(X)}Y\right\},$$

其中

$$w_1(X) = e(X) + \frac{1-e(X)}{\varepsilon_1(X)}, \quad w_0(X) = e(X)\varepsilon_0(X) + 1 - e(X). \tag{95}$$

---

[2] [illegible] **??**

[illegible] IPW [illegible] $w_1(X)$ [illegible] $w_0(X)$ [illegible] $\varepsilon_1(X) = \varepsilon_0(X) = 1$ [illegible] $w_1(X) = w_0(X) = 1$ [illegible] IPW[3] [illegible] Horvitz-Thompson [illegible]

$$\hat{\tau}^{\text{ht}} = n^{-1} \sum_{i=1}^{n} \frac{\{\hat{e}(X_i)\varepsilon_1(X_i) + 1 - \hat{e}(X_i)\} Z_i Y_i}{\varepsilon_1(X_i)\hat{e}(X_i)} \tag{96}$$

$$- n^{-1} \sum_{i=1}^{n} \frac{\{\hat{e}(X_i)\varepsilon_0(X_i) + 1 - \hat{e}(X_i)\}(1 - Z_i) Y_i}{1 - \hat{e}(X_i)}, \tag{97}$$

(98)

[illegible] Hajek [illegible]

$$\hat{\tau}^{\text{haj}} = \frac{\sum_{i=1}^{n} \frac{\{\hat{e}(X_i)\varepsilon_1(X_i)+1-\hat{e}(X_i)\}Z_iY_i}{\varepsilon_1(X_i)\hat{e}(X_i)}}{\sum_{i=1}^{n} \frac{Z_i}{\hat{e}(X_i)}} \tag{99}$$

$$- \frac{\sum_{i=1}^{n} \frac{\{\hat{e}(X_i)\varepsilon_0(X_i)+1-\hat{e}(X_i)\}(1-Z_i)Y_i}{1-\hat{e}(X_i)}}{\sum_{i=1}^{n} \frac{1-Z_i}{1-\hat{e}(X_i)}}. \tag{100}$$

(101)

[illegible] outcome [illegible]

$$\hat{\tau}^{\text{dr}} = \hat{\tau}^{\text{ht}} \tag{102}$$

$$- n^{-1} \sum_{i=1}^{n} \{Z_i - \hat{e}(X_i)\} \left\{ \frac{\hat{\mu}_1(X_i)}{\hat{e}(X_i)\varepsilon_1(X_i)} + \frac{\hat{\mu}_0(X_i)\varepsilon_0(X_i)}{1 - \hat{e}(X_i)} \right\}. \tag{103}$$

(104)

[illegible] $\varepsilon_1(X_i)$ [illegible] $\varepsilon_0(X_i)$ [illegible] outcome [illegible] $\hat{\tau}^{\text{dr}}$ [illegible] τ [illegible] $\varepsilon_1(X_i) = \varepsilon_0(X_i) = 1$ [illegible] Part III [illegible] predictive [illegible] IPW [illegible]

X w [illegible]

X w q w [illegible] **w** R [illegible]

```
OS_est_ta = function(z, y, x, out.family = gaussian,
                     truncps = c(0, 1), e1 = 1, e0 = 1) {
    # fitted propensity score
    pscore = glm(z ~ x, family = binomial)$fitted.values
    pscore = pmax(truncps[1], pmin(truncps[2], pscore))

    # fitted potential outcomes
    outcome1 = glm(y ~ x, weights = z,
                   family = out.family)$fitted.values
    outcome0 = glm(y ~ x, weights = (1 - z),
                   family = out.family)$fitted.values
```

---

[3] [illegible] ??

```
    # outcome regression estimator
    ace.reg = mean(z*y) + mean((1-z)*outcome1/e1) -
              mean(z*outcome0*e0) - mean((1-z)*y)

    # IPW estimators
    w1 = pscore + (1-pscore)/e1
    w0 = pscore*e0 + (1-pscore)

    ace.ipw0 = mean(z*y*w1/pscore) - mean((1-z)*y*w0/(1-pscore))
    ace.ipw = mean(z*y*w1/pscore)/mean(z/pscore) -
              mean((1-z)*y*w0/(1-pscore))/mean((1-z)/(1-pscore))

    # doubly robust estimator
    aug = outcome1/pscore/e1 + outcome0*e0/(1-pscore)
    ace.dr = ace.ipw0 + mean((z-pscore)*aug)

    return(c(ace.reg, ace.ipw0, ace.ipw, ace.dr))
}
```

[illegible] NHANES [illegible] BMI [illegible]

$$\varepsilon_1(X) = \varepsilon_0(X) \in \{1/2, 1/1.7, 1/1.5, 1/1.3, 1, 1.3, 1.5, 1.7, 2\},$$

[illegible]
[illegible]

TABLE 1. [illegible] BMI [illegible]

ε	1/2	1/1.7	1/1.5	1/1.3	1	1.3	1.5	1.7	2
1/2	11.62	10.44	9.40	8.03	4.96	0.97	−1.69	−4.35	−8.34
1/1.7	9.22	8.05	7.00	5.64	2.57	−1.42	−4.08	−6.75	−10.74
1/1.5	7.63	6.45	5.41	4.05	0.97	−3.02	−5.68	−8.34	−12.33
1/1.3	6.03	4.86	3.81	2.45	−0.62	−4.61	−7.27	−9.94	−13.93
1	3.64	2.47	1.42	0.06	−3.01	−7.01	−9.67	−12.33	−16.32
1.3	1.80	0.63	−0.42	−1.78	−4.85	−8.85	−11.51	−14.17	−18.16
1.5	0.98	−0.19	−1.24	−2.60	−5.67	−9.66	−12.33	−14.99	−18.98
1.7	0.36	−0.82	−1.86	−3.23	−6.30	−10.29	−12.95	−15.61	−19.60
2	−0.35	−1.52	−2.57	−3.93	−7.00	−10.99	−13.65	−16.32	−20.31

[illegible] $\varepsilon_1(X) > 1$ [illegible] $\varepsilon_0(X) > 1$ [illegible] BMI [illegible] BMI [illegible] $\varepsilon_1(X) < 1$ [illegible] $\varepsilon_0(X) < 1$ [illegible]

[illegible]

[illegible]

(1) [illegible]

(2) i ; l 。 e a [illegible] outcomes [illegible]

(3) [illegible] $\varepsilon_1(X)$ [illegible] $\varepsilon_0(X)$ [illegible]

(4) [illegible] outcome regression~ IPW [illegible]

(5) [illegible] predictive~ projective~ IPW [illegible]

(6) [illegible] ε [illegible]

[illegible]

18.1 (18.1). 证明定理 [illegible]

18.2 (18.2). 证明定理 [illegible]

18.3 (18.3). 第 [illegible]节只给出了点估计。报告相应的 *Bootstrap* 标准误差。

18.4 (18.4). （处理组平均因果效应的灵敏度分析）本习题将第 [illegible]章的结果推广到允许未测混杂的情形，估计

$$\tau_{\mathrm{T}} = \mathbb{E}\{Y(1) - Y(0) \mid Z = 1\} = \mathbb{E}(Y \mid Z = 1) - \mathbb{E}\{Y(0) \mid Z = 1\}.$$

我们容易用样本矩估计 $\mathbb{E}(Y \mid Z = 1)$：$\hat{\mu}_{\mathrm{T1}} = \sum_{i=1}^n Z_i Y_i / \sum_{i=1}^n Z_i$。唯一的反事实项是 $\mathbb{E}\{Y(0) \mid Z = 1\}$，因此我们只需要灵敏度参数 $\varepsilon_0(X)$。在已知 $\varepsilon_0(X)$ 的条件下，我们有以下两个识别公式。

[illegible] 18.4 ([illegible]). 在已知 $\varepsilon_0(X)$ 的条件下，

$$\mathbb{E}\{Y(0) \mid Z = 1\} = \frac{\mathbb{E}\{Z\mu_0(X)\varepsilon_0(X)\}}{e} = \frac{\mathbb{E}\left\{e(X)\varepsilon_0(X)\frac{1-Z}{1-e(X)}Y\right\}}{e},$$

其中 $e = \Pr(Z = 1)$。

证明定理 [illegible]

[illegible]：定理 [illegible]启发我们使用 $\hat{\tau}_{\mathrm{T}}^* = \hat{\mu}_{\mathrm{T1}} - \hat{\mu}_{\mathrm{T0}}^*$ 估计 τ_{T}，其中

$$\hat{\mu}_{\mathrm{T0}}^{\mathrm{reg}} = n_1^{-1} \sum_{i=1}^n Z_i \varepsilon_0(X_i) \hat{\mu}_0(X_i),$$

$$\hat{\mu}_{\mathrm{T0}}^{\mathrm{ht}} = n_1^{-1} \sum_{i=1}^n \varepsilon_0(X_i) \hat{o}(X_i)(1 - Z_i) Y_i,$$

$$\hat{\mu}_{\mathrm{T0}}^{\mathrm{haj}} = \frac{\sum_{i=1}^n \varepsilon_0(X_i) \hat{o}(X_i)(1 - Z_i) Y_i}{\sum_{i=1}^n \hat{o}(X_i)(1 - Z_i)},$$

其中 $\hat{o}(X_i) = \hat{e}(X_i)/\{1 - \hat{e}(X_i)\}$ 是给定协变量条件下治疗的条件比数比估计量。此外，我们可以构造 τ_{T} 的双重稳健估计量

$$\hat{\tau}_{\mathrm{T}}^{\mathrm{dr}} = \hat{\mu}_{\mathrm{T1}} - \hat{\mu}_{\mathrm{T0}}^{\mathrm{dr}},$$

其中

$$\hat{\mu}_{\mathrm{T}0}^{\mathrm{dr}} = \hat{\mu}_{\mathrm{T}0}^{\mathrm{ht}} - n_1^{-1} \sum_{i=1}^{n} \varepsilon_0(X_i) \frac{\hat{e}(X_i) - Z_i}{1 - \hat{e}(X_i)} \hat{\mu}_0(X_i).$$

详见 *Lu and Ding (2023)*。

[illegible]

- [illegible] Manski [illegible]
- [illegible]
- [illegible] E-value [illegible]
- [illegible]
- [illegible]

CHAPTER 19

[illegible]

[illegible]

[illegible] matching [illegible] 11 [illegible]

[illegible] omitted variable bias [illegible]

Paul Rosenbaum 1987 [illegible] sensitivity analysis [illegible]

[illegible] Rosenbaum [illegible] p [illegible] Rosenbaum [illegible]

[illegible]

[illegible]

[illegible] n [illegible] i [illegible] $j = 1$ [illegible] $j = 0$ [illegible] $(i, 1)$ [illegible] i [illegible] $(i, 0)$ [illegible]

[illegible] (i, j) [illegible] $Y_{ij}(1)$ [illegible] $Y_{ij}(0)$ [illegible] sharp null hypothesis [illegible]

$$H_{0F} : Y_{ij}(1) = Y_{ij}(0) \quad \text{for all } i, j.$$

[illegible] $\pi_{i1} = 1/2$

[illegible]

[illegible] Rosenbaum (1987b) [illegible]

[illegible] 19.1 (Rosenbaum [illegible]). 对于第 i 对中的两个单位，处理分配的 *odds* 比被一个有界常数 $\Gamma \geq 1$ 控制：

$$\frac{\pi_{i1}/\pi_{i0}}{(1-\pi_{i1})/(1-\pi_{i0})} \leq \Gamma, \quad i = 1, \ldots, n,$$

其中 $\pi_{i1} = \Pr\{Z_{i1} = 1 \mid pair\ i\}$ 是配对 i 中处理组接受处理的概率。

[illegible] Γ [illegible]

- [illegible] $\Gamma = 1$ [illegible] $\pi_{i1} = \pi_{i0} = 1/2$ [illegible] MPE [illegible]

- [illegible] $\Gamma > 1$ [illegible] Γ [illegible]
- Γ [illegible]

[illegible] odds [illegible] odds [illegible] Γ [illegible] p [illegible]

[illegible]

[illegible] sharp null hypothesis H_{0F} [illegible] ” ” [illegible] Fisher randomization test [illegible] q_i [illegible] i [illegible] choices [illegible]

(1) [illegible] $q_i = Z_{i1} - Z_{i0}$ [illegible] vs [illegible]
(2) [illegible] $q_i = Y_{i1} - Y_{i0}$ [illegible]
(3) [illegible] $|Y_{i1} - Y_{i0}|$ [illegible]

[illegible]

$$T = \sum_{i=1}^{n} S_i q_i,$$

[illegible] $S_i \in \{0, 1\}$ [illegible] i [illegible] $S_i = 1$ [illegible] q_i [illegible] i [illegible]

[illegible] S_i [illegible] S_i [illegible]

[illegible]

[illegible] p [illegible] 在 [illegible] 下，零假设 H_{0F} 成立时，配对 i 的处理指示 S_i 服从最坏情况的 *Bernoulli* 分布：

$$S_i \sim \mathit{Bernoulli}\left(\frac{\Gamma}{1+\Gamma}\right),$$

且各配对的 S_i 之间相互独立。

[illegible] odds ratio [illegible] Γ [illegible] S_i [illegible] $[\pi_{\min}, \pi_{\max}]$ [illegible]

$$\pi_{\max} = \frac{\Gamma}{1+\Gamma}, \quad \pi_{\min} = \frac{1}{1+\Gamma}.$$

[illegible] p [illegible] S_i [illegible] $\pi_{\max}$ [illegible] ” [illegible] ” [illegible] p[1]

[illegible]

[illegible] p [illegible] T^{obs} [illegible] p [illegible]

$$\mathrm{p}_\Gamma = \Pr_\Gamma\left(|T - T^{\text{obs}}| \geq |T^{\text{obs}}|\right),$$

[illegible] T [illegible] S_i [illegible]

---

[1] [illegible] ?? [illegible] ??

[illegible] $q_i \geq 0$ [illegible]

$$T = \sum_{i=1}^{n} S_i q_i, \quad S_i \sim \text{Bernoulli}\left(\frac{\Gamma}{1+\Gamma}\right).$$

q_i [illegible] T [illegible] n [illegible] Bernoulli [illegible]

[illegible]

[illegible] LaLonde (1986) [illegible] National Supported Work Demonstration [illegible]

[illegible]

$$T = \sum_{i=1}^{n} S_i |\hat{\tau}_i|,$$

[illegible] $\hat{\tau}_i$ [illegible] i [illegible]

- [illegible] MPE [illegible] $\Gamma = 1$ [illegible] p [illegible] 0.002 [illegible]
- [illegible] $\Gamma = 1.1$ [illegible] 10% [illegible] odds [illegible] p [illegible] 0.011
- [illegible] $\Gamma = 1.3$ [illegible] p [illegible] 0.05 [illegible]

[illegible] $\Gamma = 1.233$ [illegible] odds [illegible] 1.233 [illegible] 0.05 [illegible]

[illegible] 19.1 (LaLonde [illegible]). 图 [illegible] 展示了 p 值随 Γ 变化的灵敏度曲线。曲线的陡峭程度告诉我们结论对未测量混杂的敏感程度——曲线越平缓，结论越稳健。

[illegible] Rosenbaum [illegible] R [illegible] `sensitivity2x2` [illegible] `sensitivitymv` [illegible]

[illegible] 2×2 [illegible] Γ [illegible]

[illegible]

[illegible]

[illegible] 19.1. *Rosenbaum 灵敏度模型的基本性质* 证明：当 $\Gamma \to \infty$ 时，[illegible] 中的约束变得无效，此时 S_i 的最坏情况分布趋近于 *Bernoulli*$(1/2)$。这对应着什么实际含义？

[illegible] 19.2. *最坏情况 p 值的单调性* 设 T^{obs} 为观测到的检验统计量。证明：最坏情况 p 值 p_Γ 是 Γ 的单调递增函数。解释这在直觉上意味着什么。

[illegible] 19.3. *Sign statistic 的灵敏度分析* 对于 *sign statistic* $q_i = 1$（所有配对权重相等），推导 T 在 [illegible] 下的分布，并给出 p 值的显式计算公式。

[illegible] 19.4. *LaLonde 数据的计算* 使用 *LaLonde* 数据集，计算在不同 Γ 值$(1, 1.1, 1.2, 1.3, 1.5)$下的最坏情况 p 值，并绘制灵敏度曲线。

[illegible]

[illegible] Rosenbaum [illegible] [illegible]

[illegible]

1. (1) [illegible] odds ratio [illegible] $\Gamma \geq 1$ [illegible]
2. (2) $\Gamma = 1$ [illegible] $\Gamma > 1$ [illegible]
3. (3) [illegible] S_i [illegible] p
4. (4) [illegible] ” [illegible] ” [illegible]

[illegible] Rosenbaum (1987b) [illegible] Rosenbaum [illegible] (2002, 2010) [illegible]

[illegible] 11 [illegible] 3 Fisher randomization test [illegible] 12 [illegible] ” [illegible] ” [illegible]

CHAPTER 20

[illegible]

[illegible]

“If we could intervene on a unit and set its treatment, then we would know its potential outcome under that treatment. But what if the treatment is deterministic for some units? Can we still conceive their counterfactual outcomes?” — (, adapted from King and Zeng, 2006)

[illegible] ** [illegible] ** [illegible] ** [illegible] ** [illegible] Rubin [illegible] Rubin, 1974, 1975 [illegible] ” [illegible] ” [illegible] { {[illegible] **[illegible]**

$$0 < e(X) < 1. \tag{20.1}$$

[illegible] $e(X) = 0$ [illegible] $e(X) = 1$ [illegible] ” [illegible] ” [illegible] **[illegible]**

[illegible]

[illegible] { {[illegible] { [illegible] D’Amour [illegible] 2021 [illegible]

- [illegible] ignorability [illegible] { { [illegible] M-bias [illegible]
- [illegible]

[illegible]

[illegible] 20.1 ([illegible]). 存在某个 $\eta \in (0, 1/2)$ 使得

$$\eta \le e(X) \le 1 - \eta.$$

D’Amour [illegible] 2021 [illegible] p [illegible][1] [illegible]

[illegible] (D’Amour et al., 2021) 在 { {下，设 $e = \Pr(Z = 1)$ 为治疗组比例，λ_1 和 λ_0 分别为 $X|Z = 1$ 和 $X|Z = 0$ 条件下协变量矩阵的最大特征值。则

$$\frac{1}{p}\sum_{k=1}^{p}\bigl|\mathbb{E}(X_k \mid Z = 1) - \mathbb{E}(X_k \mid Z = 0)\bigr| \le p^{-1/2}C^{1/2}\bigl\{e\lambda_1^{1/2} + (1-e)\lambda_0^{1/2}\bigr\}, \tag{20.2}$$

---

[1] [illegible] **??**

其中 $C = \dfrac{(e-\eta)(1-e-\eta)}{e^2(1-e)^2\eta(1-\eta)}$ 是仅依赖于 (e,η) 的正常数。

[illegible]

[illegible]

[illegible], 2009; Yang and Ding, 2018b[illegible]

[illegible] D’Amour [illegible] 2021[illegible]

[illegible] X [illegible] $r(X)$[illegible]

$$\mathbb{E}\{Y(z) \mid X\} = f_z(r(X)), \quad (z = 0, 1), \tag{20.3}$$

[illegible] $r(X)$ [illegible] $r(X)$ [illegible] X [illegible][2] [illegible]

[illegible]

[illegible] X [illegible]

$$Z = \mathbb{I}(X \geq x_0), \tag{20.4}$$

[illegible] x_0 [illegible]

$$Z \perp \{Y(1), Y(0)\} \mid X, \tag{20.5}$$

Z [illegible] X [illegible]

$$e(x) = \Pr(Z = 1 \mid X = x) = \mathbb{I}(x \geq x_0) = \begin{cases} 1 & \text{[illegible]} \ x \geq x_0, \\ 0 & \text{[illegible]} \ x < x_0. \end{cases} \tag{20.6}$$

[illegible]

---

[2] [illegible] ??

[illegible] [illegible] Thistlethwaite [illegible] Campbell [illegible] 1960 [illegible] "[illegible]" [illegible] "[illegible]" [illegible]

[illegible] HIV [illegible] Bor [illegible] 2014 [illegible] HIV [illegible] CD4 [illegible] 200 cells/μL [illegible]

[illegible] Carpenter [illegible] Dobkin [illegible] 2009 [illegible]

[illegible] X [illegible] x_0 [illegible]

$$\tau(x_0) = \mathbb{E}\{Y(1) - Y(0) \mid X = x_0\}. \tag{20.7}$$

[illegible] (RD [illegible]) 假设治疗由 $Z = \mathbb{I}(X \geq x_0)$ 确定，其中 x_0 是预设阈值。假设 $\mathbb{E}\{Y(1) \mid X = x\}$ 在 x_0 处右连续，$\mathbb{E}\{Y(0) \mid X = x\}$ 在 x_0 处左连续。则在 $X = x_0$ 处的局部平均因果效应由下式识别：

$$\tau(x_0) = \lim_{\varepsilon \to 0^+} \mathbb{E}(Y \mid Z = 1, X = x_0 + \varepsilon) - \lim_{\varepsilon \to 0^+} \mathbb{E}(Y \mid Z = 0, X = x_0 - \varepsilon). \tag{20.8}$$

[illegible] $\tau(x_0)$ [illegible][3]

[illegible] $\mathbb{E}(Y \mid Z = 1, X = x) = \gamma_1 + \beta_1 x$ [illegible] $\mathbb{E}(Y \mid Z = 0, X = x) = \gamma_0 + \beta_0 x$ [illegible] x [illegible] x_0 [illegible]

$$\hat{\tau}(x_0) = (\hat{\gamma}_1 - \hat{\gamma}_0) + (\hat{\beta}_1 - \hat{\beta}_0)x_0. \tag{20.9}$$

[illegible] $\hat{\tau}(x_0)$ OLS [illegible]

$$Y_i \sim \{1, Z_i, X_i - x_0, Z_i(X_i - x_0)\} \tag{20.10}$$

[illegible] Z_i [illegible][4]

[illegible] `rdrobust` [illegible] "[illegible]" [illegible]

[illegible] RD [illegible]

[illegible] 21 [illegible]

McCrary [illegible] 2008 [illegible]

[illegible]

[illegible]

(1) [illegible]

(2) [illegible]

---

[3] [illegible] **??**

[4] [illegible] **??**

(3) [illegible] RD [illegible]
[illegible]

RD [illegible]
[illegible] ” [illegible] ” [illegible]
[illegible]

[illegible]

[illegible] [illegible][5]
[illegible] $\Sigma_1 = \mathrm{Cov}(X \mid Z = 1)$ [illegible] $\Sigma_0 = \mathrm{Cov}(X \mid Z = 0)$ [illegible]

$$\Sigma = \mathbb{E}\{\mathrm{Cov}(X \mid Z)\} + \mathrm{Cov}\{\mathbb{E}(X \mid Z)\}. \tag{20.11}$$

[illegible] Σ_1 [illegible] Σ_0 [illegible] C [illegible] Cauchy-Schwarz [illegible]
[illegible]

$$\frac{1}{p}\sum_{k=1}^{p} |\mathbb{E}(X_k \mid Z = 1) - \mathbb{E}(X_k \mid Z = 0)| \le \frac{1}{\sqrt{p}}\|\Sigma^{1/2}\| \cdot \|d\|, \tag{20.12}$$

[illegible] $d = \mathbb{E}(X \mid Z = 1) - \mathbb{E}(X \mid Z = 0)$ [illegible] $\|\cdot\|$ [illegible]
[illegible] ☐

[illegible] [illegible] $\tau = \mathbb{E}\{Y(1) - Y(0)\}$ [illegible]

$$\tau = \mathbb{E}\big\{\mathbb{E}(Y \mid Z = 1, r(X)) - \mathbb{E}(Y \mid Z = 0, r(X))\big\} \tag{20.13}$$

[illegible]

$$\tau = \mathbb{E}\left\{\frac{ZY}{e(r(X))}\right\} - \mathbb{E}\left\{\frac{(1-Z)Y}{1-e(r(X))}\right\}, \tag{20.14}$$

[illegible] $e(r(X)) = \Pr\{Z = 1 \mid r(X)\}$[6]

[illegible] [illegible]

$$\mathbb{E}\{Y(1) \mid X = x_0\} = \lim_{\varepsilon \to 0^+} \mathbb{E}\{Y(1) \mid X = x_0 + \varepsilon\}. \tag{20.15}$$

Z [illegible] $X = x_0 + \varepsilon$ [illegible] $Z = 1$ [illegible]

$$\mathbb{E}\{Y(1) \mid X = x_0 + \varepsilon\} = \mathbb{E}(Y \mid Z = 1, X = x_0 + \varepsilon). \tag{20.16}$$

---

[5] [illegible] D’Amour et al. (2021, Corollary 1)

[6] [illegible] $r(X)$ [illegible]

[illegible] $\mathbb{E}\{Y(0) \mid X = x_0\}$ [illegible] { { □

WwXw [illegible] **w**[illegible] $R_i = \max(X_i - x_0, 0)$ [illegible] $L_i = \min(X_i - x_0, 0)$

$$Z_i(X_i - x_0) = Z_iR_i - (1 - Z_i)L_i = R_i - L_i - (1 - Z_i)R_i + (1 - Z_i)L_i. \tag{20.17}$$

[illegible] $Y_i \sim \{1, Z_i, X_i - x_0, Z_i(X_i - x_0)\}$ [illegible] Z_i [illegible]

$$\hat{\tau} = (\hat{\gamma}_1 - \hat{\gamma}_0) + (\hat{\beta}_1 - \hat{\beta}_0)x_0. \tag{20.18}$$

{ { [illegible] $Y_i \sim \{1, Z_i, R_i, L_i\}$. [illegible] □

CHAPTER 21

CRE

(1) Jobs treatment

(2)

(3)

Z D D

Z D Y D

D Z

“如果我无法做随机实验，能否找到一个自然的随机化来源？”

- $Z \in \{0,1\}$ $Z = 1$
- $D \in \{0,1,2,\ldots\}$
- Y

21.1 ( ). 治疗分配 Z 与所有潜在结果独立：

$$Z \perp\!\!\!\perp \{D(1), D(0), Y(1), Y(0)\},$$

---

[1] ??

[2] ??

其中 $D(z)$ 是在分配 z 下的潜在治疗接受情况，$Y(z)$ 是相应的潜在结局。

[illegible] Z [illegible] D [illegible] τ_Y [illegible] ITT [illegible][3] [illegible] ITT [illegible]

$$\hat{\tau}_Y = \bar{Y}_1 - \bar{Y}_0$$

[illegible] IV [illegible]

[illegible]

[illegible] $D_i(1)$ [illegible] $D_i(0)$ [illegible] $U_i = (D_i(1), D_i(0))$ [illegible]

(1) [illegible] $U_i = a$ [illegible] $D_i(1) = D_i(0) = 1$ [illegible]

(2) [illegible] $U_i = c$ [illegible] $D_i(1) = 1, D_i(0) = 0$ [illegible]

(3) [illegible] $U_i = d$ [illegible] $D_i(1) = 0, D_i(0) = 1$ [illegible]

(4) [illegible] $U_i = n$ [illegible] $D_i(1) = D_i(0) = 0$ [illegible]

[illegible] $D(1)$ [illegible] $D(0)$ [illegible] U_i [illegible]

[illegible][4]

$$
\begin{aligned}
\tau_Y &= \mathbb{E}\{Y(1) - Y(0)\} && (105)\\
&= \mathbb{E}\{Y(1) - Y(0) \mid U = a\} \Pr(U = a) && (106)\\
&\quad + \mathbb{E}\{Y(1) - Y(0) \mid U = c\} \Pr(U = c) && (107)\\
&\quad + \mathbb{E}\{Y(1) - Y(0) \mid U = d\} \Pr(U = d) && (108)\\
&\quad + \mathbb{E}\{Y(1) - Y(0) \mid U = n\} \Pr(U = n). && (109)
\end{aligned}
$$

[illegible] complier [illegible]

[illegible]

[illegible] complier [illegible] CACE [illegible]

---

[3] [illegible] ITT [illegible] **??**

[4] [illegible] **??**

[illegible]

[illegible] 21.2 ([illegible]). 没有 *defier*: $\Pr(U_i = d) = 0$，或者等价地，$D_i(1) \geq D_i(0)$ 对所有 i 成立。

[illegible] Defier [illegible]

$$\Pr(D = 1 \mid Z = 1) \geq \Pr(D = 1 \mid Z = 0), \tag{110}$$

[illegible][5] [illegible] $D_i(1) \geq D_i(0)$ [illegible] $D_i(0) = 0$ [illegible]

[illegible]

[illegible] 21.3 ([illegible]). 对 *always taker* 和 *never taker*，治疗分配不影响结局：

$$Y_i(1) = Y_i(0) \quad 当 \quad U_i = a \text{ 或} U_i = n.$$

[illegible] always taker [illegible] never taker [illegible] Z [illegible] Z [illegible] Z [illegible] Y [illegible] $D_i(1) = D_i(0)$ [illegible] $Y_i(1) = Y_i(0)$ [illegible] i [illegible] Z [illegible] D [illegible] Y

[illegible] Z [illegible]

[illegible]

[illegible] always taker [illegible] never taker [illegible] $Y(1) = Y(0)$ [illegible] defier [illegible]

$$\tau_Y = \mathbb{E}\{Y(1) - Y(0) \mid U = c\} \Pr(U = c). \tag{111}$$

[illegible]

$$\tau_D = \mathbb{E}\{D(1) - D(0)\} = \Pr(U = c), \tag{112}$$

[illegible] complier [illegible] $Z = 1$ [illegible] $Z = 0$ [illegible]

---

[5] [illegible] ?? [illegible] ??

[illegible] CACE [illegible]) [illegible],

$$\tau_c \overset{def}{=} \mathbb{E}\{Y(1) - Y(0) \mid U = c\} = \frac{\tau_Y}{\tau_D}. \tag{113}$$

[illegible] CACE/LATE [illegible]) [illegible],

$$\tau_c = \frac{\mathbb{E}(Y \mid Z = 1) - \mathbb{E}(Y \mid Z = 0)}{\mathbb{E}(D \mid Z = 1) - \mathbb{E}(D \mid Z = 0)}. \tag{114}$$

[illegible] ITT [illegible] ITT [illegible] τ_Y/τ_D [illegible] complier average causal effect [illegible] CACE [illegible][6] [illegible]

[illegible] CACE [illegible] complier [illegible] complier [illegible] $D(1) > D(0)$ [illegible] $D(1) = 1, D(0) = 0$ [illegible] complier [illegible] $D = Z$ [illegible] CACE [illegible] complier [illegible] CACE [illegible] complier [illegible] always taker [illegible] never taker[5] [illegible] defier [illegible] IV [illegible]

[illegible]

1 [illegible] τ_c [illegible]

$$\hat{\tau}_c = \frac{\hat{\tau}_Y}{\hat{\tau}_D} = \frac{\bar{Y}_1 - \bar{Y}_0}{\bar{D}_1 - \bar{D}_0}, \tag{115}$$

[illegible] $\bar{Y}_z$ [illegible] $\bar{D}_z$ [illegible] $Z = z$ [illegible] 5 [illegible] [7] $\tau_c \neq 0$ [illegible]

$$\hat{V}(\hat{\tau}_c) = \frac{\hat{V}_Y}{\hat{\tau}_D^2},$$

[illegible] $\hat{V}_Y$ [illegible] $\hat{\tau}_Y$ [illegible] Neyman [illegible]

$$\left(\frac{\hat{\tau}_Y}{\hat{\tau}_D} - z_{1-\alpha/2}\sqrt{\frac{\hat{V}_Y}{\hat{\tau}_D^2}},\ \frac{\hat{\tau}_Y}{\hat{\tau}_D} + z_{1-\alpha/2}\sqrt{\frac{\hat{V}_Y}{\hat{\tau}_D^2}} \right).$$

[illegible] $\tau_c = 0$ [illegible] $\hat{V}_Y/\hat{\tau}_D^2$ [illegible] Wald [illegible] ITT [illegible] $1/\hat{\tau}_D > 1$ [illegible]

[illegible]

[illegible] IV [illegible] X

---

[6] [illegible] ??

[7] [illegible] ??

[illegible]

$$\tau_c(x) = \frac{\mathbb{E}(Y \mid Z=1, X=x) - \mathbb{E}(Y \mid Z=0, X=x)}{\mathbb{E}(D \mid Z=1, X=x) - \mathbb{E}(D \mid Z=0, X=x)}. \tag{116}$$

[illegible] X [illegible] CACE [illegible] CRE [illegible] $X = x$ [illegible] IV [illegible]

[illegible] X [illegible]

$$Z \perp\!\!\!\perp \{D(1), D(0), Y(1), Y(0)\} \mid X,$$

[illegible] X [illegible]

[illegible]

[illegible] $\tau_D > 0$ [illegible] complier [illegible] τ_D [illegible] 0 [illegible] Wald [illegible] $\tau_D > 0$ [illegible] $\hat{\tau}_D = 0$ [illegible] $\hat{\tau}_c = \hat{\tau}_Y / \hat{\tau}_D$ [illegible][8] [illegible] IV [illegible]

(1) $\hat{\tau}_c$ [illegible]
(2) $\hat{\tau}_c$ [illegible]
(3) Wald [illegible]

[illegible] Y [illegible] $\tau_Y \in [-1, 1]$ [illegible] $\hat{\tau}_c = \hat{\tau}_Y / \hat{\tau}_D$ [illegible]

[illegible] IV [illegible][9] FAR [illegible] τ_c [illegible] τ_c [illegible] IV [illegible] CRE [illegible] FAR [illegible]

$$(\tau - \hat{\tau}_Y)^2 \leq z_{1-\alpha/2}^2 \hat{V}_Y,$$

[illegible]

[illegible]

notation $\{D(1), D(0), Y(1), Y(0)\}$ [illegible] Z [illegible] notation [illegible] notation [illegible] τ_c [illegible] complier [illegible] complier [illegible] $D(1) = 1$ [illegible] $D(0) = 0$ [illegible] $D = Z$ [illegible] τ_c [illegible] complier [illegible] $Y_i(z, d)$ [illegible] i [illegible] z [illegible] d [illegible] notation [illegible]

[illegible] 21.4 ([illegible] Angrist notation[illegible]). $Y_i(z, d) = Y_i(d)$ 对所有 i, z, d 成立，即潜在结局只是 d 的函数。

---

[8] [illegible] $\hat{\tau}_D$ [illegible] 0 [illegible] ??

[9] [illegible] FAR [illegible] ??

notation [illegible] Z [illegible] Y [illegible] Z [illegible] D [illegible]
[illegible]

$$\tau_c = \frac{\mathbb{E}\{Y(D(1)) - Y(D(0))\}}{\mathbb{E}\{D(1) - D(0)\}},$$

[illegible] CACE [illegible]
[illegible] τ_c [illegible] complier [illegible] ” [illegible] ”[illegible]
[illegible] Z [illegible] Z [illegible] complier [illegible]
[illegible] LATE [illegible]

21.1. [illegible] ***CACE*** [illegible] “[illegible]” [illegible]**：** 因为它是总体中一个特定子群体（*complier*）的平均因果效应。这个子群体恰好是那些“被 Z 影响而改变治疗状态”的人。不同的工具变量可能识别不同的子群体。

[illegible]

[illegible] 21.1. [illegible]*Wald* 估计量的方差 证明 *Wald* 估计量 $\hat{\tau}_c = \hat{\tau}_Y/\hat{\tau}_D$ 的渐近方差为 $\hat{V}_Y/\tau_D^2$，其中 $\hat{V}_Y$ 是 *ITT* 估计量 $\hat{\tau}_Y$ 的方差。

[illegible] 21.2. [illegible]强单调性下的 *IV* 估计 在强单调性（所有人都是 *complier* 或 *never taker*，且 $\Pr(U = n) = 0$）下，证明 *IV* 估计量 $\hat{\tau}_c$ 的一致性。

[illegible] 21.3. [illegible]单侧非依从性 在单侧非依从性（*never taker* 存在，但没有 *always taker*，且 $D_i(0) = 0$ 对所有人成立）下，推导 *CACE* 的识别公式，并解释其含义。

[illegible] 21.4. [illegible]违反排除限制的后果 假设排除限制不成立，即存在 *always taker* 使得 $Y_i(1) \neq Y_i(0)$。证明此时 [illegible]估计量是有偏的，并讨论偏误的方向。

CHAPTER 22

[illegible]

[illegible]

[illegible]

“The essential problem of causal inference is that we can observe at most one of the potential outcomes for any unit.”

[illegible]

[illegible]

[illegible]

- $Z_i \in \{0,1\}$ [illegible] $1 =$ [illegible] $0 =$ [illegible]
- $D_i \in \{0,1\}$ [illegible] $1 =$ [illegible] $0 =$ [illegible]
- Y_i [illegible]

[illegible] $Z_i \neq D_i$ [illegible]

[illegible]

$$\{D_i(1), D_i(0), Y_i(1), Y_i(0)\},$$

[illegible] $D_i(z)$ [illegible] z [illegible] $Y_i(z)$ [illegible] z [illegible]

$$D_i = Z_i D_i(1) + (1 - Z_i) D_i(0), \quad Y_i = Z_i Y_i(1) + (1 - Z_i) Y_i(0).$$

[illegible]

[illegible] 22.1 ([illegible]). 治疗分配 Z 与潜在结果独立：

$$Z \perp\!\!\!\perp \{D(1), D(0), Y(1), Y(0)\}.$$

[illegible] D [illegible] Y [illegible][1]

$$\tau_D = \mathbb{E}\{D(1) - D(0)\} = \mathbb{E}(D \mid Z = 1) - \mathbb{E}(D \mid Z = 0),$$

---

[1] [illegible] **??**

$$\tau_Y = \mathbb{E}\{Y(1) - Y(0)\} = \mathbb{E}(Y \mid Z = 1) - \mathbb{E}(Y \mid Z = 0).$$

[illegible] $\hat{\tau}_D$ [illegible] $\hat{\tau}_Y$ [illegible] τ_D [illegible] τ_Y

[illegible] $\hat{\tau}_Y$ [illegible][2] [illegible]
ITT [illegible]

$$\underbrace{\hat{\tau}_Y}_{\text{ITT [illegible]}} \xrightarrow{p} \underbrace{\tau_Y}_{\text{[illegible]}},$$

[illegible]

$$\underbrace{\tau_{\text{treatment}}}_{\text{[illegible]}} \neq \underbrace{\tau_Y}_{\text{[illegible]}}.$$

[illegible]

[illegible]

[illegible] $\{D_i(1), D_i(0)\}$ [illegible] D [illegible]

22.1 ([illegible]). 令 $U_i = \{D_i(1), D_i(0)\}$，则：

$$U_i = \begin{cases} a \text{（}\textit{always taker}\text{）}, & \text{若} D_i(1) = 1 \text{ 且} D_i(0) = 1; \\ c \text{（}\textit{complier}\text{）}, & \text{若} D_i(1) = 1 \text{ 且} D_i(0) = 0; \\ d \text{（}\textit{defier}\text{）}, & \text{若} D_i(1) = 0 \text{ 且} D_i(0) = 1; \\ n \text{（}\textit{never taker}\text{）}, & \text{若} D_i(1) = 0 \text{ 且} D_i(0) = 0. \end{cases}$$

[illegible]

- [illegible]
- [illegible]
- [illegible]
- [illegible]

[illegible] $D_i(1)$ [illegible] $D_i(0)$ [illegible] U_i [illegible]

[illegible] U [illegible] τ_Y [illegible][3]

$$\begin{aligned} \tau_Y = {} & \mathbb{E}\{Y(1) - Y(0) \mid U = \mathrm{a}\} \Pr(U = \mathrm{a}) \\ & + \mathbb{E}\{Y(1) - Y(0) \mid U = \mathrm{c}\} \Pr(U = \mathrm{c}) \\ & + \mathbb{E}\{Y(1) - Y(0) \mid U = \mathrm{d}\} \Pr(U = \mathrm{d}) \\ & + \mathbb{E}\{Y(1) - Y(0) \mid U = \mathrm{n}\} \Pr(U = \mathrm{n}). \end{aligned}$$

[illegible] τ_Y [illegible]

---

[2] ITT [illegible]

[3] [illegible] **??**

[illegible] τ_D [illegible][4]

$$\begin{aligned}\tau_D = {} & \mathbb{E}\{D(1) - D(0) \mid U = \mathrm{a}\} \Pr(U = \mathrm{a}) \\ & + \mathbb{E}\{D(1) - D(0) \mid U = \mathrm{c}\} \Pr(U = \mathrm{c}) \\ & + \mathbb{E}\{D(1) - D(0) \mid U = \mathrm{d}\} \Pr(U = \mathrm{d}) \\ & + \mathbb{E}\{D(1) - D(0) \mid U = \mathrm{n}\} \Pr(U = \mathrm{n}) \\ = {} & 0 \cdot \Pr(U = \mathrm{a}) + 1 \cdot \Pr(U = \mathrm{c}) + (-1) \cdot \Pr(U = \mathrm{d}) + 0 \cdot \Pr(U = \mathrm{n}) \\ = {} & \Pr(U = \mathrm{c}) - \Pr(U = \mathrm{d}).\end{aligned}$$

X w [illegible]

[illegible] τ_Y [illegible] τ_Y [illegible] τ_D [illegible]

X w [illegible] w

[illegible] 22.2 ([illegible]). $\Pr(U = d) = 0$，或等价地，$D_i(1) \geq D_i(0)$ 对所有单元成立。即，[illegible] ***defiers***。

[illegible] $D_i(0) = 0$ [illegible] $D_i(1) \geq D_i(0)$ [illegible][5]

$$\Pr(D = 1 \mid Z = 1) \geq \Pr(D = 1 \mid Z = 0).$$

[illegible] $D_i(1)$ [illegible] $D_i(0)$ [illegible]

$$\tau_D = \Pr(U = \mathrm{c}) \equiv \pi_c.$$

compliers [illegible] π_c [illegible]

X w [illegible] w

[illegible] 22.3 ([illegible]). 对于 *always takers* ($U = a$) 和 *never takers* ($U = n$)，有 $Y_i(1) = Y_i(0)$。

[illegible]

$$\tau_Y = \mathbb{E}\{Y(1) - Y(0) \mid U = \mathrm{c}\} \Pr(U = \mathrm{c}).$$

[illegible]

---

[4] [illegible] ??

[5] [illegible] ??

[illegible]

[illegible] (CACE/LATE [illegible]) 在 [illegible] 和 [illegible] 下，如果 $\tau_D \neq 0$，则

$$\mathbb{E}\{Y(1) - Y(0) \mid U = c\} = \frac{\tau_Y}{\tau_D}.$$

[illegible][6] [illegible] τ_Y/τ_D [illegible] compliers [illegible] compliers [illegible] ” [illegible] ” [illegible] always takers [illegible] never takers [illegible] defiers [illegible]

[illegible]

[illegible]

[illegible] 22.3 (CACE [illegible] LATE). [illegible] ***complier average causal effect, CACE*** 或 [illegible] ***local average treatment effect, LATE*** 定义为：

$$\tau_c = \mathbb{E}\{Y(1) - Y(0) \mid U = c\}.$$

CACE [illegible]

$$\tau_c = \mathbb{E}\{Y(1) - Y(0) \mid D(1) = 1, D(0) = 0\} = \mathbb{E}\{Y(1) - Y(0) \mid D(1) > D(0)\}.$$

1 [illegible]

$$\tau_c = \frac{\tau_Y}{\tau_D}.$$

[illegible] **[illegible]** [illegible]

$$\tau_c = \frac{\mathbb{E}(Y \mid Z = 1) - \mathbb{E}(Y \mid Z = 0)}{\mathbb{E}(D \mid Z = 1) - \mathbb{E}(D \mid Z = 0)}.$$

[illegible] **[illegible]** [illegible]

[illegible]

[illegible]

[illegible] CACE [illegible]

$$\hat{\tau}_c = \frac{\hat{\tau}_Y}{\hat{\tau}_D},$$

[illegible]

$$\hat{\tau}_Y = \frac{1}{n_1} \sum_{i:Z_i=1} Y_i - \frac{1}{n_0} \sum_{i:Z_i=0} Y_i, \quad \hat{\tau}_D = \frac{1}{n_1} \sum_{i:Z_i=1} D_i - \frac{1}{n_0} \sum_{i:Z_i=0} D_i.$$

[illegible] Z [illegible] D [illegible]

[illegible] Wald [illegible] ” [illegible] ” ITT [illegible] $\hat{\tau}_Y$ [illegible] $\hat{\tau}_D$ [illegible] ” [illegible] ” [illegible] compliers [illegible] ” [illegible] ” [illegible] compliers [illegible]

---

[6] [illegible] ??

[illegible] $\hat{\tau}_c$ [illegible][7] [illegible] $\hat{\tau}_c$ [illegible][8] [illegible]

$$\hat{\tau}_c - \tau_c \approx \frac{\hat{\tau}_Y - \tau_c \hat{\tau}_D}{\tau_D}.$$

[illegible] $A_i = Y_i - \tau_c D_i$ [illegible] $\hat{\tau}_A / \tau_D$ [illegible] $\hat{\tau}_A$ [illegible]

[illegible] $\hat{\tau}_c$ [illegible] $\hat{V}_A / \tau_D^2$ [illegible] $\hat{V}_A$ [illegible] Neyman [illegible]

$$\hat{V}_A = \frac{\hat{S}_A^2(1)}{n_1} + \frac{\hat{S}_A^2(0)}{n_0},$$

[illegible] $\hat{S}_A^2(1)$ [illegible] $\hat{S}_A^2(0)$ [illegible]

$$\hat{V}_{\hat{\tau}_c} = \frac{\hat{V}_A}{\hat{\tau}_D^2}$$

$\hat{\tau}_c$ [illegible] τ_c [illegible] $1 - \alpha$ [illegible]

$$\hat{\tau}_c \pm z_{1-\alpha/2} \sqrt{\hat{V}_{\hat{\tau}_c}}.$$

[illegible]

[illegible] 5% [illegible] Wald [illegible]

[illegible] $\tau_D > 0$ [illegible] compliers [illegible] $\hat{\tau}_D$ [illegible] Wald [illegible] $\tau_D \to 0$ [illegible] $\operatorname{var}(\hat{\tau}_c) \to \infty$[9] [illegible]

- $\hat{\tau}_c$ [illegible]
- [illegible]
- [illegible]

[illegible] F [illegible] 5 [illegible] $\hat{\tau}_D$ [illegible] $\hat{\tau}_D$ [illegible] Wald [illegible]

[illegible] [illegible] $\tau_c = \tau_Y / \tau_D$ [illegible] $\tau_c = 0$ [illegible] $\tau_Y = 0$ [illegible][10] [illegible] τ_c [illegible] b [illegible] $H_0(b) : \tau_c = b$ [illegible] $\tau_Y - b\tau_D = 0$ [illegible] $A(b)_i = Y_i - bD_i$ [illegible]

---

[7] [illegible] ??

[8] [illegible] ??

[9] [illegible] Fieller-Creasy [illegible]

[10] FAR [illegible] ??

[illegible] Wald [illegible]

$$\left\{ b : \frac{\hat{\tau}_A^2(b)}{\hat{V}_A(b)} \le z_{1-\alpha/2}^2 \right\},$$

[illegible] $\hat{\tau}_A(b)$ [illegible] $\hat{V}_A(b)$ [illegible] $A(b)_i = Y_i - bD_i$ [illegible] FAR [illegible] Wald [illegible]

[illegible]

[illegible]

[illegible] [illegible]

$$Z \perp \{D(1), D(0), Y(1), Y(0), X\}.$$

[illegible]

$$D_i = \beta_0 + \beta_1 Z_i + \beta_2 X_i + \beta_3 Z_i X_i + \epsilon_i,$$

$$Y_i = \alpha_0 + \alpha_1 Z_i + \alpha_2 X_i + \alpha_3 Z_i X_i + \eta_i,$$

[illegible] $\hat{\tau}_{D,\mathrm{L}} = \hat{\beta}_1$ [illegible] $\hat{\tau}_{Y,\mathrm{L}} = \hat{\alpha}_1$ [illegible] CACE [illegible]

$$\hat{\tau}_{c,\mathrm{L}} = \frac{\hat{\tau}_{Y,\mathrm{L}}}{\hat{\tau}_{D,\mathrm{L}}}.$$

[illegible] Bootstrap [illegible]

[illegible]

[illegible] JOBS II [illegible] jobsdata.csv [illegible]

- `treat` [illegible] Z [illegible]
- `comply` [illegible] D [illegible]
- `jobseek` [illegible] Y [illegible] 1-5 [illegible]

[illegible][11]

[illegible] 22.1 (JOBS II [illegible] IV [illegible]). 我们使用 *Wald* 估计量分析 *JOBS II* 数据。结果如下：

方法	估计量	标准误	*95% CI* 下限	*95% CI* 上限
Delta 方法	*0.109*	*0.081*	*-0.050*	*0.268*
Bootstrap	*0.109*	*0.083*	*-0.054*	*0.271*
含协变量调整	*0.118*	*0.082*	*-0.042*	*0.278*

[illegible]：点估计表明，对于 *compliers*，参加培训项目使求职自我效能得分提高约 *0.11* 分（满分 *5* 分）。但 *95%* 置信区间包含零，这意味着在常规显著性水平下，我们无法拒绝”培训对 *compliers* 无效果”的原假设。

---

[11] [illegible] **??**

[illegible]

[illegible] CACE [illegible] CACE [illegible]

[illegible] τ_c [illegible]

(1) [illegible] $\tau_c = E\{Y(1) - Y(0) \mid U = c\}$ [illegible] compliers [illegible]
(2) [illegible] compliers [illegible] $D(1) = 1$ [illegible] $D(0) = 0$ [illegible] $D = Z$ [illegible] τ_c [illegible] compliers [illegible]

[illegible]

[illegible] CACE [illegible] compliers [illegible] always takers、never takers [illegible] defiers [illegible] compliers [illegible] CACE [illegible]

[illegible]

[illegible]

(1) [illegible]
(2) [illegible] always takers、compliers、defiers、never takers
(3) [illegible] defiers [illegible]
(4) [illegible] CACE $= \tau_Y / \tau_D$
(5) [illegible] Wald [illegible] $\hat{\tau}_c = \hat{\tau}_Y / \hat{\tau}_D$
(6) [illegible] compliers [illegible] Wald [illegible] FAR [illegible]
(7) [illegible]

[illegible]

[illegible] 22.1. *ITT 与 CACE 的关系* 证明在 [illegible]下，$\tau_Y = \tau_c \cdot \pi_c$，其中 $\pi_c = \Pr(U = c)$。

[illegible] 22.2. *单调性的可检验含义* 证明在完全随机化下，[illegible]蕴含 [illegible]。

[illegible] 22.3. *Wald 估计量的方差* 证明当 $\tau_D = 0$ 时，$var(\hat{\tau}_c) = \infty$。

[illegible] 22.4. *单侧依从性* 考虑单侧依从性情形：对照组被试无法获得治疗，即 $D_i(0) = 0$ 对所有 i 成立。

(1) 证明此时 [illegible]自动成立。
(2) 在这种情况下，*CACE* 如何简化为 *ITT* 的函数？

[illegible] 22.5. *FAR 置信区间的形状* 在 [illegible]中，*FAR* 置信集可能是单个闭区间、两个不相连的区间、空集或整个实数线。找出每种情况对应的精确条件。

[illegible] 22.6. *协变量调整的 CACE* 使用 *JOBS II* 数据进行以下分析：

(1) 计算原始 *Wald* 估计量及其标准误。
(2) 加入协变量（*sex, age, marital, nonwhite, educ, income*）后重新估计。
(3) 比较两种方法的估计量和标准误。

Ǘ 22.7. *Always takers* 与 *Never takers* 假设我们想估计 *always takers* 的平均因果效应。

(1) 在当前框架下，这个量能否被识别？为什么？
(2) 如果加入额外假设”*always takers* 和 *never takers* 的潜在结果分布相同”，结果会如何改变？

Ǘ 22.8. 模拟研究 进行模拟研究，比较以下三种情况：

(1) 强工具变量（$\pi_c = 0.5$）
(2) 中等强度工具变量（$\pi_c = 0.2$）
(3) 弱工具变量（$\pi_c = 0.05$）

对每种情况，报告 *Wald* 估计量的偏差、均方误差和置信区间的覆盖概率。

CHAPTER 23

[illegible]

[illegible]

[illegible] just-identified [illegible] IV [illegible] Z [illegible] D [illegible] $E(ZY) = E(ZD^\top)\beta$ [illegible] IV [illegible] over-identified [illegible]

[illegible] Angrist and Krueger (1991) [illegible]

[illegible] ” [illegible] ” [illegible] Theil (1953) [illegible] Basmann (1957) [illegible] Two-Stage Least Squares, TSLS [illegible] TSLS [illegible]

[illegible]

[illegible]

$$Y = D^\top\beta + \varepsilon, \tag{117}$$

[illegible] Y [illegible] D [illegible] p [illegible] β [illegible] p [illegible] q [illegible] Z [illegible] restrictions [illegible] $E(\varepsilon Z) = 0$ [illegible] $q > p$ [illegible]

$$n^{-1}\sum_{i=1}^{n} Z_iY_i = n^{-1}\sum_{i=1}^{n} Z_iD_i^\top\beta \tag{118}$$

[illegible] q [illegible] p [illegible] $q > p$ [illegible] ” [illegible] ” [illegible]

[illegible]

TSLS [illegible] [[illegible]] [illegible] Z [illegible] D [illegible] β [illegible] TSLS [illegible]

[illegible] D [illegible] Z OLS [illegible] $\hat{D}_i$ [illegible] $i = 1, \ldots, n$ [illegible] D_i [illegible] OLS [illegible] $\hat{D}$ [illegible] i [illegible] $\hat{D}_i^\top$ [illegible]

[illegible] Y [illegible] $\hat{D}$ OLS [illegible] $\hat{\beta}_{\text{TSLS}}$

[illegible]

[illegible] TSLS [illegible][1] [illegible] OLS [illegible]

$$\hat{D}_i = \hat{\Gamma}^\top Z_i, \tag{119}$$

[illegible] $\hat{\Gamma} = \left(\sum_{i=1}^n Z_i Z_i^\top\right)^{-1} \sum_{i=1}^n Z_i D_i^\top$ [illegible] OLS [illegible]

$$\hat{\beta}_{\text{TSLS}} = \left(\sum_{i=1}^n \hat{D}_i \hat{D}_i^\top\right)^{-1} \sum_{i=1}^n \hat{D}_i Y_i. \tag{120}$$

[illegible] $Y_i = D_i^\top \beta + \varepsilon_i$ [illegible]

$$\hat{\beta}_{\text{TSLS}} = \beta + \left\{\hat{\Gamma}^\top \left(n^{-1} \sum_{i=1}^n Z_i Z_i^\top\right) \hat{\Gamma}\right\}^{-1} \hat{\Gamma}^\top \left(n^{-1} \sum_{i=1}^n Z_i \varepsilon_i\right). \tag{121}$$

[illegible] TSLS [illegible] $E(Z\varepsilon) = 0$ [illegible] 0 [illegible] $\hat{\beta}_{\text{TSLS}}$ [illegible] β [illegible] $q = p$ [illegible] TSLS [illegible] IV [illegible] $\hat{\beta}_{\text{IV}}$ [illegible]

[illegible]

[illegible] TSLS [illegible] $\hat{\varepsilon}_i = Y_i - \hat{\beta}_{\text{TSLS}}^\top D_i$ [illegible] Eicker-Huber-White [illegible]

$$\hat{V}_{\text{TSLS}} = \left(\sum_{i=1}^n \hat{D}_i \hat{D}_i^\top\right)^{-1} \left(\sum_{i=1}^n \hat{\varepsilon}_i^2 \hat{D}_i \hat{D}_i^\top\right) \left(\sum_{i=1}^n \hat{D}_i \hat{D}_i^\top\right)^{-1}. \tag{122}$$

[illegible] $\hat{\varepsilon}_i$ [illegible] Y_i on D_i [illegible] 而不是 [illegible] OLS [illegible] Y_i on $\hat{D}_i$ [illegible] $\hat{V}_{\text{TSLS}}$ [illegible]

[illegible]

[illegible] Z [illegible] D [illegible]

$$\begin{cases} Y_i = \beta_0 + \beta_1 D_i + \beta_2^\top X_i + \varepsilon_i, \\ D_i = \gamma_0 + \gamma_1 Z_i + \gamma_2^\top X_i + \varepsilon_{2i}, \end{cases} \tag{123}$$

[illegible] D_i [illegible] Z_i [illegible] $E(\varepsilon_i Z_i) = 0$ [illegible] X_i [illegible] $E(\varepsilon_i X_i) = 0$ [illegible]

---

[1] [illegible] **??**

[illegible] [illegible] TSLS [illegible]

[illegible] TSLS] [illegible] TSLS [illegible]

[illegible] D [illegible] $(1, Z, X)$ OLS [illegible] $\hat{D}_i$ [illegible] $i = 1, \ldots, n$ [illegible]

[illegible] Y [illegible] $(1, \hat{D}, X)$ OLS [illegible] $\hat{\beta}_{\text{TSLS}}$ [illegible] $\hat{\beta}_{1,\text{TSLS}}$ [illegible] $\hat{D}$ [illegible]

[illegible]

[illegible] [illegible] reduced form [illegible]

$$
\begin{cases} Y_i = \Gamma_0 + \Gamma_1 Z_i + \Gamma_2^\top X_i + \varepsilon_{1i}, \\ D_i = \gamma_0 + \gamma_1 Z_i + \gamma_2^\top X_i + \varepsilon_{2i}, \end{cases} \tag{124}
$$

[illegible] $\Gamma_0 = \beta_0 + \beta_1\gamma_0$ [illegible] $\Gamma_1 = \beta_1\gamma_1$ [illegible] $\Gamma_2 = \beta_2 + \beta_1\gamma_2$ [illegible] $\varepsilon_{1i} = \varepsilon_i + \beta_1\varepsilon_{2i}$ [illegible] (Z, X) [illegible] OLS [illegible] $\hat{\Gamma}_1$ [illegible] $\hat{\gamma}_1$ [illegible] β_1 [illegible]

$$
\beta_1 = \frac{\Gamma_1}{\gamma_1}. \tag{125}
$$

[illegible] Indirect Least Squares, ILS [illegible]

$$
\hat{\beta}_{1,\text{ILS}} = \frac{\hat{\Gamma}_1}{\hat{\gamma}_1}. \tag{126}
$$

[illegible] ILS [illegible] TSLS [illegible] [TSLS ILS [illegible]] [illegible]

$$
\hat{\beta}_{1,\text{ILS}} = \hat{\beta}_{1,\text{TSLS}}. \tag{127}
$$

[illegible] TSLS [illegible] ILS [illegible][2] [illegible]

[illegible] TSLS [illegible] γ_1 [illegible]

[illegible]

[illegible] TSLS [illegible] $\hat{\gamma}_1$ [illegible] $\hat{\Gamma}_1$ [illegible] γ_1 [illegible]

[illegible] Fieller-Anderson-Rubin (FAR) [illegible] b [illegible] $H_0(b) : \beta_1 = b$ [illegible] $Y_i(b) = Y_i - bD_i$ [illegible] $Y_i(b)$ [illegible] Z_i [illegible] b [illegible] $E(Z\varepsilon) = 0$ [illegible]

[illegible] FAR [illegible]

$$
\left\{ b : |t_Z(b)| \le z_{1-\alpha/2} \right\}, \tag{128}
$$

---

[2] [illegible] **??** [illegible] **??**

[illegible] $t_Z(b)$ [illegible] Z [illegible] $Y_i(b)$ on $(1, Z, X)$ [illegible] t [illegible] EHW [illegible] $z_{1-\alpha/2}$ [illegible] $1-\alpha/2$ [illegible]

[illegible] FAR [illegible] 1 [illegible] TSLS [illegible] Wald [illegible] TSLS [illegible]

[illegible]

[illegible] TSLS [illegible] Card (1993) [illegible]

[illegible] Z [illegible] ” [illegible] ” [illegible] D [illegible] Y [illegible] 1976 [illegible] X [illegible]

[illegible] TSLS [illegible] 95% [illegible]

	TSLS [illegible]	[illegible]	[illegible]
[illegible]	0.132	0.026	0.237

[illegible] 13.2% [illegible] FAR [illegible] $[0.028, 0.282]$ [illegible] TSLS [illegible]

[illegible]

[illegible] TSLS [illegible] TSLS [illegible] OLS [illegible]

[illegible] TSLS [illegible] IV [illegible] TSLS [illegible] ILS [illegible] FAR [illegible] TSLS [illegible]

[illegible]

[illegible] 23.1 (TSLS [illegible] IV [illegible]). 证明在恰好识别情形下（Z 与 D 维度相同），*TSLS* 估计量 $\hat{\beta}_{\text{TSLS}}$ 与 *IV* 估计量 $\hat{\beta}_{\text{IV}}$ 数值相同。

[illegible]：利用第一阶段 *OLS* 的正交性条件 $\sum_{i=1}^n \hat{D}_i \check{D}_i^\top = 0$。

[illegible] 23.2 (TSLS [illegible] ILS [illegible]). 证明定理 [illegible] 在单一内生处理变量和单一工具变量情形下，*TSLS* 估计量等于约化形式系数比值 $\hat{\Gamma}_1/\hat{\gamma}_1$。

[illegible]：利用第一阶段拟合值的表达式 $\hat{D}_i = \hat{\gamma}_0 + \hat{\gamma}_1 Z_i + \hat{\gamma}_2^\top X_i$，验证第二阶段 *OLS* 中 $\hat{D}$ 的系数恰好等于比值形式。

[illegible] 23.3 ([illegible]). 考虑 *TSLS* 估计量 $\hat{\beta}_{1,\text{TSLS}} = \hat{\Gamma}_1/\hat{\gamma}_1$。当工具变量较弱（即 γ_1 接近于零）时，讨论：

(1) 为什么 $\hat{\gamma}_1$ 的小误差会导致 $\hat{\beta}_{1,\text{TSLS}}$ 的巨大偏差？
(2) *FAR* 置信区间相比 *Wald* 型置信区间的优势是什么？

CHAPTER 24

[illegible]

[illegible]

[illegible] Regression Discontinuity Design, RDD[illegible] Instrumental Variables, IV[illegible]

[illegible] Fuzzy Regression Discontinuity, FRDD[illegible] ” [illegible] ”[illegible]

[illegible]

[illegible] [illegible] Sharp RDD[illegible] $Z_i = I(X_i \geq x_0)$ [illegible] running variable[illegible] X_i [illegible] x_0 [illegible]

[illegible] RDD [illegible] $\{Y_i(1), Y_i(0)\}$ [illegible]

$$\tau_Y(x_0) = \lim_{\varepsilon \to 0+} \mathbb{E}(Y \mid X = x_0 + \varepsilon) - \lim_{\varepsilon \to 0+} \mathbb{E}(Y \mid X = x_0 - \varepsilon). \tag{129}$$

[illegible] [illegible] RDD [illegible] $Z_i = I(X_i \geq x_0)$ [illegible]

(1) [illegible] Z_i [illegible] D_i [illegible] $X_i \geq x_0$ [illegible]
(2) [illegible] Z_i [illegible] D_i [illegible] Y_i[illegible]

[illegible]

[illegible]

[illegible] 24.1 ([illegible]). [illegible] 研究了印度政府 *2000* 年启动的”总理乡村道路计划”。该计划根据 *2001* 年人口普查的阈值来优先考虑规模较大的村庄。

关键点：村庄人口规模与阈值的差并不能 [illegible] 是否获得道路——它只改变了获得道路的[illegible]。在断点附近，获得道路的概率发生了跳跃，但不是从 *0* 到 *1* 的确定性跳跃。

[illegible] 24.2 ([illegible]). [illegible] 研究了意大利大学助学金对辍学率的影响。学生如果标准化家庭收入低于 *15,000* 欧元，则有资格获得助学金。

为获得助学金，学生必须先申请。因此，资格状态和申请状态共同决定了最终治疗状态。运行变量（*15,000* 减去标准化家庭收入）单独并不能决定治疗状态，但它在断点处改变了治疗概率。

Figure 1. [illegible]

[illegible]

[illegible] X_i [illegible] x_0 [illegible]

$$Z_i = I(X_i \ge x_0).$$

[illegible] D_i [illegible] Z_i [illegible] $\mathbb{P}(D_i = 1 \mid X_i = x)$ [illegible] x_0 [illegible] RDD [illegible] RDD [illegible]

[illegible]

[illegible] Y_i [illegible] Z_i [illegible] $\{D_i(1), D_i(0), Y_i(1), Y_i(0)\}$ [illegible] RDD [illegible]

$$\tau_D(x_0) = \mathbb{E}\{D(1) - D(0) \mid X = x_0\}$$

[illegible]

$$\tau_Y(x_0) = \mathbb{E}\{Y(1) - Y(0) \mid X = x_0\}. \tag{130}$$

[illegible] Z [illegible] D [illegible] $X = x_0$ [illegible] Local Complier Average Causal Effect, LATE[illegible]

[illegible] **[illegible]** RDD [illegible]) 设治疗由 $Z = I(X \ge x_0)$ 决定，其中 x_0 是预定的阈值。

假设[illegible]：

$$D_i(1) \ge D_i(0), \tag{131}$$

和[illegible]：

$$D_i(1) = D_i(0) \Longrightarrow Y_i(1) = Y_i(0), \tag{132}$$

在 x_0 的无穷小邻域内成立。

[illegible]：$E\{D(1) \mid X = x\}$ 和 $E\{Y(1) \mid X = x\}$ 在 $x = x_0$ 处右连续，$E\{D(0) \mid X = x\}$ 和 $E\{Y(0) \mid X = x\}$ 在 $x = x_0$ 处左连续。

局部依从者平均因果效应（*Local Average Treatment Effect, LATE*）定义为：

$$\tau_{\text{LATE}}(x_0) = \mathbb{E}\{Y(1) - Y(0) \mid D(1) > D(0), X = x_0\}, \tag{133}$$

可以由下式识别：

$$\tau_{\text{LATE}}(x_0) = \frac{\mathbb{E}\{Y(1) - Y(0) \mid X = x_0\}}{\mathbb{E}\{D(1) - D(0) \mid X = x_0\}}. \tag{134}$$

如果 $E(D \mid X = x)$ 在 $x = x_0$ 处有非零跳跃，则局部依从者平均因果效应可以由下式识别：

$$\tau_{\text{LATE}}(x_0) = \frac{\lim_{\varepsilon \to 0+} \mathbb{E}(Y \mid Z = 1, X = x_0 + \varepsilon) - \lim_{\varepsilon \to 0+} \mathbb{E}(Y \mid Z = 0, X = x_0 - \varepsilon)}{\lim_{\varepsilon \to 0+} \mathbb{E}(D \mid Z = 1, X = x_0 + \varepsilon) - \lim_{\varepsilon \to 0+} \mathbb{E}(D \mid Z = 0, X = x_0 - \varepsilon)}. \tag{135}$$

[illegible] RDD [illegible] ({) [illegible] ({) [illegible] RDD [illegible] Z [illegible] IV [illegible] IV [illegible]

[illegible]

[illegible]

[illegible] local randomization [illegible] [illegible]

[illegible] RDD [illegible] h [illegible]

[illegible]

[illegible]

[illegible]

(1) [illegible] RDD [illegible]

(2) [illegible] $Z = I(X \geq x_0)$ [illegible] LATE

(3) [illegible] ILS [illegible] TSLS [illegible]

(4) [illegible]

(5) [illegible] ~**MM** [illegible] RDD [illegible] RDD [illegible] IV [illegible]

[illegible] RDD [illegible]

[illegible]

[illegible] 24.1. [illegible]模糊 *RDD* 识别定理的证明 证明 [illegible]中的识别公式。

[illegible] 24.2. [illegible]清晰 *RDD* 与模糊 *RDD* 的比较

(1) 在清晰 *RDD* 中，治疗分配 D 和工具变量 Z 有什么关系？

(2) 在模糊 *RDD* 中，这种关系如何变化？

(3) 给出清晰 *RDD* 和模糊 *RDD* 各一个现实生活中的例子。

[illegible] 24.3. [illegible]带宽敏感性分析 对于 [illegible]中的印度道路数据，使用不同的带宽 h 进行敏感性分析。报告点估计和标准误如何随 h 变化。

[illegible] 24.4. [illegible]*ILS* 与 *TSLS* 的等价性 证明在模糊 *RDD* 中，间接最小二乘估计器（比率形式）和两阶段最小二乘估计器在数值上是等价的。

CHAPTER 25

[illegible]

[illegible]

[illegible] IV [illegible] IV [illegible]

[illegible] 20 [illegible] 80 [illegible] Katan [illegible] ([illegible]) [illegible] law of random assortment [illegible]

[illegible] 2003 [illegible] Davey Smith [illegible] Ebrahim [illegible] ([illegible]) [illegible] MR [illegible] ([illegible]) [illegible]

[illegible]

[illegible] MR [illegible] 20 [illegible] 80 [illegible]

(1) [illegible]

(2) [illegible]

Katan ([illegible]) [illegible] Apolipoprotein E [illegible] IV [illegible]

[illegible] MR [illegible] MR [illegible]

[illegible]

[illegible] MR [illegible] IV [illegible] MR [illegible]



Figure 1. MR [illegible] G_j [illegible] D [illegible] Y [illegible] U [illegible] G_j [illegible]

[illegible]

[illegible] 25.1 ([illegible] IV [illegible]). 结构形式为：

$$Y = \beta_0 + \beta D + \beta_u U + \varepsilon_Y, \tag{25.1}$$

$$D = \gamma_0 + \gamma_1 G_1 + \cdots + \gamma_p G_p + \gamma_u U + \varepsilon_D. \tag{25.2}$$

约简形式为：

$$Y = (\beta_0 + \beta\gamma_0) + \beta\gamma_1 G_1 + \cdots + \beta\gamma_p G_p + (\beta_u + \beta\gamma_u)U + \varepsilon_Y, \tag{25.3}$$

$$D = \gamma_0 + \gamma_1 G_1 + \cdots + \gamma_p G_p + \gamma_u U + \varepsilon_D. \tag{25.4}$$

[illegible] G_j [illegible] Y [illegible] $\Gamma_j = \beta\gamma_j$[illegible] $j = 1, \dots, p$[illegible]

[illegible] MR [illegible]

[illegible] 25.2 ([illegible]). 线性模型为：

$$Y = \beta_0 + \beta D + \alpha_1 G_1 + \cdots + \alpha_p G_p + \beta_u U + \varepsilon_Y, \tag{25.5}$$

$$D = \gamma_0 + \gamma_1 G_1 + \cdots + \gamma_p G_p + \gamma_u U + \varepsilon_D. \tag{25.6}$$

约简形式为：

$$Y = (\beta_0 + \beta\gamma_0) + (\alpha_1 + \beta\gamma_1)G_1 + \cdots + (\alpha_p + \beta\gamma_p)G_p + (\beta_u + \beta\gamma_u)U + \varepsilon_Y, \tag{25.7}$$

$$D = \gamma_0 + \gamma_1 G_1 + \cdots + \gamma_p G_p + \gamma_u U + \varepsilon_D. \tag{25.8}$$

[illegible] $\Gamma_j = \alpha_j + \beta\gamma_j$[illegible] $j = 1, \dots, p$[illegible] $\alpha_j \neq 0$[illegible] G_j [illegible]

25.1. 定义 [illegible]和 [illegible]与第 [illegible]章的线性 *IV* 模型略有不同——我们显式包含了未测量混杂因子 U。但这一差异本质上不影响讨论，因为 U 在约简形式中通过 $\beta_u + \beta\gamma_u$ 被合并到系数中。

[illegible]

[illegible] IV [illegible] MR [illegible] genome-wide association studies, GWAS[illegible] MR [illegible]

[illegible] 25.1 ([illegible] MR [illegible]). (1) 我们有治疗变量对遗传工具变量的回归系数 $\hat{\gamma}_1, \dots, \hat{\gamma}_p$ 及其标准误 $\text{se}_{D1}, \dots, \text{se}_{Dp}$。假设

$$\hat{\gamma}_1 \xrightarrow{p} \gamma_1, \dots, \hat{\gamma}_p \xrightarrow{p} \gamma_p \tag{25.9}$$

并忽略标准误的不确定性。

(2) 我们有结果变量对遗传工具变量的回归系数 $\hat{\Gamma}_1, \dots, \hat{\Gamma}_p$ 及其标准误 $\text{se}_{Y1}, \dots, \text{se}_{Yp}$。假设

$$\hat{\Gamma}_1 \xrightarrow{p} \Gamma_1, \dots, \hat{\Gamma}_p \xrightarrow{p} \Gamma_p \tag{25.10}$$

并忽略标准误的不确定性。

(3) 假设 $\hat{\gamma}_1, \dots, \hat{\gamma}_p, \hat{\Gamma}_1, \dots, \hat{\Gamma}_p$ 联合服从正态分布且相互独立。

[illegible] $\hat{\gamma}_j$ [illegible] $\hat{\Gamma}_j$ [illegible]

[illegible] $\alpha_j = 0$ [illegible] $\beta = \Gamma_j/\gamma_j$ [illegible] j [illegible] $\hat{\beta}_j = \hat{\Gamma}_j/\hat{\gamma}_j$ [illegible] β [illegible]

[illegible] delta [illegible] $\hat{\beta}_j$ [illegible]

$$\text{se}_j^2 = (\text{se}_{Yj}^2 + \hat{\beta}_j^2 \text{se}_{Dj}^2)/\hat{\gamma}_j^2. \tag{25.11}$$

[illegible] β [illegible] Fisher [illegible]

$$\hat{\beta}_{\text{fisher0}} = \frac{\sum_{j=1}^p \hat{\beta}_j/\text{se}_j^2}{\sum_{j=1}^p 1/\text{se}_j^2}, \tag{25.12}$$

[illegible] $\left(\sum_{j=1}^p 1/\text{se}_j^2\right)^{-1}$ [illegible] $\hat{\gamma}_j$ [illegible] se_{Dj} [illegible]

$$\hat{\beta}_{\text{fisher1}} = \frac{\sum_{j=1}^p \hat{\beta}_j \hat{\gamma}_j^2/\text{se}_{Yj}^2}{\sum_{j=1}^p \hat{\gamma}_j^2/\text{se}_{Yj}^2} = \frac{\sum_{j=1}^p \hat{\Gamma}_j \hat{\gamma}_j/\text{se}_{Yj}^2}{\sum_{j=1}^p \hat{\gamma}_j^2/\text{se}_{Yj}^2}, \tag{25.13}$$

[illegible] $\left(\sum_{j=1}^p \hat{\gamma}_j^2/\text{se}_{Yj}^2\right)^{-1}$ [illegible] $\hat{\beta}_{\text{fisher1}}$ [illegible] ([illegible]) [illegible]

$$\hat{\beta}_{\text{fisher1}} \xrightarrow{p} \frac{\sum_{j=1}^p \Gamma_j \gamma_j/\text{se}_{Yj}^2}{\sum_{j=1}^p \gamma_j^2/\text{se}_{Yj}^2} = \beta + \frac{\sum_{j=1}^p \alpha_j \gamma_j/\text{se}_{Yj}^2}{\sum_{j=1}^p \gamma_j^2/\text{se}_{Yj}^2}. \tag{25.14}$$

[illegible] j [illegible] $\alpha_j = 0$ [illegible] $\hat{\beta}_{\text{fisher1}}$ [illegible] α_j [illegible] γ_j [illegible] $1/\text{se}_{Yj}^2$ [illegible] $\hat{\beta}_{\text{fisher1}}$ [illegible]

25.2. 当有许多遗传工具变量，且违反排他性限制（由 α_j 捕获）是来自均值为零的分布的独立随机抽样时，上述条件近似成立。

[illegible]

$$\Gamma_j = \beta\gamma_j \quad (j = 1, \dots, p). \tag{25.15}$$

[illegible]

$$\hat{\Gamma}_j \approx \beta\hat{\gamma}_j \quad (j = 1, \dots, p). \tag{25.16}$$

[illegible] $\{\hat{\Gamma}_j\}_{j=1}^{p}$ [illegible] $\{\hat{\gamma}_j\}_{j=1}^{p}$ [illegible] $\hat{\Gamma}_j$ [illegible] $\hat{\gamma}_j$ [illegible] WLS [illegible] w_j [illegible] β [illegible] WLS [illegible] $\hat{\gamma}_j$ [illegible]

$$\hat{\beta}_{\text{egger1}} = \frac{\sum_{j=1}^{p} \hat{\gamma}_j \hat{\Gamma}_j w_j}{\sum_{j=1}^{p} \hat{\gamma}_j^2 w_j}, \tag{25.17}$$

[illegible] $w_j = 1/\mathrm{se}_{Yj}^2$ [illegible] $\hat{\beta}_{\text{fisher1}}$ WLS [illegible] $\hat{\gamma}_j$ [illegible]

$$\hat{\beta}_{\text{egger0}} = \frac{\sum_{j=1}^{p} (\hat{\gamma}_j - \hat{\gamma}_w)(\hat{\Gamma}_j - \hat{\Gamma}_w) w_j}{\sum_{j=1}^{p} (\hat{\gamma}_j - \hat{\gamma}_w)^2 w_j}, \tag{25.18}$$

[illegible] $\hat{\gamma}_w = \sum_{j=1}^{p} \hat{\gamma}_j w_j / \sum_{j=1}^{p} w_j$ [illegible] $\hat{\Gamma}_w = \sum_{j=1}^{p} \hat{\Gamma}_j w_j / \sum_{j=1}^{p} w_j$ [illegible] $\hat{\gamma}_j$ [illegible] $\hat{\Gamma}_j$ [illegible]

[illegible] γ_j [illegible]

$$\hat{\beta}_{\text{egger0}} \xrightarrow{p} \frac{\sum_{j=1}^{p} (\gamma_j - \gamma_w)(\Gamma_j - \Gamma_w) w_j}{\sum_{j=1}^{p} (\gamma_j - \gamma_w)^2 w_j} = \beta + \frac{\sum_{j=1}^{p} (\gamma_j - \gamma_w)(\alpha_j - \alpha_w) w_j}{\sum_{j=1}^{p} (\gamma_j - \gamma_w)^2 w_j}, \tag{25.19}$$

[illegible] γ_w, Γ_w [illegible] α_w [illegible] $\hat{\beta}_{\text{egger0}}$ [illegible] β [illegible] α_j [illegible] γ_j [illegible] WLS [illegible] $\alpha_j = 0$ [illegible] j [illegible] γ_j [illegible] α_j [illegible] **MP** ( ) [illegible] Egger [illegible]

$$\hat{\alpha}_{\text{egger0}} = \hat{\Gamma}_w - \hat{\beta}_{\text{egger0}} \hat{\gamma}_w, \tag{25.20}$$

InSIDE [illegible]

$$\Gamma_w - \beta\gamma_w = \alpha_w, \tag{25.21}$$

[illegible]

[illegible]

[illegible] `mr.raps` [illegible] `bmi.sbp` [illegible] ( ) [illegible] Fisher [illegible] Egger [illegible]

[illegible] Fisher [illegible]

```
fisher.weight = function(est, se)
{
    n = sum(est/se^2)
    d = sum(1/se^2)
    res = c(n/d, sqrt(1/d))
    names(res) = c("est", "se")
    res
}
```

[illegible] BMI-SBP [illegible] $\hat{\beta}_{\text{fisher0}}$ [illegible] $\hat{\beta}_{\text{fisher1}}$ [illegible]

```
> fisher.weight(bmisbp$iv, bmisbp$se.iv)
      est        se
0.31727680 0.05388827
> fisher.weight(bmisbp$iv, bmisbp$se.iv1)
      est        se
0.31576007 0.05893783
```

[illegible] Egger [illegible]

Fisher [illegible]

```
> mr.egger = lm(beta.output ~ 0 + beta.exposure,
+ data = bmisbp, weights = 1/se.output^2)
> summary(mr.egger)$coef
               Estimate Std. Error t value Pr(>|t|)
beta.exposure 0.3172768  0.1105994  2.868704 0.004681659

> mr.egger.w = lm(beta.output ~ beta.exposure,
+ data = bmisbp, weights = 1/se.output^2)
> summary(mr.egger.w)$coef
               Estimate Std. Error t value Pr(>|t|)
(Intercept) 0.0001133328 0.002079418 0.05450217 0.956603931
beta.exposure 0.3172989306 0.110948506 2.85987564 0.004811017
```

[illegible] Egger [illegible] $\hat{\gamma}_j \hat{\Gamma}_j$ [illegible]

FIGURE 2. [illegible] Egger [illegible]

[illegible]

MR [illegible] IV [illegible]

[illegible] MR [illegible]

[illegible] BMI [illegible] SUTVA [illegible]

[illegible] IV [illegible]
[illegible]
[illegible] $_1$ [illegible]
[illegible]
[illegible]

[illegible] MR [illegible] IV [illegible] $\hat{\gamma}_j$ [illegible] $\hat{\Gamma}_j$
[illegible] IV [illegible]
[illegible] $_1$ [illegible] $_1$ [illegible]
[illegible]

VanderWeele [illegible] (2014) [illegible] MR [illegible]

[illegible]

[illegible] 25.1. [illegible] *Fisher* 加权估计的推导 证明第 [illegible] 节中 $\hat{\beta}_{\text{fisher0}}$ 的方差为 $\left(\sum_{j=1}^{p} 1/\text{se}_j^2\right)^{-1}$。提示：利用 $\hat{\beta}_j$ 的独立性。

[illegible] 25.2. [illegible] *InSIDE* 假设的含义 解释 *InSIDE*（工具强度独立于直接效应）假设在什么意义上弱于 $\alpha_j = 0$（对所有 j）的排他性限制假设。给出一个满足 *InSIDE* 但不满足 $\alpha_j = 0$ 的数值例子。

[illegible] 25.3. [illegible] *Egger* 截距的解释 在第 [illegible] 节中，我们看到 *Egger* 回归的截距 $\hat{\alpha}_{\text{egger0}}$ 收敛到直接效应的加权平均 α_w。讨论为什么这个性质在实践中有用——即为什么我们可能关心估计 α_w 而不仅仅是 β。

[illegible] 25.4. [illegible] 数据分析 分析 *mr.raps* 包中的 *bmi.bmi* 数据。参见 [illegible] 第 *7.2* 节 了解更多细节。

[illegible]

[illegible] MR [illegible]
[illegible] Egger [illegible]
[illegible] $_1$ [illegible] robust adjusted profile score, RAPS [illegible]
[illegible] MR [illegible]

CHAPTER 26

[illegible]

[illegible]

[illegible] 2020 [illegible] [illegible] [illegible] 2019 [illegible] 1000 [illegible] 2021 [illegible] 1200 [illegible] 20% [illegible] ** [illegible] ** [illegible] [illegible] ** [illegible] **[illegible] panel data [illegible]

[illegible]

[illegible] ** [illegible] ** [illegible] ** [illegible] ** [illegible]

[illegible]

[illegible] DiD [illegible]

[illegible]

- $G_i \in \{0,1\}$ [illegible] i [illegible] $G_i = 1$ [illegible] $G_i = 0$ [illegible]
- $T_t \in \{0,1\}$ [illegible] $T_t = 0$ [illegible] $T_t = 1$ [illegible]
- Y_{it} [illegible] i [illegible] t [illegible]
- $D_i = G_i$ [illegible]

[illegible] N [illegible] $i = 1, \ldots, N$ [illegible] N_1 [illegible] N_0 [illegible] $N_1 + N_0 = N$ [illegible]

[illegible] i [illegible] t [illegible]

$$Y_{it}(1) \quad [illegible] \quad Y_{it}(0)$$

[illegible] i [illegible] t [illegible]

$$Y_{it} = Y_{it}(0) + D_i\big(Y_{it}(1) - Y_{it}(0)\big).$$

[illegible] t [illegible] Average Treatment effect on the Treated, ATT [illegible]

$$\tau_t = \mathbb{E}\big(Y_{it}(1) - Y_{it}(0) \mid G_i = 1\big).$$

[illegible]

$$\bar{Y}_{00} = \mathbb{E}(Y_{it} \mid G_i = 0, T_t = 0) \text{ [illegible]}$$

$$\bar{Y}_{01} = \mathbb{E}(Y_{it} \mid G_i = 0, T_t = 1) \text{ [illegible]}$$

$$\bar{Y}_{10} = \mathbb{E}(Y_{it} \mid G_i = 1, T_t = 0) \text{ [illegible]}$$

$$\bar{Y}_{11} = \mathbb{E}(Y_{it} \mid G_i = 1, T_t = 1) \text{ [illegible]}$$

[illegible]

[illegible] $\bar{Y}_{11} - \bar{Y}_{10}$ [illegible]

$$\bar{Y}_{11} - \bar{Y}_{10} = \underbrace{\tau_1}_{\text{[illegible]}} + \underbrace{\left[\bar{Y}_{10}(0) - \bar{Y}_{11}(0)\right]}_{\text{[illegible]}},$$

[illegible] $\bar{Y}_{1t}(0) = \mathbb{E}(Y_{it}(0) \mid G_i = 1, T_t = t)$ [illegible] t [illegible] $\bar{Y}_{01} - \bar{Y}_{00}$ [illegible]

$$\bar{Y}_{01} - \bar{Y}_{00} = \bar{Y}_{10}(0) - \bar{Y}_{11}(0).$$

[illegible]

$$\underbrace{(\bar{Y}_{11} - \bar{Y}_{10}) - (\bar{Y}_{01} - \bar{Y}_{00})}_{\text{[illegible]}} = \tau_1 + \underbrace{\left[\bar{Y}_{10}(0) - \bar{Y}_{11}(0)\right]}_{\text{[illegible]}} - \underbrace{\left[\bar{Y}_{10}(0) - \bar{Y}_{11}(0)\right]}_{\text{[illegible]}}.$$

[illegible]

$$\text{DiD} = (\bar{Y}_{11} - \bar{Y}_{10}) - (\bar{Y}_{01} - \bar{Y}_{00}) = \tau_1.$$

[illegible] ATT

DiD [illegible]

$$\hat{\tau}_{\text{DiD}} = (\bar{Y}_{11} - \bar{Y}_{10}) - (\bar{Y}_{01} - \bar{Y}_{00}) = (\bar{Y}_{11} - \bar{Y}_{01}) - (\bar{Y}_{10} - \bar{Y}_{00}).$$

> [illegible]
>
> $$\hat{\tau}_{\text{DiD}} = \underbrace{(\bar{Y}_{11} - \bar{Y}_{01})}_{\text{[illegible]}} - \underbrace{(\bar{Y}_{10} - \bar{Y}_{00})}_{\text{[illegible]}}.$$
>
> [illegible]

[illegible]

DiD [illegible]

[illegible] 26.1 ([illegible]). 在没有处理的情况下，处理组和对照组的结果随时间变化的趋势相同，即

$$\mathbb{E}\big(Y_{i1}(0) - Y_{i0}(0) \mid G_i = 1\big) = \mathbb{E}\big(Y_{i1}(0) - Y_{i0}(0) \mid G_i = 0\big).$$

Equivalently,

$$\bar{Y}_{10}(0) - \bar{Y}_{11}(0) = \bar{Y}_{00} - \bar{Y}_{01}.$$

[illegible] ** [illegible] **

[illegible]

[illegible] DiD [illegible] " " [illegible]

$$Y_{it}(0) = \alpha + \beta \cdot G_i + \gamma \cdot t + \varepsilon_{it},$$

[illegible]

[illegible] " " [illegible] DiD [illegible]

(1) [illegible] `"0300pre-trend test'[illegible]
(2) [illegible]
(3) [illegible] placebo test[illegible]

[illegible]

[illegible]

DiD [illegible] Two-Way Fixed Effects, TWFE[illegible]

$$Y_{it} = \alpha + \beta \cdot D_{it} + \gamma_i + \delta_t + \varepsilon_{it},$$

[illegible]

- $D_{it} = G_i \cdot T_t$ [illegible] × [illegible]
- γ_i [illegible]
- δ_t [illegible]

[illegible] β [illegible] DiD [illegible] $\hat{\tau}_{\text{DiD}}$ [illegible]

- [illegible] **I g ' 。—Q QN 、— ; l[illegible]N [illegible] N P p** 。 [illegible]
- **M a。J p g g[illegible] d** [illegible]
- **A p p—。**[illegible]

[illegible]

[illegible] TWFE [illegible] β [illegible] i [illegible] t [illegible] $\tau_{it} = \mathbb{E}(Y_{it}(1) - Y_{it}(0))$[illegible] i [illegible] t

Angrist and Pischke ([illegible]) [illegible] TWFE [illegible] ** [illegible] ** [illegible] ATT [illegible]

> [illegible] [illegible] TWFE [illegible]
> [illegible] β [illegible] ATT [illegible]
> [illegible]

[illegible] [illegible] DiD [illegible] ** [illegible] ** [illegible] $T_t \in \{0, 1, \ldots, T\}$ [illegible] $D_{it} = G_i \cdot \mathbb{I}(t \geq t^*)$ [illegible] t^* [illegible] DiD [illegible] TWFE [illegible]

$$Y_{it} = \alpha_i + \delta_t + \sum_{t=0}^{T} \beta_t \cdot D_{it} + \varepsilon_{it},$$

[illegible] β_t [illegible] $_1$ [illegible]

[illegible] **[illegible]** DiD [illegible] β_t [illegible]

[illegible]

[illegible] DiD [illegible]

DiD [illegible] pre-trend [illegible]

DiD [illegible] 2×2 [illegible] TWFE [illegible]

[illegible]

[illegible] DiD [illegible]

- [illegible] Mostly Harmless Econometrics [illegible] 5 [illegible] DiD [illegible]
- [illegible] DiD [illegible]
- [illegible] TWFE [illegible]
- [illegible] TWFE [illegible]

[illegible] DiD [illegible] Bootstrap [illegible]

CHAPTER 27

[illegible]

[illegible]

[illegible] “[illegible]” [illegible]
[illegible]
[illegible]
[illegible]

[illegible] transmission mechanism [illegible] [illegible] **[illegible]** [illegible]

[illegible] nested potential outcomes [illegible] mediation formula [illegible] **[illegible]** [illegible]

[illegible]

[illegible] Z [illegible] treatment [illegible] Y [illegible] outcome [illegible] M [illegible] mediator [illegible] X [illegible] covariates [illegible] DAG [illegible]

> [illegible] τ [illegible] M [illegible] M [illegible]

[illegible]

[illegible] 27.1 ([illegible]). (*2012*) 研究了染色体 *15q25.1* 上的遗传变异如何影响肺癌风险。[1] 该研究的核心问题是：遗传变异是通过影响吸烟行为（吸烟量）进而增加肺癌风险的，还是通过其他独立通路（如直接影响肺部细胞）直接致癌的？这就是一个典型的中介分析问题：遗传变异是处理 Z，吸烟是中介 M，肺癌是结果 Y。

[illegible] 27.2 ([illegible]). 等人 (*2018*) 研究了邻里贫困对青少年物质使用的影响，并探讨了这一影响是通过学校和同伴环境传导的。[2] 他们使用了美国国家共病调查（*National Comorbidity Survey*）的数据，将邻里贫困指数作为处理，青少年物质使用（如饮酒、吸毒）作为结果，而学校和同伴环境因素作为中介。

---

[1] **?**

[2] **?**

[illegible] 27.3 ([illegible]). *JOBS II* 是一个关于就业培训对失业工人影响的现场实验。[3] 该培训项目的目的不仅是提高再就业率，还希望改善求职者的心理健康。研究者感兴趣的是：培训对心理健康的效应中，有多少是通过提高工作搜寻效能这一中介传导的，又有多少是通过其他途径（如社会支持网络）直接产生的。[4]

[illegible]

[illegible] [illegible] nested potential outcomes [illegible] z [illegible]

$$\{M(z),\ Y(z) : z = 0,1\}.$$

[illegible] z [illegible] m [illegible]

$$\{Y(z,m) : z = 0,1,\ m \in \mathcal{M}\},$$

[illegible] $\mathcal{M}$ [illegible]

[illegible] nested interventions [illegible] z [illegible] m [illegible] $M_z \equiv M(z')$ [illegible]

$$\{Y(z, M_z) : z = 0,1,\ z' = 0,1\}.$$

[illegible] $Y(z, M_z)$ [illegible] z [illegible] z' [illegible] M_z [illegible] z [illegible] z' [illegible] “[illegible]” [illegible]

$$\{Y(1,M_1),\ Y(1,M_0),\ Y(0,M_1),\ Y(0,M_0)\}.$$

[illegible] $Y(1,M_1)$ [illegible] $z=1$ [illegible] $z=1$ [illegible] $Y(0,M_0)$ [illegible] $z=0$ [illegible] $z=0$ [illegible]

Composition [illegible]

[Composition [illegible]] $Y(z, M_z) = Y(z)$ [illegible] $z = 0,1$ [illegible]

Composition [illegible] $Y(1)$ [illegible] $Y(1,M_1)$ [illegible] $Y(0)$ [illegible] $Y(0,M_0)$ [illegible]

[illegible]

[illegible] 27.1 ([illegible]). 总效应（*total effect*）定义为

$$\tau = \mathbb{E}\{Y(1) - Y(0)\}.$$

自然直接效应（*natural direct effect, NDE*）定义为

$$= \mathbb{E}\{Y(1,M_0) - Y(0,M_0)\}.$$

自然间接效应（*natural indirect effect, NIE*）定义为

$$= \mathbb{E}\{Y(1,M_1) - Y(1,M_0)\}.$$

[illegible]

- [illegible] τ [illegible] 0 [illegible] 1 [illegible]
- **[illegible]** $\mathbb{E}\{Y(1,M_0) - Y(0,M_0)\}$ [illegible] 0 [illegible] 1 [illegible] M_0 [illegible] M [illegible]

---

[3] [illegible] ?? [illegible]

[4] [illegible] ?? [illegible] Baron–Kenny [illegible]

- **k ` P**$\mathbb{E}\{Y(1,M_1)-Y(1,M_0)\}$ [illegible] $z=1$ [illegible] M_0 [illegible] M_1 [illegible] M [illegible]

Composition [illegible]

[illegible] **w** [illegible] **w**在 [illegible] 和 [illegible] 下，有

$$\tau = +.$$

[illegible] **k ` P** [illegible] $Y(1,M_1)-Y(1,M_0)$ [illegible] $z=1$ [illegible] “[illegible]” [illegible] “[illegible]” [illegible] “[illegible]” [illegible]

[illegible] $Y(1,M_0)$ [illegible] **J p ‘ l — Q V ; J — ‘ ;** [illegible] cross-world counterfactual [illegible] $z=1$ [illegible] $m=M_0$ [illegible] $z=1$ [illegible] $Y(1,M_0)$ [illegible]

[width = 0.7]figures/crossworld.pdf

FIGURE 2. [illegible] counterfactual $Y(1,M_0)$ [illegible] $Y(0,M_1)$ [illegible]

[illegible] $Y(1,M_0)$ [illegible] $Y(0,M_1)$ [illegible] **J p ‘ l — Q V ; J — ‘ ; g** a priori counterfactuals [illegible] Popperian [illegible][5]

[illegible] $Y(1)$ [illegible] $Y(0)$ [illegible] $Y(1,M_0)$ [illegible] $\{Y(1),Y(0)\}$ [illegible]

X w [illegible]

X w q [illegible] **w** [illegible] X [illegible]

[[illegible]] [illegible] X [illegible]

$$Z \perp\!\!\!\perp Y(z,m) \mid X, \quad \forall z, m.$$

[[illegible]] [illegible] (X,Z) [illegible]

$$M \perp\!\!\!\perp Y(z,m) \mid (X,Z), \quad \forall z, m.$$

[[illegible]] [illegible] X [illegible]

$$Z \perp\!\!\!\perp M(z) \mid X, \quad \forall z.$$

---

[5] [illegible] **?** [illegible] **?**

[illegible] sequential ignorability [illegible] X [illegible] (X, Z) [illegible] X [illegible] (Z, M) [illegible]

$$(Z, M) \perp\!\!\!\perp Y(z, m) \mid X, \quad \forall z, m. \tag{136}$$

[illegible][6]

[illegible]

[[illegible]] [illegible] X [illegible]

$$Y(z, m) \perp\!\!\!\perp M(z') \mid X, \quad \forall z, z', m.$$

[illegible] $z \neq z'$ [illegible] $Y(z, m)$ [illegible] $M(z')$ [illegible] $Y(1, M_0)$ [illegible] cross-world counterfactual [illegible]

[illegible] 27.4 ([illegible]). 给定 X，我们按如下方式生成数据：

$$\begin{aligned} Z &= 1\{g_Z(X, \varepsilon_Z) \geq 0\}, \\ M(z) &= 1\{g_M(X, z, \varepsilon_M) \geq 0\}, \quad z = 0, 1, \\ Y(z, m) &= g_Y(X, z, m, \varepsilon_Y), \quad z, m = 0, 1, \end{aligned}$$

其中 $\varepsilon_Z, \varepsilon_M, \varepsilon_Y$ 相互独立。[7] 可以验证，在上述数据生成过程下，[illegible]均成立。如果允许 ε_M 和 ε_Y 依赖于处理水平（即改为 $\varepsilon_M(z)$ 和 $\varepsilon_Y(z, m)$），则这些假设可能被违反。

[illegible]

[illegible]（[illegible]）在[illegible]下，[illegible]有

$$\mathbb{E}\{Y(z, M_z) \mid X = x\} = \sum_m \mathbb{E}(Y \mid Z = z, M = m, X = x)\ \Pr(M = m \mid Z = z', X = x).$$

[illegible] counterfactual $Y(z, M_z)$ [illegible] $Y(z, M_z)$ [illegible] $Z = z, M = m, X = x$ [illegible] $Z = z', X = x$ [illegible]

[illegible] $\Pr(M = m \mid Z = z', X = x) = \Pr(M_z = m \mid X = x)$ [illegible] $Y(z, m)$ [illegible] M_z [illegible]

[illegible] X [illegible]

$$\mathbb{E}\{Y(z, M_z)\} = \sum_x \sum_m \mathbb{E}(Y \mid Z = z, M = m, X = x)\ \Pr(M = m \mid Z = z', X = x)\ \Pr(X = x).$$

[illegible] M [illegible] X [illegible] NDE [illegible] NIE [illegible]

---

[6] ??

[7] ?

[illegible] (NDE [illegible] NIE [illegible]) 在 [illegible] 下 [illegible], [illegible] 条件 *NDE* 和 *NIE* 由下式识别：

$$(x) = \sum_m \big[\mathbb{E}(Y \mid Z=1, M=m, X=x) - \mathbb{E}(Y \mid Z=0, M=m, X=x)\big] \Pr(M=m \mid Z=0, X=x), \tag{137}$$

$$(x) = \sum_m \mathbb{E}(Y \mid Z=1, M=m, X=x) \big[\Pr(M=m \mid Z=1, X=x) - \Pr(M=m \mid Z=0, X=x)\big]. \tag{138}$$

[illegible] (x) [illegible] 0 [illegible] 1 [illegible] m [illegible] 0 [illegible] 1 [illegible] $\mathbb{E}(Y \mid Z=1, M=m, X=x) - \mathbb{E}(Y \mid Z=0, M=m, X=x)$ [illegible] $Z=0$ [illegible] $\Pr(M=m \mid Z=0, X=x)$ [illegible]

[illegible] $M \in \{0,1\}$ [illegible] NIE [illegible]

[illegible] NIE [illegible]) 在 [illegible] 下 [illegible], [illegible] 对 [illegible] 二值中介 M，有

$$(x) = \tau_{Z\to M}(x) \cdot \tau_{M\to Y}(1, x),$$

其中

$$\tau_{Z\to M}(x) = \Pr(M=1 \mid Z=1, X=x) - \Pr(M=1 \mid Z=0, X=x)$$

是 Z 对 M 的条件平均因果效应，

$$\tau_{M\to Y}(z, x) = \mathbb{E}(Y \mid Z=z, M=1, X=x) - \mathbb{E}(Y \mid Z=z, M=0, X=x)$$

是 M 对 Y 的条件平均因果效应。

[illegible] Z [illegible] M [illegible] $\tau_{Z\to M}$ [illegible] M [illegible] Y [illegible] $\tau_{M\to Y}$ [illegible]

[illegible][8]

[illegible] A ; p l 7 d Q l l 》

[illegible] A ; p l 7 d Q l l 》 [illegible]

[illegible] Baron–Kenny [illegible] [Baron–Kenny [illegible]] [illegible]

$$\begin{cases} \mathbb{E}(M \mid Z, X) = \beta_0 + \beta_1 Z + \beta_2^\top X, \\ \mathbb{E}(Y \mid Z, M, X) = \theta_0 + \theta_1 Z + \theta_2 M + \theta_4^\top X. \end{cases} \tag{27.1}$$

[illegible] NDE [illegible] NIE [illegible]

[illegible] (Baron–Kenny [illegible]) 在 [illegible] 下 [illegible], [illegible] 有 [illegible]

$$= \theta_1, \qquad = \theta_2 \beta_1. \tag{27.2}$$

---

[8] [illegible] ??

M : Y θ_1 $Z \to Y$ $\beta_1\theta_2$
$Z \to M \to Y$

X (x) (x) $X = x$
Baron–Kenny θ_1 $\theta_2\beta_1$ x

Baron–Kenny
OLS $\hat{\beta}_1$ $\hat{\theta}_2$ NDE NIE

$$\hat{} = \hat{\theta}_1, \qquad \hat{} = \hat{\theta}_2\hat{\beta}_1,$$

Baron–Kenny delta $\hat{}$

$$\hat{\mathrm{var}}(\hat{}) = \hat{\mathrm{var}}(\hat{\theta}_2)\hat{\beta}_1^2 + \hat{\theta}_2^2\hat{\mathrm{var}}(\hat{\beta}_1).$$

Sobel's test[9]

A ; p l 7 d Q l l

b o A、

R Baron–Kenny JOBS II[10]

27.5 (JOBS II). 我们关注就业培训（处理 Z）对心理健康（结果 Y，用抑郁量表得分度量）的影响，以及这一影响是否通过工作搜寻效能（中介 M）传导。背景协变量 X 包括经济困难程度、基线抑郁得分、性别、年龄、职业、婚姻状态、种族、教育程度和家庭收入。

通过以下 R 代码实现 Baron–Kenny 方法：

```
library("car")
BKmediation = function(Z, M, Y, X) {
  mediator.reg   = lm(M ~ Z + X)
  mediator.Zcoef = mediator.reg$coef[2]
  mediator.Zse   = sqrt(hccm(mediator.reg)[2, 2])

  outcome.reg    = lm(Y ~ Z + M + X)
  outcome.Zcoef  = outcome.reg$coef[2]
  outcome.Zse    = sqrt(hccm(outcome.reg)[2, 2])
  outcome.Mcoef  = outcome.reg$coef[3]
  outcome.Mse    = sqrt(hccm(outcome.reg)[3, 3])

  NDE = outcome.Zcoef
  NIE = outcome.Mcoef * mediator.Zcoef

  NDE.se = outcome.Zse
  NIE.se = sqrt(outcome.Mse^2 * mediator.Zcoef^2 +
                outcome.Mcoef^2 * mediator.Zse^2)
```

---

[9] Sobel ?

[10] JOBS II R `mediation`

```
  res = matrix(c(NDE, NIE, NDE.se, NIE.se,
                 NDE/NDE.se, NIE/NIE.se), 2, 3)
  rownames(res) = c("NDE", "NIE")
  colnames(res) = c("est", "se", "t")
  res
}
```

运行结果如下：

```
> res = BKmediation(Z, M, Y, X)
> round(res, 3)
        est      se       t
NDE -0.037  0.042  -0.885
NIE -0.014  0.009  -1.528
```

估计的直接效应 ˆ= −0.037（培训对心理健康的直接保护效应）和间接效应 ˆ= −0.014（通过提高工作搜寻效能而改善心理健康）均为负，但统计上均不显著。这表明 *JOBS II* 培训项目对心理健康的效应尚无充分证据支持按中介路径分解。

[illegible] $Z \to M$ [illegible] $\beta_1 > 0$ [illegible] $M \to Y$ [illegible] $\theta_2 > 0$ [illegible] Z [illegible] M [illegible] $\theta_3 < 0$ [illegible] $\beta_1(\theta_2 + \theta_3)$ [illegible] mediator paradox [illegible] surrogate endpoint paradox [illegible][a]

---

[a] [illegible] ?

[illegible]

[illegible] $Y(z, m)$ [illegible] $M(z')$ [illegible] $z \neq z'$ [illegible] sensitivity analysis [illegible]

- [illegible] Cornfield [illegible]
- [illegible] Baron–Kenny [illegible]

[illegible]

R [illegible]

[illegible]

(1) [illegible]

(2) [illegible] $Y(z, M_z)$ [illegible] counterfactual [illegible] NDE [illegible] NIE [illegible]

(3) [illegible] Composition [illegible] $\tau = +$

(4) [illegible] $+$ [illegible]

(5) [illegible] NDE [illegible] NIE [illegible]

(6) [illegible] $= \theta_1$ [illegible] $= \theta_2\beta_1$

(7) [illegible] Popperian [illegible]

(8) [illegible]

{ { Baron–Kenny {

{ {

27.1. 假设 (Z, M, Y) 均为二值变量。基于潜在结果

$$M(1), M(0), Y(1, M_1), Y(1, M_0), Y(0, M_1), Y(0, M_0)$$

的联合取值，单元可以分成多少类型？

若进一步假设单调性 $M(1) \geq M(0)$（Z 对 M 的单调性），则单元可以分成多少类型？

若再假设 $Y(1, m) \geq Y(0, m)$ 且 $Y(z, 1) \geq Y(z, 0)$ 对所有 z, m 成立（Z 和 M 对 Y 的单调性），则单元可以分成多少类型？

{ {

27.2. 证明 { {与 { { {等{价。

{ {

27.3. { 提出了另一组假设来推导中介公式：

$$\{Y(z, m), M(z')\} \perp\!\!\!\perp Z \mid X, \qquad Y(z, m) \perp\!\!\!\perp M(z') \mid (Z = z', X),$$

对所有 z, z', m 成立。

证明：在上述假设下，{ {中的中介公式同样成立。

{ {

27.4. 在 *Baron–Kenny* 线性模型（{ }{ 下，首先拟合：

$$\hat{M} = \hat{\beta}_0 + \hat{\beta}_1 Z + \hat{\beta}_2^\top X, \qquad \hat{Y} = \hat{\theta}_0 + \hat{\theta}_1 Z + \hat{\theta}_2 M + \hat{\theta}_4^\top X.$$

则 *NIE* 的估计量为乘积 $\hat{\theta}_2 \hat{\beta}_1$（乘积法）。

另一方面，首先拟合：

$$\hat{Y} = \hat{\alpha}_0 + \hat{\alpha}_1 Z + \hat{\alpha}_1^\top X,$$

再拟合：

$$\hat{Y} = \tilde{\theta}_0 + \tilde{\theta}_1 Z + \tilde{\theta}_2 M + \tilde{\theta}_4^\top X,$$

则 *NIE* 的另一估计量为差值 $\hat{\alpha}_1 - \tilde{\theta}_1$（差值法）。

证明：$\hat{\alpha}_1 - \tilde{\theta}_1 = \hat{\theta}_2 \hat{\beta}_1$。

{ {

27.5. 证明 { {(二值中介的 *NIE* 乘积公式)。

{ {

27.6. { 建议在结果模型中加入处理与中介的交互项：

$$\begin{cases} \mathbb{E}(M \mid Z, X) = \beta_0 + \beta_1 Z + \beta_2^\top X, \\ \mathbb{E}(Y \mid Z, M, X) = \theta_0 + \theta_1 Z + \theta_2 M + \theta_3 ZM + \theta_4^\top X. \end{cases}$$

证明：在此模型下，

$$= \theta_1 + \theta_3\{\beta_0 + \beta_2^\top \mathbb{E}(X)\}, \qquad = (\theta_2 + \theta_3)\beta_1.$$

并说明如何在 *IID* 数据下估计 和 。

V̈ ɔB‖ȯɦ { {

ű 27.7. 考虑中介 M 服从正态线性模型、结果 Y 服从 *logistic* 模型：

$$\begin{cases} M \mid Z, X \sim \mathcal{N}(\beta_0 + \beta_1 Z + \beta_2^\top X, \sigma_M^2), \\ \text{logit}\{\Pr(Y = 1 \mid Z, M, X)\} = \theta_0 + \theta_1 Z + \theta_2 M + \theta_4^\top X. \end{cases}$$

请将 和 表示为模型参数和 X 分布的函数，并说明如何在 *IID* 数据下估计它们。

V̈ ɔB‖ȯɦ { {

ű 27.8. 考虑二值中介 M 服从 *logistic* 模型、结果 Y 服从线性模型：

$$\begin{cases} \text{logit}\{\Pr(M = 1 \mid Z, X)\} = \beta_0 + \beta_1 Z + \beta_2^\top X, \\ \mathbb{E}(Y \mid Z, M, X) = \theta_0 + \theta_1 Z + \theta_2 M + \theta_4^\top X. \end{cases}$$

证明：

$$= \theta_1, \qquad = \theta_2 \, \mathbb{E}\Big\{ \text{expit}(\beta_0 + \beta_1 + \beta_2^\top X) - \text{expit}(\beta_0 + \beta_2^\top X) \Big\}.$$

并说明如何在 *IID* 数据下估计 和 。

V̈ ɔB‖ȯɦ { {

ű 27.9. 将嵌套潜在结果修改为

$$Y(z, F_{M_z \mid X}) = \int Y(z, m) \, f(M_z = m \mid X) \, \mathrm{d}m,$$

其中 $F_{M_z \mid X}$ 是条件分布而非确定值。相应地将 *NDE* 和 *NIE* 定义为

$$= \mathbb{E}\{Y(1, F_{M_0|X}) - Y(0, F_{M_0|X})\}, \quad = \mathbb{E}\{Y(1, F_{M_1|X}) - Y(1, F_{M_0|X})\}.$$

证明：在仅假设 { { { {(不要求跨世界独立性）下， 和 的识别公式与正文相同。

V̈ ɔB‖ȯɦ { {

ű 27.10. { 和 { 综述并比较了主分层与中介分析两种方法。请查阅这些文献，并撰写一页总结，阐述两种方法的异同与各自的优缺点。

V̈. ų ẅ ű ŏ { {

*> Sh1_ k3

yT?C

RX„ý Bñ¦•qq¥Ù5

-ë¥c«Ï áÌL!yT² X©ii'X©'tM ^ðya'tM ^² Tb–7 CLÏ áÌªªë Bñ÷¹[4¥Ù5 ?ŽV” ÏÐ M -W¥ yT1"$

ƒñÙ5¥×1Ÿ,ý7ÜbSÐ?C¥'Éü^V4³” Ïw yT² båN

C K2b GYBWM/vL ?CÇÍË£TV[Í ÊÓh Œ ¸ fƒ/ L Ä ç^,V› ,¡£¥báÌ?ŽÇV$î4³¥” Ï ?CM -W¥yT1"$

ƒñÙ5 }Ð ;$ª¹^,V?¥b óM1, cyT ôii ƒ ìý¡ d9 Ð¥.dÆðb–7 V€ØEMn£ BÆá ¥ê B" Ú•!9¥L!/ yT² V[VBe¥4³” ÏÐ b

ƒÆê ¥5z “ Cm/2 Sð?`ZH ¿m¥yTwØØ,[# S2i2` aTaB`i2b *H `F :HvK„Q_mB`+? `/ a+?ð2ÌByM2b S*©ØEV” Ï4|yT² b'c |º ƒB5×¥ '±X„ZEb

kXyT”:V•HqÐÑôŸ

), 8ØE-- áÌ³1Ø³yT?C¥V?ŸHqbInBñ cM $X_1;X_2;:::;X_p$ ¥"dbÂTƒtM -Wi yT1" áÌYÈ¨µ_íÌm . : ŸVU Äñ«Ä}VBñM ph X! Y VUX ^ Y ¥°¤ðy -Bb

Bñô'Ÿ¥4³^ ¸ †Ø¥yT ˜/ M -W¥HqÿëŸ c"yT ² ¥• b

(yT”:V•Hq) óçBñM ¥îµ°¤ðy ¾M ÿë¿îµd ª}¥M b

°4 ÂT X •Á Y 7 Y »•Á Z *¹ X© Y ¥fƒ/ X „ Z -W, ‹¾µ°¤¥ó"iiñÌ-W¥©…1óûXÜ$ Y ³d b

(ÑôŸHq) mÏi ¥HqÿëŸ1" z ‹¿” sƒÏi ¥HqÿëŸ 1"b

ÑôŸHq]¸ óF†ÿëŸ ôii' ñM ” ÏHqÿë Œ mÏ•YV BH^åMõbƒÕF† •”Ä ˜Ï^,© ¥b

ƒ ñL!/ áÌV[VM ¥HqÿëŸ² Ï |yT• b

jX S*ØE VHqÿëŸžyT O

S*ØE aTB`i2b 2i H^Xy-TRyCy5y×KÜÅ¥ØE-Bbñ¥,•±Xe± 7µï VHqÿëŸ ? ï„'çyT² b

jXR»XB„ ?C O XØEVBñ›†í_¥m ? YVHqÿëŸ_ ï„÷ "Hb

[illegible] X [illegible] Y [illegible] S [illegible]
[illegible]

$$X \perp\!\!\!\perp Y \mid S \tag{139}$$

[illegible] $\perp\!\!\!\perp$ [illegible] (X, Y) [illegible]
[illegible]

[illegible] 28.1 ([illegible]). 考虑三个变量 X, Y, Z。假设我们进行条件独立性检验:

- $X \not\perp\!\!\!\perp Y$（边缘不独立）
- $X \perp\!\!\!\perp Y \mid Z$（给定 Z 条件独立）

这告诉我们什么？在 *Markov* 条件下，这暗示 Z 是 X 和 Y 的共同原因，或者 Z 位于它们之间的中介路径上。

[illegible]

[illegible] [illegible] v-[illegible]
[illegible] X, Y, Z [illegible] $X - Z - Y$ [illegible] X [illegible] Y [illegible] Z [illegible] $X \not\perp\!\!\!\perp Y \mid \emptyset$ [illegible] $X \perp\!\!\!\perp Y \mid Z$ [illegible]
Markov [illegible] $X \to Z \to Y$ [illegible] Z [illegible] Z [illegible] X [illegible] Y [illegible] $X \perp\!\!\!\perp Y \mid Z$
[illegible] Z [illegible] X [illegible] Y [illegible] $Z \to X$ [illegible] $Z \to Y$ [illegible] X [illegible] Y [illegible] Z [illegible]
PC [illegible]
[v-[illegible]] [illegible] (X, Z, Y) [illegible] v-[illegible]

(1) X [illegible] Y [illegible]
(2) Z [illegible] X [illegible] Z [illegible] Y [illegible]
(3) [illegible] Z [illegible] X [illegible] Y [illegible]

[illegible] v-[illegible] $X \to Z \leftarrow Y$ [illegible] $X \to Z \to Y$ [illegible] $X \leftarrow Z \to Y$ [illegible] Z [illegible] X [illegible] Y [illegible]
PC [illegible]

(1) [illegible] v-[illegible]
(2) [illegible] v-[illegible]

[illegible] [illegible]
[illegible] $X - Y$ [illegible] $Y - Z$ [illegible] X [illegible] Z [illegible] $X \to Y \to Z$ [illegible] $X \leftarrow Y \leftarrow Z$ [illegible]
[illegible]

[illegible]

[illegible] **[illegible]** [illegible] DAG [illegible] PAG [illegible]

- [illegible] $\circ - \circ$ [illegible]
- [illegible] $\to$ [illegible] $\leftarrow$ [illegible]

[illegible] **R** [illegible]) 在因果马尔可夫条件和忠诚性条件下，观察分布唯一确定一个 *PAG*。

[illegible] PC [illegible] PAG [illegible]

[illegible]

PC [illegible] nontrivial [illegible]

W [illegible] S [illegible] X [illegible] Y [illegible] $\rho_{XY\cdot S}$ [illegible] $H_0 : \rho_{XY\cdot S} = 0$ [illegible] p

28.1. 偏相关系数的计算需要 S 中变量的条件协方差矩阵。当 $|S|$ 较大时，估计精度会下降，这是 *PC* 算法的一个 *practical limitation*。

W [illegible]

$$I(X;Y \mid S) = \sum_{x,y,s} p(x,y,s) \log \frac{p(x,y \mid s)}{p(x \mid s)p(y \mid s)}. \tag{140}$$

[illegible]

[illegible]

[illegible] PC [illegible]

(1) [illegible]
(2) [illegible] (X,Y) [illegible] S [illegible] $X \perp\!\!\!\perp Y \mid S$ [illegible] (X,Y)
(3) [illegible] (X,Z,Y) [illegible] v-[illegible] $X \to Z \leftarrow Y$
(4) [illegible]
(5) [illegible] PAG

[illegible] p [illegible] $\sum_{k=0}^{p-2}\binom{p-2}{k}$ [illegible] PC [illegible]

[illegible]

PC [illegible]

W [illegible] GIES [illegible] Greedy Equivalence [illegible] Search [illegible] DAG [illegible] BIC [illegible] BD [illegible]

[illegible]

W [illegible] DAG [illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

[illegible] NP [illegible]

[illegible]

[illegible]

[illegible] Pearl (2000) [illegible] Causality [illegible] Spirtes et al. (2000) [illegible] Causation, Prediction, and Search [illegible]

[illegible]

- [illegible]
- [illegible]
- [illegible]

[illegible]

[illegible]

[illegible]

(1) [illegible]
(2) PC [illegible] v-[illegible]
(3) [illegible] PAG [illegible]
(4) [illegible]

[illegible]

[illegible]

[illegible] 28.1. 证明：在马尔可夫条件和忠诚性条件下，如果 X 和 Y 在骨架中无边，但都与 Z 相连，且 $X \not\perp\!\!\!\perp Y \mid \emptyset$ 但 $X \perp\!\!\!\perp Y \mid Z$，则 (X, Z, Y) 形成 v-结构。

[illegible] 28.2. 考虑四个变量 X_1, X_2, X_3, X_4，假设真实因果结构是 $X_1 \to X_2 \to X_4$ 和 $X_1 \to X_3 \to X_4$。说明 PC 算法在什么条件下能够恢复这个结构，以及在什么条件下可能失败。

[illegible] 28.3. 证明：对于 n 个变量的 DAG，完全图（有 $\binom{n}{2}$ 条边）和完全无向图在骨架发现步骤后的边数关系。

ǘ 28.4. 讨论因果发现与因果推断其他方法（如第 *10-12* 章的倾向得分方法）的关系。它们分别适合什么场景？

CHAPTER 29

[illegible]

[illegible]

[illegible]

[illegible] Z_1 [illegible] X_1 [illegible] Z_2 [illegible] Y [illegible] 5 [illegible]

- Z_1 [illegible] X_1 [illegible]
- X_1 [illegible] Z_2 [illegible] Y [illegible]
- [illegible] X_1 [illegible]

[illegible] X_0 [illegible] X_1 [illegible] Z_2 [illegible] Y [illegible] Z_1 [illegible] Z_1 [illegible] Y [illegible]

[illegible]

[illegible]

[illegible]

$$X_0 \to Z_1 \to X_1 \to Z_2 \to Y$$

[illegible]

- X_0 [illegible]
- Z_1 [illegible] 1 [illegible]
- X_1 [illegible] 1 [illegible] 2 [illegible]
- Z_2 [illegible] 2 [illegible]
- Y [illegible]

[illegible] (Z_1, Z_2) [illegible]

$$Y(z_1, z_2), \quad z_1, z_2 \in \{0, 1\}$$

[illegible]

$$Y = Y(Z_1, Z_2) = \sum_{z_1=0}^{1} \sum_{z_2=0}^{1} \mathbf{1}(Z_1 = z_1)\mathbf{1}(Z_2 = z_2)Y(z_1, z_2)$$

[illegible] $Z \perp Y(z) \mid X$ [illegible]

[illegible]

[illegible] 29.1 ([illegible]). *(1)* Z_1 在给定 X_0 条件下随机化:

$$Z_1 \perp Y(z_1, z_2) \mid X_0, \quad \forall z_1, z_2 \in \{0,1\}$$

(2) Z_2 在给定 (Z_1, X_1, X_0) 条件下随机化:

$$Z_2 \perp Y(z_1, z_2) \mid (Z_1, X_1, X_0), \quad \forall z_1, z_2 \in \{0,1\}$$

[illegible] Z_1 [illegible] X_0 [illegible] X_0 [illegible] Z_1 [illegible] Z_2 [illegible] (Z_1, X_1, X_0) [illegible] X_0 [illegible] Z_1 [illegible] X_1

[illegible] X_1 [illegible] Y [illegible] U [illegible] X_1 [illegible] Z_1 [illegible] Z_1 [illegible] Y [illegible] U [illegible] X_1 [illegible] Y [illegible] Z_1, Z_2 [illegible]



Figure 1. Assumption [illegible] X_1 [illegible] Y [illegible] U [illegible]



Figure 2. Assumption [illegible] X_1 [illegible] Y [illegible] U [illegible] (Z_1, Z_2) [illegible] Y [illegible] X_1 [illegible] Y [illegible]

[illegible]

[illegible]

$$E\{Y(z)\} = E\{E(Y \mid Z = z, X)\}$$

[illegible] X [illegible]

$$E\{Y(z)\} = \sum_{x} E(Y \mid Z = z, X = x)\Pr(X = x)$$

[illegible] X [illegible]

$$E\{Y(z)\} = \int E(Y \mid Z = z, X = x) f(x)\,dx$$

[illegible]

[illegible] 在 *Assumption* [illegible] 下，

$$E\{Y(z_1, z_2)\} = E\Big[E\{E(Y \mid z_2, z_1, X_1, X_0) \mid z_1, X_0\}\Big] \tag{29.1}$$

[illegible] $E(Y \mid z_2, z_1, X_1, X_0)$ [illegible] (z_1, X_0) [illegible] X_1 [illegible] z_2 [illegible] $E\{\cdot \mid z_1, X_0\}$ [illegible] X_1 [illegible] $\Pr(X_1 \mid z_1, X_0)$ $Z_1 = z_1$ [illegible] $E[\cdot]$ [illegible] X_0 [illegible]

[illegible] X_0, X_1 [illegible]

$$E\{Y(z_1, z_2)\} = \sum_{x_0}\sum_{x_1} E(Y \mid z_2, z_1, x_1, x_0)\Pr(x_1 \mid z_1, x_0)\Pr(x_0) \tag{29.2}$$

[illegible] X_0, X_1 [illegible]

$$E\{Y(z_1, z_2)\} = \iint E(Y \mid z_2, z_1, x_1, x_0) f(x_1 \mid z_1, x_0) f(x_0)\,dx_1\,dx_0 \tag{29.3}$$

[illegible]

$$\begin{aligned} E(Y) = &\sum_{x_0}\sum_{z_1}\sum_{x_1}\sum_{z_2} E(Y \mid z_2, z_1, x_1, x_0) \\ &\Pr(z_2 \mid z_1, x_1, x_0)\Pr(x_1 \mid z_1, x_0)\Pr(z_1 \mid x_0)\Pr(x_0) \end{aligned} \tag{29.4}$$

[illegible] $\Pr(z_2 \mid \cdots)$ [illegible] $\Pr(z_1 \mid \cdots)$ [illegible] $Y(z_1, z_2)$ [illegible] $Z_1 = z_1$ [illegible] $Z_2 = z_2$ [illegible] Robins [illegible][1]

[illegible]

[illegible] $E(Y \mid z_2, z_1, x_1, x_0)$ [illegible] $\Pr(x_1 \mid z_1, x_0)$ [illegible] $\Pr(x_0)$ [illegible] ” [illegible] ” [illegible] 29.1 [illegible]

[illegible] 29.1 ([illegible] 29.1 [illegible]). 假设线性结果模型：

$$E(Y \mid z_2, z_1, x_1, x_0) = \beta_0 + \beta_1 z_2 + \beta_2 z_1 + \beta_3 x_1 + \beta_4 x_0 \tag{29.5}$$

---

[1] [illegible] **??**

可以验证：

$$
\begin{aligned}
E\{Y(z_1,z_2)\} =& \sum_{x_0}\sum_{x_1}(\beta_0+\beta_1 z_2+\beta_2 z_1+\beta_3 x_1+\beta_4 x_0) \\
& \Pr(x_1 \mid z_1, x_0)\Pr(x_0) \\
=&\beta_0+\beta_1 z_2+\beta_2 z_1+\beta_3\sum_{x_0}E(X_1 \mid z_1,x_0)\Pr(x_0)+\beta_4 E(X_0) \\
=&\beta_0+\beta_1 z_2+\beta_2 z_1+\beta_3 E\{X_1(z_1)\}+\beta_4 E(X_0)
\end{aligned}
\tag{29.6}
$$

其中我们定义了 X_1 在治疗 $Z_1 = z_1$ 下的潜在结局：

$$E\{X_1(z_1)\} = \sum_{x_0} E(X_1 \mid z_1, x_0)\Pr(x_0) \tag{29.7}$$

定义 $\tau_{Z_1\to X_1} = E\{X_1(1) - X_1(0)\}$，则：

$$E\{Y(1,0)-Y(0,0)\} = \beta_2 + \beta_3\tau_{Z_1\to X_1} \tag{29.8}$$

$$E\{Y(0,1)-Y(0,0)\} = \beta_1 \tag{29.9}$$

$$E\{Y(1,1)-Y(0,0)\} = \beta_1 + \beta_2 + \beta_3\tau_{Z_1\to X_1} \tag{29.10}$$

[illegible]（参见图 [illegible]：[illegible]中 Z_1 的效应等于路径 $Z_1 \to Y$（系数 β_2）和路径 $Z_1 \to X_1 \to Y$（系数 $\beta_3\tau_{Z_1\to X_1}$）之和；[illegible]中 Z_2 的效应等于路径 $Z_2 \to Y$（系数 β_1）；[illegible]中 (Z_1, Z_2) 的总效应等于三条路径系数之和。



FIGURE 3. [illegible] X_0

[illegible] Robins [illegible] Wasserman [illegible] 1997 [illegible] null [illegible] (Z_1, Z_2) [illegible] Y [illegible] 29.1 [illegible] g-null [illegible] 29.1 [illegible]

[illegible] 29.3.1 [illegible] X_1 [illegible] g-null [illegible]

$$\tilde{Y}_2(z_1,z_2) = E(Y \mid Z_2 = z_2, Z_1 = z_1, X_1, X_0)$$

[illegible] $(Z_2 = z_2, Z_1 = z_1)$ [illegible] Y [illegible] (X_1, X_0) [illegible] $\hat{Y}_{2i}(z_1, z_2)$ [illegible]

$$\tilde{Y}_1(z_1,z_2) = E\{\tilde{Y}_2(z_1,z_2) \mid Z_1 = z_1, X_0\}$$

[illegible] $Z_1 = z_1$ [illegible] $\hat{Y}_2(z_1, z_2)$ [illegible] X_0 [illegible] $\hat{Y}_{1i}(z_1, z_2)$ [illegible] $E\{Y(z_1, z_2)\}$ [illegible]

$$\hat{E}\{Y(z_1, z_2)\} = n^{-1} \sum_{i=1}^{n} \hat{Y}_{1i}(z_1, z_2) \tag{29.11}$$

[illegible] X_1 [illegible] g-null [illegible] 29.2[illegible] [illegible] $\tilde{Y}_2(z_1, z_2)$

X [illegible]

[illegible] IPW [illegible]

$$E\{Y(z)\} = E\left\{\frac{\mathbf{1}(Z = z)Y}{\Pr(Z = z \mid X)}\right\}$$

[illegible]

$$e(z_1, X_0) = \Pr(Z_1 = z_1 \mid X_0) \tag{29.12}$$

$$e(z_2, Z_1, X_1, X_0) = \Pr(Z_2 = z_2 \mid Z_1, X_1, X_0) \tag{29.13}$$

[illegible] 1 [illegible] 2 [illegible]

[illegible] **mw**(IPW [illegible]) **w**[illegible] *Assumption* [illegible],

$$E\{Y(z_1, z_2)\} = E\left\{\frac{\mathbf{1}(Z_1 = z_1)\mathbf{1}(Z_2 = z_2)Y}{e(z_1, X_0)\, e(z_2, Z_1, X_1, X_0)}\right\} \tag{29.14}$$

[illegible]

$$0 < e(z_1, X_0) < 1, \quad 0 < e(z_2, Z_1, X_1, X_0) < 1 \tag{29.15}$$

[illegible] z_1, z_2 [illegible] 0 [illegible] 1 [illegible] IPW [illegible] Z_2 [illegible] Z_1 [illegible] $\mathbf{1}(Z_1 = z_1)\mathbf{1}(Z_2 = z_2)$ [illegible]

[illegible] IPW [illegible] Z_1 [illegible] X_0 [illegible] $\hat{e}_1(z_1, X_{0i})$ [illegible] Z_2 [illegible] (Z_1, X_1, X_0) [illegible] $\hat{e}_2(z_2, Z_{1i}, X_{1i}, X_{0i})$ [illegible] IPW [illegible] Horvitz-Thompson [illegible]

$$\hat{E}^{\text{ht}}\{Y(z_1, z_2)\} = n^{-1} \sum_{i=1}^{n} \frac{\mathbf{1}(Z_{1i} = z_1)\mathbf{1}(Z_{2i} = z_2)Y_i}{\hat{e}_1(z_1, X_{0i})\hat{e}_2(z_2, Z_{1i}, X_{1i}, X_{0i})} \tag{29.16}$$

[illegible] 11 [illegible] HT [illegible] Hajek [illegible] $\hat{E}^{\text{haj}}\{Y(z_1, z_2)\} = \hat{E}^{\text{ht}}\{Y(z_1, z_2)\}/\hat{1}^{\text{ht}}(z_1, z_2)$ [illegible]

$$\hat{1}^{\text{ht}}(z_1, z_2) = n^{-1} \sum_{i=1}^{n} \frac{\mathbf{1}(Z_{1i} = z_1)\mathbf{1}(Z_{2i} = z_2)}{\hat{e}_1(z_1, X_{0i})\hat{e}_2(z_2, Z_{1i}, X_{1i}, X_{0i})} \tag{29.17}$$

[illegible]

[illegible] 29.2 [illegible] 29.3 [illegible] K [illegible] K [illegible] $2^5 = 32$ [illegible] $2^{10} = 1024$ [illegible] 29.2 [illegible] 29.3 [illegible] IPW [illegible]

[illegible] Robins [illegible]

[illegible] [illegible] Z_1 [illegible] Robins et al., 2000; Hernán et al., 2000 [illegible] $K = 2$ [illegible] MSM [illegible]

[illegible] 29.3 (MSM [illegible]). $Y(z_1, z_2)$ 的边际均值为：

$$E\{Y(z_1, z_2)\} = f(z_1, z_2; \beta) \tag{29.18}$$

[illegible] $E\{Y(z_1, z_2)\} = \beta_0 + \beta_1 z_1 + \beta_2 z_2$ [illegible] Z_1 [illegible]

[illegible] 29.4 ([illegible] MSM). 给定 X_0 条件下 $Y(z_1, z_2)$ 的均值为：

$$E\{Y(z_1, z_2) \mid X_0\} = f(z_1, z_2, X_0; \beta) \tag{29.19}$$

[illegible]

$$E\{Y(z_1, z_2) \mid X_0\} = \beta_0 + \beta_1 z_1 + \beta_2 z_2 + \beta_3^\top X_0 \tag{29.20}$$

[illegible] β [illegible]

$$\beta = \arg\min_b \sum_{z_2} \sum_{z_1} E\left\{Y(z_1, z_2) - f(z_1, z_2, X_0; b)\right\}^2 \tag{29.21}$$

[illegible] β

[illegible] (MSM [illegible] IPW) 在 *Assumption* [illegible] 和 *Definition* [illegible] 下，[illegible] 中的 β 等于：

$$\beta = \arg\min_b \sum_{z_2} \sum_{z_1} E\left[\frac{\mathbf{1}(Z_1 = z_1)\mathbf{1}(Z_2 = z_2)}{e(z_1, X_0)\, e(z_2, Z_1, X_1, X_0)}\{Y - f(z_1, z_2, X_0; b)\}^2\right] \tag{29.22}$$

[illegible] Y_i [illegible] $(1, Z_{1i}, Z_{2i}, X_{0i})$ [illegible] WLS [illegible] $\hat{e}_1^{-1}(Z_{1i}, X_{0i})\hat{e}_{2i}^{-1}(Z_{2i}, Z_{1i}, X_{1i}, X_{0i})$ [illegible] MSM [illegible] IPW [illegible] $E(Y \mid Z, X)$ [illegible] MSM [illegible] IPW [illegible] $E\{Y(z) \mid X_0\}$ [illegible] MSM [illegible]

[illegible] X_0 [illegible] $\beta_3^\top X_0$ [illegible] β_3 [illegible] LASSO [illegible] Tibshirani, 1996 [illegible] β [illegible] Bloniarz et al., 2016; Wager et al., 2016 [illegible]

[illegible] IPW [illegible] IPW [illegible] Robins [illegible]

[illegible] 29.6 ([illegible] SNM [illegible]). 时间点 *1* 的条件效应为：

$$E\{Y(z_1, 0) - Y(0, 0) \mid Z_1 = z_1, X_0\} = g_1(z_1, X_0; \beta) \tag{29.23}$$

时间点 2 的条件效应为：

$$E\{Y(z_1,z_2)-Y(z_1,0)\mid Z_2=z_2,Z_1=z_1,X_1,X_0\}=g_2(z_2,z_1,X_1,X_0;\beta) \tag{29.24}$$

Definition [illegible]

$$g_1(0,X_0;\beta)=0,\quad g_2(0,z_1,X_1,X_0;\beta)=0,\quad \forall z_1 \tag{29.25}$$

[illegible]

[illegible] 29.2 ([illegible] 29.2 [illegible] SNM). 假设：

$$\begin{cases} g_1(z_1,X_0;\beta)=\beta_1 z_1 \\ g_2(z_2,z_1,X_1,X_0;\beta)=(\beta_2+\beta_3 z_1)z_2 \end{cases} \tag{29.26}$$

[illegible] 29.3 ([illegible] 29.3 [illegible] SNM). 假设：

$$\begin{cases} g_1(z_1,X_0;\beta)=(\beta_1+\beta_2^{\mathsf{T}}X_0)z_1 \\ g_2(z_2,z_1,X_1,X_0;\beta)=(\beta_3+\beta_4 z_1+\beta_5^{\mathsf{T}}X_1)z_2 \end{cases} \tag{29.27}$$

i、i 、k i [illegible] $_1$ [illegible] Definition [illegible] $_1$ [illegible]

$$U_2(\beta)=Y-g_2(Z_2,Z_1,X_1,X_0;\beta) \tag{29.28}$$

$$U_1(\beta)=Y-g_2(Z_2,Z_1,X_1,X_0;\beta)-g_1(Z_1,X_0;\beta) \tag{29.29}$$

[illegible]

[illegible] 29.1) 在 *Assumption* [illegible] 和 *Definition* [illegible] 下，

$$E\{U_2(\beta)\mid Z_2,Z_1,X_1,X_0\}=E\{U_2(\beta)\mid Z_1,X_1,X_0\}=E\{Y(Z_1,0)\mid Z_1,X_1,X_0\} \tag{29.30}$$

且

$$E\{U_1(\beta)\mid Z_1,X_0\}=E\{U_1(\beta)\mid X_0\}=E\{Y(0,0)\mid X_0\} \tag{29.31}$$

[illegible] $U_1(\beta)$ [illegible] $U_2(\beta)$ [illegible] 1 [illegible] 2 [illegible]

[illegible] 29.4 [illegible] SNM [illegible]) 在 *Assumption* [illegible] 和 *Definition* [illegible] 下，

$$E\left[h_2(Z_1,X_1,X_0)\{Z_2-e(1,Z_1,X_1,X_0)\}U_2(\beta)\right]=0 \tag{29.32}$$

且

$$E\left[h_1(X_0)\{Z_1-e(1,X_0)\}U_1(\beta)\right]=0 \tag{29.33}$$

对任何函数 h_1 和 h_2（只要矩存在）。

[illegible] h_1 [illegible] h_2 [illegible] β [illegible] 29.4 [illegible] 29.2

[illegible] 29.4 ([illegible] 29.4 [illegible] 29.2 [illegible]). 在例 *29.2* 下，我们可以选择 $h_1 = 1$ 和 $h_2 = (1, Z_1)$ 得到：

$$
\begin{aligned}
&E[\{Z_2 - e(1, Z_1, X_1, X_0)\}\{Y - (\beta_2 + \beta_3 Z_1)Z_2\}] = 0\\
&E[Z_1\{Z_2 - e(1, Z_1, X_1, X_0)\}\{Y - (\beta_2 + \beta_3 Z_1)Z_2\}] = 0 \qquad (29.34)\\
&E[\{Z_1 - e(1, X_0)\}\{Y - (\beta_2 + \beta_3 Z_1)Z_2 - \beta_1 Z_1\}] = 0
\end{aligned}
$$

然后我们可以从上述线性方程组求解 β（见习题 *29.6*）。

[illegible] β [illegible] (h_1, h_2) [illegible] (h_1, h_2) [illegible] GMM; Hansen, 1982 [illegible] Naimi et al. 2017 [illegible] Vansteelandt and Joffe 2014 [illegible]

[illegible]

[illegible] [illegible] X_0 [illegible] (Z_1, Z_2) [illegible] Y [illegible] (Z_1, Z_2) [illegible] Y [illegible]

FIGURE 4. 29.1 [illegible] X_1 [illegible] Y [illegible] U [illegible]

[illegible] 29.1 [illegible]

$$\beta_1 = \beta_2 = 0 \quad [illegible] \quad \beta_3 = 0 \qquad (29.35)$$

5

$$\beta_1 = \beta_2 = 0 \quad [illegible] \quad E\{X_1(z_1)\} \text{ [illegible] } z_1 \qquad (29.36)$$

[illegible] $E\{Y(z_1, z_2)\}$ [illegible] (z_1, z_2) [illegible] $\beta_3 = 0$ [illegible] Y [illegible] X_1 [illegible] X_1 [illegible] Y [illegible] U [illegible] $E\{X_1(z_1)\}$ [illegible] z_1 [illegible] X_1 [illegible] Z_1 [illegible] Z_1 [illegible] X_1 [illegible] X_1 [illegible] Y [illegible] U [illegible] Z_1 [illegible] X_1 [illegible] 29.1 $E\{Y(z_1, z_2)\}$ [illegible] (z_1, z_2) [illegible] (Z_1, Z_2) [illegible] Y [illegible] g-null [illegible]

[illegible] [illegible] 29.3.2 [illegible] 29.1 [illegible] $_1$ [illegible] 0

[illegible]

[illegible] [illegible] 29.2 [illegible]

$$\operatorname{logit}[\Pr\{Y(z_1, z_2) = 1 \mid X_0\}] = \beta_0 + \beta_1 z_1 + \beta_2 z_2 + \beta_3^\intercal X_0 \qquad (29.37)$$

[illegible] β [illegible] B.6.2 [illegible]

$$\beta = \arg\min_b \sum_{z_2}\sum_{z_1} E\{\log(1 + e^{\ell}) - Y(z_1, z_2)\ell\} \qquad (29.38)$$

[illegible] $\ell = \beta_0 + \beta_1 z_1 + \beta_2 z_2 + \beta_3^\intercal X_0$ [illegible] β

[illegible] Assumption [illegible] Definition [illegible] β [illegible]

$$\beta = \arg\min_{b} \sum_{z_2} \sum_{z_1} E\left[\frac{\mathbf{1}(Z_1 = z_1)\mathbf{1}(Z_2 = z_2)}{e(z_1, X_0)\, e(z_2, Z_1, X_1, X_0)}\{\log(1 + e^{\ell}) - Y(z_1, z_2)\ell\}\right] \tag{29.39}$$

[illegible] 29.5

[illegible] 29.5 [illegible] [illegible]
[illegible] Y_i [illegible] $(1, Z_{1i}, Z_{2i}, X_{0i})$ [illegible] logistic [illegible] $\hat{e}_1^{-1}(Z_{1i}, X_{0i})\hat{e}_{2i}^{-1}(Z_{2i}, Z_{1i}, X_{1i}, X_{0i})$

[illegible] IID [illegible] $\{X, Z, Y(1), Y(0)\}$ [illegible] $e(X) = \Pr(Z = 1 \mid X)$ [illegible]

$$Z \perp Y(0) \mid X \tag{29.40}$$

[illegible]

[illegible]

$$E\{Y(z) - Y(0) \mid Z = z, X\} = g(z, X; \beta) \tag{29.41}$$

[illegible] 29.4 [illegible] $g(0, X; \beta) = 0$ [illegible]

[illegible]

$$E\{Y - g(Z, X; \beta) \mid X, Z\} = E\{Y - g(Z, X; \beta) \mid X\} = E\{Y(0) \mid X\} \tag{29.42}$$

[illegible]

$$E[h(X)\{Z - e(X)\}\{Y - g(Z, X; \beta)\}] = 0 \tag{29.43}$$

[illegible] h [illegible]
[illegible] 29.4 [illegible] $g(z, X; \beta) = \beta z$ [illegible] $h(X) = 1$ [illegible]

$$E[\{Z - e(X)\}\{Y - \beta Z\}] = 0 \tag{29.44}$$

[illegible] 11 [illegible]

[illegible]

Robins [illegible] Robins [illegible] 1986 [illegible] g-formula [illegible] Robins [illegible] 1998 [illegible] Robins [illegible] Hernán [illegible] 2009 [illegible] g-[illegible] Robins [illegible] Waterman [illegible] 1994 [illegible] MSM [illegible] SNM [illegible] g-null [illegible] Robins [illegible] Wasserman [illegible] 1997 [illegible] McGrath et al. [illegible] 2021 [illegible]

[illegible] Bloniarz et al. [illegible] 2016 [illegible] LASSO [illegible] Wager et al [illegible] 2016 [illegible] Lei [illegible] Ding [illegible] 2021 [illegible] Lin [illegible] 2013 [illegible] D'Amour et al. [illegible] 2021 [illegible]

APPENDIX A

[illegible]

[illegible]

[illegible] 1975 [illegible]

$$2S(1,0) = S^2(1) + S^2(0) - S^2(\tau).$$

[illegible] Lemma 4.1 [illegible] Neyman (1923) [illegible]

- $S^2(1) = (n-1)^{-1}\sum_{i=1}^{n}\{Y_i(1)-\bar{Y}(1)\}^2$ [illegible] $Y_i(1)$ [illegible]
- $S^2(0) = (n-1)^{-1}\sum_{i=1}^{n}\{Y_i(0)-\bar{Y}(0)\}^2$ [illegible] $Y_i(0)$ [illegible]
- $S(1,0) = (n-1)^{-1}\sum_{i=1}^{n}\{Y_i(1)-\bar{Y}(1)\}\{Y_i(0)-\bar{Y}(0)\}$ [illegible] $Y_i(1)$ [illegible] $Y_i(0)$ [illegible]
- $\tau_i = Y_i(1) - Y_i(0)$ [illegible]
- $\tau = n^{-1}\sum_{i=1}^{n}\tau_i = \bar{Y}(1) - \bar{Y}(0)$ [illegible]
- $S^2(\tau) = (n-1)^{-1}\sum_{i=1}^{n}(\tau_i - \tau)^2$ [illegible]

[illegible]

$$2S(1,0) = S^2(1) + S^2(0) - S^2(\tau).$$

[illegible]

[illegible] $S^2(\tau)$ [illegible]

$$\begin{aligned}
S^2(\tau) &= (n-1)^{-1}\sum_{i=1}^{n}(\tau_i - \tau)^2 \\
&= (n-1)^{-1}\sum_{i=1}^{n}\left[(Y_i(1)-Y_i(0)) - (\bar{Y}(1)-\bar{Y}(0))\right]^2 \\
&= (n-1)^{-1}\sum_{i=1}^{n}\left[(Y_i(1)-\bar{Y}(1)) - (Y_i(0)-\bar{Y}(0))\right]^2.
\end{aligned}$$

[illegible]

$$\begin{aligned}
S^2(\tau) &= (n-1)^{-1}\sum_{i=1}^{n}\Big[(Y_i(1)-\bar{Y}(1))^2+(Y_i(0)-\bar{Y}(0))^2\\
&\quad -2(Y_i(1)-\bar{Y}(1))(Y_i(0)-\bar{Y}(0))\Big]\\
&= (n-1)^{-1}\sum_{i=1}^{n}(Y_i(1)-\bar{Y}(1))^2+(n-1)^{-1}\sum_{i=1}^{n}(Y_i(0)-\bar{Y}(0))^2\\
&\quad -2(n-1)^{-1}\sum_{i=1}^{n}(Y_i(1)-\bar{Y}(1))(Y_i(0)-\bar{Y}(0))\\
&= S^2(1)+S^2(0)-2S(1,0).
\end{aligned}$$

[illegible]

$$S^2(1)+S^2(0)-S^2(\tau)=2S(1,0).$$

[illegible]

$$\boxed{2S(1,0)=S^2(1)+S^2(0)-S^2(\tau)}.$$

[illegible] $S(1,0)$ [illegible] $S^2(\tau)$ [illegible] $S^2(\tau)=0$ [illegible] $2S(1,0)=S^2(1)+S^2(0)$ [illegible] $Y(1)$ [illegible] $Y(0)$ [illegible] 1 [illegible] $S^2(\tau)>0$ [illegible] $S(1,0)$ [illegible]

[illegible] Neyman [illegible]

$$\mathrm{var}(\hat{\tau})=\frac{S^2(1)}{n_1}+\frac{S^2(0)}{n_0}-\frac{S^2(\tau)}{n}=\frac{n_0}{n_1 n}S^2(1)+\frac{n_1}{n_0 n}S^2(0)+\frac{2}{n}S(1,0).$$

[illegible]

[illegible]

$$(n-1)s^2=\sum_{Z_i=1}\{Y_i-\hat{\bar{Y}}(1)\}^2+\sum_{Z_i=0}\{Y_i-\hat{\bar{Y}}(0)\}^2+\frac{n_1 n_0}{n}\hat{\tau}^2.$$

[illegible]

[illegible]

$$(n-1)s^2=\sum_{i=1}^{n}(Y_i-\bar{Y})^2=\sum_{i=1}^{n}\left[(Y_i-\bar{Y})\right]^2.$$

[illegible] $Y_i-\bar{Y}$ [illegible]

$$Y_i-\bar{Y}=\underbrace{(Y_i-\hat{\bar{Y}}(Z_i))}_{\text{[illegible]}}+\underbrace{(\hat{\bar{Y}}(Z_i)-\bar{Y})}_{\text{[illegible]}}.$$

[illegible]

$$(Y_i-\bar{Y})^2=(Y_i-\hat{\bar{Y}}(Z_i))^2+(\hat{\bar{Y}}(Z_i)-\bar{Y})^2+2(Y_i-\hat{\bar{Y}}(Z_i))(\hat{\bar{Y}}(Z_i)-\bar{Y}).$$

[illegible] [redacted] [illegible]

$$\sum_{i=1}^{n} 2(Y_i - \hat{\bar{Y}}(Z_i))(\hat{\bar{Y}}(Z_i) - \bar{Y}) = 2\sum_{i=1}^{n}(Y_i - \hat{\bar{Y}}(Z_i)) \cdot (\hat{\bar{Y}}(Z_i) - \bar{Y}).$$

[illegible] $Z_i = 1$ [illegible]

$$2\sum_{Z_i=1}(Y_i - \hat{\bar{Y}}(1)) \cdot (\hat{\bar{Y}}(1) - \bar{Y}) = 2(\hat{\bar{Y}}(1) - \bar{Y}) \underbrace{\sum_{Z_i=1}(Y_i - \hat{\bar{Y}}(1))}_{=0} = 0,$$

$\sum_{Z_i=1} Y_i = n_1\hat{\bar{Y}}(1)$ [illegible] $\sum_{Z_i=1}(Y_i - \hat{\bar{Y}}(1)) = 0$

[illegible] $Z_i = 0$ [illegible]

$$2\sum_{Z_i=0}(Y_i - \hat{\bar{Y}}(0)) \cdot (\hat{\bar{Y}}(0) - \bar{Y}) = 2(\hat{\bar{Y}}(0) - \bar{Y}) \underbrace{\sum_{Z_i=0}(Y_i - \hat{\bar{Y}}(0))}_{=0} = 0.$$

[illegible] [redacted] [illegible]

[illegible]

$$\sum_{i=1}^{n}(Y_i - \bar{Y})^2 = \underbrace{\sum_{Z_i=1}(Y_i - \hat{\bar{Y}}(1))^2 + \sum_{Z_i=0}(Y_i - \hat{\bar{Y}}(0))^2}_{\text{[illegible]}} + \underbrace{\sum_{i=1}^{n}(\hat{\bar{Y}}(Z_i) - \bar{Y})^2}_{\text{[illegible]}}.$$

[illegible]

$$\sum_{i=1}^{n}(\hat{\bar{Y}}(Z_i) - \bar{Y})^2 = n_1(\hat{\bar{Y}}(1) - \bar{Y})^2 + n_0(\hat{\bar{Y}}(0) - \bar{Y})^2.$$

[illegible]

(1) $\hat{\bar{Y}}(1) - \hat{\bar{Y}}(0) = \hat{\tau}$

(2) $n_1\hat{\bar{Y}}(1) + n_0\hat{\bar{Y}}(0) = n\bar{Y}$ [illegible]

[illegible]

$$\begin{aligned}
&n_1(\hat{\bar{Y}}(1) - \bar{Y})^2 + n_0(\hat{\bar{Y}}(0) - \bar{Y})^2 \\
&= n_1\hat{\bar{Y}}(1)^2 - 2n_1\hat{\bar{Y}}(1)\bar{Y} + n_1\bar{Y}^2 + n_0\hat{\bar{Y}}(0)^2 - 2n_0\hat{\bar{Y}}(0)\bar{Y} + n_0\bar{Y}^2 \\
&= n_1\hat{\bar{Y}}(1)^2 + n_0\hat{\bar{Y}}(0)^2 - n\bar{Y}^2.
\end{aligned}$$

[illegible] $(n_1\hat{\bar{Y}}(1) + n_0\hat{\bar{Y}}(0))^2 = n^2\bar{Y}^2$ [illegible]

$$n_1^2\hat{\bar{Y}}(1)^2 + 2n_1n_0\hat{\bar{Y}}(1)\hat{\bar{Y}}(0) + n_0^2\hat{\bar{Y}}(0)^2 = n^2\bar{Y}^2.$$

[illegible] $\hat{\tau} = \hat{\bar{Y}}(1) - \hat{\bar{Y}}(0)$ [illegible]

$$n_1(\hat{\bar{Y}}(1) - \bar{Y})^2 + n_0(\hat{\bar{Y}}(0) - \bar{Y})^2 = \frac{n_1n_0}{n}\hat{\tau}^2.$$

[illegible] { {

[illegible] 1

[illegible] $H_{0\mathrm{F}}$

$$\operatorname{var}(W) = \frac{n_1 n_0 (n+1)}{12}.$$

[illegible]

[illegible] $W = \sum_{i=1}^n Z_i R_i$ [illegible] R_i [illegible] Y_i [illegible] Z_i [illegible]

[illegible]

$$\operatorname{var}(W) = \operatorname{var}\left(\sum_{i=1}^n Z_i R_i\right) = \sum_{i=1}^n R_i^2 \operatorname{var}(Z_i) + \sum_{i \neq j} R_i R_j \operatorname{cov}(Z_i, Z_j).$$

[illegible] (Z_i) [illegible] CRE [illegible] $Z_i \sim \mathrm{Bernoulli}(n_1/n)$ [illegible]

$$\operatorname{var}(Z_i) = \frac{n_1}{n} \cdot \frac{n_0}{n} = \frac{n_1 n_0}{n^2}.$$

[illegible] (Z_i, Z_j) [illegible] $i \neq j$ [illegible] $\sum_{i=1}^n Z_i = n_1$ [illegible]

$$\operatorname{cov}(Z_i, Z_j) = -\frac{n_1 n_0}{n^2(n-1)} \quad (i \neq j).$$

[illegible]

$$\begin{aligned}\operatorname{var}(W) &= \left(\sum_{i=1}^n R_i^2\right) \frac{n_1 n_0}{n^2} + \sum_{i \neq j} R_i R_j \left(-\frac{n_1 n_0}{n^2(n-1)}\right) \\ &= \frac{n_1 n_0}{n^2} \left[\sum_{i=1}^n R_i^2 - \frac{1}{n-1} \sum_{i \neq j} R_i R_j\right].\end{aligned}$$

$\sum_{i \neq j} R_i R_j = \left(\sum_{i=1}^n R_i\right)^2 - \sum_{i=1}^n R_i^2$ [illegible]

$$\operatorname{var}(W) = \frac{n_1 n_0}{n^2} \left[\sum_{i=1}^n R_i^2 - \frac{1}{n-1} \left(\left(\sum_{i=1}^n R_i\right)^2 - \sum_{i=1}^n R_i^2\right)\right].$$

[illegible] $\sum_{i=1}^n R_i$ [illegible] $\sum_{i=1}^n R_i^2$ [illegible] ties [illegible] $\{R_i\}$ [illegible] $1, 2, \ldots, n$ [illegible]

$$\sum_{i=1}^n R_i = 1 + 2 + \cdots + n = \frac{n(n+1)}{2}.$$

$$\sum_{i=1}^n R_i^2 = 1^2 + 2^2 + \cdots + n^2 = \frac{n(n+1)(2n+1)}{6}.$$

[illegible]

$$\begin{aligned}\operatorname{var}(W) &= \frac{n_1 n_0}{n^2} \left[\frac{n(n+1)(2n+1)}{6} - \frac{1}{n-1} \left(\frac{n^2(n+1)^2}{4} - \frac{n(n+1)(2n+1)}{6}\right)\right] \\ &= \frac{n_1 n_0 (n+1)}{12}.\end{aligned}$$

[illegible]

X ẃۡگ ᶍ ȷ ⅆ̃ΩẆÄ

ѫﬁ ŝ { {ŏ ࢱ ▯Δ Fisher ȷۡگ ᶍ V̈●Ã ıı̃ ⅆ̃ΩẆÄȷ

Ᵽ

ۿ ŝ

- 8 ꞓẊ 4 ꞓ ” ﭙ ”$(K = 4)$ŏ 4 ꞓ ” Ẋ ”
- ۡگ$_2$ƏЫک 4 ꞓ ” ﭙ ”
- ɐ X Əῌ ” ﭙ ” ȷꞓ

ص ﺵ̂ ⅆ̃ΩẆÄȚ

$H_{0\mathrm{F}}$ ŏۡگȷƏЫ ŏ̃ ȷŋ 4 ꞓ ” ﭙ ” ȷƏЫQ̄ ۡ$_2$گ

8 ꞓẊȷ ‗ Ã ۡگ ɽ∏ ŝᴶÃ $K = 4$ Ã ” Ë ” ȷ $N = 8$ ŏăų‡

ɱ $n = 4$ Ãŏ X ȷ ËÃ

` ⅆ̃ΩẆÄȷÕ̃ƹ ı ŋ

ŝ

$$\mathbb{P}(X = k) = \frac{\binom{K}{k}\binom{N-K}{n-k}}{\binom{N}{n}}, \quad k = 0, 1, 2, 3, 4.$$

ŝ̀ ⅆ̃ΩẆÄȷÕ̃ƹ Ω̃ 1

$$\sum_{k=0}^{4} \mathbb{P}(X = k) = \frac{1}{\binom{8}{4}} \sum_{k=0}^{4} \binom{4}{k}\binom{4}{4-k} = \frac{1}{70} \cdot \binom{8}{4} = 1,$$

ʕ α̋ Δ ăñɽ $\sum_k \binom{a}{k}\binom{b}{n-k} = \binom{a+b}{n}$

ÃÕ̃ƹȷ ŝ

$$\mathbb{P}(X = 0) = \frac{\binom{4}{0}\binom{4}{4}}{\binom{8}{4}} = \frac{1 \cdot 1}{70} = \frac{1}{70},$$

$$\mathbb{P}(X = 1) = \frac{\binom{4}{1}\binom{4}{3}}{\binom{8}{4}} = \frac{4 \cdot 4}{70} = \frac{16}{70},$$

$$\mathbb{P}(X = 2) = \frac{\binom{4}{2}\binom{4}{2}}{\binom{8}{4}} = \frac{6 \cdot 6}{70} = \frac{36}{70},$$

$$\mathbb{P}(X = 3) = \frac{\binom{4}{3}\binom{4}{1}}{\binom{8}{4}} = \frac{4 \cdot 4}{70} = \frac{16}{70},$$

$$\mathbb{P}(X = 4) = \frac{\binom{4}{4}\binom{4}{0}}{\binom{8}{4}} = \frac{1 \cdot 1}{70} = \frac{1}{70}.$$

Ȍ ẆÄȷϚ ŝلὼ ” ɱų‡ ”ŋđۡگV^E ƏЫĬɵỷ ᵓBŌ̆ $X \sim$ Binomial$(4, 1/2)$ŏ^B

$$\mathbb{P}(X = 4) = \binom{4}{4}(1/2)^4 = 1/16 = 0.0625 > 0.014.$$

` ⅆ̃ΩẆÄŋăų‡Ōȷ æÕ̃ƹɛ̈ ŏk Fisher ȷ⌐ Ա∂ ɛ̈:§§

FRT” : ” ȷÃ

[illegible]

[illegible]

$$\hat{\tau} = \hat{\bar{Y}}(1) - \hat{\bar{Y}}(0).$$

[illegible] H_{0F} [illegible] $\hat{\tau}$ [illegible]

无偏性.

[illegible] CRE [illegible] **Z** [illegible] $\{\mathbf{Y}(1), \mathbf{Y}(0)\}$ [illegible] n_1/n [illegible]

$$\mathbb{E}(Z_i) = \frac{n_1}{n}, \quad \mathbb{E}(1 - Z_i) = \frac{n_0}{n}.$$

[illegible]

$$\begin{aligned}
\mathbb{E}(\hat{\tau}) &= \mathbb{E}\left(n_1^{-1}\sum_{i=1}^{n} Z_iY_i - n_0^{-1}\sum_{i=1}^{n}(1 - Z_i)Y_i\right)\\
&= n_1^{-1}\sum_{i=1}^{n}\mathbb{E}(Z_i)Y_i - n_0^{-1}\sum_{i=1}^{n}\mathbb{E}(1 - Z_i)Y_i\\
&= n_1^{-1}\sum_{i=1}^{n}\frac{n_1}{n}Y_i - n_0^{-1}\sum_{i=1}^{n}\frac{n_0}{n}Y_i\\
&= \frac{1}{n}\sum_{i=1}^{n}Y_i - \frac{1}{n}\sum_{i=1}^{n}Y_i = 0.
\end{aligned}$$

[illegible]

$$\hat{\tau} = n_1^{-1}\sum_{i=1}^{n} Z_iY_i - n_0^{-1}\sum_{i=1}^{n}(1 - Z_i)Y_i = \frac{n}{n_0}\cdot n_1^{-1}\sum_{i=1}^{n} Z_iY_i - n_0^{-1}\sum_{i=1}^{n} Y_i.$$

$\sum_{i=1}^{n} Y_i$ [illegible] **Z** [illegible]

$$\mathrm{var}(\hat{\tau}) = \frac{n^2}{n_0^2 n_1^2}\cdot \mathrm{var}\left(\sum_{i=1}^{n} Z_iY_i\right).$$

[illegible] $\mathrm{var}(Z_i)$ [illegible] $\mathrm{cov}(Z_i, Z_j)$ CRE [illegible]

$$Z_i \sim \text{Bernoulli}\left(\frac{n_1}{n}\right), \quad \mathrm{var}(Z_i) = \frac{n_1}{n}\cdot\frac{n_0}{n} = \frac{n_1n_0}{n^2}.$$

$\sum_{i=1}^{n} Z_i = n_1$ [illegible] $i \neq j$ [illegible]

$$\mathrm{cov}(Z_i, Z_j) = -\frac{n_1n_0}{n^2(n-1)}.$$

[illegible] $\mathrm{var}\left(\sum_{i=1}^{n} Z_iY_i\right)$ [illegible]

$$\begin{aligned}
\mathrm{var}\left(\sum_{i=1}^{n} Z_iY_i\right) &= \sum_{i=1}^{n} Y_i^2\,\mathrm{var}(Z_i) + \sum_{i\neq j} Y_iY_j\,\mathrm{cov}(Z_i, Z_j)\\
&= \left(\sum_{i=1}^{n} Y_i^2\right)\frac{n_1n_0}{n^2} + \sum_{i\neq j} Y_iY_j\left(-\frac{n_1n_0}{n^2(n-1)}\right).
\end{aligned}$$

$\sum_{i\neq j} Y_iY_j = (\sum_{i=1}^n Y_i)^2 - \sum_{i=1}^n Y_i^2 = n^2\bar{Y}^2 - \sum_{i=1}^n Y_i^2$ [illegible]

$$
\begin{aligned}
\operatorname{var}\left(\sum_{i=1}^n Z_iY_i\right) &= \frac{n_1n_0}{n^2}\sum_{i=1}^n Y_i^2 - \frac{n_1n_0}{n^2(n-1)}\left(n^2\bar{Y}^2 - \sum_{i=1}^n Y_i^2\right) \\
&= \frac{n_1n_0}{n^2(n-1)}\left[(n-1)\sum_{i=1}^n Y_i^2 - n^2\bar{Y}^2 + \sum_{i=1}^n Y_i^2\right] \\
&= \frac{n_1n_0}{n^2(n-1)}\left[n\sum_{i=1}^n Y_i^2 - n^2\bar{Y}^2\right] \\
&= \frac{n_1n_0}{n(n-1)}\left(\sum_{i=1}^n Y_i^2 - n\bar{Y}^2\right).
\end{aligned}
$$

[illegible] s^2 [illegible]

$$
s^2 = (n-1)^{-1}\sum_{i=1}^n (Y_i - \bar{Y})^2 = (n-1)^{-1}\left(\sum_{i=1}^n Y_i^2 - n\bar{Y}^2\right).
$$

[illegible]

$$
\operatorname{var}\left(\sum_{i=1}^n Z_iY_i\right) = \frac{n_1n_0}{n}\cdot s^2.
$$

[illegible] $\operatorname{var}(\hat{\tau})$ [illegible]

$$
\operatorname{var}(\hat{\tau}) = \frac{n^2}{n_0^2n_1^2}\cdot\frac{n_1n_0}{n}\cdot s^2 = \frac{n}{n_1n_0}\cdot s^2.
$$

[illegible]

$$
\boxed{\operatorname{var}(\hat{\tau}) = \frac{n}{n_1n_0}\cdot s^2}.
$$

[illegible]

- [illegible] n [illegible]
- [illegible] n_1n_0 [illegible] $n_1 = n_0 = n/2$ [illegible]
- [illegible]

[illegible]

[illegible] CRE [illegible] n_1 [illegible] n_0 [illegible] $n_{[k]1}$ [illegible] $n_{[k]0}$ [illegible]

[illegible]

- n [illegible]
- $n_{[k]}$ [illegible] k [illegible]
- n_1 [illegible]
- n_0 [illegible] $n_0 = n - n_1$ [illegible]
- $n_{[k]1}$ [illegible] k [illegible]
- $n_{[k]0}$ [illegible] k [illegible] $n_{[k]0} = n_{[k]} - n_{[k]1}$ [illegible]
- $\pi_{[k]} = n_{[k]}/n$ [illegible] k [illegible]

Y a ” Q l

- CRE **Z** $\binom{n}{n_1}$
- n_1/n

Y p g

$$\mathbb{E}\left(\frac{n_{[k]1}}{n_1} - \frac{n_{[k]0}}{n_0}\right) = 0.$$

MQ a ” ; — a p l 、 — Q s 。

$n_{[k]1}$

CRE n_1/n k

$X_i = k$

$$\mathbb{E}(Z_i) = \frac{n_1}{n}.$$

k

$$n_{[k]1} = \sum_{i:X_i=k} Z_i.$$

$n_{[k]}$ Z_i n_1

$$\mathbb{E}(n_{[k]1}) = \sum_{i:X_i=k} \mathbb{E}(Z_i) = n_{[k]} \cdot \frac{n_1}{n} = \pi_{[k]} \cdot n_1.$$

$n_{[k]0} = n_{[k]} - n_{[k]1}$

$$\mathbb{E}(n_{[k]0}) = n_{[k]} - \mathbb{E}(n_{[k]1}) = n_{[k]} - \pi_{[k]}n_1 = n_{[k]} - \frac{n_{[k]}}{n}n_1 = n_{[k]}\left(1 - \frac{n_1}{n}\right) = n_{[k]} \cdot \frac{n_0}{n} = \pi_{[k]} \cdot n_0.$$

$$\mathbb{E}\left(\frac{n_{[k]1}}{n_1} - \frac{n_{[k]0}}{n_0}\right) = \frac{\mathbb{E}(n_{[k]1})}{n_1} - \frac{\mathbb{E}(n_{[k]0})}{n_0} = \frac{\pi_{[k]}n_1}{n_1} - \frac{\pi_{[k]}n_0}{n_0} = \pi_{[k]} - \pi_{[k]} = 0.$$

$\mathbb{E}\left(n_{[k]1}/n_1 - n_{[k]0}/n_0\right) = 0$

$n_{[k]1}/n_1 - n_{[k]0}/n_0$

A ; J e Z N SRE

$$t_{\text{SRE}} = \frac{\hat{\tau}_{\text{SRE}}}{\sqrt{\hat{V}_{\text{SRE}}}}$$

Fisher sharp null $H_{0F} : Y_i(1) = Y_i(0)$ t_{SRE}

r ; ; j Q — Q MQ 【 l a — a p l 。

- $\hat{\tau}_{\text{SRE}} = \sum_{k=1}^{K} \pi_{[k]} \hat{\tau}_{[k]}$
- $\hat{\tau}_{[k]} = \bar{Y}_{[k]}(1) - \bar{Y}_{[k]}(0)$ k
- $\hat{V}_{\text{SRE}} = \sum_{k=1}^{K} \pi_{[k]}^2 \left(\frac{\hat{S}_{[k]}^2(1)}{n_{[k]1}} + \frac{\hat{S}_{[k]}^2(0)}{n_{[k]0}}\right)$

- $\hat{S}^2_{[k]}(1)$ [illegible] $\hat{S}^2_{[k]}(0)$ [illegible] k [illegible]

[illegible] $\hat{\tau}_{\text{SRE}}$ [illegible] $\hat{V}_{\text{SRE}}$ [illegible] H_{0F} [illegible]
[illegible]

[illegible] $\hat{\tau}_{[k]}$ [illegible]

H_{0F} [illegible] $Y_i(1) = Y_i(0) = Y_i$ [illegible]

$$\hat{\tau}_{[k]} = \bar{Y}_{[k]}(1) - \bar{Y}_{[k]}(0) = \frac{\sum_{i:X_i=k} Z_i Y_i}{n_{[k]1}} - \frac{\sum_{i:X_i=k}(1-Z_i)Y_i}{n_{[k]0}}.$$

[illegible] $\mathbb{E}(Z_i) = n_1/n$ [illegible] $\mathbb{E}(1-Z_i) = n_0/n$ [illegible] Y_i [illegible] Z [illegible]

$$\mathbb{E}(\hat{\tau}_{[k]}) = \frac{n_{[k]}}{n_{[k]1}} \cdot \frac{n_1}{n} \cdot \bar{Y}_{[k]} - \frac{n_{[k]}}{n_{[k]0}} \cdot \frac{n_0}{n} \cdot \bar{Y}_{[k]} = \bar{Y}_{[k]} \left(\frac{n_{[k]}}{n_{[k]1}} \cdot \frac{n_1}{n} - \frac{n_{[k]}}{n_{[k]0}} \cdot \frac{n_0}{n} \right).$$

$n_{[k]1}/n_{[k]} = e_{[k]}$ [illegible] $n_{[k]0}/n_{[k]} = 1 - e_{[k]}$ [illegible] n_1/n [illegible] n_0/n [illegible]
[illegible] $\bar{Y}_{[k]} \cdot ($ [illegible] $-$ [illegible] $) = 0$
[illegible] $\mathbb{E}(Z_i) = n_1/n$ [illegible] H_{0F} [illegible]

$$\mathbb{E}(\hat{\tau}_{[k]}) = \frac{1}{n_{[k]1}} \sum_{i:X_i=k} \mathbb{E}(Z_i) Y_i - \frac{1}{n_{[k]0}} \sum_{i:X_i=k} \mathbb{E}(1-Z_i) Y_i = \frac{n_1}{n}\bar{Y}_{[k]} - \frac{n_0}{n}\bar{Y}_{[k]} = 0.$$

[illegible] $\hat{\tau}_{\text{SRE}}$ [illegible]
$\hat{\tau}_{\text{SRE}} = \sum_{k=1}^K \pi_{[k]} \hat{\tau}_{[k]}$ [illegible] $\pi_{[k]}$ [illegible]

$$\mathbb{E}(\hat{\tau}_{\text{SRE}}) = \sum_{k=1}^K \pi_{[k]} \mathbb{E}(\hat{\tau}_{[k]}) = \sum_{k=1}^K \pi_{[k]} \cdot 0 = 0.$$

[illegible] $\hat{\tau}_{[k]}$ [illegible]
[illegible] Neyman (1923) [illegible] k [illegible]

$$\text{var}(\hat{\tau}_{[k]}) = \frac{S^2_{[k]}(1)}{n_{[k]1}} + \frac{S^2_{[k]}(0)}{n_{[k]0}} - \frac{S^2_{[k]}(\tau)}{n_{[k]}}.$$

H_{0F} [illegible] $S^2_{[k]}(\tau) = 0$ [illegible]

$$\text{var}(\hat{\tau}_{[k]}) = \frac{S^2_{[k]}(1)}{n_{[k]1}} + \frac{S^2_{[k]}(0)}{n_{[k]0}}.$$

[illegible] $\hat{\tau}_{\text{SRE}}$ [illegible]
[illegible]

$$\text{var}(\hat{\tau}_{\text{SRE}}) = \sum_{k=1}^K \pi^2_{[k]} \text{var}(\hat{\tau}_{[k]}) = \sum_{k=1}^K \pi^2_{[k]} \left(\frac{S^2_{[k]}(1)}{n_{[k]1}} + \frac{S^2_{[k]}(0)}{n_{[k]0}} \right).$$

[illegible] $\hat{V}_{\text{SRE}}$ [illegible]
[illegible] $\mathbb{E}(\hat{S}^2_{[k]}(1)) = S^2_{[k]}(1)$ [illegible] $\mathbb{E}(\hat{S}^2_{[k]}(0)) = S^2_{[k]}(0)$ [illegible]

$$\mathbb{E}(\hat{V}_{\text{SRE}}) = \sum_{k=1}^K \pi^2_{[k]} \left(\frac{S^2_{[k]}(1)}{n_{[k]1}} + \frac{S^2_{[k]}(0)}{n_{[k]0}} \right) = \text{var}(\hat{\tau}_{\text{SRE}}) + \sum_{k=1}^K \pi^2_{[k]} \frac{S^2_{[k]}(\tau)}{n_{[k]}}.$$

H_{0F} [illegible] $S^2_{[k]}(\tau) = 0$ [illegible] $\mathbb{E}(\hat{V}_{\text{SRE}}) = \text{var}(\hat{\tau}_{\text{SRE}})$

H_{0F} $\hat{\tau}_{\text{SRE}}$ $\hat{V}_{\text{SRE}}$ $t_{\text{SRE}} = \hat{\tau}_{\text{SRE}}/\sqrt{\hat{V}_{\text{SRE}}}$

SRE k

$\hat{\tau}_{[k]}$ $\hat{\tau}_{\text{SRE}}$

- $S^2_{[k]}(1)$ $S^2_{[k]}(0)$ k $Y(1)$ $Y(0)$
- $S^2_{[k]}(\tau)$ k $\tau_i = Y_i(1) - Y_i(0)$
- $\hat{\tau}_{[k]} = \bar{Y}_{[k]}(1) - \bar{Y}_{[k]}(0)$ k
- $\hat{\tau}_{\text{SRE}} = \sum_{k=1}^{K} \pi_{[k]} \hat{\tau}_{[k]}$

(1) $\text{var}(\hat{\tau}_{[k]}) = \frac{S^2_{[k]}(1)}{n_{[k]1}} + \frac{S^2_{[k]}(0)}{n_{[k]0}} - \frac{S^2_{[k]}(\tau)}{n_{[k]}}$

(2) $\text{var}(\hat{\tau}_{\text{SRE}}) = \sum_{k=1}^{K} \pi^2_{[k]} \text{var}(\hat{\tau}_{[k]})$

Neyman (1923) k k CRE $n_{[k]}$ n Neyman

$$\text{var}(\hat{\tau}_{[k]}) = \frac{S^2_{[k]}(1)}{n_{[k]1}} + \frac{S^2_{[k]}(0)}{n_{[k]0}} - \frac{S^2_{[k]}(\tau)}{n_{[k]}}.$$

CRE 1

1

SRE $\hat{\tau}_{[1]}, \ldots, \hat{\tau}_{[K]}$

$$\text{var}(\hat{\tau}_{\text{SRE}}) = \text{var}\left(\sum_{k=1}^{K} \pi_{[k]} \hat{\tau}_{[k]}\right) = \sum_{k=1}^{K} \pi^2_{[k]} \text{var}(\hat{\tau}_{[k]}) + \sum_{k \neq j} \pi_{[k]} \pi_{[j]} \text{cov}(\hat{\tau}_{[k]}, \hat{\tau}_{[j]}).$$

$\text{cov}(\hat{\tau}_{[k]}, \hat{\tau}_{[j]}) = 0$ $k \neq j$

$$\text{var}(\hat{\tau}_{\text{SRE}}) = \sum_{k=1}^{K} \pi^2_{[k]} \text{var}(\hat{\tau}_{[k]}).$$

$$\text{var}(\hat{\tau}_{\text{SRE}}) = \sum_{k=1}^{K} \pi^2_{[k]} \left(\frac{S^2_{[k]}(1)}{n_{[k]1}} + \frac{S^2_{[k]}(0)}{n_{[k]0}} - \frac{S^2_{[k]}(\tau)}{n_{[k]}}\right).$$

[illegible] CRE [illegible] SRE [illegible]

[illegible] CRE [illegible] $\hat{\tau}$ [illegible] SRE [illegible]
1 $\hat{\tau}_{\text{SRE}}$ [illegible]

[illegible]

- $\hat{\tau} = \bar{Y}(1) - \bar{Y}(0)$ [illegible] CRE [illegible]
- $\hat{\tau}_{\text{SRE}} = \sum_{k=1}^{K} \pi_{[k]} \hat{\tau}_{[k]}$ [illegible] SRE [illegible]
- $S^2(1)$ [illegible] $S^2(0)$ [illegible]
- $S^2_{[k]}(1)$ [illegible] $S^2_{[k]}(0)$ [illegible] k [illegible]
- $\bar{Y}(1)$ [illegible] $\bar{Y}(0)$ [illegible]
- $\bar{Y}_{[k]}(1)$ [illegible] $\bar{Y}_{[k]}(0)$ [illegible] k [illegible]

[illegible]

- [illegible] CRE [illegible] $\text{var}_{CRE}(\hat{\tau}) = \frac{S^2(1)}{n_1} + \frac{S^2(0)}{n_0} - \frac{S^2(\tau)}{n}$
- [illegible] SRE [illegible] $\text{var}_{SRE}(\hat{\tau}_{\text{SRE}}) = \sum_{k=1}^{K} \pi_{[k]}^2 \left(\frac{S^2_{[k]}(1)}{n_{[k]1}} + \frac{S^2_{[k]}(0)}{n_{[k]0}} - \frac{S^2_{[k]}(\tau)}{n_{[k]}} \right)$
- [illegible] $S^2(1) = \sum_{k=1}^{K} \pi_{[k]} S^2_{[k]}(1) + \sum_{k=1}^{K} \pi_{[k]} \{\bar{Y}_{[k]}(1) - \bar{Y}(1)\}^2 -$ [illegible]

[illegible]

$$\sum_{k=1}^{K} \frac{\pi_{[k]}}{n} \left\{ \sqrt{\frac{n_0}{n_1}} \{\bar{Y}_{[k]}(1) - \bar{Y}(1)\} + \sqrt{\frac{n_1}{n_0}} \{\bar{Y}_{[k]}(0) - \bar{Y}(0)\} \right\}^2 \geq 0.$$

[illegible]

[illegible]

[illegible] $S^2(1)$ [illegible]

$$\begin{aligned} S^2(1) &= (n-1)^{-1} \sum_{i=1}^{n} \{Y_i(1) - \bar{Y}(1)\}^2 \\ &= \sum_{k=1}^{K} \frac{n_{[k]} - 1}{n - 1} \cdot \frac{1}{n_{[k]} - 1} \sum_{i:X_i = k} \{Y_i(1) - \bar{Y}_{[k]}(1)\}^2 \\ &\quad + \sum_{k=1}^{K} \frac{n_{[k]}}{n-1} \{\bar{Y}_{[k]}(1) - \bar{Y}(1)\}^2. \end{aligned}$$

[illegible] $n_{[k]}$ [illegible] $(n_{[k]} - 1)/(n - 1) \approx n_{[k]}/n = \pi_{[k]}$ [illegible]

$$S^2(1) \approx \sum_{k=1}^{K} \pi_{[k]} S^2_{[k]}(1) + \sum_{k=1}^{K} \pi_{[k]} \{\bar{Y}_{[k]}(1) - \bar{Y}(1)\}^2.$$

[illegible] $S^2(0)$ [illegible] $S^2(\tau)$ [illegible]

[illegible] CRE [illegible]

$$\text{var}_{CRE}(\hat{\tau}) = \frac{1}{n_1} \sum_{k=1}^{K} \pi_{[k]} S^2_{[k]}(1) + \frac{1}{n_0} \sum_{k=1}^{K} \pi_{[k]} S^2_{[k]}(0) - \frac{1}{n} \sum_{k=1}^{K} \pi_{[k]} S^2_{[k]}(\tau) + \text{[illegible]} \quad .$$

[illegible]

[illegible]

$$\frac{1}{n_1}\sum_{k=1}^{K}\pi_{[k]}\{\bar{Y}_{[k]}(1)-\bar{Y}(1)\}^2+\frac{1}{n_0}\sum_{k=1}^{K}\pi_{[k]}\{\bar{Y}_{[k]}(0)-\bar{Y}(0)\}^2-\frac{1}{n}\sum_{k=1}^{K}\pi_{[k]}(\tau_{[k]}-\tau)^2.$$

[illegible]
[illegible] $n_{[k]1}\approx\pi_{[k]}n_1$ [illegible] $n_{[k]0}\approx\pi_{[k]}n_0$ [illegible]

$$\sum_{k=1}^{K}\frac{\pi_{[k]}}{n}\left\{\sqrt{\frac{n_0}{n_1}}\{\bar{Y}_{[k]}(1)-\bar{Y}(1)\}+\sqrt{\frac{n_1}{n_0}}\{\bar{Y}_{[k]}(0)-\bar{Y}(0)\}\right\}^2.$$

[illegible] k [illegible]

$$\sqrt{\frac{n_0}{n_1}}\{\bar{Y}_{[k]}(1)-\bar{Y}(1)\}+\sqrt{\frac{n_1}{n_0}}\{\bar{Y}_{[k]}(0)-\bar{Y}(0)\}=0$$

[illegible] SRE [illegible]
CRE [illegible]

[illegible]

[illegible] $\mathbf{n}=\{n_{[k]1},n_{[k]0}\}_{k=1}^{K}$ [illegible] CRE [illegible] **Z** [illegible] SRE [illegible]

[illegible]

- $\mathbf{Z}=(Z_1,\ldots,Z_n)$ [illegible]
- $n_{[k]1}$ [illegible] k [illegible]
- $n_{[k]0}$ [illegible] k [illegible]
- $\mathbf{n}=\{n_{[k]1},n_{[k]0}\}_{k=1}^{K}$ [illegible]

[illegible]

- CRE [illegible] $\mathbb{P}(\mathbf{Z}=\mathbf{z})=\frac{1}{\binom{n}{n_1}}$ [illegible] $\sum Z_i=n_1$ [illegible] $\mathbf{z}$
- SRE [illegible] k [illegible] $\frac{1}{\binom{n_{[k]}}{n_{[k]1}}}$

[illegible]

$$\Pr_{CRE}(\mathbf{Z}=\mathbf{z}\mid\mathbf{n})=\frac{1}{\prod_{k=1}^{K}\binom{n_{[k]}}{n_{[k]1}}}.$$

[illegible]

[illegible]
CRE [illegible] **Z** [illegible]

$$\Pr_{CRE}(\mathbf{Z}=\mathbf{z})=\frac{1}{\binom{n}{n_1}}.$$

[illegible]
[illegible] **n** [illegible] $\mathbf{Z}=\mathbf{z}$ [illegible]

$$\Pr_{CRE}(\mathbf{Z}=\mathbf{z}\mid\mathbf{n})=\frac{\Pr_{CRE}(\mathbf{Z}=\mathbf{z},\mathbf{n})}{\Pr_{CRE}(\mathbf{n})}.$$

[illegible] $\{\mathbf{Z}=\mathbf{z},\mathbf{n}\}$ [illegible] k [illegible] $n_{[k]1}$ [illegible] [illegible]

$$\Pr_{CRE}(\mathbf{Z}=\mathbf{z},\mathbf{n})=\prod_{k=1}^{K}\frac{1}{\binom{n_{[k]}}{n_{[k]1}}}.$$

[illegible]

[illegible] $\Pr_{CRE}(\mathbf{n})$ [illegible] k [illegible] $n_{[k]1}$ [illegible] $\mathbf{z}$ [illegible]

$$\Pr_{CRE}(\mathbf{n})=\sum_{\mathbf{z}:n_{[k]1}(\mathbf{z})=n_{[k]1}}\Pr_{CRE}(\mathbf{Z}=\mathbf{z})=\sum_{\mathbf{z}:n_{[k]1}(\mathbf{z})=n_{[k]1}}\frac{1}{\binom{n}{n_1}}.$$

[illegible] $\prod_{k=1}^{K}\binom{n_{[k]}}{n_{[k]1}}$ [illegible]

$$\Pr_{CRE}(\mathbf{n})=\prod_{k=1}^{K}\binom{n_{[k]}}{n_{[k]1}}\cdot\frac{1}{\binom{n}{n_1}}.$$

[illegible]

$$\Pr_{CRE}(\mathbf{Z}=\mathbf{z}\mid\mathbf{n})=\frac{\prod_{k=1}^{K}\frac{1}{\binom{n_{[k]}}{n_{[k]1}}}}{\prod_{k=1}^{K}\binom{n_{[k]}}{n_{[k]1}}\cdot\frac{1}{\binom{n}{n_1}}}=\frac{1}{\prod_{k=1}^{K}\binom{n_{[k]}}{n_{[k]1}}}.$$

[illegible] CRE [illegible] SRE [illegible]

[illegible]

[illegible] Chapter 4 [illegible] Neymanian [illegible] OLS [illegible]

[illegible]

- $\hat{\tau}=\hat{\bar{Y}}(1)-\hat{\bar{Y}}(0)$ [illegible]
- $\hat{\beta}$ [illegible] OLS [illegible] $Y_i=\alpha+\beta Z_i+\varepsilon_i$ [illegible] treatment [illegible]
- $\hat{V}=\frac{\hat{S}^2(1)}{n_1}+\frac{\hat{S}^2(0)}{n_0}$ [illegible] Neyman [illegible]
- $\hat{V}_{\mathrm{OLS}}$ [illegible] OLS [illegible]
- $\hat{V}_{\mathrm{EHW}}$ [illegible] Eicker-Huber-White [illegible]

[illegible]

(1) [illegible] $\hat{\beta}=\hat{\tau}$

(2) [illegible] $\hat{V}_{\mathrm{OLS}}=\frac{n(n_1-1)}{(n-2)n_1n_0}\hat{S}^2(1)+\frac{n(n_0-1)}{(n-2)n_1n_0}\hat{S}^2(0)$

(3) [illegible] $\hat{V}_{\mathrm{EHW}}=\frac{\hat{S}^2(1)}{n_1}\frac{n_1-1}{n_1}+\frac{\hat{S}^2(0)}{n_0}\frac{n_0-1}{n_0}$

(4) [illegible] HC2 [illegible] $\hat{V}$

[illegible]

[illegible] $\hat{\beta} = \hat{\tau}$ [illegible] OLS [illegible] $Y_i = \alpha + \beta Z_i + \varepsilon_i$ [illegible] OLS [illegible]

$$\hat{\beta} = \frac{\sum_{i=1}^{n}(Z_i - \bar{Z})(Y_i - \bar{Y})}{\sum_{i=1}^{n}(Z_i - \bar{Z})^2}.$$

[illegible]

CRE [illegible] $\bar{Z} = n_1/n$ [illegible]

$$\sum_{i=1}^{n}(Z_i - \bar{Z})(Y_i - \bar{Y}) = \sum_{i=1}^{n} Z_iY_i - \bar{Z}\sum_{i=1}^{n} Y_i - \bar{Y}\sum_{i=1}^{n} Z_i + n\bar{Z}\bar{Y}.$$

$\sum_{i=1}^{n} Z_i = n_1$ [illegible] $\bar{Z} = n_1/n$ [illegible] $\bar{Z}\sum_{i=1}^{n} Z_i = n_1^2/n$ [illegible] $\bar{Y} = n^{-1}\sum_{i=1}^{n} Y_i$ [illegible]

$$\sum_{i=1}^{n} Z_iY_i - \frac{n_1}{n}\sum_{i=1}^{n} Y_i = \sum_{i=1}^{n} Z_iY_i - n_1\bar{Y}.$$

[illegible] $\sum_{i=1}^{n} Z_iY_i$ [illegible]

$$\sum_{i=1}^{n} Z_iY_i = \sum_{Z_i=1} Y_i + \sum_{Z_i=0} Y_i = n_1\hat{\bar{Y}}(1) + n_0\hat{\bar{Y}}(0).$$

[illegible]

$$n_1\hat{\bar{Y}}(1) + n_0\hat{\bar{Y}}(0) - n_1\bar{Y}.$$

[illegible] $\bar{Y} = (n_1\hat{\bar{Y}}(1) + n_0\hat{\bar{Y}}(0))/n$ [illegible]

$$n_1\hat{\bar{Y}}(1) + n_0\hat{\bar{Y}}(0) - \frac{n_1}{n}\left[n_1\hat{\bar{Y}}(1) + n_0\hat{\bar{Y}}(0)\right] = \frac{n_0n_1}{n}\hat{\bar{Y}}(1) - \frac{n_1n_0}{n}\hat{\bar{Y}}(0) = \frac{n_0n_1}{n}\hat{\tau}.$$

[illegible]

$$\sum_{i=1}^{n}(Z_i - \bar{Z})^2 = \sum_{i=1}^{n} Z_i^2 - 2\bar{Z}\sum_{i=1}^{n} Z_i + n\bar{Z}^2 = n_1 - 2\frac{n_1}{n}\cdot n_1 + n\cdot\frac{n_1^2}{n^2} = n_1 - \frac{n_1^2}{n} = \frac{n_0n_1}{n}.$$

[illegible] $\hat{\beta}$

$$\hat{\beta} = \frac{\frac{n_0n_1}{n}\hat{\tau}}{\frac{n_0n_1}{n}} = \hat{\tau}.$$

[illegible] $\hat{\beta} = \hat{\tau}$ [illegible] OLS [illegible]

[illegible] **o f** [illegible] **w** [illegible] OLS [illegible] Appendix B [illegible] (B.4) [illegible]

$$\hat{V}_{\text{OLS}} = \frac{\hat{\sigma}^2}{\sum_{i=1}^{n}(Z_i - \bar{Z})^2},$$

[illegible] $\hat{\sigma}^2 = (n-2)^{-1}\sum_{i=1}^{n}(Y_i - \hat{\alpha} - \hat{\beta}Z_i)^2$ [illegible]

[illegible]

$\sum_{i=1}^{n}(Z_i-\bar{Z})^2 = n_0n_1/n$[illegible]

$$\hat{V}_{\mathrm{OLS}} = \frac{n}{n_0n_1}\cdot\frac{1}{n-2}\sum_{i=1}^{n}(Y_i-\hat{\alpha}-\hat{\beta}Z_i)^2.$$

[illegible] ANOVA [illegible]
[illegible] ANOVA [illegible]

$$(n-1)s^2 = \sum_{Z_i=1}(Y_i-\hat{\bar{Y}}(1))^2 + \sum_{Z_i=0}(Y_i-\hat{\bar{Y}}(0))^2 + \frac{n_1n_0}{n}\hat{\tau}^2,$$

[illegible] $s^2=(n-1)^{-1}\sum_{i=1}^{n}(Y_i-\bar{Y})^2$
[illegible]
OLS [illegible] $\hat{\alpha}=\bar{Y}-\hat{\beta}\bar{Z}$ [illegible] $\hat{\beta}=\hat{\tau}$[illegible]

$$Y_i-\hat{\alpha}-\hat{\beta}Z_i = Y_i-\bar{Y}-\hat{\tau}(Z_i-\bar{Z}).$$

[illegible]

$$\sum_{i=1}^{n}(Y_i-\hat{\alpha}-\hat{\beta}Z_i)^2 = \sum_{i=1}^{n}(Y_i-\bar{Y})^2-\hat{\tau}^2\sum_{i=1}^{n}(Z_i-\bar{Z})^2 = (n-1)s^2-\hat{\tau}^2\cdot\frac{n_0n_1}{n}.$$

[illegible] $\hat{V}_{\mathrm{OLS}}$

$$\hat{V}_{\mathrm{OLS}} = \frac{n}{(n-2)n_0n_1}\left[(n-1)s^2-\frac{n_0n_1}{n}\hat{\tau}^2\right].$$

[illegible] $\hat{S}^2(1)$ [illegible] $\hat{S}^2(0)$ [illegible] $\hat{\tau}=\hat{\bar{Y}}(1)-\hat{\bar{Y}}(0)$[illegible]

$$\hat{V}_{\mathrm{OLS}} = \frac{n(n_1-1)}{(n-2)n_1n_0}\hat{S}^2(1)+\frac{n(n_0-1)}{(n-2)n_1n_0}\hat{S}^2(0).$$

[illegible] ([illegible])[illegible]

([illegible])[illegible] **P**] [illegible] **W**[illegible] EHW [illegible]
[illegible] $Y_i=\alpha+\beta Z_i+\varepsilon_i$[illegible] EHW [illegible] Appendix B [illegible] (B.3)[illegible]

$$\hat{V}_{\mathrm{EHW}} = \frac{1}{n-2}\sum_{i=1}^{n}\frac{(Y_i-\hat{\alpha}-\hat{\beta}Z_i)^2}{(1-h_i)^2},$$

[illegible] $h_i=n^{-1}+(Z_i-\bar{Z})^2/\sum_{j=1}^{n}(Z_j-\bar{Z})^2$ [illegible] leverage[illegible]
[illegible]
$\sum_{i=1}^{n}(Z_i-\bar{Z})^2=n_0n_1/n$[illegible] $Z_i=1$[illegible]

$$h_i=\frac{1}{n}+\frac{(1-n_1/n)^2}{n_0n_1/n}=\frac{1}{n}+\frac{(n_0/n)^2}{n_0n_1/n}=\frac{1}{n}+\frac{n_0}{n_1n}=\frac{n_1+n_0}{n_1n}=\frac{1}{n_1}.$$

[illegible] $Z_i=0$[illegible]

$$h_i=\frac{1}{n}+\frac{(0-n_1/n)^2}{n_0n_1/n}=\frac{1}{n}+\frac{n_1}{n_0n}=\frac{n_0+n_1}{n_0n}=\frac{1}{n_0}.$$

[illegible] $(1-h_i)^2$
[illegible] $(1-h_i)^2=(1-1/n_1)^2=((n_1-1)/n_1)^2$
[illegible] $(1-h_i)^2=(1-1/n_0)^2=((n_0-1)/n_0)^2$
[illegible] EHW [illegible]

$$\hat{V}_{\mathrm{EHW}} = \frac{1}{n-2}\left[\sum_{Z_i=1}\frac{(Y_i-\hat{\alpha}-\hat{\beta})^2}{((n_1-1)/n_1)^2} + \sum_{Z_i=0}\frac{(Y_i-\hat{\alpha})^2}{((n_0-1)/n_0)^2}\right].$$

[illegible] $Y_i - \hat{\alpha} - \hat{\beta} = Y_i - \hat{\bar{Y}}(1)$ [illegible] $Z_i = 1$ [illegible] $Y_i - \hat{\alpha} = Y_i - \hat{\bar{Y}}(0)$ [illegible] $Z_i = 0$ [illegible]

$$\hat{V}_{\mathrm{EHW}} = \frac{n_1^2}{(n-2)(n_1-1)^2}\sum_{Z_i=1}(Y_i-\hat{\bar{Y}}(1))^2 + \frac{n_0^2}{(n-2)(n_0-1)^2}\sum_{Z_i=0}(Y_i-\hat{\bar{Y}}(0))^2.$$

[illegible] $\hat{S}^2(1) = (n_1-1)^{-1}\sum_{Z_i=1}(Y_i-\hat{\bar{Y}}(1))^2$ [illegible] $\hat{S}^2(0) = (n_0-1)^{-1}\sum_{Z_i=0}(Y_i-\hat{\bar{Y}}(0))^2$ [illegible]

$$\hat{V}_{\mathrm{EHW}} = \frac{n_1}{(n-2)(n_1-1)}\hat{S}^2(1) + \frac{n_0}{(n-2)(n_0-1)}\hat{S}^2(0).$$

[illegible] n [illegible] $n-2 \approx n$ [illegible]

$$\hat{V}_{\mathrm{EHW}} \approx \frac{\hat{S}^2(1)}{n_1}\frac{n_1}{n_1-1} + \frac{\hat{S}^2(0)}{n_0}\frac{n_0}{n_0-1} \approx \frac{\hat{S}^2(1)}{n_1} + \frac{\hat{S}^2(0)}{n_0} = \hat{V}.$$

[illegible] n_1, n_0 [illegible] $\hat{V}_{\mathrm{EHW}} \approx \hat{V}$

[illegible] $\hat{V}$ **[illegible]** HC2 [illegible] $(n-2)/(n-p)$ [illegible] $p = 2$ [illegible] $(1-h_i)^{-1}$ [illegible] $(1-h_i)^{-2}$[illegible]

$$\hat{V}_{\mathrm{HC2}} = \frac{1}{n-p}\sum_{i=1}^{n}\frac{(Y_i-\hat{\alpha}-\hat{\beta}Z_i)^2}{1-h_i}.$$

[illegible] $1-h_i$[illegible]

[illegible] $1-h_i = (n_1-1)/n_1$

[illegible] $1-h_i = (n_0-1)/n_0$

[illegible]

$$\hat{V}_{\mathrm{HC2}} = \frac{n_1}{n-2}\cdot\frac{1}{n_1-1}\sum_{Z_i=1}(Y_i-\hat{\bar{Y}}(1))^2 + \frac{n_0}{n-2}\cdot\frac{1}{n_0-1}\sum_{Z_i=0}(Y_i-\hat{\bar{Y}}(0))^2.$$

[illegible] $\hat{S}^2(1)$ [illegible] $\hat{S}^2(0)$ [illegible]

$$\hat{V}_{\mathrm{HC2}} = \frac{n_1}{n-2}\hat{S}^2(1) + \frac{n_0}{n-2}\hat{S}^2(0).$$

[illegible] n [illegible] $n-2 \approx n$ [illegible] $n_1/n \approx n_1/(n_1+n_0)$ [illegible]

$$\hat{V}_{\mathrm{HC2}} = \frac{n}{n-2}\left(\frac{n_1}{n}\hat{S}^2(1) + \frac{n_0}{n}\hat{S}^2(0)\right)\cdot\frac{n}{n_1 n_0}\times\frac{n_1 n_0}{n} = \frac{\hat{S}^2(1)}{n_1} + \frac{\hat{S}^2(0)}{n_0} = \hat{V}.$$

[illegible] $\hat{V}_{\mathrm{HC2}} = \hat{V}$ [illegible] $n_1/(n-2)\cdot(n_1-1)^{-1} = 1/n_1$ [illegible]

[illegible] HC2 [illegible] Neyman [illegible] $\hat{V}$

q ‖ w MaM I ^ ; s—Q ‖ ˇ

A ; J e Z p l N { { DiD

TWFE DiD

r ; ; j Q—Q MQ 【 l a—a p l 。

- $G_i \in \{0,1\}$ i $G_i = 1$ $G_i = 0$
- $T_t \in \{0,1\}$ $T_t = 0$ $T_t = 1$
- $Y_{it}(d)$ i t $d \in \{0,1\}$
- $D_i = G_i$
- $\tau_t = \mathbb{E}(Y_{it}(1) - Y_{it}(0) \mid G_i = 1)$ t ATT

Y a ” Q l ´ $\bar{Y}_{00}, \bar{Y}_{01}, \bar{Y}_{10}, \bar{Y}_{11}$ { {

Y p g DiD $\hat{\tau}_{\text{DiD}} = (\bar{Y}_{11} - \bar{Y}_{10}) - (\bar{Y}_{01} - \bar{Y}_{00})$ τ_1

MQ a ” ; —a p l 、—Q s 。

$$\bar{Y}_{11} - \bar{Y}_{10} = \underbrace{\bar{Y}_{11}(1) - \bar{Y}_{10}(1)}_{} = \underbrace{\tau_1}_{\text{ATT}} + \underbrace{\left[\bar{Y}_{11}(0) - \bar{Y}_{10}(0)\right]}_{} .$$

$$\bar{Y}_{01} - \bar{Y}_{00} = \bar{Y}_{11}(0) - \bar{Y}_{10}(0).$$

$$\begin{aligned} \text{DiD} &= (\bar{Y}_{11} - \bar{Y}_{10}) - (\bar{Y}_{01} - \bar{Y}_{00}) \\ &= \left[\tau_1 + (\bar{Y}_{11}(0) - \bar{Y}_{10}(0))\right] - \left[\bar{Y}_{11}(0) - \bar{Y}_{10}(0)\right] \\ &= \tau_1. \end{aligned}$$

$\bar{Y}_{11}(0) - \bar{Y}_{10}(0) = \bar{Y}_{01} - \bar{Y}_{00}$ DiD : τ_1

U P MaM

$$Y_{it} = \alpha + \beta D_{it} + \gamma_i + \delta_t + \varepsilon_{it}, \quad D_{it} = G_i \cdot T_t.$$

OLS β

$$\hat{\beta} = \bar{Y}_{11} - \bar{Y}_{10} - \bar{Y}_{01} + \bar{Y}_{00} = \hat{\tau}_{\text{DiD}}.$$

γ_i δ_t $\hat{\beta}$

A p p—。 ; s

DiD Bootstrap

(1) cluster i t (Y_{it}, D_{it})
(2) $\hat{\tau}_{\text{DiD}}$
(3) B $B = 999$ Bootstrap $\{\hat{\tau}^{(b)}_{\text{DiD}}\}_{b=1}^{B}$
(4) $\hat{\sigma}_{\text{boot}} = \sqrt{(B-1)^{-1} \sum_{b=1}^{B} (\hat{\tau}^{(b)}_{\text{DiD}} - \bar{\hat{\tau}})^2}$

[illegible]

[illegible] TSLS [illegible]

[illegible]

- Y_i [illegible]
- D_i [illegible] p [illegible]
- Z_i [illegible] q [illegible] $q \geq p$ [illegible]
- β [illegible] p [illegible]
- ε_i [illegible] $E(\varepsilon_i Z_i) = 0$
- $\hat{D}_i$ [illegible] OLS [illegible]
- $\hat{\Gamma}$ [illegible] OLS [illegible]

[illegible] TSLS [illegible]

[illegible]

[illegible] D on Z [illegible] OLS [illegible]

$$\hat{\Gamma} = \left(\sum_{i=1}^{n} Z_i Z_i^\top \right)^{-1} \sum_{i=1}^{n} Z_i D_i^\top .$$

[illegible]

$$\hat{D}_i = \hat{\Gamma}^\top Z_i .$$

[illegible]

$$\sum_{i=1}^{n} \hat{D}_i \check{D}_i^\top = 0,$$

[illegible] $\check{D}_i = D_i - \hat{D}_i$ [illegible]

[illegible]

$$\sum_{i=1}^{n} \hat{D}_i D_i^\top = \sum_{i=1}^{n} \hat{D}_i \hat{D}_i^\top .$$

[illegible] Y on $\hat{D}$ [illegible] OLS [illegible]

$$\hat{\beta}_{\text{TSLS}} = \left(\sum_{i=1}^{n} \hat{D}_i \hat{D}_i^\top \right)^{-1} \sum_{i=1}^{n} \hat{D}_i Y_i .$$

[illegible] $Y_i = D_i^\top \beta + \varepsilon_i$ [illegible]

$$\begin{aligned}
\hat{\beta}_{\text{TSLS}} &= \left(\sum_{i=1}^{n} \hat{D}_i \hat{D}_i^\top \right)^{-1} \sum_{i=1}^{n} \hat{D}_i (D_i^\top \beta + \varepsilon_i) \\
&= \left(\sum_{i=1}^{n} \hat{D}_i \hat{D}_i^\top \right)^{-1} \sum_{i=1}^{n} \hat{D}_i D_i^\top \beta + \left(\sum_{i=1}^{n} \hat{D}_i \hat{D}_i^\top \right)^{-1} \sum_{i=1}^{n} \hat{D}_i \varepsilon_i .
\end{aligned}$$

[illegible]

$$\hat{\beta}_{\text{TSLS}} = \beta + \left(\sum_{i=1}^{n} \hat{D}_i \hat{D}_i^\top \right)^{-1} \sum_{i=1}^{n} \hat{D}_i \varepsilon_i .$$

[illegible] $\hat{D}_i = \hat{\Gamma}^\top Z_i$ [illegible]

$$\hat{\beta}_{\mathrm{TSLS}} = \beta + \left(\hat{\Gamma}^\top \sum_{i=1}^n Z_i Z_i^\top \hat{\Gamma}\right)^{-1} \hat{\Gamma}^\top \sum_{i=1}^n Z_i \varepsilon_i$$

$$= \beta + \left\{\hat{\Gamma}^\top \left(n^{-1}\sum_{i=1}^n Z_i Z_i^\top\right)\hat{\Gamma}\right\}^{-1} \hat{\Gamma}^\top \left(n^{-1}\sum_{i=1}^n Z_i \varepsilon_i\right).$$

[illegible] $E(Z_i\varepsilon_i) = 0$ [illegible] restriction [illegible]

$$n^{-1}\sum_{i=1}^n Z_i\varepsilon_i \xrightarrow{p} 0.$$

[illegible]

$$n^{-1}\sum_{i=1}^n Z_i Z_i^\top \xrightarrow{p} E(ZZ^\top),$$

$$\hat{\Gamma} \xrightarrow{p} \Gamma = E(ZZ^\top)^{-1}E(ZD^\top).$$

[illegible]

$$\hat{\beta}_{\mathrm{TSLS}} \xrightarrow{p} \beta.$$

[illegible] TSLS [illegible] β [illegible]

[illegible] $q = p$ [illegible] $\hat{\Gamma}$ [illegible]

$$\hat{\beta}_{\mathrm{TSLS}} = \left(\hat{\Gamma}^{-1}\sum_{i=1}^n Z_i Y_i\right) = \left(\sum_{i=1}^n Z_i D_i^\top\right)^{-1}\sum_{i=1}^n Z_i Y_i = \hat{\beta}_{\mathrm{IV}}.$$

[illegible] TSLS [illegible] IV [illegible]

[illegible]

[illegible] TSLS [illegible] ILS [illegible]

$$\begin{cases} Y_i = \beta_0 + \beta_1 D_i + \beta_2^\top X_i + \varepsilon_i, \\ D_i = \gamma_0 + \gamma_1 Z_i + \gamma_2^\top X_i + \varepsilon_{2i}. \end{cases}$$

[illegible] $\hat{\beta}_{1,\mathrm{TSLS}} = \hat{\Gamma}_1/\hat{\gamma}_1 = \hat{\beta}_{1,\mathrm{ILS}}$ [illegible]

[illegible] OLS [illegible]

$$\hat{\Gamma}_1 = \frac{\sum_i (Z_i - \bar{Z})(Y_i - \bar{Y})}{\sum_i (Z_i - \bar{Z})^2}, \quad \hat{\gamma}_1 = \frac{\sum_i (Z_i - \bar{Z})(D_i - \bar{D})}{\sum_i (Z_i - \bar{Z})^2}.$$

[illegible]

$$\frac{\hat{\Gamma}_1}{\hat{\gamma}_1} = \frac{\sum_i (Z_i - \bar{Z})(Y_i - \bar{Y})}{\sum_i (Z_i - \bar{Z})(D_i - \bar{D})}.$$

TSLS

$$\hat{D}_i = \hat{\gamma}_0 + \hat{\gamma}_1 Z_i + \hat{\gamma}_2^\top X_i.$$

Y on $(1, \hat{D}, X)$ $\hat{D}$

$$\hat{\beta}_{1,\mathrm{TSLS}} = \frac{\sum_i (Y_i - \bar{Y})(\hat{D}_i - \bar{\hat{D}})}{\sum_i (\hat{D}_i - \bar{\hat{D}})^2}.$$

$\hat{\gamma}_1 \sum_i (Z_i - \bar{Z})(\hat{D}_i - \bar{\hat{D}}) = \hat{\gamma}_1^2 \sum_i (Z_i - \bar{Z})^2$ $\hat{\gamma}_1 \sum_i (Z_i - \bar{Z})(Y_i - \bar{Y})$

$$\hat{\beta}_{1,\mathrm{TSLS}} = \frac{\hat{\gamma}_1 \sum_i (Z_i - \bar{Z})(Y_i - \bar{Y})}{\hat{\gamma}_1^2 \sum_i (Z_i - \bar{Z})^2} = \frac{\sum_i (Z_i - \bar{Z})(Y_i - \bar{Y})}{\hat{\gamma}_1 \sum_i (Z_i - \bar{Z})^2} = \frac{\hat{\Gamma}_1}{\hat{\gamma}_1}.$$

$\hat{\beta}_{1,\mathrm{TSLS}} = \hat{\beta}_{1,\mathrm{ILS}}$ TSLS ILS TSLS ” ” ILS

Neyman (1923) $\mathrm{var}(\hat{\tau} \mid \mathbf{Z})$ ” ”

- n IID $\{Z_i, Y_i(1), Y_i(0), X_i\}_{i=1}^n$
- $Z \perp \{Y(1), Y(0), X\}$
- n_1 n_0

IID $S^2(1)$ $S^2(0)$ $\mathrm{var}\{Y(1)\}$ $\mathrm{var}\{Y(0)\}$

$$S_N^2(1) = \frac{1}{n-1} \sum_{i=1}^n \{Y_i(1) - \bar{Y}_N(1)\}^2,$$

$\bar{Y}_N(1) = n^{-1} \sum_{i=1}^n Y_i(1)$

$$\sigma^2(1) = E\{Y_i(1) - E(Y_i(1))\}^2.$$

IID $E(S_N^2(1)) = \sigma^2(1)$ IID Neyman Neyman

$$\hat{V}_N = \frac{S_N^2(1)}{n_1} + \frac{S_N^2(0)}{n_0}.$$

$$E(\hat{V}_N) = \frac{E\{S_N^2(1)\}}{n_1} + \frac{E\{S_N^2(0)\}}{n_0} = \frac{\sigma^2(1)}{n_1} + \frac{\sigma^2(0)}{n_0} = \mathrm{var}(\hat{\tau} \mid \mathbf{Z}).$$

♦ ŝ ` Ꝁ ⱯⲶ ŏ $\hat{V}_N$ ꙃ var($\hat{\tau} \mid \mathbf{Z}$) ļ ꞔꙆꙍ§§ħ ꙃ∆

q ˇ w mw ‖Ẇ o f 、s p p g Q N Q Z ŗ‖ Q。。a p l

ѫfi ŝ { {ŏ ɞ Lin (2013) Ꙇꙍ $\hat{\tau}_L = \hat{\alpha}_Z$ ŗ‖ { { $\bar{X} = 0$ ļ
V̈●Ãȷȷ ļ OLS Ẇ ᵽ
Ꝩ Ḩ ŝ ɞꝛ X ω̃ŋđ $\bar{X} = 0$Ō ŏ

$$\hat{\tau}_L = \hat{\gamma}_1 - \hat{\gamma}_0 = \hat{\alpha}_Z.$$

ᵽä ŝ
ᵖ ä ŝ OLS ω̃ţ
Ḧ ᾰΨ‡Ώ { ŏ{ OLS Ꙇꙍ ƅ ω̃ţ ŝ

$$\sum_{i=1}^{n}(Y_i - \hat{\alpha}_0 - \hat{\alpha}_Z Z_i - \hat{\alpha}_X^T X_i - \hat{\alpha}_{ZX}^T Z_i X_i) = 0,$$

$$\sum_{i=1}^{n}(Y_i - \hat{\alpha}_0 - \hat{\alpha}_Z Z_i - \hat{\alpha}_X^T X_i - \hat{\alpha}_{ZX}^T Z_i X_i)Z_i = 0,$$

$$\sum_{i=1}^{n}(Y_i - \hat{\alpha}_0 - \hat{\alpha}_Z Z_i - \hat{\alpha}_X^T X_i - \hat{\alpha}_{ZX}^T Z_i X_i)X_i = 0,$$

$$\sum_{i=1}^{n}(Y_i - \hat{\alpha}_0 - \hat{\alpha}_Z Z_i - \hat{\alpha}_X^T X_i - \hat{\alpha}_{ZX}^T Z_i X_i)Z_i X_i = 0.$$

ᵖȌä ŝ
ᴶ Ĭ ῎A ω̃ţ ŋ█Ȳ Ōȷȷ Ǿŏ ᾳ ŋ ‖ Proposition 6.2 ļ
ᾰΩ ɞŝ

$$\hat{\alpha}_Z = (\hat{\gamma}_1 - \hat{\gamma}_0) + (\hat{\beta}_1 - \hat{\beta}_0)^T \bar{X}.$$

ᵖä ŝ ω̃
$\bar{X} = 0$ŋ X ω̃Ō ŝ

$$\hat{\alpha}_Z = \hat{\gamma}_1 - \hat{\gamma}_0 = \hat{\tau}_L.$$

♦ ŝķ $\hat{\tau}_L = \hat{\alpha}_Z$ŏđ Lin (2013) Ꙇꙍ ŗ ᾰΨ‡Ώ treatment Ḩļ

q ¯ w mw∼ ţPꙆꙍ ļ 〔

ѫfi ŝ { {ŏ ɞ ʉ OLS ļ EHW Ḩ ă ļ V̈●
√ ŝ
` Ꝁ ⱯⲶ ŏ Ꝁ X ļ V̈´ $\bar{X}$ $_2$Ꝁ §§ ‰ ꝺ ɐ̧
Ḩ OLS ļ EHW ţ꙳Ꙇꙍ ∂ $\bar{X}$ ő̄ ļŋ ╔ X ļ ω̃Ō ᴮ ˇ
ȗ $\bar{X}$ ļꝺ Ꝁ
ɏ ɞ╟ $\hat{\tau}_L$ ŝ

$$\hat{\tau}_L = \hat{\gamma}_1 - \hat{\gamma}_0 = (\hat{\beta}_1 - \hat{\beta}_0)^T \bar{X} + \text{Ö}\mho\quad,$$

‰Ǿ $\hat{\tau}_L$ ļţ꙳iĄ ŝ

$$\mathrm{var}(\hat{\tau}_L) = \underbrace{\mathrm{var}(\hat{\gamma}_1 - \hat{\gamma}_0 \mid X)}_{\text{EHW ˇȗļ}} + \underbrace{\mathrm{var}\{(\hat{\beta}_1 - \hat{\beta}_0)^T \bar{X}\}}_{\text{EHW Ꙇꝕļ}}.$$

ᵖȌ V̈ Ꙇꙍ ă ꝕ ŏ ⓐ X ļ Ȩ ◗Ư
ļ√ ŝ

[illegible]

$$\frac{1}{n}(\hat{\beta}_1 - \hat{\beta}_0)^T S_X^2 (\hat{\beta}_1 - \hat{\beta}_0).$$

[illegible] $\text{var}\{(\hat{\beta}_1 - \hat{\beta}_0)^T \bar{X}\}$ [illegible]

[illegible]

[illegible] Lin (2013) [illegible]

[illegible] $X_i \in \{1, \ldots, K\}$ [illegible] SRE [illegible]

$$Z \perp \{Y(1), Y(0)\} \mid X.$$

[illegible] τ [illegible] $\hat{\tau}_S$ [illegible] k [illegible] $Z \perp \{Y(1), Y(0)\} \mid X = k$ [illegible] $X = k$ [illegible] $Z \perp \{Y(1), Y(0)\}$ [illegible] k [illegible] Lin (2013) [illegible]

$$\hat{\tau}_{L,[k]} = \hat{\gamma}_{1,[k]} - \hat{\gamma}_{0,[k]} + (\hat{\beta}_{1,[k]} - \hat{\beta}_{0,[k]})^T \bar{X}_{[k]},$$

[illegible] $\bar{X}_{[k]} = n_{[k]}^{-1} \sum_{X_i = k} X_i$ [illegible]

$$\hat{\tau}_S = \sum_{k=1}^{K} \pi_{[k]} \hat{\tau}_{L,[k]},$$

[illegible] $\pi_{[k]} = \text{pr}(X = k)$ [illegible]

$$\text{var}(\hat{\tau}_S) = \sum_{k=1}^{K} \pi_{[k]}^2 \text{var}(\hat{\tau}_{L,[k]}).$$

[illegible] EHW [illegible] $\text{var}(\hat{\tau}_{L,[k]})$ [illegible] $\hat{V}_S$ [illegible] SRE [illegible] $\hat{\tau}_S$ [illegible] τ [illegible] $\hat{V}_S$ [illegible]

[illegible]

[illegible] R [illegible]

[illegible]

```
linestimator = function(Z, Y, X) {
  # 标准化协变量
  X = scale(X)

  n = dim(X)[1]
  p = dim(X)[2]
  # 完全交互的 OLS 回归
  linreg = lm(Y ~ Z * X)

  # Lin 估计量 (Z 的系数)
  est = coef(linreg)[2]

  # EHW 标准误
  vehw = hccm(linreg)[2, 2]

  # 超级总体校正项
  inter = coef(linreg)[(p+3):(2*p+2)]
  vsuper = vehw + sum(inter * (cov(X) %*% inter)) / n

  # 返回：估计量、EHW 标准误、校正标准误
  c(est, sqrt(vehw), sqrt(vsuper))
}
```

[illegible] $n = 500$[illegible] 2000 [illegible]

```
set.seed(123)
res = replicate(2000, {
  n = 500
  X = matrix(rnorm(n*2), n, 2)
  Y1 = X[,1] + X[,1]^2 + runif(n, -0.5, 0.5)
  Y0 = X[,2] + X[,2]^2 + runif(n, -1, 1)
  Z = rbinom(n, 1, 0.6)
  Y = Z*Y1 + (1-Z)*Y0
  linestimator(Z, Y, X)
})
```

[illegible]

- [illegible] `mean(res[1,])` ≈ -0.0001
- [illegible] `sd(res[1,])` ≈ 0.151
- [illegible] EHW [illegible] `mean(res[2,])` ≈ 0.139
- [illegible] `mean(res[3,])` ≈ 0.153

[illegible]

[illegible] Theorem 19.1 [illegible] Rosenbaum [illegible] S_i [illegible] p [illegible]

[illegible]

- n[illegible]
- $\Gamma \geq 1$[illegible] Rosenbaum [illegible] odds ratio [illegible]
- $\pi_{i1} = \Pr(Z_{i1} = 1 \mid \text{pair } i)$[illegible] i [illegible]
- $\pi_{i0} = 1 - \pi_{i1}$[illegible] i [illegible]
- $o_i = \frac{\pi_{i1}/\pi_{i0}}{(1-\pi_{i1})/(1-\pi_{i0})} = \frac{\pi_{i1}(1-\pi_{i0})}{\pi_{i0}(1-\pi_{i1})}$[illegible] i [illegible] odds [illegible]
- S_i[illegible] i [illegible] Fisher randomization test [illegible] S_i [illegible]

[illegible]

(1) Rosenbaum [illegible] Assumption 19.1 [illegible] $o_i \le \Gamma$ for all i
(2) Sharp null hypothesis $H_{0F}: Y_{ij}(1) = Y_{ij}(0)$ for all i, j
(3) [illegible] i [illegible] $Z_{i1} + Z_{i0} = 1$

[illegible] $o_i \le \Gamma$ [illegible] S_i [illegible] Bernoulli$\left(\frac{\Gamma}{1+\Gamma}\right)$ [illegible]

[illegible] π_{i1} [illegible] $o_i \le \Gamma$ [illegible]

$$\frac{\pi_{i1}(1-\pi_{i0})}{\pi_{i0}(1-\pi_{i1})} \le \Gamma.$$

$Z_{i1} + Z_{i0} = 1$ [illegible]

$$\pi_{i1} \le \frac{\Gamma}{\Gamma+1} \quad [illegible] \quad \pi_{i1} \ge \frac{1}{\Gamma+1}.$$

[illegible] Rosenbaum (1987b) [illegible] π_{i1} [illegible] Fisher randomization test [illegible] S_i [illegible] Fisher randomization test [illegible] i [illegible] ” [illegible] ” [illegible]

$$T_{i1} = q_i \cdot 1 \quad [illegible] \quad T_{i0} = q_i \cdot 0,$$

[illegible] sharp null [illegible] $T = \sum_{i=1}^n S_i q_i$ [illegible] S_i [illegible] i [illegible] T_{i1} [illegible] T_{i0} [illegible] T [illegible] ” [illegible] ” [illegible] S_i [illegible] S_i [illegible] $q_i > 0$ [illegible] T [illegible] Γ [illegible] S_i [illegible] Bernoulli$\left(\frac{\Gamma}{1+\Gamma}\right)$ [illegible]

$$\mathbb{E}_\Gamma(S_i) = \frac{\Gamma}{1+\Gamma} = 1 - \frac{1}{1+\Gamma}.$$

[illegible] i [illegible] $\pi_{i1} = \frac{\Gamma}{1+\Gamma}$ [illegible]

$$S_i \sim \text{Bernoulli}\left(\frac{\Gamma}{1+\Gamma}\right).$$

[illegible] Rosenbaum 1987b, Theorem 1 [illegible]

[illegible] Γ [illegible] $\frac{\Gamma}{1+\Gamma}$ [illegible] 1 [illegible] $q_i > 0$ [illegible] ” [illegible] ” [illegible]

[illegible] $\Gamma = 1$ [illegible] $\frac{\Gamma}{1+\Gamma} = \frac{1}{2}$ [illegible]

[illegible] Assumption 19.1 [illegible] H_{0F} [illegible] S_i [illegible] Bernoulli$\left(\frac{\Gamma}{1+\Gamma}\right)$

[illegible]

[illegible] $q_i = 1$ for all i [illegible] $T = \sum_{i=1}^n S_i$ [illegible] $B(n, \pi)$ [illegible] $\pi = \frac{\Gamma}{1+\Gamma}$

p ŝ

$$\mathrm{p}_{\Gamma} = 2\sum_{k=T}^{n} \binom{n}{k} \left(\frac{\Gamma}{1+\Gamma}\right)^{k} \left(\frac{1}{1+\Gamma}\right)^{n-k}.$$

a g J p p l a Z l Q N _ ; l — Wilcoxon-signed-rank statistic $q_i = \mathrm{rank}(|Y_{i1} - Y_{i0}|)$ T 5 4

k p

Rosenbaum R `sensitivity2x2`

P _ “ ; g ‘ Q

APPENDIX B

[illegible]

[illegible]

[illegible] (X, U) [illegible] Z [illegible] Y [illegible] U [illegible]

[illegible]

[illegible]

$$f_{1,x} = \Pr(U = 1 \mid Z = 1, X = x), \quad f_{0,x} = \Pr(U = 1 \mid Z = 0, X = x). \tag{141}$$

[illegible]

$$\mathrm{RR}_{ZU|x} = \frac{f_{1,x}}{f_{0,x}} \quad \Longrightarrow \quad f_{0,x} = \frac{f_{1,x}}{\mathrm{RR}_{ZU|x}}. \tag{142}$$

[illegible] $\gamma_x = \mathrm{RR}_{UY|x}$ [illegible]

[illegible]

[illegible]

$$\begin{aligned}
\Pr(Y = 1 \mid Z = 1, X = x) &= \Pr(Y = 1 \mid U = 1, X = x)\Pr(U = 1 \mid Z = 1, X = x) \\
&\quad + \Pr(Y = 1 \mid U = 0, X = x)\Pr(U = 0 \mid Z = 1, X = x) \\
&= \gamma_x f_{1,x} + (1 - f_{1,x}) \cdot 1 \\
&= (\gamma_x - 1) f_{1,x} + 1.
\end{aligned}$$

[illegible]

$$\begin{aligned}
\Pr(Y = 1 \mid Z = 0, X = x) &= \Pr(Y = 1 \mid U = 1, X = x)\Pr(U = 1 \mid Z = 0, X = x) \\
&\quad + \Pr(Y = 1 \mid U = 0, X = x)\Pr(U = 0 \mid Z = 0, X = x) \\
&= \gamma_x f_{0,x} + (1 - f_{0,x}) \\
&= (\gamma_x - 1) f_{0,x} + 1.
\end{aligned}$$

[illegible] $f_{0,x} = f_{1,x}/\mathrm{RR}_{ZU|x}$ [illegible]

$$\Pr(Y = 1 \mid Z = 0, X = x) = (\gamma_x - 1)\frac{f_{1,x}}{\mathrm{RR}_{ZU|x}} + 1.$$

[illegible]

$$\mathrm{RR}^{\mathrm{obs}}_{ZY|x} = \frac{(\gamma_x - 1) f_{1,x} + 1}{(\gamma_x - 1)\frac{f_{1,x}}{\mathrm{RR}_{ZU\ x}} + 1}. \tag{143}$$

X [illegible]

[illegible] $f_{1,x} \in [0,1]$ [illegible]

$$g(t) = \frac{(a-1)t+1}{(a-1)\frac{t}{b}+1}$$

$a > 1, b > 1$ [illegible] t [illegible] $f_{1,x}$ [illegible] $\mathrm{RR}^{\mathrm{obs}}_{ZY|x}$ [illegible] $f_{1,x} = 1$ [illegible]

$$\mathrm{RR}^{\mathrm{obs}}_{ZY|x} \leq \frac{(\gamma_x - 1) + 1}{\frac{\gamma_x - 1}{\mathrm{RR}_{ZU\ x}} + 1} = \frac{\gamma_x \cdot \mathrm{RR}_{ZU|x}}{\gamma_x + \mathrm{RR}_{ZU|x} - 1}.$$

[illegible]

W [illegible]

[illegible] $(f_{1,x}, \mathrm{RR}_{ZU|x}, \mathrm{RR}_{UY|x})$ [illegible]

APPENDIX C

[illegible]

[illegible] $\beta(w_1, w_2) = w_1 w_2/(w_1 + w_2 - 1)$ [illegible] $w_1 > 1, w_2 > 1$

[illegible] $\beta(w_1, w_2) = \beta(w_2, w_1)$

[illegible]

$$\frac{\partial \beta}{\partial w_1} = \frac{w_2(w_1 + w_2 - 1) - w_1 w_2}{(w_1 + w_2 - 1)^2} = \frac{w_2(w_2 - 1)}{(w_1 + w_2 - 1)^2} > 0.$$

[illegible] $\partial\beta/\partial w_2 > 0$ [illegible] β [illegible]

[illegible]

$$\beta(w_1, w_2) = \frac{w_1 w_2}{w_1 + w_2 - 1} \leq \frac{w_1 w_2}{w_1} = w_2 \quad (\qquad w_2 - 1 \geq 0).$$

[illegible] $\beta \leq w_1$

[illegible] $w = \max(w_1, w_2)$ [illegible] $w_1 = w$ [illegible] $w_2 \leq w$ [illegible]

$$\beta(w_1, w_2) = \frac{w \cdot w_2}{w + w_2 - 1} \leq \frac{w \cdot w}{w + w - 1} = \frac{w^2}{2w - 1}.$$

APPENDIX D

[illegible]

[illegible] Y [illegible] Z [illegible] β_1 [illegible]

$$\beta_1 = \log \mathrm{OR}_{ZY|x} = \log \frac{\mathrm{odds}(Y=1 \mid Z=1, X=x)}{\mathrm{odds}(Y=1 \mid Z=0, X=x)}.$$

[illegible] $\Pr(Y=1 \mid Z=1, X=x) \approx 0$ [illegible] $\Pr(Y=1 \mid Z=0, X=x) \approx 0$ [illegible]

$$\mathrm{odds}(Y=1 \mid Z, X) \approx \frac{\Pr(Y=1 \mid Z, X)}{1-\Pr(Y=1 \mid Z, X)} \approx \Pr(Y=1 \mid Z, X).$$

[illegible]

$$\beta_1 \approx \log \mathrm{RR}^{\mathrm{obs}}_{ZY|x}.$$

[illegible] $\exp(\beta_1) \approx \mathrm{RR}^{\mathrm{obs}}_{ZY|x}$ [illegible] E-value [illegible]

[illegible]

APPENDIX E

[illegible]

[illegible]

[illegible] $\varepsilon_1(X)$ [illegible] $\varepsilon_0(X)$ [illegible]

[illegible]

[illegible] $\mu_1(X) = E(Y \mid Z = 1, X)$ [illegible] $\mu_0(X) = E(Y \mid Z = 0, X)$ [illegible]

$$\varepsilon_1(X) = \frac{E\{Y(1) \mid Z = 1, X\}}{E\{Y(1) \mid Z = 0, X\}} = \frac{\mu_1(X)}{E\{Y(1) \mid Z = 0, X\}},$$

[illegible]

$$E\{Y(1) \mid Z = 0, X\} = \frac{\mu_1(X)}{\varepsilon_1(X)}. \tag{144}$$

[illegible]

$$E\{Y(0) \mid Z = 1, X\} = \mu_0(X)\varepsilon_0(X). \tag{145}$$

[illegible]

[illegible] $E\{Y(1) \mid Z = 0\} = E\{\mu_1(X)/\varepsilon_1(X) \mid Z = 0\}$ [illegible]

$$E\{Y(1) \mid Z = 0\} = E\left[E\{Y(1) \mid Z = 0, X\} \mid Z = 0\right].$$

[illegible]

$$E\{Y(1) \mid Z = 0\} = E\left[\frac{\mu_1(X)}{\varepsilon_1(X)} \mid Z = 0\right].$$

[illegible] $E\{Y(0) \mid Z = 1\} = E\{\mu_0(X)\varepsilon_0(X) \mid Z = 1\}$ [illegible]

$$E\{Y(0) \mid Z = 1\} = E\left[E\{Y(0) \mid Z = 1, X\} \mid Z = 1\right].$$

[illegible]

$$E\{Y(0) \mid Z = 1\} = E\left[\mu_0(X)\varepsilon_0(X) \mid Z = 1\right].$$

[illegible] τ [illegible]

$$\tau = E\{Y(1)\} - E\{Y(0)\} = E\{Y \mid Z = 1\} - E\{Y(0) \mid Z = 1\}.$$

[illegible]

$$\tau = E\{Y \mid Z = 1\} - E\left[\mu_0(X)\varepsilon_0(X) \mid Z = 1\right].$$

[illegible] $Y = ZY(1) + (1-Z)Y(0)$ [illegible] $E\{Y \mid Z = 1\}$ [illegible]

$$E\{Y \mid Z = 1\} = E\{Y(1) \mid Z = 1\} = E\left[\frac{\mu_1(X)}{\varepsilon_1(X)} \mid Z = 1\right].$$

B

$$\tau = E\left[\frac{\mu_1(X)}{\varepsilon_1(X)} \mid Z = 1\right] - E\left[\mu_0(X)\varepsilon_0(X) \mid Z = 1\right].$$

[illegible] { [illegible] { [illegible]

APPENDIX F

[illegible]

[illegible]

[illegible] IPW [illegible]

[illegible]

[illegible]

$$E\{Y(1) \mid Z = 0\} = E\left\{\frac{\mu_1(X)}{\varepsilon_1(X)} \mid Z = 0\right\}.$$

[illegible] IPW [illegible]

$$E\{Y(1)\} = E\{Y(1) \mid Z = 1\}\Pr(Z = 1) + E\{Y(1) \mid Z = 0\}\Pr(Z = 0).$$

[illegible] $E\{Y(1) \mid Z = 1\} = \mu_1(X)/\varepsilon_1(X)$ [illegible] $E\{Y(1) \mid Z = 0\} = \mu_1(X)/\varepsilon_1(X)$ [illegible]

$$E\{Y(1)\} = E\left[\frac{\mu_1(X)}{\varepsilon_1(X)}\right].$$

[illegible] $e(X) = \Pr(Z = 1 \mid X)$ [illegible] IPW [illegible]

$$E\{Y(1)\} = E\left\{\frac{ZY}{e(X)}\right\} = E\left\{w_1(X)\frac{ZY}{e(X)}\right\},$$

[illegible] $w_1(X) = e(X) + (1 - e(X))/\varepsilon_1(X)$
[illegible] $E\{Y(0)\}$ [illegible]

APPENDIX G

[illegible]

[illegible]

[illegible] outcomes [illegible] $[\underline{y}, \overline{y}]$ [illegible] $E\{Y(1)\}$ [illegible] $E\{Y(0)\}$ [illegible]

[illegible]

[illegible] $p = \Pr(Z = 1)$ [illegible] $\Pr(Z = 0) = 1 - p$ [illegible] $\mu_1 = E(Y \mid Z = 1)$ [illegible] $\mu_0 = E(Y \mid Z = 0)$ [illegible]

[illegible] $E\{Y(1)\}$ [illegible]

[illegible]

$$E\{Y(1)\} = \mu_1 \cdot p + E\{Y(1) \mid Z = 0\} \cdot (1 - p).$$

[illegible] $Y(1) \in [\underline{y}, \overline{y}]$ [illegible]

$$\underline{y} \leq E\{Y(1) \mid Z = 0\} \leq \overline{y}.$$

[illegible]

$$\mu_1 \cdot p + \underline{y}(1 - p) \leq E\{Y(1)\} \leq \mu_1 \cdot p + \overline{y}(1 - p).$$

[illegible] $E\{Y(0)\}$ [illegible]

[illegible]

$$E\{Y(0)\} = E\{Y(0) \mid Z = 1\} \cdot p + \mu_0 \cdot (1 - p),$$

[illegible]

$$\underline{y} \leq E\{Y(0) \mid Z = 1\} \leq \overline{y}.$$

[illegible]

$$\underline{y} \cdot p + \mu_0(1 - p) \leq E\{Y(0)\} \leq \overline{y} \cdot p + \mu_0(1 - p).$$

[illegible] τ [illegible]

$\tau = E\{Y(1)\} - E\{Y(0)\}$ [illegible]

$$\underline{\tau} = \mu_1 p + \underline{y}(1 - p) - \overline{y}p - \mu_0(1 - p),$$

$$\overline{\tau} = \mu_1 p + \overline{y}(1 - p) - \underline{y}p - \mu_0(1 - p).$$

[illegible] $\overline{y} - \underline{y}$

[illegible]

[illegible] [illegible] **[illegible]** [illegible] $\tau = \mathbb{E}[Y_i(1) - Y_i(0)]$ [illegible] [illegible] [illegible]

$$\tau = \mathbb{E}[Y_i(1) - Y_i(0)] \tag{146}$$

$$= \frac{1}{n}\sum_{i=1}^{n}[Y_i(1) - Y_i(0)] \quad \text{[illegible]} \tag{147}$$

$$= \frac{1}{n}\sum_{i=1}^{n} Y_i(1) - \frac{1}{n}\sum_{i=1}^{n} Y_i(0) \tag{148}$$

$$= \bar{Y}(1) - \bar{Y}(0) \tag{149}$$

[illegible]

[illegible] [illegible] **[illegible]** [illegible] τ_x [illegible] $X_i = x$ [illegible] [illegible] [illegible] [illegible]

$$\tau_x = \mathbb{E}[Y_i(1) - Y_i(0) \mid X_i = x] \tag{150}$$

$$= \frac{\sum_{i=1}^{n} \mathbb{I}(X_i = x)[Y_i(1) - Y_i(0)]}{\sum_{i=1}^{n} \mathbb{I}(X_i = x)} \quad \text{[illegible]} \tag{151}$$

[illegible] $\mathbb{I}(X_i = x)$ [illegible] $X_i = x$ [illegible] 1 [illegible] 0 [illegible]

[illegible] [illegible] **[illegible]** [illegible] [illegible] [illegible]

[illegible]

$$\tau = \frac{1}{n}\sum_{i=1}^{n}\{Y_i(1) - Y_i(0)\} \tag{152}$$

$$= \frac{1}{n}\sum_{i=1}^{n}\big[\mathbb{I}(X_i = 1) + \mathbb{I}(X_i = 0)\big]\{Y_i(1) - Y_i(0)\} \tag{153}$$

(154) [illegible] $\mathbb{I}(X_i = 1) + \mathbb{I}(X_i = 0) = 1$

$$= \underbrace{\frac{1}{n}\sum_{i=1}^{n}\mathbb{I}(X_i = 1)\{Y_i(1) - Y_i(0)\}}_{\text{treatment}} + \underbrace{\frac{1}{n}\sum_{i=1}^{n}\mathbb{I}(X_i = 0)\{Y_i(1) - Y_i(0)\}}_{\text{control}} \tag{155}$$

$$= \frac{n_1}{n}\cdot\underbrace{\frac{\sum_{i=1}^{n}\mathbb{I}(X_i = 1)\{Y_i(1) - Y_i(0)\}}{n_1}}_{\tau_1} + \frac{n_0}{n}\cdot\underbrace{\frac{\sum_{i=1}^{n}\mathbb{I}(X_i = 0)\{Y_i(1) - Y_i(0)\}}{n_0}}_{\tau_0} \tag{156}$$

$$= \pi_1\tau_1 + \pi_0\tau_0 \tag{157}$$

[illegible] $n_1 = \sum_{i=1}^{n}\mathbb{I}(X_i = 1)$ [illegible] $n_0 = \sum_{i=1}^{n}\mathbb{I}(X_i = 0)$ [illegible] $\pi_x = n_x/n$ [illegible]

[illegible] [illegible]

[illegible] $\delta_1 \neq \delta_2$ [illegible]

[illegible] $n = 4$ [illegible]

i	$Y_i(1)$	$Y_i(0)$
1	10	2
2	8	3
3	6	4
4	4	5

[illegible]

- $\delta_1 = \text{median}\{Y_i(1)\} - \text{median}\{Y_i(0)\} = 7 - 3.5 = 3.5$
- $\delta_2 = \text{median}\{Y_i(1) - Y_i(0)\} = \text{median}\{8, 5, 2, -1\} = 3.5$

[illegible] $\delta_1 = \delta_2$

[illegible] $n = 5$ [illegible]

i	$Y_i(1)$	$Y_i(0)$
1	100	1
2	90	2
3	50	3
4	10	4
5	5	5

[illegible]

- $\delta_1 = \text{median}\{Y_i(1)\} - \text{median}\{Y_i(0)\} = 50 - 3 = 47$
- $\delta_2 = \text{median}\{Y_i(1) - Y_i(0)\} = \text{median}\{99, 88, 47, 6, 0\} = 47$

[illegible] $\delta_1 > \delta_2$ [illegible]

i	$Y_i(1)$	$Y_i(0)$
1	100	0
2	1	0
3	1	0
4	1	0

[illegible]

- $\delta_1 = 50.5 - 0 = 50.5$
- $\delta_2 = \text{median}\{100, 1, 1, 1\} = 1$

[illegible] $\delta_1 = 50.5 > \delta_2 = 1$ [illegible] δ_1 [illegible] δ_2 [illegible]

[illegible]

[illegible] [illegible] $\hat{\tau}_X$ [illegible]
[illegible] $\text{cov}(\hat{\tau}_X) = \frac{n}{n_1 n_0} S_X^2$
[illegible] …[1]

[illegible] R^2 [illegible] [illegible] R^2 [illegible]
[illegible] { {
[illegible] …

[illegible] [illegible]
[illegible] $\hat{\tau}(\beta_1, \beta_0)$ [illegible]
[illegible] …

[illegible] [illegible] $\hat{\tau}(\hat{\beta}_1, \hat{\beta}_0)$ [illegible] OLS [illegible] Z_i [illegible]
[illegible] OLS [illegible]
[illegible] …

[illegible] [illegible] $\hat{\tau}_{\text{pred}} = \hat{\tau}_{\text{L}}$
[illegible] Lin [illegible]
[illegible] …

[illegible] [illegible] $\hat{\tau}(\beta_1, \beta_0)$ [illegible]
[illegible] $\hat{\tau}(\beta_1, \beta_0) = \hat{\tau} - \gamma^\top \hat{\tau}_X$
[illegible] …

---

[1] [illegible]

R w [illegible]

R w q w [illegible] $\sum_{i=1}^n(a_i - \bar{a})^2 = \sum_{i=1}^n a_i^2 - n\bar{a}^2$ [illegible] $\mathrm{var}(W) = \mathbb{E}(W^2) - (\mathbb{E}W)^2$ [illegible]

$$
\begin{aligned}
n(n-1)\mathbb{E}(\hat{V}) &= \mathbb{E}\left\{\sum_{i=1}^n(\hat{\tau}_i - \hat{\tau})^2\right\} \\
&= \mathbb{E}\left(\sum_{i=1}^n \hat{\tau}_i^2 - n\hat{\tau}^2\right) \\
&= \sum_{i=1}^n \left\{\mathrm{var}(\hat{\tau}_i) + \tau_i^2\right\} - n\left\{\mathrm{var}(\hat{\tau}) + \tau^2\right\} \\
&= \sum_{i=1}^n \mathrm{var}(\hat{\tau}_i) - n\mathrm{var}(\hat{\tau}) + \sum_{i=1}^n \tau_i^2 - n\tau^2 \\
&= n^2\mathrm{var}(\hat{\tau}) - n\mathrm{var}(\hat{\tau}) + \sum_{i=1}^n(\tau_i - \tau)^2.
\end{aligned}
$$

[illegible]

$$
\mathbb{E}(\hat{V}) = \mathrm{var}(\hat{\tau}) + \{n(n-1)\}^{-1}\sum_{i=1}^n(\tau_i - \tau)^2 \geq \mathrm{var}(\hat{\tau}).
$$

□

R w ‖ w [illegible] $(\hat{\tau}_1, \ldots, \hat{\tau}_n)^\top$ [illegible] OLS [illegible]

$$
\hat{\beta}_0 = \bar{\hat{\tau}} = n^{-1}\sum_{i=1}^n \hat{\tau}_i = \hat{\tau},
$$

[illegible] $\hat{\tau}$ [illegible] $\hat{\varepsilon}_i = \hat{\tau}_i - \hat{\tau}$ [illegible]

$$
\hat{\sigma}^2 = (n-1)^{-1}\sum_{i=1}^n \hat{\varepsilon}_i^2 = (n-1)^{-1}\sum_{i=1}^n(\hat{\tau}_i - \hat{\tau})^2 = \hat{V}.
$$

[illegible] $\hat{\tau}$ [illegible] $\hat{V}$ [illegible] OLS [illegible] □

m w [illegible]

m w q̈ w t [illegible] **w** H_{0F} [illegible] $\hat{\tau}_i = S_iD_i$ [illegible] $S_i = 2Z_i - 1 \in \{-1, 1\}$ [illegible] $D_i = Y_{i1} - Y_{i2}$ [illegible] S_i [illegible] 0 [illegible] 1 [illegible]

$$
\mathrm{var}(\hat{\tau}_i) = \mathbb{E}(S_i^2 D_i^2) - \mathbb{E}(S_i)^2\mathbb{E}(D_i)^2 = D_i^2 - 0 = D_i^2 = \hat{\tau}_i^2.
$$

[illegible]

$$
\frac{\hat{\tau} - \mathbb{E}\hat{\tau}}{\sqrt{\mathrm{var}(\hat{\tau})}} = \frac{\hat{\tau}}{\sqrt{n^{-2}\sum_{i=1}^n \hat{\tau}_i^2}} \to \mathrm{N}(0,1)
$$

[illegible] □

[illegible] H_{0F} [illegible] $|\hat{\tau}_i|$ [illegible] R_i [illegible] $U_i = \mathbb{I}(\hat{\tau}_i > 0)$ [illegible] $W = \sum_{i=1}^n U_i R_i$ [illegible] $U_i \overset{\text{IID}}{\sim} \text{Bernoulli}(1/2)$ [illegible] R_i [illegible]

$$\mathbb{E}(W) = \sum_{i=1}^n \mathbb{E}(U_i) R_i = \frac{1}{2}\sum_{i=1}^n i = \frac{n(n+1)}{4},$$

$$\text{var}(W) = \sum_{i=1}^n \text{var}(U_i) R_i^2 = \frac{1}{4}\sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{24}.$$

[illegible] □

[illegible] H_{0F} [illegible] $\Delta = \sum_{i=1}^n \mathbb{I}(\hat{\tau}_i > 0) \sim \text{Binomial}(n, 1/2)$ [illegible]

$$\frac{\Delta - n/2}{\sqrt{n/4}} \to \text{N}(0,1)$$

[illegible] □

MPE [illegible] i [illegible] Z_i [illegible] $1 - Z_i$ [illegible]

$$\hat{\tau}_i = Z_i Y_{i1} + (1 - Z_i) Y_{i2} - [(1 - Z_i) Y_{i1} + Z_i Y_{i2}] = (2Z_i - 1)(Y_{i1} - Y_{i2}) = S_i D_i,$$

[illegible] $S_i = 2Z_i - 1$ [illegible] $D_i = Y_{i1} - Y_{i2}$

$Z_i \overset{\text{IID}}{\sim} \text{Bernoulli}(1/2)$ [illegible] $\mathbb{E}(Z_i) = 1/2$ [illegible] $\text{var}(Z_i) = 1/4$ [illegible] $\mathbb{E}(S_i) = 0$ [illegible] $\text{var}(S_i) = 1$ [illegible] S_i [illegible] D_i [illegible] D_i [illegible] S_i [illegible]

$$\mathbb{E}(\hat{\tau}_i) = \mathbb{E}(S_i)\mathbb{E}(D_i) = 0,$$

$$\text{var}(\hat{\tau}_i) = \mathbb{E}(S_i^2)\mathbb{E}(D_i^2) - \mathbb{E}(S_i)^2 \mathbb{E}(D_i)^2 = 1 \cdot \mathbb{E}(D_i^2) = \mathbb{E}(D_i^2).$$

$D_i = Y_{i1} - Y_{i2}$ [illegible]

$$\mathbb{E}(D_i^2) = \mathbb{E}\left[\{Z_i Y_{i1}(1) + (1 - Z_i) Y_{i1}(0) - Z_i Y_{i2}(0) - (1 - Z_i) Y_{i2}(1)\}^2\right].$$

[illegible] $Z_i^2 = Z_i$ [illegible] $(1 - Z_i)^2 = 1 - Z_i$ [illegible] $Z_i(1 - Z_i) = 0$ [illegible] $\mathbb{E}(Z_i) = 1/2$ [illegible] $\mathbb{E}(Z_i^2) = 1/2$ [illegible][2] [illegible] $\text{var}(\hat{\tau}_i) = \mathbb{E}(Y_{i1}(1) - Y_{i2}(1) - Y_{i1}(0) + Y_{i2}(0))^2/4 = \tau_i^2$ [illegible]

[illegible] $\hat{\tau} = n^{-1}\sum_{i=1}^n \hat{\tau}_i$ [illegible]

$$\text{var}(\hat{\tau}) = n^{-2}\sum_{i=1}^n \text{var}(\hat{\tau}_i) + [illegible] .$$

Z_i [illegible] $\hat{\tau}_i$ [illegible]

$$\text{var}(\hat{\tau}) = n^{-2}\sum_{i=1}^n \text{var}(\hat{\tau}_i).$$

□

---

[2] [illegible]

q q [illegible]

[illegible] (i, j) [illegible] X_{ij} [illegible]

$$\hat{\tau}_{X,i} = X_{i1} - X_{i2},$$

[illegible] $\hat{\tau}_X = n^{-1} \sum_{i=1}^{n} \hat{\tau}_{X,i}$ MPE [illegible] Z_i [illegible] X_{ij} [illegible] [illegible]

$$\mathbb{E}(\hat{\tau}_{X,i}) = \mathbb{E}(X_{i1} - X_{i2}) = \mathbb{E}(X_{i1}) - \mathbb{E}(X_{i2}) = 0,$$

[illegible] $\mathbb{E}(\hat{\tau}_X) = 0$ [illegible]

$$\operatorname{cov}(\hat{\tau}_X) = n^{-2} \sum_{i=1}^{n} \operatorname{cov}(\hat{\tau}_{X,i}) = n^{-2} \sum_{i=1}^{n} \hat{\tau}_{X,i} \hat{\tau}_{X,i}^{\top},$$

[illegible] $\hat{\tau}_{X,i}$ [illegible] $\hat{\tau}(\gamma) = \hat{\tau} - \gamma^{\top} \hat{\tau}_X$ [illegible]

$$\operatorname{var}(\hat{\tau}(\gamma)) = \operatorname{var}(\hat{\tau}) + \gamma^{\top} \operatorname{cov}(\hat{\tau}_X) \gamma - 2\gamma^{\top} \operatorname{cov}(\hat{\tau}_X, \hat{\tau}).$$

[illegible]

$$\tilde{\gamma} = \operatorname{cov}(\hat{\tau}_X)^{-1} \operatorname{cov}(\hat{\tau}_X, \hat{\tau}).$$

$\operatorname{cov}(\hat{\tau}_X)$ [illegible] $\operatorname{cov}(\hat{\tau}_X, \hat{\tau})$ [illegible] { { [illegible] $\hat{\theta}$ [illegible] [illegible] $\hat{\gamma}$ [illegible] $\hat{\tau}_{\text{adj}} = \hat{\tau} - \hat{\gamma}^{\top} \hat{\tau}_X$ [illegible] $\hat{V}_{\text{adj}}$ □

q [illegible] p e [illegible] p[illegible]s

1 8 [illegible]

[illegible]	[illegible] $\hat{\tau}_{X,i}$	[illegible] $\hat{\tau}_i$
1	-1	-5
2	2	-1
3	0	3
4	1	2
5	2	5
6	0	-3
7	0	-1
8	2	9

[illegible]

$$\hat{\tau} = \frac{1}{8} \sum_{i=1}^{8} \hat{\tau}_i = \frac{-5 + (-1) + 3 + 2 + 5 + (-3) + (-1) + 9}{8} = \frac{9}{8} = 1.125.$$

[illegible]

$$\hat{V} = \frac{1}{8 \times 7} \sum_{i=1}^{8} (\hat{\tau}_i - \hat{\tau})^2 = \frac{1}{56} \left[(-6.125)^2 + (-2.125)^2 + (1.875)^2 + \cdots \right] \approx 4.98.$$

95% [illegible] $\hat{\tau} \pm 1.96\sqrt{\hat{V}} = 1.125 \pm 1.96 \times 2.23 = [-3.25, 5.50]$

[illegible] $\hat{\gamma}$ [illegible] $\hat{\tau}_{\text{adj}}$ [3]

q～[illegible]

q～w q w { { [illegible] **w** [illegible] ignorability [illegible]

$$\Pr\{Z=1 \mid Y(1),Y(0),e(X)\} = \Pr\{Z=1 \mid e(X)\}.$$

[illegible] tower property[illegible]

$$\Pr\{Z=1 \mid Y(1),Y(0),e(X)\} = E\{Z \mid Y(1),Y(0),e(X)\}.$$

[illegible] tower property[illegible] X [illegible] $Y(1),Y(0),e(X)$ [illegible]

$$E\{Z \mid Y(1),Y(0),e(X)\} = E\big[E\{Z \mid Y(1),Y(0),X\} \mid Y(1),Y(0),e(X)\big].$$

[illegible] ignorability [illegible] $Z \perp \{Y(1),Y(0)\} \mid X$[illegible]

$$E\{Z \mid Y(1),Y(0),X\} = E\{Z \mid X\} = e(X).$$

[illegible]

$$E\{Z \mid Y(1),Y(0),e(X)\} = E\{e(X) \mid Y(1),Y(0),e(X)\} = e(X).$$

[illegible]

$$\Pr\{Z=1 \mid e(X)\} = E\{Z \mid e(X)\} = E\big[E\{Z \mid X\} \mid e(X)\big] = E\{e(X) \mid e(X)\} = e(X).$$

[illegible]

q～w ‖ w k] [illegible] **P、** ～ [illegible] **w** [illegible] R [illegible]

```
# 读取数据
nhanes'bmi = Q ; JN。("nhanes'bmi.csv")[, −1]
z = nhanes'bmiOSchoolmeal
y = nhanes'bmiOBMI
x = ; 。j ; — (nhanes'bmi[, −J(1, 2)])
x = 。 J ;(g)Q # 标准化协变量

# 估计倾向得分
pscore = Z g(jz > x, V ; j a=g》B a l p j) O y g a —.−valQNes

# 倾向得分分层
n.strata = J(5, 10, 20, 50, 80)
strat.res = 。 ; s s(n.g》strata, FUN = V ' l J −(nn) p {
   z.pscore = z ' ; l −(pascgoQe, (1:(nn−1)) ¨ nn)
   s。strata = J '(−pscore, breaks = J(0, z.pscore, 1),
                  g ; B Q=g 1:nn)
  Neyman'SRE(z, y, s。strata)
})

  p〕 l ;(jstQrat.res) = J("est", "se")
J p g l ;(jstQrat.res) = n.strata
```

---

[3] [illegible]

p ‘ (lsNrat.res, 3)

ʕ `Neyman_SRE` ŋ̇ṁ̄ ᴅ { {ŏ Ωᵩ SRE ⅧΣ ḽ Neyman ʽΙΩ ʰˢʕ

Ḫ

q Xὀẅ̄ ɹ ‖ mẅỵڨq

̄ ɹ { {ڨ ŏ

$$E\{Y(z_1,z_2)\} = E[E\{Y(z_1,z_2) \mid X_0\}]$$

ᴮ ۏ̇ʲ▌ $E\{Y(z_1,z_2) \mid X_0\}$ Assumption { (1) ῼ ŏ

$$\begin{aligned}E\{Y(z_1,z_2) \mid X_0\} &= E\{Y(z_1,z_2) \mid z_1, X_0\} \\ &= E[E\{Y(z_1,z_2) \mid z_1, X_1, X_0\} \mid z_1, X_0]\end{aligned}$$

Assumption { (2)ŏ

$$\begin{aligned}E\{Y(z_1,z_2) \mid X_0\} &= E[E\{Y \mid z_2, z_1, X_1, X_0\} \mid z_1, X_0] \\ &= E\{E(Y \mid z_2, z_1, X_1, X_0) \mid z_1, X_0\}\end{aligned}$$

ڨڤ ḏķ { { □

APPENDIX A

[illegible]

[illegible]

[illegible] [illegible]

[illegible] subgroup [illegible] X_i [illegible]

[illegible] $\tau = \pi_1\tau_1 + \pi_0\tau_0$ [illegible] $\tau = \pi_1\tau_1 + \pi_0\tau_0$ [illegible]

[illegible] $\mathbb{I}(X_i = 1) + \mathbb{I}(X_i = 0) = 1$ [illegible]

[illegible] H_{0F} $\mathbf{Y}$ [illegible] Fisher's sharp null ($H_{0F}: Y_i(1) = Y_i(0)$) [illegible] $\mathbf{Y}$ [illegible]

[illegible] H_{0F} [illegible] treatment [illegible] $Y_i(1) = Y_i(0) = Y_i$ [illegible] i [illegible] $\mathbf{Y}(1) = \mathbf{Y}(0) = \mathbf{Y}$ [illegible] FRT [illegible] condition on [illegible] $\mathbf{Y}$ [illegible] $\mathbf{Z}$ [illegible] $M = \binom{n}{n_1}$ [illegible] FRT" [illegible] " [illegible]

[illegible] [illegible] $\tilde{p}_{frt}$ [illegible]

[illegible]

$$\hat{p}_{frt} = \frac{1}{R}\sum_{r=1}^{R} \mathbb{I}\{T(\mathbf{z}^r, \mathbf{Y}) \geq T(\mathbf{Z}, \mathbf{Y})\}$$

[illegible] R [illegible] p [illegible]

[illegible]

$$\tilde{p}_{frt} = \frac{1 + \sum_{r=1}^{R} \mathbb{I}\{T(\mathbf{z}^r, \mathbf{Y}) \geq T(\mathbf{Z}, \mathbf{Y})\}}{1 + R}$$

[illegible] 1 [illegible] " [illegible] " [illegible]

[illegible] $R = 99$ [illegible] 0 [illegible]

$$\hat{p}_{frt} = \frac{0}{99} = 0$$

[illegible] M [illegible] p [illegible] $1/M$ [illegible] 0

[illegible] 1 [illegible]

$$\tilde{p}_{frt} = \frac{1}{100}$$

[illegible]

(1) p [illegible] 1[illegible]
(2) [illegible] R [illegible] $\mathbb{P}(p \le u) \le u$
(3) [illegible] $R \to \infty$ [illegible] $\tilde{p}_{frt} \to \hat{p}_{frt}$

[illegible] [illegible] FRT [illegible] p [illegible] ” [illegible] ” [illegible]

$$\mathbb{P}(p_{frt} \le u) \le u \quad \text{for all } u \in [0,1]$$

[illegible] finite-sample validity [illegible] p_{frt} [illegible] FRT [illegible]

[illegible] [illegible] Fisher [illegible]

- [illegible] “[illegible]” [illegible]
- **[illegible]** “[illegible]” [illegible]

Fisher [illegible] sharp null [illegible] FRT [illegible] justify [illegible]

[illegible] [illegible] justify [illegible] FRT [illegible] FRT [illegible]

- [illegible] Fisher [illegible] $Y_i \sim \text{i.i.d. } \mathcal{N}(\mu, \sigma^2)$ [illegible]
- [illegible] CLT [illegible]

[illegible] FRT [illegible] ” [illegible] ” [illegible]

- [illegible] ” [illegible] ” [illegible]
- FRT [illegible] ” [illegible] ” [illegible]

[illegible] CRE [illegible] p [illegible] ” [illegible] ” [illegible] FRT [illegible]

[illegible] [illegible] FRT

[illegible]

- [illegible]
- [illegible] CLT [illegible]

[illegible] [illegible] + sharp null [illegible] Monte Carlo [illegible]

- [illegible]
- FRT [illegible] treatment [illegible]

FRT [illegible] CLT [illegible] FRT [illegible] p [illegible]

[illegible] FRT [illegible] FRT [illegible] p

[illegible] Fisher [illegible]

[illegible] Potential Outcomes [illegible] treatment [illegible]

- $Y_i(1)$ [illegible] i [illegible] treatment [illegible]
- $Y_i(0)$ [illegible] i [illegible] control [illegible]

[illegible] treatment [illegible] counterfactual [illegible]

$$Y_i^{obs} = Z_i Y_i(1) + (1 - Z_i) Y_i(0)$$

[illegible] $Z_i = 1$ [illegible] treatment [illegible] $Y_i(1)$ [illegible] $Z_i = 0$ [illegible] $Y_i(0)$ [illegible] $Y_i(1) - Y_i(0)$ [illegible] i [illegible]

[illegible] Fisher [illegible]

[illegible] treatment [illegible] control [illegible] selection bias [illegible]

[illegible] randomization mechanism [illegible] p [illegible]

(1) [illegible] H_{0F} [illegible] treatment [illegible] $Y_i(1) = Y_i(0)$ [illegible]
(2) [illegible]
(3) [illegible] Monte Carlo [illegible]
(4) [illegible]

[illegible]

Fisher Randomization Test

Fisher

Fisher Randomization Test (FRT) Ronald Fisher 1935 Design of Experiments

(1) $H_{0F}: Y_i(1) = Y_i(0)$

Y

(2) **Z**

(3) CRE **Z** $\binom{n}{n_1}$

(4)

(5) p

FRT condition on **Y** **Z** condition on FRT condition on p FRT finite-sample exactness

$\{\mathbf{z}^1, \ldots, \mathbf{z}^M\}$

CRE **Z** $\{\mathbf{z}^1, \ldots, \mathbf{z}^M\}$

Z $\mathbf{Z} = (Z_1, \ldots, Z_n)$ $Z_i = 1$ i $Z_i = 0$ i

$\{\mathbf{z}^1, \ldots, \mathbf{z}^M\}$ $n = 3, n_1 = 1$

$$\mathbf{z}^1 = (1,0,0), \mathbf{z}^2 = (0,1,0), \mathbf{z}^3 = (0,0,1)$$

$M = \binom{3}{1} = 3$

CRE

$\mathbf{z}^m$

$$\mathbb{P}(\mathbf{Z} = \mathbf{z}^m) = \frac{1}{M} = \frac{1}{\binom{n}{n_1}}$$

$n = 100, n_1 = 50$ $M \approx 10^{29}$ 10^{-29}

CRE treatment control Fisher "randomization provides a valid basis for inference"

Z

Z **Z**

Z Fisher treatment

Z

Z n $\mathbf{Z} = (Z_1, Z_2, \ldots, Z_n)$ n

- $Z_i = 1$ [illegible] i [illegible] treatment [illegible]
- $Z_i = 0$ [illegible] i [illegible] control [illegible]

[illegible] $n = 5, n_1 = 2$ [illegible] $\mathbf{Z} = (1,0,0,1,0)$ [illegible] 1 [illegible] 4 [illegible] 2 [illegible] 3 [illegible] 5 [illegible]

[illegible] CRE [illegible] $\mathbf{Z}$ [illegible] $M = \binom{n}{n_1}$ [illegible] $1/M$ [illegible]

- $n = 3, n_1 = 1$ [illegible] $M = 3$ [illegible] $(1,0,0)$ [illegible] $(0,1,0)$ [illegible] $(0,0,1)$ [illegible] $1/3$
- $n = 100, n_1 = 50$ [illegible] $M \approx 10^{29}$ [illegible] 10^{-29}

$\mathbf{z}^r$ [illegible] Monte Carlo [illegible] M [illegible] R [illegible] $\mathbf{z}^r$ [illegible] $\mathbf{z}^r$ [illegible] $\mathbf{Z}$ [illegible] ⋆ [illegible] $\mathbf{Z}$ [illegible] $\mathbf{Z}$ [illegible] indicator vector [illegible] Monte Carlo [illegible]

[illegible]

[illegible] Potential Outcomes [illegible] Potential Outcomes [illegible] ** [illegible] ** [illegible] i [illegible]

- $Y_i(1)$ [illegible] i [illegible] treatment [illegible]
- $Y_i(0)$ [illegible] i [illegible] control [illegible]

[illegible] ** [illegible] ** [illegible] counterfactual [illegible] Fundamental Problem of Causal Inference [illegible]

[illegible] First Fundamental Form [illegible]

[illegible]

[illegible] far confidence set [illegible]

[illegible]

[illegible] ITT [illegible] Intent-to-Treat [illegible]

[illegible] ITT [illegible] Intent-to-Treat [illegible]

[illegible] Instrumental Variable [illegible]

[illegible] Z [illegible] 1 [illegible] treatment [illegible] 2 [illegible] 3 [illegible] treatment [illegible] variation [illegible] variation [illegible] treatment [illegible]

[illegible] Monotonicity [illegible]

[illegible] $Y_i(1) \geq Y_i(0)$ [illegible]

Randomization Assumption

Z $(Y_i(0), Y_i(1))$ $(D_i(0), D_i(1))$

$Z \perp (Y_i(0), Y_i(1), D_i(0), D_i(1))$

Weak IV Problem

treatment F

F

Staiger-Stock 10

LATE ATE

Compliers” variation

Compliers $D_i(1) > D_i(0)$ Defiers Always-takers

Compliers ATE LATE

m Y

$m(X) = \mathbb{E}[Y|X]$

m Logit

m

Robins mediation formula latent groups

Y

cross-world independence

mediation analysis

Surrogate Sr Jr

当你在阅读过程中提出问题时，回答将被记录在此处。

在这里记录你的学习心得和思考。

A a B g a p Z ; s ^⟫

Joshua D. Angrist, Guido W. Imbens, and Donald B. Rubin. Identification of causal effects using instrumental variables. *Journal of the American Statistical Association*, 91(434): 444–472, 1996. Cited in Chapter 21.

Peter J. Bickel, Eugene A. Hammel, and J. William O'Connell. Sex bias in graduate admissions: Data from berkeley. *Science*, 187(4175):398–404, 1975. Cited in Chapter 1.

Marie-Abele C. Bind and Lea M. Rattay. When possible, report a fisher-exact p-value and display its underlying null randomization distribution. *Journal of the American Statistical Association*, 115(532):1644–1646, 2020. Cited in Chapter 3.

George E. P. Box, J. Stuart Hunter, and William G. Hunter. *Statistics for Experimenters: An Introduction to Experimental Design*. John Wiley & Sons, 1978. Cited in Chapter 5.

Peng Ding. A paradox from randomization-based causal inference. *Statistical Science*, 32 (3):331–345, 2017. Cited in Chapter 4.

Peng Ding and Tyler J. VanderWeele. Sensitivity analysis without assumptions. *Epidemiology*, 27(3):368–377, 2016. Cited in Chapters 17, 18.

Herman F. Dorn. The planner's dilemma. *Journal of the American Statistical Association*, 48(264):601–604, 1953. Cited in Chapters 18, 21.

Ronald A. Fisher. The design of experiments. *Statistical Research Memoirs*, 1:1–26, 1935. Cited in Chapter 3.

J. L. Hodges and E. L. Lehmann. Rank methods for combination of independent experiments in analysis of variance. *The Annals of Mathematical Statistics*, 33(2):482–497, 1962. Cited in Chapter 5.

Paul W. Holland. Statistics and causal inference. *Journal of the American Statistical Association*, 81(396):945–960, 1986. Cited in Chapters 1, 2, 3.

Guido W. Imbens. Sensitivity to exogeneity assumptions in program evaluation. *American Economic Review*, 93(2):126–132, 2003. Cited in Chapter 18.

Guido W. Imbens and Joshua D. Angrist. Identification and estimation of local average treatment effects. *Econometrica*, 62(2):467–475, 1994. Cited in Chapter 21.

Guido W. Imbens and Donald B. Rubin. *Causal Inference for Statistics, Social, and Biomedical Sciences: An Introduction*. Cambridge University Press, 2015. Cited in Chapters 4, 5.

Gary King and Langche Zeng. The dangers of extreme counterfactuals. *Political Analysis*, 14(2):131–159, 2006. Cited in Chapter 20.

Winston Lin. Agnostic notes on regression adjustments to experimental data: Reexamining freedman's critique. *The Annals of Applied Statistics*, 7(1):295–318, 2013. Cited in Chapter 3.

Bo Lu and Peng Ding. Sensitivity analysis for causal inference: Theory and applications. *Unpublished manuscript*, 2023. Cited in Chapter 18.

Charles F. Manski. Nonparametric bounds on treatment effects. *American Economic Review*, 80(2):319–323, 1990. Cited in Chapter 18.

Charles F. Manski. *Monitoring the dissolving of the Soviet Union*, volume 21. 2003. Cited in Chapter 18.

Jerzy Neyman. Sur les applications de la théorie des probabilités aux experiences agricoles: Essai d'application de la méthode "chia". *Roczniki Nauk Rolniczych*, 10:1–51, 1923. Cited in Chapters 2, 4.

Judea Pearl. On the consistency rule in causal inference. *Biometrika*, 97(2):490–492, 2010. Cited in Chapter 2.

Judea Pearl and Dana Mackenzie. *The Book of Why: The New Science of Cause and Effect*. Basic Books, 2018. Cited in Chapter 1.

Paul R. Rosenbaum and Donald B. Rubin. Assessing sensitivity to an unobserved binary covariate in an observational study with binary outcome. *Journal of the Royal Statistical Society: Series B*, 45(2):212–218, 1983. Cited in Chapters 17, 18, 19.

Donald B. Rubin. Comment: Bayesian inference for causal effects: The role of randomization. *The Annals of Statistics*, 8(1):44–45, 1980. Cited in Chapter 3.

Donald B. Rubin. Causal inference using potential outcomes: Design, modeling, decisions. *Journal of the American Statistical Association*, 100(469):322–331, 2005. Cited in Chapter 2.

Donald B. Rubin. For objective causal inference, design trumps analysis. *The Annals of Applied Statistics*, 2(3):808–840, 2008. Cited in Chapter 20.

P. H. Van Elteren. On the combination of independent statistical tests. *Statistica Neerlandica*, 14(1):11–24, 1960. Cited in Chapter 5.

Tyler J. VanderWeele and Peng Ding. Introducing the e-value. *Annals of Internal Medicine*, 167(4):285, 2017. Cited in Chapters 17, 18.