

PROOF OF POSITIVITY IN QUANTUM TRIPLE SCHUBERT CALCULUS

YUE YIHUA AND XIAO TONGREN

ABSTRACT. We prove the positivity conjecture for the product of quantum double Schubert polynomials with three sets of variables. Specifically, for any $u, v \in S_n$, the expansion coefficients

$$c_{u,v}^{w,(d)}(\mathbf{y}, \mathbf{z})$$

in

$$\tilde{\mathfrak{S}}_u(\mathbf{x}; \mathbf{y}) \cdot \tilde{\mathfrak{S}}_v(\mathbf{x}; \mathbf{z}) = \sum_d \sum_w q^d c_{u,v}^{w,(d)}(\mathbf{y}, \mathbf{z}) \tilde{\mathfrak{S}}_w(\mathbf{x}; \mathbf{z}).$$

are polynomials in $\mathbf{y}$ and $\mathbf{z}$ with nonnegative integer coefficients in the negative roots $y_i - z_j$. Our approach combines the equivariant quantum cohomology framework of Mihalcea [?] with the triple Schubert calculus of Gao–Xiong [?], extending the Graham positivity theorem [?] to the quantum setting.

1. INTRODUCTION

The theory of Schubert polynomials, initiated by Lascoux and Schützenberger, provides polynomial representatives for Schubert classes in the cohomology of flag varieties. In the classical setting, the Littlewood–Richardson coefficients $c_{u,v}^w$ appearing in

$$\mathfrak{S}_u(\mathbf{x}) \cdot \mathfrak{S}_v(\mathbf{x}) = \sum_w c_{u,v}^w \mathfrak{S}_w(\mathbf{x})$$

are nonnegative integers, reflecting the geometric fact that they count intersection points of Schubert varieties. Graham’s seminal theorem [?] established that in the equivariant setting these coefficients become polynomials in the negative roots x_i with nonnegative integer coefficients.

The quantum deformation of this story was developed by Kirillov and Maeno [?], who introduced quantum double Schubert polynomials $\tilde{\mathfrak{S}}_w(\mathbf{x}; \mathbf{y})$. In the quantum setting the product involves formal variables q^β encoding curve-counting information:

$$(1) \quad \tilde{\mathfrak{S}}_u(\mathbf{x}; \mathbf{y}) \cdot \tilde{\mathfrak{S}}_v(\mathbf{x}; \mathbf{z}) = \sum_d \sum_w q^d c_{u,v}^{w,(d)}(\mathbf{y}, \mathbf{z}) \tilde{\mathfrak{S}}_w(\mathbf{x}; \mathbf{z}).$$

Mihalcea [?] proved that when $\mathbf{y} = \mathbf{z}$ (the double case), the coefficients $c_{u,v}^{w,(d)}(\mathbf{y}, \mathbf{y})$ are polynomials in the negative roots $y_i - y_j$ with nonnegative integer coefficients, establishing the Peterson conjecture for equivariant quantum cohomology.

In this paper we prove the corresponding positivity for the *triple* case, where $\mathbf{y} \neq \mathbf{z}$ are independent variable sets.

The proof strategy follows the method of Mihalcea, combined with the triple Schubert calculus established by Gao and Xiong in [?]. The key idea is a reduction from the quantum triple case to the already-established double positivity of Mihalcea, via a specialization argument using the separated descents condition of [?].

2. NOTATIONS AND PRELIMINARIES

2.1. Flag variety and Weyl group. Let $G = SL_n(\mathbb{C})$ and $B \subset G$ a Borel subgroup of upper triangular matrices. The **flag variety** is $Fl_n = G/B$, a smooth complex algebraic variety of dimension $\binom{n}{2}$. Its Weyl group is the symmetric group S_n , generated by simple reflections s_i ($i = 1, \dots, n-1$) satisfying the Coxeter relations. For $w \in S_n$ the **length** $\ell(w)$ equals the number of inversions of w , and $w_0 = (n, n-1, \dots, 1)$ denotes the longest element.

The set of **descents** of w is

$$\text{Des}(w) = \{ i : w(i) > w(i+1) \},$$

and we say (u, v) satisfies the **separated descents condition** if $\max \text{Des}(u) \leq \min \text{Des}(v)$; see [? , Definition 2.1].

2.2. Divided difference operators. For $i = 1, \dots, n-1$, the **divided difference operator** ∂_i acts on polynomials in variables $x_1, \dots, x_n$ by

$$\partial_i f = \frac{f(\dots, x_i, x_{i+1}, \dots) - f(\dots, x_{i+1}, x_i, \dots)}{x_i - x_{i+1}},$$

and these operators satisfy $\partial_i^2 = 0$ and the braid relations, so for any $w \in S_n$ the operator $\partial_w = \partial_{i_1} \cdots \partial_{i_k}$ is well-defined when $(i_1, \dots, i_k)$ is a reduced word for w .

The **skew divided difference operator** for $v \leq w$ in Bruhat order is

$$\partial_{w/v} = \sum_{\substack{u \\ v \leq u \leq w}} (-1)^{\ell(w) - \ell(u)} \partial_u.$$

2.3. Schubert polynomials. The (single) Schubert polynomials of Lascoux–Schützenberger are defined by

$$\mathfrak{S}_w(\mathbf{x}) = \partial_{w^{-1}w_0}(x_1^{n-1}x_2^{n-2} \cdots x_n^0).$$

The **double Schubert polynomials** introduce a second variable set:

$$\mathfrak{S}_w(\mathbf{x}; \mathbf{y}) = \partial_{w^{-1}w_0}^{(\mathbf{x})} \prod_{i+j \leq n} (x_i + y_j).$$

The **quantum double Schubert polynomials** $\tilde{\mathfrak{S}}_w(\mathbf{x}; \mathbf{y})$ were introduced in [?]. They depend on n variables $\mathbf{x} = (x_1, \dots, x_n)$, another n variables $\mathbf{y} = (y_1, \dots, y_n)$, and $n-1$ quantum variables $\mathbf{q} = (q_1, \dots, q_{n-1})$. The top polynomial is

$$\tilde{\mathfrak{S}}_{w_0}(\mathbf{x}; \mathbf{y}) = \prod_{i=1}^{n-1} \Delta_i(y_{n-i} \mid x_1, \dots, x_i),$$

where

$$\Delta_k(t \mid x_1, \dots, x_k) = \sum_{j=0}^k t^{k-j} e_j(x_1, \dots, x_k \mid q_1, \dots, q_{k-1}),$$

and $e_j(\cdot \mid \mathbf{q})$ are the **quantum elementary symmetric functions**. For any $w \in S_n$,

$$\tilde{\mathfrak{S}}_w(\mathbf{x}; \mathbf{y}) = \partial_{w w_0}^{(\mathbf{y})} \tilde{\mathfrak{S}}_{w_0}(\mathbf{x}; \mathbf{y}).$$

When $q = 0$ we recover the classical double Schubert polynomial: $\tilde{\mathfrak{S}}_w(\mathbf{x}; \mathbf{y})|_{q=0} = \mathfrak{S}_w(\mathbf{x}; \mathbf{y})$.