

Differential Geometry of Curves and Surfaces

Manfredo P. do Carmo

March 24, 2026

Contents

引言：文献概述与背景知识	5
0.1 总述	5
0.2 文献概述	5
0.3 背景知识安排	6
0.4 学习路径	6
0.5 与其他教材的关系	6
1 Curves	7
1.1 Introduction	7
1.2 Parametrized Curves	8
1.3 Regular Curves; Arc Length	12
1.4 The Vector Product in $\mathbb{R}^3$	18
1.5 The Local Theory of Curves Parametrized by Arc Length	21
1.5.1 The Curvature	21
1.5.2 The Torsion (Or Bending)	22
1.5.3 The Frenet-Serret Formulas	24
1.6 The Local Canonical Form	30
1.7 Global Properties of Plane Curves	32

引言：文献概述与背景知识

0.1 总述

本笔记学习的主题是 **微分几何 (differential geometry)** ——研究曲线和曲面几何性质的数学分支。这门学科发端于微积分诞生之时，由 Gauss、Riemann 等数学家发展成熟。

本笔记以 Manfredo P. do Carmo 的经典教材《Differential Geometry of Curves and Surfaces》为主线，系统学习曲线与曲面的局部理论。

0.2 文献概述

Differential Geometry of Curves and Surfaces

作者: Manfredo P. do Carmo

出版社: Dover Publications (Revised & Updated Second Edition, 2016)

原版: Prentice-Hall, 1976

页数: 约 500 页

内容简介: 这是微分几何领域最经典的入门教材之一。do Carmo 来自巴西 IMPA (数学与应用数学研究所)，他的这本书以清晰、严谨的风格著称。

全书分为五章：

- **第 1 章** 曲线理论—局部几何 (曲率、挠率、Frenet-Serret 公式)
- **第 2 章** 正则曲面—曲面的定义、切平面、第一基本形式
- **第 3 章** Gauss 映射—曲面的曲率、第二基本形式
- **第 4 章** 曲面内在几何—等距映射、测地线、Gauss-Bonnet 定理
- **第 5 章** 整体微分几何—曲面的整体性质

写作特点:

- 每章开头有引言，说明该章内容和学习路线
- 大量详细的例子和图示
- 习题丰富，配有提示和答案
- 强调几何直观与代数方法的结合

前置要求:

- 线性代数（向量、矩阵、行列式）
- 多变量微积分（偏导数、链式法则、积分）
- 微分方程（初步了解即可）

0.3 背景知识安排

第 ?? 章收集学习曲线理论所需的**纯数学基础**，包括：

1. **欧氏空间** $\mathbb{R}^n$ — 向量、范数、内积的基本性质
2. **向量值函数** — 求导法则、链式法则
3. **向量积（叉积）** — $\mathbb{R}^3$ 中特有的运算
4. **行列式与有向体积** — 标量三重积的几何意义

这些内容大部分来自本科线性代数和微积分课程。这里统一整理，便于后续查阅。

0.4 学习路径

根据 do Carmo 在 Preface 中的建议，本笔记采用以下学习顺序：

1. **第 1 步**: 阅读第 ?? 章的背景知识
2. **第 2 步**: 学习第 1 章（曲线）— 1-2 节到 1-5 节是核心，1-6、1-7 节是拓展
3. **第 3 步**: 学习第 ?? 章（曲面）— 2-2、2-3 节是基础，2-4、2-5 节是核心
4. **第 4 步**: 深入第 3-4 章（Gauss 映射、内在几何）
5. **第 5 步**: 选读第 5 章（整体微分几何）

0.5 与其他教材的关系

微分几何的标准教材还有：

- do Carmo（本书）— 注重几何直观，例子丰富
- Klingenberg — 更代数化，强调抽象理论
- O'Neill — 入门友好，风格接近 do Carmo
- Morgan — 更加入门，适合本科生

do Carmo 的这本书适合作为**第一本**微分几何教材。

注 0.1. 主文献: *do Carmo - Differential Geometry of Curves and Surfaces*

注 0.2. 学习顺序: 先曲线，再曲面，最后整体理论

Chapter 1

Curves

1.1 Introduction

The problem we face. 我们生活在一个三维空间中，而视觉是我们感知世界的主要方式之一。当我们观察一条曲线——无论是一条道路、一条河流的轮廓、还是行星在天空中的运动轨迹——我们的大脑本能地识别它的“弯曲程度”。这种对弯曲的直观感知，引导我们提出一个根本性的问题：

如何用精确的数学语言来描述曲线的弯曲程度？


这个问题看似简单，却贯穿了整个微分几何的发展历程。从欧几里得几何中直线与圆的基本定义，到高斯研究曲面的内在几何，再到达布（Darboux）建立曲线与曲面的局部理论，这个问题的答案逐渐清晰：曲线的弯曲程度可以通过其在一点处的加速度来度量。

The main theme. 微分几何研究的是曲线和曲面的几何性质。与微积分研究函数变化不同，微分几何关注的是空间中曲线和曲面的“形状”——弯曲程度、扭曲程度等内在几何量。


经典微分几何始于微积分诞生之时，其核心是研究曲线和曲面的**局部性质**（local properties）——即那些仅依赖于曲线或曲面在一点附近行为的几何性质。使用的工具正是微分计算。我们将会看到，一个出人意料的事实是：空间曲线的形状完全由两个函数决定——曲率（curvature）和挠率（torsion）。

A guiding example. 在学习曲线的理论时，一个例子将贯穿始终，这就是圆柱螺旋线（cylindrical helix）。想象一条弹簧从空中掉落，它在空中划出的轨迹就是螺旋线。螺旋线的非凡之处在于：它在弯曲的同时保持恒定的扭转率。这种“均匀弯曲 + 均匀扭曲”的特性，使其成为理解曲率与挠率概念的完美载体。


Organization of this chapter. 本章的组织方式如下。1-2 节到 1-4 节包含基础内容（参数曲线、弧长、向量积），可作为复习；1-5 节是核心，引入曲率和挠率的概念；1-6 节和 1-7 节讨论局部标准形式和整体性质。若只想学习曲面，可以跳过 1-6、1-7 节。



(a) Figure 1-1. 圆柱螺旋线 (helix) — 弯曲与扭曲的完美结合



(b) Figure 1-2. 奇异点例子—曲线在尖点处不可微



(a) Figure 1-3. 非单射曲线例子—同一点被多次经过

1.2 Parametrized Curves

Why parametrization? 在我们提出“如何描述曲线的弯曲程度”这个问题之后，自然的下一步是找到描述曲线的数学工具。我们已经知道，微积分是研究函数变化的有力武器——那么，是否可以将曲线视为某种“函数”，从而运用微分学的工具？

这个想法引导我们引入**参数化**的概念。其核心思想简单而优雅：将曲线看作一个从区间到空间的映射。例如，当我们说“质点沿曲线运动”时，时间 t 是参数，位置 $\alpha(t)$ 是 t 的函数。这种视角不仅统一了古代对圆、椭圆等特殊曲线的研究，更为重要的是，它让我们能够使用导数来研究曲线的局部几何性质。

接下来，我们首先回顾 $\mathbb{R}^3$ 中向量内积的基本性质，这对后续讨论至关重要。

注 1.1. 设 $u = (u_1, u_2, u_3) \in \mathbb{R}^3$ ，定义其范数（或长度）为

$$|u| = \sqrt{u_1^2 + u_2^2 + u_3^2}.$$


几何上， $|u|$ 是点 (u_1, u_2, u_3) 到原点 $O = (0, 0, 0)$ 的距离。

设 $u = (u_1, u_2, u_3)$ 和 $v = (v_1, v_2, v_3)$ 是 $\mathbb{R}^3$ 中的向量， θ ($0 \leq \theta \leq \pi$) 是线段 Ou 和 Ov 之间的夹角。内积 $u \cdot v$ 定义为（见 fig. 1.3a, 教材 Figure 1-6）:

$$u \cdot v = |u||v| \cos \theta.$$

内积的基本性质：

1. 若 u 和 v 为非零向量，则 $u \cdot v = 0$ 当且仅当 u 与 v 正交；
2. $u \cdot v = v \cdot u$;
3. $\lambda(u \cdot v) = \lambda u \cdot v = u \cdot \lambda v$;
4. $u \cdot (v + w) = u \cdot v + u \cdot w$ 。



(a) Figure 1-6. 内积的几何意义

The central notion. 我们用 $\mathbb{R}^3$ 表示所有三元实数有序组 (x, y, z) 的集合。现在，我们正式引入参数化曲线的概念。

定义 1.1 (可微参数曲线). **可微参数曲线**是一个可微映射

$$\alpha : \mathbf{I} \rightarrow \mathbb{R}^3,$$

其中 $\mathbf{I} = (\mathbf{a}, \mathbf{b})$ 是实数直线 $\mathbb{R}$ 上的一个开区间。

可微意味着 α 将每个 $t \in I$ 映射到点 $\alpha(t) = (x(t), y(t), z(t)) \in \mathbb{R}^3$, 其中函数 $x(t), y(t), z(t)$ 都是可微的。变量 t 称为曲线的**参数**。

如果记 $x(t)$ 在 t 处的一阶导数为 $x'(t)$, 等等, 则向量

$$(x'(t), y'(t), z'(t)) = \alpha'(t) \in \mathbb{R}^3$$

称为曲线 α 在 t 处的**切向量** (tangent vector) 或**速度向量** (velocity vector)。像集 $\alpha(I) \subset \mathbb{R}^3$ 称为 α 的**迹** (trace)。如下面的例 5 所示, 我们必须注意区分参数曲线 (它是一个映射) 和它的迹 (它是 $\mathbb{R}^3$ 的一个子集)。


如果记 $x(t)$ 在 t 处的一阶导数为 $x'(t)$, 等等, 则向量

$$(x'(t), y'(t), z'(t)) = \alpha'(t) \in \mathbb{R}^3$$


称为曲线 α 在 t 处的**切向量** (tangent vector) 或**速度向量** (velocity vector)。像集 $\alpha(I) \subset \mathbb{R}^3$ 称为 α 的**迹** (trace)。如下面的例 5 所示, 我们必须注意区分参数曲线 (它是一个映射) 和它的迹 (它是 $\mathbb{R}^3$ 的一个子集)。

注 1.2. 术语说明: 许多人用“无限可微”来表示具有各阶导数 (且自动连续) 的函数, 而用“可微”仅表示一阶导数的存在性。我们不采用这种用法——本书中“可微”意味着所有阶导数都存在。

Examples. 为了建立直觉, 我们来看几个关键的例子, 它们将贯穿本章的讨论。



(a) Figure 1-1. 圆柱螺旋线 (helix)



(b) Figure 1-2. 奇异点例子

例 1.1 (圆柱螺旋线). 参数方程


$$\alpha(t) = (a \cos t, a \sin t, bt), \quad t \in \mathbb{R}$$

在 $\mathbb{R}^3$ 中的迹是圆柱面 $x^2 + y^2 = a^2$ 上的螺旋线 (helix), 螺距 (pitch) 为 $2\pi b$ 。这里参数 t 测量的是 x 轴与从原点 O 到点 $\alpha(t)$ 在 xy 平面上投影点的连线之间的夹角 (见 fig. 1.4a, 教材 Figure 1-1)。

正如本章引言所述, 螺旋线将成为我们理解曲率与挠率的完美载体——它的”均匀弯曲 + 均匀扭曲”特性, 使其在后续讨论中反复出现。

例 1.2 (奇异点). 映射 $\alpha: \mathbb{R} \rightarrow \mathbb{R}^2$, $\alpha(t) = (t^3, t^2)$, 是一条可微参数曲线 (见 fig. 1.4b, 教材 Figure 1-2)。注意 $\alpha'(0) = (0, 0)$ ——即 $t = 0$ 处的速度向量为零。这样的点称为**奇异点** (singular point)。

奇异点的出现提醒我们: 可微的参数方程并不保证曲线在每一点都有”良好”的行为。





(a) Figure 1-3. 非单射曲线例子

例 1.3 (非单射曲线). 映射 $\alpha: \mathbb{R} \rightarrow \mathbb{R}^2$, $\alpha(t) = (t^3 - 4t, t^2 - 4)$, 是一条可微参数曲线 (见 fig. 1.5a, 教材 Figure 1-3)。注意 $\alpha(2) = \alpha(-2) = (0, 0)$ ——即映射 α 不是单射。

这个例子说明一个重要事实: 参数化与曲线本身是不同的概念。同一条曲线可以有多种参数化方式。

例 1.4 (不可微曲线). 映射 $\alpha: \mathbb{R} \rightarrow \mathbb{R}^2$, $\alpha(t) = (t, |t|)$, $t \in \mathbb{R}$, 不是可微参数曲线, 因为 $|t|$ 在 $t = 0$ 处不可微 (见 fig. 1.6a, 教材 Figure 1-4)。

这个例子表明我们的定义排除了”带尖角”的曲线——这正是微分几何研究平滑曲线的自然选择。

(a) Figure 1-4. 不可微曲线 $(t, |t|)$ 

(b) Figure 1-5. 相同圆的不同参数化

例 1.5 (相同迹的不同参数化). 两条不同的参数曲线

$$\alpha(t) = (\cos t, \sin t), \quad \beta(t) = (\cos 2t, \sin 2t),$$

其中 $t \in (-\epsilon, 2\pi + \epsilon)$, $\epsilon > 0$, 具有相同的迹——即单位圆 $x^2 + y^2 = 1$ 。但第二条曲线的速度向量是第一条的两倍 (见 *fig. 1.6b*, 教材 *Figure 1-5*)。

这引出一个关键观察: 曲线的“形状”(迹)与描述它的“方式”(参数化)是不同的概念。这种区别在后续引入弧长参数化时将变得尤为重要。

1-2 节练习

练习 1-2, 1 —do Carmo, Exercise 1-2, 1 Find a parametrized curve $\alpha(t)$ whose trace is the circle $x^2 + y^2 = 1$ such that $\alpha(t)$ runs clockwise around the circle with $\alpha(0) = (0, 1)$.

练习 1-2, 2 —do Carmo, Exercise 1-2, 2 Let $\alpha(t)$ be a parametrized curve which does not pass through the origin. If $\alpha(t_0)$ is a point of the trace of α closest to the origin and $\alpha'(t_0) \neq 0$, show that the position vector $\alpha(t_0)$ is orthogonal to $\alpha'(t_0)$.

练习 1-2, 3 —do Carmo, Exercise 1-2, 3 A parametrized curve $\alpha(t)$ has the property that its second derivative $\alpha''(t)$ is identically zero. What can be said about α ?

练习 1-2, 4 —do Carmo, Exercise 1-2, 4 Let $\alpha : I \rightarrow \mathbb{R}^3$ be a parametrized curve and let $v \in \mathbb{R}^3$ be a fixed vector. Assume that $\alpha'(t)$ is orthogonal to v for all $t \in I$ and that $\alpha(0)$ is also orthogonal to v . Prove that $\alpha(t)$ is orthogonal to v for all $t \in I$.

练习 1-2, 5 —do Carmo, Exercise 1-2, 5 Let $\alpha : I \rightarrow \mathbb{R}^3$ be a parametrized curve, with $\alpha'(t) \neq 0$ for all $t \in I$. Show that $|\alpha(t)|$ is a nonzero constant if and only if $\alpha(t)$ is orthogonal to $\alpha'(t)$ for all $t \in I$.

1.3 Regular Curves; Arc Length

The need for regular curves. 当我们引入参数化曲线的概念后, 自然会问: 是否每个可微参数曲线都能用微分学的工具来研究?

答案并非总是肯定的。我们已经看到，奇异点（如 $\alpha(t) = (t^3, t^2)$ 在 $t = 0$ 处）的出现会导致切向量为零，这会给后续的微分运算带来麻烦。更根本的是：如果 $\alpha'(t) = 0$ ，那么在该点处就不存在确定的切线——这对于研究曲线的“弯曲程度”是致命的。

因此，在研究曲线的微分几何时，我们必须把注意力限制在每一点都有良好定义的切线的曲线上。这引导我们引入正则曲线的概念。

设 $\alpha: I \rightarrow \mathbb{R}^3$ 是一条可微参数曲线。对于每个满足 $\alpha'(t) \neq 0$ 的 $t \in I$ ，存在一条唯一定义直线，它经过点 $\alpha(t)$ 且与向量 $\alpha'(t)$ 平行。这条直线称为曲线 α 在 t 处的**切线** (tangent line)。

定义 1.2 (正则曲线). 如果 $\alpha'(t) \neq 0$ 对所有 $t \in I$ 成立，则称可微参数曲线 $\alpha: I \rightarrow \mathbb{R}^3$ 为**正则曲线** (regular curve)。

Why arc length? 在有了正则曲线的概念之后，我们转向一个更基本的问题：如何测量曲线的长度？


这个问题看似简单，却揭示了微分几何的一个核心思想。直观上，曲线的长度应该是其“内在”属性——它不应该依赖于我们如何参数化这条曲线。果真如此吗？

答案是肯定的，这正是弧长公式的非凡之处。


定义 1.3 (弧长). 设 $\alpha: [a, b] \rightarrow \mathbb{R}^3$ 是一条可微曲线。 α 从 $t = a$ 到 t 的**弧长** (arc length) 定义为

$$s(t) = \int_a^t \|\alpha'(u)\| du.$$


弧长是曲线的内在不变量，与参数化选择无关。



(a) Figure 1-7. 旋轮线 (cycloid)



(a) Figure 1-8. Diocles 割舌线 (cissoid)



(b) Figure 1-9. 曳物线 (tractrix)

命题 1.4 (弧长参数化). 设 $\alpha: I \rightarrow \mathbb{R}^3$ 是正则曲线。定义

$$s(t) = \int_{t_0}^t \|\alpha'(u)\| du,$$

其中 $t_0 \in I$ 是固定点。由于 α 是正则的, $s'(t) = \|\alpha'(t)\| > 0$, 所以 s 是严格递增的, 可以取逆函数 $t(s)$ 。则 $\beta(s) = \alpha(t(s))$ 是单位速度曲线。

定义 1.5 (单位速度曲线). 如果 $\|\alpha'(t)\| = 1$ 对所有 t 成立, 则称 α 为**单位速度曲线** (unit speed curve) 或**弧长参数化曲线**。

从现在开始, 除非特别说明, 我们将使用弧长 s 作为参数, 即 $\alpha(s)$ 是单位速度曲线。

注 1.3. 弧长参数化后, 速度向量恒为 1, 这是最自然的参数化方式。此时 $\alpha'(s)$ 是单位切向量。

例 1.6 (圆的弧长). 半径为 a 的圆可以参数化为 $\alpha(t) = (a \cos t, a \sin t, 0)$, $t \in [0, 2\pi]$ 。则弧长


$$s(t) = \int_0^t \|\alpha'(u)\| du = \int_0^t a du = at,$$

所以从 $t = 0$ 到 $t = 2\pi$ 的全长是 $2\pi a$ 。


例 1.7 (圆柱螺旋线的弧长). 圆柱螺旋线 $\alpha(t) = (a \cos t, a \sin t, bt)$ 的速度向量为 $\alpha'(t) = (-a \sin t, a \cos t, b)$, 模长为

$$\|\alpha'(t)\| = \sqrt{a^2 + b^2}.$$

因此弧长为 $s(t) = \sqrt{a^2 + b^2} t$.



(a) Figure 1-10. 笛卡尔叶形线 (folium of Descartes)



(b) Figure 1-11. 对数螺旋线 (logarithmic spiral)

1-3 节练习

练习 1-3, 1 —do Carmo, Exercise 1-3, 1 Show that the tangent lines to the regular parametrized curve $\alpha(t) = (3t, 3t^2, 2t^3)$ make a constant angle with the line $y = 0, z = x$.

练习 1-3, 2 —do Carmo, Exercise 1-3, 2 A circular disk of radius 1 in the plane xy rolls without slipping along the x axis. The figure described by a point of the circumference of the disk is called a cycloid (见 fig. 1.7a, 教材 Figure 1-7).

- Obtain a parametrized curve $\alpha : \mathbb{R} \rightarrow \mathbb{R}^2$ the trace of which is the cycloid, and determine its singular points.
- Compute the arc length of the cycloid corresponding to a complete rotation of the disk.

练习 1-3, 3 —do Carmo, Exercise 1-3, 3 Let $OA = 2a$ be the diameter of a circle S^1 . Prove that:

- The trace of $\alpha(t) = \left(\frac{2at^2}{1+t^2}, \frac{2at^3}{1+t^2}\right)$, $t \in \mathbb{R}$, is the cissoid of Diocles (见 fig. 1.8a, 教材 Figure 1-8).
- The origin $(0,0)$ is a singular point of the cissoid.
- As $t \rightarrow \infty$, $\alpha(t)$ approaches the line $x = 2a$ (asymptote).

练习 1-3, 4 —do Carmo, Exercise 1-3, 4 Let $\alpha : (0, \pi) \rightarrow \mathbb{R}^2$ be given by $\alpha(t) = (\sin t, \cos t + \log \tan \frac{t}{2})$. The trace of α is called the tractrix (见 fig. 1.8b, 教材 Figure 1-9). Show that:

- a. α is a differentiable parametrized curve, regular except at $t = \pi/2$.
- b. The length of the segment of the tangent of the tractrix between the point of tangency and the y axis is constantly equal to 1.

练习 1-3, 5 —do Carmo, Exercise 1-3, 5 Let $\alpha : (-1, +\infty) \rightarrow \mathbb{R}^2$ be given by

$$\alpha(t) = \left(\frac{3at}{1+t^3}, \frac{3at^2}{1+t^3} \right).$$

Prove that:

- a. For $t = 0$, α is tangent to the x axis.
- b. As $t \rightarrow +\infty$, $\alpha(t) \rightarrow (0, 0)$ and $\alpha'(t) \rightarrow (0, 0)$.
- c. Take the curve with the opposite orientation. Now, as $t \rightarrow -1$, the curve and its tangent approach the line $x + y + a = 0$.

The figure obtained by completing the trace of α in such a way that it becomes symmetric relative to the line $y = x$ is called the **folium of Descartes** (笛卡尔叶形线, 见 fig. 1.9a, 教材 Figure 1-10).

练习 1-3, 6 —do Carmo, Exercise 1-3, 6 Let $\alpha(t) = (ae^{bt} \cos t, ae^{bt} \sin t)$, $t \in \mathbb{R}$, a and b constants, $a > 0$, $b < 0$, be a parametrized curve.

- a. Show that as $t \rightarrow +\infty$, $\alpha(t)$ approaches the origin $\mathbf{0}$, spiraling around it (because of this, the trace of α is called the **logarithmic spiral**, 对数螺旋线, 见 fig. 1.9b, 教材 Figure 1-11).
- b. Show that $\alpha'(t) \rightarrow (0, 0)$ as $t \rightarrow +\infty$ and that

$$\lim_{t \rightarrow +\infty} \int_{t_0}^t \|\alpha'(t)\| dt$$

is finite; that is, α has finite arc length in $[t_0, \infty)$.

练习 1-3, 7 —do Carmo, Exercise 1-3, 7 A map $\alpha : I \rightarrow \mathbb{R}^3$ is called a curve of class C^k if each of the coordinate functions in the expression $\alpha(t) = (x(t), y(t), z(t))$ has continuous derivatives up to order k . If α is merely continuous, we say that α is of class C^0 . A curve α is called **simple** if the map α is one-to-one.

Let $\alpha : I \rightarrow \mathbb{R}^3$ be a simple curve of class C^0 . We say that α has a **weak tangent** at $t = t_0 \in I$ if the line determined by $\alpha(t_0 + h)$ and $\alpha(t_0)$ has a limit position when $h \rightarrow 0$. We say that α has a **strong tangent** at $t = t_0$ if the line determined by $\alpha(t_0 + h)$ and $\alpha(t_0 + k)$ has a limit position when $h, k \rightarrow 0$. Show that

- a. $\alpha(t) = (t^3, t^2)$, $t \in \mathbb{R}$, has a weak tangent but not a strong tangent at $t = 0$.
- b. If $\alpha : I \rightarrow \mathbb{R}^3$ is of class C^1 and regular at $t = t_0$, then it has a strong tangent at $t = t_0$.

c. The curve given by

$$\alpha(t) = \begin{cases} (t^2, t^2), & t \geq 0, \\ (t^2, -t^2), & t \leq 0, \end{cases}$$

is of class C^1 but not of class C^2 . Draw a sketch of the curve and its tangent vectors.

练习 1-3, 8* —do Carmo, Exercise 1-3, 8 Let $\alpha : I \rightarrow \mathbb{R}^3$ be a differentiable curve and let $[a, b] \subset I$ be a closed interval. For every partition

$$a = t_0 < t_1 < \cdots < t_n = b$$

of $[a, b]$, consider the sum $\sum_{i=1}^n |\alpha(t_i) - \alpha(t_{i-1})| = l(\alpha, P)$, where P stands for the given partition. The norm $|P|$ of a partition P is defined as

$$|P| = \max(t_i - t_{i-1}), \quad i = 1, \dots, n.$$

Geometrically, $l(\alpha, P)$ is the length of a polygon inscribed in $\alpha([a, b])$ with vertices in $\alpha(t_i)$. Prove that given $\epsilon > 0$ there exists $\delta > 0$ such that if $|P| < \delta$ then

$$\left| \int_a^b |\alpha'(t)| dt - l(\alpha, P) \right| < \epsilon.$$

练习 1-3, 9 —do Carmo, Exercise 1-3, 9 a. Let $\alpha : I \rightarrow \mathbb{R}^3$ be a curve of class C^0 .

Use the approximation by polygons described in Exercise 8 to give a reasonable definition of arc length of α .

b. (**A Nonrectifiable Curve.**) The following example shows that, with any reasonable definition, the arc length of a C^0 curve in a closed interval may be unbounded. Let $\alpha : [0, 1] \rightarrow \mathbb{R}^2$ be given as $\alpha(t) = (t, t \sin(\pi/t))$ if $t \neq 0$, and $\alpha(0) = (0, 0)$. Show, geometrically, that the arc length of the portion of the curve corresponding to $1/(n+1) \leq t \leq 1/n$ is at least $2/(n+1/2)$. Use this to show that the length of the curve in the interval $1/N \leq t \leq 1$ is greater than $2 \sum_{n=1}^N 1/(n+1)$, and thus it tends to infinity as $N \rightarrow \infty$.

练习 1-3, 10* —do Carmo, Exercise 1-3, 10 (Straight Lines as Shortest.) Let $\alpha : I \rightarrow \mathbb{R}^3$ be a parametrized curve. Let $[a, b] \subset I$ and set $\alpha(a) = p$, $\alpha(b) = q$.

a. Show that, for any constant vector v , $|v| = 1$,

$$(q - p) \cdot v = \int_a^b \alpha'(t) \cdot v dt \leq \int_a^b |\alpha'(t)| dt.$$

b. Set

$$v = \frac{q - p}{|q - p|}$$

and show that

$$|\alpha(b) - \alpha(a)| \leq \int_a^b |\alpha'(t)| dt;$$

that is, the curve of shortest length from $\alpha(a)$ to $\alpha(b)$ is the straight line joining these points.

1.4 The Vector Product in $\mathbb{R}^3$

A new operation unique to three dimensions. 在我们研究 $\mathbb{R}^3$ 中的曲线时，内积已经给了我们测量长度和角度的工具。然而，仅仅有内积是不够的。考虑一个问题：给定曲线的切向量 $\alpha'(t)$ 和二阶导数 $\alpha''(t)$ ，我们如何构造一个与两者都垂直的向量？这个向量对于建立曲线的局部标架（如后续将引入的 Frenet 标架）至关重要。

在 $\mathbb{R}^2$ 中，这是不可能的——两个不平行的向量张成整个平面，不存在第三个同时垂直于两者的非零向量。但在 $\mathbb{R}^3$ 中，一个新的运算应运而生：向量积。

向量积（或叉积、外积）是 $\mathbb{R}^3$ 中特有的运算，它的引入不是人为的，而是由几何需求驱动的。我们将会看到，向量积在曲线与曲面的理论中起着关键作用。

定义 1.6 (向量积). 设 $\mathbf{u} = (u_1, u_2, u_3)$ 和 $\mathbf{v} = (v_1, v_2, v_3)$ 是 $\mathbb{R}^3$ 中的向量。定义**向量积** (cross product) $\mathbf{u} \times \mathbf{v} \in \mathbb{R}^3$ 为

$$\mathbf{u} \times \mathbf{v} = (u_2v_3 - u_3v_2, u_3v_1 - u_1v_3, u_1v_2 - u_2v_1).$$

向量积可以用行列式简洁地表示为

$$\mathbf{u} \times \mathbf{v} = \det \begin{pmatrix} \mathbf{e}_1 & \mathbf{e}_2 & \mathbf{e}_3 \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{pmatrix}.$$

向量积的基本性质：


命题 1.7 (向量积的性质). 设 $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathbb{R}^3$, $\lambda \in \mathbb{R}$, 则：

1. (反交换律) $\mathbf{u} \times \mathbf{v} = -\mathbf{v} \times \mathbf{u}$;
2. (数乘结合律) $(\lambda \mathbf{u}) \times \mathbf{v} = \lambda(\mathbf{u} \times \mathbf{v}) = \mathbf{u} \times (\lambda \mathbf{v})$;
3. (分配律) $\mathbf{u} \times (\mathbf{v} + \mathbf{w}) = \mathbf{u} \times \mathbf{v} + \mathbf{u} \times \mathbf{w}$;
4. (标量三重积) $(\mathbf{u} \times \mathbf{v}) \cdot \mathbf{w} = \det(\mathbf{u}, \mathbf{v}, \mathbf{w})$ 。

向量积的几何意义：

命题 1.8 (向量积的几何意义). 1. $\mathbf{u} \times \mathbf{v}$ 垂直于 $\mathbf{u}$ 和 $\mathbf{v}$ 所在的平面；


2. $\mathbf{u} \times \mathbf{v}$ 的方向由右手定则确定；
3. $\|\mathbf{u} \times \mathbf{v}\| = \|\mathbf{u}\| \|\mathbf{v}\| \sin \theta$, 其中 θ 是 $\mathbf{u}$ 和 $\mathbf{v}$ 之间的夹角；
4. $\mathbf{u} \times \mathbf{v} = \mathbf{0}$ 当且仅当 $\mathbf{u}$ 和 $\mathbf{v}$ 平行。



(a) Figure 1-13. 向量积几何意义

注 1.4. 向量积在曲线理论中的应用:

- 构造副法向量: $\mathbf{b} = \mathbf{t} \times \mathbf{n}$
- 计算挠率: $\tau = \frac{\det(\alpha', \alpha'', \alpha''')}{k^2}$



(a) Figure 1-12. 弧长逼近的几何意义

1-4 节练习

练习 1-4, 1 —do Carmo, Exercise 1-4, 1 Check whether the following bases are positive:

- The basis $\{(1, 3), (4, 2)\}$ in $\mathbb{R}^2$.
- The basis $\{(1, 3, 5), (2, 3, 7), (4, 8, 3)\}$ in $\mathbb{R}^3$.

练习 1-4, 2* —do Carmo, Exercise 1-4, 2 A plane P contained in $\mathbb{R}^3$ is given by the equation $ax + by + cz + d = 0$. Show that the vector $v = (a, b, c)$ is perpendicular to the plane and that $|d|/\sqrt{a^2 + b^2 + c^2}$ measures the distance from the plane to the origin $(0, 0, 0)$.

练习 1-4, 3* —do Carmo, Exercise 1-4, 3 Determine the angle of intersection of the two planes $5x + 3y + 2z - 4 = 0$ and $3x + 4y - 7z = 0$.

练习 1-4, 4* —do Carmo, Exercise 1-4, 4 Given two planes $a_ix + b_iy + c_iz + d_i = 0$, $i = 1, 2$, prove that a necessary and sufficient condition for them to be parallel is

$$\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2},$$

where the convention is made that if a denominator is zero, the corresponding numerator is also zero.

练习 1-4, 5 —do Carmo, Exercise 1-4, 5 Show that an equation of a plane passing through three noncollinear points $p_1 = (x_1, y_1, z_1)$, $p_2 = (x_2, y_2, z_2)$, $p_3 = (x_3, y_3, z_3)$ is given by

$$(p - p_1) \wedge (p - p_2) \cdot (p - p_3) = 0,$$

where $p = (x, y, z)$ is an arbitrary point of the plane.

练习 1-4, 6* —do Carmo, Exercise 1-4, 6 Given two nonparallel planes $a_ix + b_iy + c_iz + d_i = 0$, $i = 1, 2$, show that their line of intersection may be parametrized as

$$x - x_0 = u_1t, \quad y - y_0 = u_2t, \quad z - z_0 = u_3t,$$

where (x_0, y_0, z_0) belongs to the intersection and $u = (u_1, u_2, u_3)$ is the vector product $u = v_1 \wedge v_2$, $v_i = (a_i, b_i, c_i)$, $i = 1, 2$.

练习 1-4, 7* —do Carmo, Exercise 1-4, 7 Prove that a necessary and sufficient condition for the plane $ax + by + cz + d = 0$ and the line $x - x_0 = u_1t$, $y - y_0 = u_2t$, $z - z_0 = u_3t$ to be parallel is $au_1 + bu_2 + cu_3 = 0$.

练习 1-4, 8* —do Carmo, Exercise 1-4, 8 Prove that the distance ρ between the non-parallel lines

$$\begin{aligned} x - x_0 &= u_1t, & y - y_0 &= u_2t, & z - z_0 &= u_3t, \\ x - x_1 &= v_1t, & y - y_1 &= v_2t, & z - z_1 &= v_3t \end{aligned}$$

is given by

$$\rho = \frac{|(u \wedge v) \cdot r|}{|u \wedge v|},$$

where $u = (u_1, u_2, u_3)$, $v = (v_1, v_2, v_3)$, $r = (x_0 - x_1, y_0 - y_1, z_0 - z_1)$.

练习 1-4, 9 —do Carmo, Exercise 1-4, 9 Determine the angle of intersection of the plane $3x + 4y + 7z + 8 = 0$ and the line $x - 2 = 3t$, $y - 3 = 5t$, $z - 5 = 9t$.

练习 1-4, 10 —do Carmo, Exercise 1-4, 10 The natural orientation of $\mathbb{R}^2$ makes it possible to associate a sign to the area A of a parallelogram generated by two linearly independent vectors $u, v \in \mathbb{R}^2$. Show that $A^2 = \begin{vmatrix} u_1 & u_2 \\ v_1 & v_2 \end{vmatrix}^2$.

练习 1-4, 11 —do Carmo, Exercise 1-4, 11 a. Show that the volume V of a parallelepiped generated by three linearly independent vectors $u, v, w \in \mathbb{R}^3$ is given by $V = |(u \wedge v) \cdot w|$, and introduce an oriented volume in $\mathbb{R}^3$.

b. Prove that

$$V^2 = \begin{vmatrix} u \cdot u & u \cdot v & u \cdot w \\ v \cdot u & v \cdot v & v \cdot w \\ w \cdot u & w \cdot v & w \cdot w \end{vmatrix}.$$

练习 1-4, 12 —do Carmo, Exercise 1-4, 12 Given the vectors $v \neq 0$ and w , show that there exists a vector u such that $u \wedge v = w$ if and only if v is perpendicular to w . Is this vector u uniquely determined? If not, what is the most general solution?

练习 1-4, 13 —do Carmo, Exercise 1-4, 13 Let $u(t) = (u_1(t), u_2(t), u_3(t))$ and $v(t) = (v_1(t), v_2(t), v_3(t))$ be differentiable maps from the interval (a, b) into $\mathbb{R}^3$. If the derivatives $u'(t)$ and $v'(t)$ satisfy the conditions

$$u'(t) = au(t) + bv(t), \quad v'(t) = cu(t) - av(t),$$

where a, b , and c are constants, show that $u(t) \wedge v(t)$ is a constant vector.

练习 1-4, 14 —do Carmo, Exercise 1-4, 14 Find all unit vectors which are perpendicular to the vector $(2, 2, 1)$ and parallel to the plane determined by the points $(0, 0, 0)$, $(1, -2, 1)$, $(-1, 1, 1)$.

1.5 The Local Theory of Curves Parametrized by Arc Length

Answering our original question. 在引言中，我们提出了一个看似简单却贯穿整个微分几何发展的问题：如何用精确的数学语言来描述曲线的弯曲程度？

现在，我们已经有了足够的工具来回答这个问题。回想一下，对于单位速度曲线 $\alpha(s)$ ，速度向量 $\alpha'(s)$ 是单位切向量。那么，加速度向量 $\alpha''(s)$ 反映了什么？

几何直觉告诉我们：如果曲线是“直的”，那么沿曲线匀速运动的质点没有加速度；如果曲线是“弯曲的”，质点必须承受加速度以改变运动方向。因此， $\alpha''(s)$ 的大小自然就度量了曲线的弯曲程度。这引导我们引入曲率的概念。


1.5.1 The Curvature

定义 1.9 (曲率). 设 $\alpha(s)$ 是单位速度曲线。定义**曲率** (*curvature*) 为

$$k(s) = \|\alpha''(s)\|.$$

即加速度向量的模长。

曲率捕捉了曲线弯曲程度的本质。直线的曲率为零——它完全不弯曲。圆的曲率是恒定的 $1/a$ (a 为半径)——圆周上每一点的弯曲程度都是均匀的。曲率越大, 曲线弯曲得越厉害。



(a) Figure 1-14. 曲率的几何意义

例 1.8 (圆的曲率). 单位圆的参数方程 $\alpha(s) = (\cos s, \sin s, 0)$, 则

$$\alpha'(s) = (-\sin s, \cos s, 0), \quad \alpha''(s) = (-\cos s, -\sin s, 0),$$

所以 $k(s) = \|\alpha''(s)\| = 1$ 。更一般地, 半径为 a 的圆的曲率为 $1/a$ 。

例 1.9 (直线的曲率). 直线 $\alpha(s) = \mathbf{p} + \mathbf{v}s$ ($\mathbf{v}$ 为单位向量), 则 $\alpha''(s) = 0$, 所以 $k(s) = 0$ 。

定义 1.10 (主法向量). 设 $\alpha(s)$ 是单位速度曲线, $k(s) > 0$ 。定义主法向量 (*principal normal*) 为

$$\mathbf{n}(s) = \frac{\alpha''(s)}{k(s)} = \frac{\alpha''(s)}{\|\alpha''(s)\|}.$$

命题 1.11 (曲率的另一个表达式). 设 $\alpha(t)$ 是正则曲线 (不一定单位速度), 则

$$k(t) = \frac{\|\alpha'(t) \times \alpha''(t)\|}{\|\alpha'(t)\|^3}.$$

1.5.2 The Torsion (Or Bending)

Beyond curvature. 曲率已经告诉我们曲线在每一点如何弯曲。然而, 这还不够。考虑一个关键问题: 给定一条曲线, 我们可以找到包含它的最小平面 (密切平面)。但如果曲线“扭曲”出这个平面呢?

这种“扭曲”的程度由挠率度量。直观上, 如果一条曲线始终位于某个平面内 (如圆、椭圆等平面曲线), 它就没有扭曲, 挠率为零。但如果曲线像螺旋线那样在空间中盘旋, 它就会持续“扭转”出每个密切平面——这就是挠率所捕捉的。

挠率描述曲线在空间中的扭曲程度: 平面曲线的挠率恒为零 (因为 $\alpha', \alpha'', \alpha'''$ 共面), 而空间曲线 (如螺旋线) 的挠率可能非零。

定义 1.12 (挠率). 设 $\alpha(s)$ 是单位速度曲线, 假设 $k(s) > 0$. 定义**挠率** (*torsion*) 为

$$\tau(s) = \frac{\det(\alpha'(s), \alpha''(s), \alpha'''(s))}{k(s)^2}.$$

注 1.5. 挠率描述曲线在空间中的扭曲程度:


- 平面曲线: 挠率恒为零 (因为 $\alpha', \alpha'', \alpha'''$ 共面);
- 空间曲线: 挠率可能非零, 描述曲线"扭转"的程度。

例 1.10 (圆柱螺旋线的挠率). 圆柱螺旋线 $\alpha(t) = (a \cos t, a \sin t, bt)$ 的曲率为 $k = \frac{a}{a^2 + b^2}$, 挠率为 $\tau = \frac{b}{a^2 + b^2}$, 两者均为常数。

命题 1.13 (挠率的另一个表达式). 设 $\alpha(s)$ 是单位速度曲线, $k(s) > 0$, 则

$$\tau(s) = -\langle \mathbf{b}'(s), \mathbf{n}(s) \rangle,$$

其中 $\mathbf{b}(s) = \mathbf{t}(s) \times \mathbf{n}(s)$ 是副法向量。



(a) Figure 1-15. 密切平面与 Frenet 标架

1.5.3 The Frenet-Serret Formulas


The fundamental theorem. 给定一个点上的曲率和挠率，我们就知道了曲线在该点的所有几何信息。Frenet-Serret 公式揭示了一个更深层的事实：曲率和挠率不仅决定了曲线的基本性质，而且给定作为弧长函数的光滑曲率 $k(s) > 0$ 和挠率 $\tau(s)$ ，存在一条（局部意义下的）唯一曲线以之为曲率和挠率。这就是曲线论的基本定理。

这个定理的核心工具是 Frenet 标架——一个随曲线移动的正交标架。


定义 1.14 (Frenet 标架). 设 $\alpha(s)$ 是单位速度曲线， $k(s) > 0$ 。定义：

1. 切向量 (*tangent vector*): $\mathbf{t}(s) = \alpha'(s)$;
2. 主法向量 (*principal normal*): $\mathbf{n}(s) = \frac{\alpha''(s)}{k(s)}$;
3. 副法向量 (*binormal*): $\mathbf{b}(s) = \mathbf{t}(s) \times \mathbf{n}(s)$ 。

$\{\mathbf{t}, \mathbf{n}, \mathbf{b}\}$ 称为 **Frenet 标架** (*Frenet frame*)。它们构成 $\mathbb{R}^3$ 的标准正交基 (右手系)。



(a) Figure 1-16 (a). 有向曲率 - 正向



(b) Figure 1-16 (b). 有向曲率 - 负向

定义 1.15 (密切平面、法平面、化直平面). • **密切平面** (*osculating plane*): 由 $\mathbf{t}(s)$ 和 $\mathbf{n}(s)$ 张成的平面;

• **法平面** (*normal plane*): 由 $\mathbf{n}(s)$ 和 $\mathbf{b}(s)$ 张成的平面;

• **化直平面** (*rectifying plane*): 由 $\mathbf{t}(s)$ 和 $\mathbf{b}(s)$ 张成的平面。

定理 1.16 (Frenet-Serret 公式). 设 $\alpha(s)$ 是单位速度曲线， $k(s) > 0$ ，挠率为 $\tau(s)$ 。则

$$\begin{cases} \mathbf{t}' = k\mathbf{n}, \\ \mathbf{n}' = -k\mathbf{t} + \tau\mathbf{b}, \\ \mathbf{b}' = -\tau\mathbf{n}. \end{cases}$$


推论 1.17 (曲率和挠率的计算). 曲率和挠率可以通过以下公式计算:

$$k = \|\mathbf{t}'\|, \quad \tau = -\langle \mathbf{b}', \mathbf{n} \rangle.$$

注 1.6. *Frenet-Serret* 公式表明, 如果我们知道曲率 $k(s)$ 和挠率 $\tau(s)$, 以及初始的 *Frenet* 标架, 就可以唯一确定曲线的形状。这两个函数完全决定了曲线 (在局部意义上)。这类似于刚体运动学: 给定线速度和角速度, 就能确定刚体的运动。

定理 1.18 (曲线存在唯一性定理). 给定两个连续函数 $k(s) > 0$ 和 $\tau(s)$, 以及 $\mathbb{R}^3$ 中一组正交标架, 存在唯一一条单位速度曲线 $\alpha(s)$, 使得其曲率为 $k(s)$ 、挠率为 $\tau(s)$, 且 *Frenet* 标架为给定的标架。

注 1.7. 这是微分几何中的一个基本存在唯一性定理。它告诉我们: 曲率和挠率完全决定了曲线的形状。



(a) Figure 1-17. 渐伸线 (evolute)

1-5 节练习

除非特别说明, $\alpha: \mathbf{I} \rightarrow \mathbf{R}^3$ 是单位速度曲线, 曲率 $k(s) \neq 0$ 。

练习 1-5, 1 —do Carmo, Exercise 1-5, 1 Given the parametrized curve (helix)

$$\alpha(s) = \left(a \cos \frac{s}{c}, a \sin \frac{s}{c}, b \frac{s}{c} \right), \quad s \in \mathbb{R},$$

where $c^2 = a^2 + b^2$,

a. Show that the parameter s is the arc length.

- b. Determine the curvature and the torsion of α .
- c. Determine the osculating plane of α .
- d. Show that the lines containing $n(s)$ and passing through $\alpha(s)$ meet the z axis under a constant angle equal to $\pi/2$.
- e. Show that the tangent lines to α make a constant angle with the z axis.

练习 1-5, 2* —do Carmo, Exercise 1-5, 2 Show that the torsion τ of α is given by

$$\tau(s) = -\frac{\alpha'(s) \wedge \alpha''(s) \cdot \alpha'''(s)}{|k(s)|^2}.$$

练习 1-5, 3 —do Carmo, Exercise 1-5, 3 Assume that $\alpha(I) \subset \mathbb{R}^2$ (i.e., α is a plane curve) and give k a sign. Transport the vectors $t(s)$ parallel to themselves so that the origins of $t(s)$ agree with the origin of $\mathbb{R}^2$; the end points of $t(s)$ then describe a parametrized curve $s \rightarrow t(s)$ called the indicatrix of tangents of α . Let $\theta(s)$ be the angle from e_1 to $t(s)$ in the orientation of $\mathbb{R}^2$. Prove:

- a. The indicatrix of tangents is a regular parametrized curve.
- b. $dt/ds = (d\theta/ds)n$, that is, $k = d\theta/ds$.

练习 1-5, 4* —do Carmo, Exercise 1-5, 4 Assume that all normals of a parametrized curve pass through a fixed point. Prove that the trace of the curve is contained in a circle.

练习 1-5, 5 —do Carmo, Exercise 1-5, 5 A regular parametrized curve α has the property that all its tangent lines pass through a fixed point.

- a. Prove that the trace of α is a (segment of a) straight line.
- b. Does the conclusion in part a still hold if α is not regular?

练习 1-5, 6 —do Carmo, Exercise 1-5, 6 A translation by a vector v in $\mathbb{R}^3$ is the map $A : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ that is given by $A(p) = p + v$, $p \in \mathbb{R}^3$. A linear map $\rho : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is an orthogonal transformation when $\rho u \cdot \rho v = u \cdot v$ for all vectors $u, v \in \mathbb{R}^3$. A rigid motion in $\mathbb{R}^3$ is the result of composing a translation with an orthogonal transformation with positive determinant.

- a. Demonstrate that the norm of a vector and the angle θ between two vectors, $0 \leq \theta \leq \pi$, are invariant under orthogonal transformations with positive determinant.
- b. Show that the vector product of two vectors is invariant under orthogonal transformations with positive determinant. Is the assertion still true if we drop the condition on the determinant?
- c. Show that the arc length, the curvature, and the torsion of a parametrized curve are (whenever defined) invariant under rigid motions.

练习 1-5, 7* —do Carmo, Exercise 1-5, 7 Let $\alpha : I \rightarrow \mathbb{R}^2$ be a regular parametrized plane curve (arbitrary parameter), and define $n = n(t)$ and $k = k(t)$. Assume that $k(t) \neq 0$, $t \in I$. In this situation, the curve

$$\beta(t) = \alpha(t) + \frac{1}{k(t)}n(t), \quad t \in I,$$

is called the **evolute** of α (渐伸线, 见 fig. 1.15a, 教材 Figure 1-17).

- Show that the tangent at t of the evolute of α is the normal to α at t .
- Consider the normal lines of α at two neighboring points t_1, t_2 , $t_1 \neq t_2$. Let t_1 approach t_2 and show that the intersection points of the normals converge to a point on the trace of the evolute of α .

练习 1-5, 8 —do Carmo, Exercise 1-5, 8 The trace of the parametrized curve (arbitrary parameter)

$$\alpha(t) = (t, \cosh t), \quad t \in \mathbb{R},$$

is called the **catenary** (悬链线).

- Show that the signed curvature of the catenary is

$$k(t) = \frac{1}{\cosh^2 t}.$$

- Show that the evolute of the catenary is

$$\beta(t) = (t - \sinh t \cosh t, 2 \cosh t).$$

练习 1-5, 9 —do Carmo, Exercise 1-5, 9 Given a differentiable function $k(s)$, $s \in I$, show that the parametrized plane curve having $k(s) = k$ as curvature is given by

$$\alpha(s) = \left(\int \cos \theta(s) ds + a, \int \sin \theta(s) ds + b \right),$$

where

$$\theta(s) = \int k(s) ds + \varphi,$$

and that the curve is determined up to a translation of the vector (a, b) and a rotation of the angle φ .

练习 1-5, 10 —do Carmo, Exercise 1-5, 10 Consider the map

$$\alpha(t) = \begin{cases} (t, 0, e^{-1/t^2}), & t > 0, \\ (t, e^{-1/t^2}, 0), & t < 0, \\ (0, 0, 0), & t = 0. \end{cases}$$

- Prove that α is a differentiable curve.

- b. Prove that α is regular for all t and that the curvature $k(t) \neq 0$, for $t \neq 0, t \neq \pm\sqrt{2/3}$, and $k(0) = 0$.
- c. Show that the limit of the osculating planes as $t \rightarrow 0, t > 0$, is the plane $y = 0$ but that the limit of the osculating planes as $t \rightarrow 0, t < 0$, is the plane $z = 0$ (this implies that the normal vector is discontinuous at $t = 0$ and shows why we excluded points where $k = 0$).
- d. Show that τ can be defined so that $\tau \equiv 0$, even though α is not a plane curve.

练习 1-5, 11 —do Carmo, Exercise 1-5, 11 One often gives a plane curve in polar coordinates by $\rho = \rho(\theta)$, $a \leq \theta \leq b$.

- a. Show that the arc length is

$$\int_a^b \sqrt{\rho^2 + (\rho')^2} d\theta,$$

where the prime denotes the derivative relative to θ .

- b. Show that the curvature is

$$k(\theta) = \frac{2(\rho')^2 - \rho\rho'' + \rho^2}{\{((\rho')^2 + \rho^2)^{3/2}\}}.$$

练习 1-5, 12* —do Carmo, Exercise 1-5, 12 Let $\alpha : I \rightarrow \mathbb{R}^3$ be a regular parametrized curve (not necessarily by arc length) and let $\beta : J \rightarrow \mathbb{R}^3$ be a reparametrization of $\alpha(I)$ by the arc length $s = s(t)$, measured from $t_0 \in I$. Let $t = t(s)$ be the inverse function of s and set $d\alpha/dt = \alpha'$, $d^2\alpha/dt^2 = \alpha''$, etc. Prove:

- a. $dt/ds = 1/|\alpha'|$, $d^2t/ds^2 = -(\alpha' \cdot \alpha''/|\alpha'|^4)$.

- b. The curvature of α at $t \in I$ is

$$k(t) = \frac{|\alpha' \wedge \alpha''|}{|\alpha'|^3}.$$

- c. The torsion of α at $t \in I$ is

$$\tau(t) = -\frac{(\alpha' \wedge \alpha'') \cdot \alpha'''}{|\alpha' \wedge \alpha''|^2}.$$

- d. If $\alpha : I \rightarrow \mathbb{R}^2$ is a plane curve $\alpha(t) = (x(t), y(t))$, the signed curvature of α at t is

$$k(t) = \frac{x'y'' - x''y'}{((x')^2 + (y')^2)^{3/2}}.$$

练习 1-5, 13* —do Carmo, Exercise 1-5, 13 Assume that $\tau(s) \neq 0$ and $k'(s) \neq 0$ for all $s \in I$. Show that a necessary and sufficient condition for $\alpha(I)$ to lie on a sphere is that

$$R^2 + (R')^2 T^2 = \text{const.},$$

where $R = 1/k$, $T = 1/\tau$, and R' is the derivative of R relative to s .

练习 1-5, 14 —do Carmo, Exercise 1-5, 14 Let $\alpha : (a, b) \rightarrow \mathbb{R}^2$ be a regular parametrized plane curve. Assume that there exists t_0 , $a < t_0 < b$, such that the distance $|\alpha(t)|$ from the origin to the trace of α will be a maximum at t_0 . Prove that the curvature k of α at t_0 satisfies $|k(t_0)| \geq 1/|\alpha(t_0)|$.

练习 1-5, 15* —do Carmo, Exercise 1-5, 15 Show that the knowledge of the vector function $b = b(s)$ (binormal vector) of a curve α , with nonzero torsion everywhere, determines the curvature $k(s)$ and the absolute value of the torsion $\tau(s)$ of α .

练习 1-5, 16* —do Carmo, Exercise 1-5, 16 Show that the knowledge of the vector function $n = n(s)$ (normal vector) of a curve α , with nonzero torsion everywhere, determines the curvature $k(s)$ and the torsion $\tau(s)$ of α .

练习 1-5, 17 —do Carmo, Exercise 1-5, 17 In general, a curve α is called a **helix** if the tangent lines of α make a constant angle with a fixed direction. Assume that $\tau(s) \neq 0$, $s \in I$, and prove that:

- a. α is a helix if and only if $k/\tau = \text{const.}$
- b.* α is a helix if and only if the lines containing $n(s)$ and passing through $\alpha(s)$ are parallel to a fixed plane.
- c.* α is a helix if and only if the lines containing $b(s)$ and passing through $\alpha(s)$ make a constant angle with a fixed direction.
- d. The curve

$$\alpha(s) = \left(\frac{a}{c} \int \sin \theta(s) ds, \frac{a}{c} \int \cos \theta(s) ds, \frac{b}{c} s \right),$$

where $c^2 = a^2 + b^2$, is a helix, and that $k/\tau = a/b$.

练习 1-5, 18* —do Carmo, Exercise 1-5, 18 Let $\alpha : I \rightarrow \mathbb{R}^3$ be a parametrized regular curve (not necessarily by arc length) with $k(t) \neq 0$, $\tau(t) \neq 0$, $t \in I$. The curve α is called a **Bertrand curve** if there exists a curve $\bar{\alpha} : I \rightarrow \mathbb{R}^3$ such that the normal lines of α and $\bar{\alpha}$ at $t \in I$ are equal. In this case, $\bar{\alpha}$ is called a **Bertrand mate** of α , and we can write

$$\bar{\alpha}(t) = \alpha(t) + rn(t).$$

Prove that:

- a. r is constant.
- b. α is a Bertrand curve if and only if there exists a linear relation

$$Ak(t) + B\tau(t) = 1, \quad t \in I,$$

where A, B are nonzero constants and k and τ are the curvature and torsion of α , respectively.

- c. If α has more than one Bertrand mate, it has infinitely many Bertrand mates. This case occurs if and only if α is a circular helix.

1.6 The Local Canonical Form

本节研究曲线在一点附近的局部形状。

定理 1.19 (曲线在一点的局部标准形式). 设 $\alpha(s)$ 是以弧长 s 为参数的可微曲线, 在 $s = s_0$ 处 $k(s_0) > 0$ 。则

$$\alpha(s) = \alpha(s_0) + (s - s_0)\mathbf{t}_0 + \frac{(s - s_0)^2}{2}k_0\mathbf{n}_0 + \frac{(s - s_0)^3}{6}k_0\tau_0\mathbf{b}_0 + o((s - s_0)^3),$$

其中 $\mathbf{t}_0 = \mathbf{t}(s_0)$, $k_0 = k(s_0)$, $\tau_0 = \tau(s_0)$ 。


注 1.8. 这个展开式揭示了曲线的几何意义:

- 线性项 $(s - s_0)\mathbf{t}_0$: 沿切方向 $\mathbf{t}$ 的运动;
- 二次项 $\frac{(s - s_0)^2}{2}k_0\mathbf{n}_0$: 沿法方向 $\mathbf{n}$ 的弯曲 (由曲率 k 控制);
- 三次项 $\frac{(s - s_0)^3}{6}k_0\tau_0\mathbf{b}_0$: 沿副法方向 $\mathbf{b}$ 的扭曲 (由挠率 τ 控制)。


注 1.9. 当 $\tau(s_0) = 0$ 但 $k(s_0) \neq 0$ 时, 曲线在 s_0 附近是平面曲线 (因为三次项为零)。

定义 1.20 (密切圆). 曲线在 s_0 处的**密切圆** (*osculating circle*) 是半径为 $1/k(s_0)$ 、圆心在 $\alpha(s_0) + \frac{1}{k(s_0)}\mathbf{n}(s_0)$ 的圆。

注 1.10. 密切圆与曲线在切点有相同的切方向和曲率。它是曲线在该点附近最好的圆近似。






(a) Figure 1-18. 局部标准形式的投影




Projection over the plane tn

(b) 投影到 tn 平面

(a) 投影到 nb 平面(b) 投影到 tn 平面

Negative torsion

(a) Figure 1-19 (a). $\tau < 0$ 的情形

Positive torsion

(b) Figure 1-19 (b). $\tau > 0$ 的情形

1-6 节练习

练习 1-6, 1 —do Carmo, Exercise 1-6, 1 Let $\alpha : I \rightarrow \mathbb{R}^3$ be a curve parametrized by arc length with curvature $k(s) \neq 0$, $s \in I$. Let P be a plane satisfying both of the following conditions:

1. P contains the tangent line at s .
2. Given any neighborhood $J \subset I$ of s , there exist points of $\alpha(J)$ in both sides of P .

Prove that P is the osculating plane of α at s .

练习 1-6, 2 —do Carmo, Exercise 1-6, 2 Let $\alpha : I \rightarrow \mathbb{R}^3$ be a curve parametrized by arc length, with curvature $k(s) \neq 0$, $s \in I$. Show that:


- The osculating plane at s is the limit position of the plane passing through $\alpha(s)$, $\alpha(s+h_1)$, $\alpha(s+h_2)$ when $h_1, h_2 \rightarrow 0$.
- The limit position of the circle passing through $\alpha(s)$, $\alpha(s+h_1)$, $\alpha(s+h_2)$ when $h_1, h_2 \rightarrow 0$ is a circle in the osculating plane at s , the center of which is on the line that contains $n(s)$ and the radius of which is the radius of curvature $1/k(s)$; this circle is called the osculating circle at s .

练习 1-6, 3 —do Carmo, Exercise 1-6, 3 Show that the curvature $k(t) \neq 0$ of a regular parametrized curve $\alpha : I \rightarrow \mathbb{R}^3$ is the curvature at t of the plane curve $\pi \circ \alpha$, where π is the normal projection of α over the osculating plane at t .


1.7 Global Properties of Plane Curves

前面我们讨论的是曲线的局部性质。现在我们讨论平面曲线的整体（全局）性质。


定义 1.21 (简单闭曲线). 一条平面曲线 $\alpha : [0, L] \rightarrow \mathbb{R}^2$ 称为**简单闭曲线** (*simple closed curve*), 如果 $\alpha(0) = \alpha(L)$, 且 α 在 $(0, L)$ 上是单射。




(a) Figure 1-20 (a). 简单闭曲线



(b) Figure 1-20 (b). 非简单闭曲线



(a) Figure 1-21 (a). 环面上的简单闭曲线



(b) Figure 1-21 (b). 正向定向

定义 1.22 (旋转指数). 设 $\alpha : [0, L] \rightarrow \mathbb{R}^2$ 是一条简单闭平面曲线, 参数化为单位速度。定义**旋转指数** (*turning number*) 为

$$n = \frac{1}{2\pi} \int_0^L k(s) ds.$$

旋转指数表示切向量沿曲线完整一圈转过的圈数。

定理 1.23 (旋转指数定理). 简单闭平面曲线的旋转指数必为整数, 且绝对值至少为 1。

定理 1.24 (四顶点定理). 设 C 是平面上一条简单闭曲线, 曲率恒为正。则 C 上至少有四个曲率极值点 (顶点)。

注 1.11. "顶点" (*vertex*) 是指曲率取得局部极值的点。四顶点定理是平面曲线理论中最著名的整体定理之一。例如, 椭圆有四个顶点 (长轴和短轴的端点)。

定理 1.25 (Fenchel 定理). 设 $\alpha : [0, L] \rightarrow \mathbb{R}^3$ 是一条简单闭曲线。则其全曲率

$$\int_0^L k(s) ds \geq 2\pi,$$

等号成立当且仅当曲线是平面上的凸曲线。

定义 1.26 (全曲率). 曲线 α 的**全曲率** (*total curvature*) 定义为


$$\int_0^L k(s) ds,$$

即曲率沿曲线的积分。


推论 1.27. 任何简单闭曲线的全曲率至少为 2π 。

定理 1.28 (Fary-Milnor 定理). 如果 α 是一条简单闭曲线的全曲率小于 4π , 则该曲线是未打结的 (*unknotted*)。


注 1.12. 这些定理连接了曲线的局部几何 (曲率) 和全局拓扑性质。四顶点定理说明闭合曲线的“总弯曲量”必须“平衡”; *Fenchel* 定理说明闭合曲线的全曲率至少是 2π 。





(a) Figure 1-22. 等周公式证明图示




(a) Figure 1-23. 等周问题的几何构型




(b) Figure 1-24. 等周不等式证明图示

(a) Figure 1-25. 分段 C^1 曲线


(a) Figure 1-26 (a). 切线指标线




(b) Figure 1-26 (b). 切线指标线详图



(a) Figure 1-27 (a). 旋转指数为 1




(b) Figure 1-27 (b). 旋转指数为 2




(a) Figure 1-27 (c). 旋转指数为 -1




(a) Figure 1-28 (a). 凸曲线




(a) Figure 1-28 (b). 非凸曲线




(a) Figure 1-29 (a). 三个不同顶点情况




(a) Figure 1-29 (b). 直线段情况



(a) Figure 1-30. 直线族的重数



(a) Figure 1-31. 直线的参数化



(a) Figure 1-32. 刚体运动

1-7 节练习

练习 1-7, 1* —do Carmo, Exercise 1-7, 1 Is there a simple closed curve in the plane with length equal to 6 feet and bounding an area of 3 square feet?

练习 1-7, 2* —do Carmo, Exercise 1-7, 2 Let $\overline{AB}$ be a segment of straight line and let $l > \text{length of } \overline{AB}$. Show that the curve C joining A and B , with length l , and such that together with $\overline{AB}$ bounds the largest possible area is an arc of a circle passing through A and B (见 fig. 1.22a, fig. 1.22b, 教材 Figure 1-23, 1-24)。

练习 1-7, 3 —do Carmo, Exercise 1-7, 3 Compute the curvature of the ellipse $x = a \cos t$, $y = b \sin t$, $t \in [0, 2\pi]$, $a \neq b$, and show that it has exactly four vertices, namely, $(a, 0)$, $(-a, 0)$, $(0, b)$, $(0, -b)$.

练习 1-7, 4* —do Carmo, Exercise 1-7, 4 Let C be a plane curve and let T be the tangent line at a point $p \in C$. Draw a line L parallel to the normal line at p and at a distance d of p . Let h be the length of the segment determined on L by C and T . Prove that $|k(p)| = \lim_{d \rightarrow 0} \frac{2h}{d^2}$.

练习 1-7, 5* —do Carmo, Exercise 1-7, 5 If a closed plane curve C is contained inside a disk of radius r , prove that there exists a point $p \in C$ such that the curvature k of C at p satisfies $|k| \geq 1/r$.

练习 1-7, 6 —do Carmo, Exercise 1-7, 6 Let $\alpha(s)$, $s \in [0, l]$ be a closed convex plane curve positively oriented. The curve $\beta(s) = \alpha(s) - r\mathbf{n}(s)$, where r is a positive constant and $\mathbf{n}$ is the normal vector, is called a parallel curve to α . Show that:

- Length of $\beta = \text{length of } \alpha + 2\pi r$.
- $A(\beta) = A(\alpha) + rl + \pi r^2$.
- $k_\beta(s) = k_\alpha(s)/(1 + rk_\alpha(s))$.

练习 1-7, 7 —do Carmo, Exercise 1-7, 7 Let $\alpha : \mathbb{R} \rightarrow \mathbb{R}^2$ be a plane curve defined in the entire real line. Assume that α does not pass through the origin and that both $\lim_{t \rightarrow +\infty} |\alpha(t)| = \infty$ and $\lim_{t \rightarrow -\infty} |\alpha(t)| = \infty$.

- Prove that there exists a point $t_0 \in \mathbb{R}$ such that $|\alpha(t_0)| \leq |\alpha(t)|$ for all $t \in \mathbb{R}$.
- Show, by an example, that the assertion in part a is false if one does not assume both limits are infinite.

练习 1-7, 8* —do Carmo, Exercise 1-7, 8 a. Let $\alpha(s)$, $s \in [0, l]$, be a plane simple closed curve. Assume that the curvature $k(s)$ satisfies $0 < k(s) \leq c$. Prove that length of $\alpha \geq 2\pi/c$.

- In part a replace the assumption of being simple by " α has rotation index N ". Prove that length of $\alpha \geq 2\pi N/c$.

练习 1-7, 9* —do Carmo, Exercise 1-7, 9 A set $K \subset \mathbb{R}^2$ is convex if given any two points $p, q \in K$ the segment of straight line $\overline{pq}$ is contained in K (见 fig. 1.34a, 教材 Figure 1-38). Prove that a simple closed convex curve bounds a convex set.

练习 1-7, 10 —do Carmo, Exercise 1-7, 10 Let C be a convex plane curve. Prove geometrically that C has no self-intersections.

练习 1-7, 11* —do Carmo, Exercise 1-7, 11 Given a nonconvex simple closed plane curve C , we can consider its convex hull H (见 fig. 1.34b, 教材 Figure 1-39). Use this to show that, in the isoperimetric problem, we can restrict ourselves to convex curves.

练习 1-7, 12* —do Carmo, Exercise 1-7, 12 Consider a unit circle S^1 in the plane. Show that the ratio $M_1/M_2 = 1/2$, where M_2 is the measure of the set of straight lines in the plane that meet S^1 and M_1 is the measure of all such lines that determine in S^1 a chord of length $> \sqrt{3}$ (见 fig. 1.34c, 教材 Figure 1-40).

练习 1-7, 13 —do Carmo, Exercise 1-7, 13 Let C be an oriented plane closed curve with curvature $k > 0$. Assume that C has at least one point p of self-intersection. Prove that:

- There is a point $p' \in C$ such that the tangent line T' at p' is parallel to some tangent at p .
- The rotation angle of the tangent in the positive arc of C made up by $pp'p$ is $> \pi$ (见 fig. 1.35a, 教材 Figure 1-41).
- The rotation index of C is ≥ 2 .


练习 1-7, 14 —do Carmo, Exercise 1-7, 14 a. Show that if a straight line L meets a closed convex curve C , then either L is tangent to C or L intersects C in exactly two points.

- b. Use part a to show that the measure of the set of lines that meet C (without multiplicities) is equal to the length of C .


练习 1-7, 15 —do Carmo, Exercise 1-7, 15 Green's theorem in the plane: Let a simple closed plane curve be given by $\alpha(t) = (x(t), y(t))$, $t \in [a, b]$. Assume that α is positively oriented, let C be its trace, and let R be the interior of C . Let $p = p(x, y)$, $q = q(x, y)$ be real functions. Then

$$\iint_R (q_x - p_y) dx dy = \int_C (p dx + q dy).$$


- a. Set $q = x$ and $p = -y$ and conclude that $A(R) = \frac{1}{2} \int_a^b (x dy - y dx) dt$.
- b. Use Green's theorem to obtain the transformation formula for double integrals.




(a) Figure 1-38. 凸集定义



(b) Figure 1-39. 凸包



(c) Figure 1-40. 单位圆与等边三角形



(a) Figure 1-41. 自交曲线旋转角