

Differential Geometry of Curves and Surfaces

Manfredo P. do Carmo

May 11, 2026

- a. φ satisfies $\varphi'' + K\varphi = 0$ and ψ is given by $\psi = \int \sqrt{1 - (\varphi')^2} dv$.
- b. All surfaces of revolution with constant curvature $K = 1$ which intersect perpendicularly the plane xOy are given by $\varphi(v) = C \cos v$, $\psi(v) = \int_0^v \sqrt{1 - C^2 \sin^2 v} dv$, where C is a constant. Determine the domain of v and draw a rough sketch of the profile of the surface in the xz plane for the cases $C = 1$, $C > 1$, $C < 1$. Observe that $C = 1$ gives a sphere.
- c. All surfaces of revolution with constant curvature $K = -1$ may be given by one of the following types:

1. $\varphi(v) = C \cosh v$, $\psi(v) = \int_0^v \sqrt{1 - C^2 \sinh^2 v} dv$.
2. $\varphi(v) = C \sinh v$, $\psi(v) = \int_0^v \sqrt{1 - C^2 \cosh^2 v} dv$.
3. $\varphi(v) = e^v$, $\psi(v) = \int_0^v \sqrt{1 - e^{2v}} dv$.

Determine the domain of v and draw a rough sketch of the profile of the surface in the xz plane.

- d. The surface of type 3 in part c is the pseudosphere of Exercise 6.
- e. The only surfaces of revolution with $K \equiv 0$ are the right circular cylinder, the right circular cone, and the plane.

练习 3-3, 8* —do Carmo, Exercise 3-3, 8 (Contact of Order ≥ 2 of Surfaces.) Two surfaces S and $\bar{S}$, with a common point p , have contact of order ≥ 2 at p if there exist parametrizations $\mathbf{x}(u, v)$ and $\bar{\mathbf{x}}(u, v)$ in p of S and $\bar{S}$, respectively, such that $\mathbf{x}_u = \bar{\mathbf{x}}_u$, $\mathbf{x}_v = \bar{\mathbf{x}}_v$, $\mathbf{x}_{uu} = \bar{\mathbf{x}}_{uu}$, $\mathbf{x}_{uv} = \bar{\mathbf{x}}_{uv}$, $\mathbf{x}_{vv} = \bar{\mathbf{x}}_{vv}$ at p . Prove the following:

- a. Let S and $\bar{S}$ have contact of order ≥ 2 at p ; $\mathbf{x}: U \rightarrow S$ and $\bar{\mathbf{x}}: U \rightarrow \bar{S}$ be arbitrary parametrizations in p of S and $\bar{S}$, respectively; and $f: V \subset R^3 \rightarrow R$ be a differentiable function in a neighborhood V of p in R^3 . Then the partial derivatives of order ≤ 2 of $f \circ \bar{\mathbf{x}}$ are zero in $\bar{\mathbf{x}}^{-1}(p)$ if and only if the partial derivatives of order ≤ 2 of $f \circ \mathbf{x}$ are zero in $\mathbf{x}^{-1}(p)$.
- b. Let S and $\bar{S}$ have contact of order ≥ 2 at p . Let $z = f(x, y)$, $z = \bar{f}(x, y)$ be the equations, in a neighborhood of p , of S and $\bar{S}$, respectively, where the xy plane is the common tangent plane at $p = (0, 0)$. Then the function $f(x, y) - \bar{f}(x, y)$ has all partial derivatives of order ≤ 2 , at $(0, 0)$, equal to zero.
- c. Let p be a point in a surface $S \subset R^3$. Let $Oxyz$ be a Cartesian coordinate system for R^3 such that $O = p$ and the xy plane is the tangent plane of S at p . Show that the paraboloid $z = \frac{1}{2}(x^2 f_{xx} + 2xy f_{xy} + y^2 f_{yy})$ has contact of order ≥ 2 at p with S (the surface is called the osculating paraboloid of S at p).
- d. If a paraboloid (the degenerate cases of plane and parabolic cylinder are included) has contact of order ≥ 2 with a surface S at p , then it is the osculating paraboloid of S at p .

- e. If two surfaces have contact of order ≥ 2 at p , then the osculating paraboloids of S and $\bar{S}$ at p coincide. Conclude that the Gaussian and mean curvatures of S and $\bar{S}$ at p are equal.
- f. The notion of contact of order ≥ 2 is invariant by diffeomorphisms of R^3 .
- g. If S and $\bar{S}$ have contact of order ≥ 2 at p , then $\lim_{r \rightarrow 0} \frac{d}{r^2} = 0$, where d is the length of the segment cut by the surfaces in a straight line normal to $T_p(S) = T_p(\bar{S})$, which is at a distance r from p .

练习 3-3, 9 —do Carmo, Exercise 3-3, 9 (Contact of Curves.) Define contact of order $\geq n$ (n integer ≥ 1) for regular curves in R^3 with a common point p and prove that

- a. The notion of contact of order $\geq n$ is invariant by diffeomorphisms.
- b. Two curves have contact of order ≥ 1 at p if and only if they are tangent at p .

练习 3-3, 10* —do Carmo, Exercise 3-3, 10 (Contact of Curves and Surfaces.) A curve C and a surface S , which have a common point p , have contact of order $\geq n$ at p if there exists a curve $\bar{C} \subset S$ passing through p such that C and $\bar{C}$ have contact of order $\geq n$ at p . Prove that

- a. If $f(x, y, z) = 0$ is a representation of a neighborhood of p in S and $\alpha(t) = (x(t), y(t), z(t))$ is a parametrization of C in p , with $\alpha(0) = p$, then C and S have contact of order $\geq n$ if and only if $f(x(0), y(0), z(0)) = 0$, $\frac{df}{dt} = 0$, $\dots$, $\frac{d^n f}{dt^n} = 0$, where the derivatives are computed for $t = 0$.
- b. If a plane has contact of order ≥ 2 with a curve C at p , then this is the osculating plane of C at p .
- c. If a sphere has contact of order ≥ 3 with a curve C at p , and $\alpha(s)$ is a parametrization by arc length of this curve, with $\alpha(0) = p$, then the center of the sphere is given by $\alpha(0) + \frac{1}{k}n + \frac{k'}{k^2\tau}b$. Such a sphere is called the osculating sphere of C at p .

练习 3-3, 11 —do Carmo, Exercise 3-3, 11 Consider the monkey saddle S of Example 2. Construct the Dupin indicatrix at $p = (0, 0, 0)$ using the definition of Sec. 3-2, and compare it with the curve obtained as the intersection of S with a plane parallel to $T_p(S)$ and close to p . Why are they not “approximately similar”?

练习 3-3, 12 —do Carmo, Exercise 3-3, 12 Consider the parametrized surface

$$\mathbf{x}(u, v) = \left(\sin u \cos v, \sin u \sin v, \cos u + \log \tan \frac{u}{2} + \varphi(v) \right),$$

where φ is a differentiable function. Prove that

- a. The curves $v = \text{const}$ are contained in planes which pass through the z axis and intersect the surface under a constant angle θ given by $\cos \theta = \frac{\varphi'}{\sqrt{1+(\varphi')^2}}$. Conclude that the curves $v = \text{const}$ are lines of curvature of the surface.

- b. The length of the segment of a tangent line to a curve $v = \text{const}$, determined by its point of tangency and the z axis, is constantly equal to 1. Conclude that the curves $v = \text{const}$ are tractrices.

练习 3-3, 13 —do Carmo, Exercise 3-3, 13 Let $F: R^3 \rightarrow R^3$ be the map (a similarity) defined by $F(p) = cp$, $p \in R^3$, c a positive constant. Let $S \subset R^3$ be a regular surface and set $F(S) = \bar{S}$. Show that $\bar{S}$ is a regular surface, and find formulas relating the Gaussian and mean curvatures, K and H , of S with the Gaussian and mean curvatures, $\bar{K}$ and $\bar{H}$, of $\bar{S}$.

练习 3-3, 14 —do Carmo, Exercise 3-3, 14 Consider the surface obtained by rotating the curve $y = x^3$, $-1 < x < 1$, about the line $x = 1$. Show that the points obtained by rotation of the origin $(0, 0)$ of the curve are planar points of the surface.

练习 3-3, 15 —do Carmo, Exercise 3-3, 15 Give an example of a surface which has an isolated parabolic point p (that is, no other parabolic point is contained in some neighborhood of p).

练习 3-3, 16 —do Carmo, Exercise 3-3, 16 Show that a surface which is compact (i.e., it is bounded and closed in R^3) has an elliptic point.

练习 3-3, 17 —do Carmo, Exercise 3-3, 17 Define Gaussian curvature for a nonorientable surface. Can you define mean curvature for a nonorientable surface?

练习 3-3, 18 —do Carmo, Exercise 3-3, 18 Show that the Möbius strip of Fig. 3-1 can be parametrized by

$$\mathbf{x}(u, v) = \left(\left(2 - v \sin \frac{u}{2} \right) \sin u, \left(2 - v \sin \frac{u}{2} \right) \cos u, v \cos \frac{u}{2} \right)$$

and that its Gaussian curvature is

$$K = -\frac{1}{\left\{ \frac{1}{4}v^2 + (2 - v \sin(u/2))^2 \right\}^2}.$$

练习 3-3, 19 —do Carmo, Exercise 3-3, 19 Obtain the asymptotic curves of the one-sheeted hyperboloid $x^2 + y^2 - z^2 = 1$.

练习 3-3, 20 —do Carmo, Exercise 3-3, 20 Determine the umbilical points of the ellipsoid

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1.$$

练习 3-3, 21* —do Carmo, Exercise 3-3, 21 Let S be a surface with orientation N . Let $V \subset S$ be an open set in S and let $f: V \subset S \rightarrow R$ be any nowhere-zero differentiable function in V . Let v_1 and v_2 be two differentiable (tangent) vector fields in V such that at each point of V , v_1 and v_2 are orthonormal and $v_1 \wedge v_2 = N$.

- a. Prove that the Gaussian curvature K of V is given by

$$K = \frac{\langle d(fN)(v_1) \wedge d(fN)(v_2), fN \rangle}{f^3}.$$

- b. Apply the above result to show that if f is the restriction of $\sqrt{\frac{x^2}{a^4} + \frac{y^2}{b^4} + \frac{z^2}{c^4}}$ to the ellipsoid $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$, then the Gaussian curvature of the ellipsoid is $K = \frac{1}{a^2 b^2 c^2} \frac{1}{f^4}$.

练习 3-3, 22* —do Carmo, Exercise 3-3, 22 (The Hessian.) Let $h: S \rightarrow R$ be a differentiable function on a surface S , and let $p \in S$ be a critical point of h (i.e., $dh_p = 0$). Let $w \in T_p(S)$ and let $\alpha: (-\epsilon, \epsilon) \rightarrow S$ be a parametrized curve with $\alpha(0) = p$, $\alpha'(0) = w$. Set $H_p h(w) = \left. \frac{d^2(h \circ \alpha)}{dt^2} \right|_{t=0}$.

- a. Let $\mathbf{x}: U \rightarrow S$ be a parametrization of S at p , and show that $H_p h(u'\mathbf{x}_u + v'\mathbf{x}_v) = h_{uu}(p)(u')^2 + 2h_{uv}(p)u'v' + h_{vv}(p)(v')^2$. Conclude that $H_p h: T_p(S) \rightarrow R$ is a well-defined quadratic form on $T_p(S)$. $H_p h$ is called the Hessian of h at p .
- b. Let $h: S \rightarrow R$ be the height function of S relative to $T_p(S)$; that is, $h(q) = \langle q - p, N(p) \rangle$, $q \in S$. Verify that p is a critical point of h and thus that the Hessian $H_p h$ is well defined. Show that if $w \in T_p(S)$, $|w| = 1$, then $H_p h(w) =$ normal curvature at p in the direction of w . Conclude that the Hessian at p of the height function relative to $T_p(S)$ is the second fundamental form of S at p .

练习 3-3, 23* —do Carmo, Exercise 3-3, 23 (Morse Functions on Surfaces.) A critical point $p \in S$ of a differentiable function $h: S \rightarrow R$ is nondegenerate if the self-adjoint linear map $A_p h$ associated to the quadratic form $H_p h$ is nonsingular. Otherwise, p is a degenerate critical point. A differentiable function on S is a Morse function if all its critical points are nondegenerate. Let $h_r: S \subset R^3 \rightarrow R$ be the distance function from S to r ; i.e., $h_r(q) = \sqrt{\langle q - r, q - r \rangle}$, $q \in S$, $r \in R^3$, $r \notin S$.

- a. Show that $p \in S$ is a critical point of h , if and only if the straight line pr is normal to S at p .
- b. Let p be a critical point of $h_r: S \rightarrow R$. Let $w \in T_p(S)$, $|w| = 1$, and let $\alpha: (-\epsilon, \epsilon) \rightarrow S$ be a curve parametrized by arc length with $\alpha(0) = p$, $\alpha'(0) = w$. Prove that $H_p h_r(w) = \frac{1}{h_r(p)} - k_n$, where k_n is the normal curvature at p along the direction of w . Conclude that the orthonormal basis $\{e_1, e_2\}$, where e_1 and e_2 are along the principal directions of $T_p(S)$, diagonalizes the self-adjoint linear map $A_p h_r$. Conclude further that p is a degenerate critical point of h_r if and only if either $h_r(p) = 1/k_1$ or $h_r(p) = 1/k_2$, where k_1 and k_2 are the principal curvatures at p .
- c. Show that the set $B = \{r \in R^3; h_r \text{ is a Morse function}\}$ is a dense set in R^3 .

练习 3-3, 24* —do Carmo, Exercise 3-3, 24 (Local Convexity and Curvature.) A surface $S \subset R^3$ is locally convex at a point $p \in S$ if there exists a neighborhood $V \subset S$ of p such that V is contained in one of the closed half-spaces determined by $T_p(S)$ in R^3 . If, in addition, V has only one common point with $T_p(S)$, then S is called strictly locally convex at p .

- a. Prove that S is strictly locally convex at p if the principal curvatures of S at p are nonzero with the same sign (that is, the Gaussian curvature $K(p)$ satisfies $K(p) > 0$).
- b. Prove that if S is locally convex at p , then the principal curvatures at p do not have different signs (thus, $K(p) \geq 0$).
- c. To show that $K \geq 0$ does not imply local convexity, consider the surface $f(x, y) = x^3(1 + y^2)$, defined in the open set $U = \{(x, y) \in \mathbb{R}^2; y^2 < \frac{1}{2}\}$. Show that the Gaussian curvature of this surface is nonnegative on U and yet the surface is not locally convex at $(0, 0) \in U$.
- d. The example of part c is also very special in the following local sense. Let p be a point in a surface S , and assume that there exists a neighborhood $V \subset S$ of p such that the principal curvatures on V do not have different signs. Prove that S is locally convex at p .

3.3 Vector Fields on Surfaces

本节简要介绍曲面上的向量场 (vector fields), 这将在后续研究测地线时发挥重要作用。

定义 3.16 (向量场). 曲面上的一个**向量场** (vector field) X 对每一点 $p \in S$ 赋予一个切向量 $X(p) \in T_p S$. 若 $X(p)$ 随 p 光滑变化, 则称 X 为光滑向量场。

向量场让我们可以在曲面上做微分运算。曲面上的协变导数 (covariant derivative) $\nabla_X Y$ 测量了向量场 Y 沿向量场 X 方向的变化率。

测地线 (geodesic) 是曲面上的一类特殊曲线, 其切向量沿自身保持平行——也就是说, 测地线是”直的” (没有加速度)。

定义 3.17 (测地线). 曲线 $\gamma(t)$ 称为**测地线** (geodesic), 如果其切向量沿曲线平行, 即

$$\nabla_{\dot{\gamma}} \dot{\gamma} = 0. \quad (3-4-1)$$

测地线的概念是欧几里得空间中”直线”概念在弯曲曲面内蕴几何中的推广。在平面上, 测地线就是直线; 在球面上, 测地线是大圆 (great circles)。

向量场和协变导数的概念将在 Chapter 4 (测地线) 中详细展开。

3-4 节练习

练习 3-4, 1 —do Carmo, Exercise 3-4, 1 Prove that the differentiability of a vector field does not depend on the choice of a coordinate system.

练习 3-4, 2 —do Carmo, Exercise 3-4, 2 Prove that the vector field obtained on the torus by parametrizing all its meridians by arc length and taking their tangent vectors (Example 1) is differentiable.

练习 3-4, 3 —do Carmo, Exercise 3-4, 3 Prove that a vector field w defined on a regular surface $S \subset \mathbb{R}^3$ is differentiable if and only if it is differentiable as a map $w: S \rightarrow \mathbb{R}^3$.

练习 3-4, 4 —do Carmo, Exercise 3-4, 4 Let S be a surface and $\mathbf{x}: U \rightarrow S$ be a parametrization of S . Then $a(u, v)u' + b(u, v)v' = 0$, where a and b are differentiable functions, determines a field of directions r on $\mathbf{x}(U)$, namely, the correspondence which assigns to each $\mathbf{x}(u, v)$ the straight line containing the vector $b\mathbf{x}_u - a\mathbf{x}_v$. Show that a necessary and sufficient condition for the existence of an orthogonal field r' on $\mathbf{x}(U)$ is that both functions $Eb - Fa$ and $Fb - Ga$ are nowhere simultaneously zero (here E, F , and G are the coefficients of the first fundamental form in $\mathbf{x}$) and that r' is then determined by $(Eb - Fa)u' + (Fb - Ga)v' = 0$.

练习 3-4, 5 —do Carmo, Exercise 3-4, 5 Let S be a surface and $\mathbf{x}: U \rightarrow S$ be a parametrization of S . If $ac - b^2 < 0$, show that $a(u, v)(u')^2 + 2b(u, v)u'v' + c(u, v)(v')^2 = 0$ can be factored into two distinct equations, each of which determines a field of directions on $\mathbf{x}(U) \subset S$. Prove that these two fields of directions are orthogonal if and only if $Ec - 2Fb + Ga = 0$.

练习 3-4, 6* —do Carmo, Exercise 3-4, 6 A straight line r meets the z axis and moves in such a way that it makes a constant angle $\alpha \neq 0$ with the z axis and each of its points describes a helix of pitch $c \neq 0$ about the z axis. The figure described by r is the trace of the parametrized surface

$$\mathbf{x}(u, v) = (v \sin \alpha \cos u, v \sin \alpha \sin u, v \cos \alpha + cu).$$

Restrict the parameters (u, v) to an open set U so that $\mathbf{x}(U) = S$ is a regular surface.

- Find the orthogonal family to the family of coordinate curves $u = \text{const}$.
- Use the curves $u = \text{const}$ and their orthogonal family to obtain an orthogonal parametrization for S . Show that in the new parameters $(\bar{u}, \bar{v})$ the coefficients of the first fundamental form are $\bar{G} = 1$, $\bar{F} = 0$, $\bar{E} = \{c^2 + (\bar{v} - c\bar{u} \cos \alpha)^2\} \sin^2 \alpha$.

练习 3-4, 7 —do Carmo, Exercise 3-4, 7 Define the derivative $\mathbf{w}(\mathbf{f})$ of a differentiable function $f: U \subset \mathbf{S} \rightarrow \mathbf{R}$ relative to a vector field $\mathbf{w}$ in $\mathbf{U}$ by $\mathbf{w}(\mathbf{f})(\mathbf{q}) = \frac{d}{dt}(f \circ \alpha)|_{t=0}$, where $\alpha: I \rightarrow S$ is a curve such that $\alpha(0) = q$, $\alpha'(0) = w(q)$. Prove that

- w is differentiable in U if and only if $w(f)$ is differentiable for all differentiable f in U .
- Let λ and μ be real numbers and $g: U \subset S \rightarrow \mathbf{R}$ be a differentiable function on U ; then $w(\lambda f + \mu g) = \lambda w(f) + \mu w(g)$ and $w(fg) = w(f)g + fw(g)$.

练习 3-4, 8 —do Carmo, Exercise 3-4, 8 Show that if w is a differentiable vector field on a surface S and $w(p) \neq 0$ for some $p \in S$, then it is possible to parametrize a neighborhood of p by $\mathbf{x}(u, v)$ in such a way that $\mathbf{x}_u = w$.

练习 3-4, 9* —do Carmo, Exercise 3-4, 9 a. Let $A: V \rightarrow W$ be a nonsingular linear map of vector spaces V and W of dimension 2 and endowed with inner products $\langle, \rangle$ and $(,)$, respectively. A is a similitude if there exists a real number $\lambda \neq 0$ such that $\langle Av_1, Av_2 \rangle = \lambda \langle v_1, v_2 \rangle$ for all vectors $v_1, v_2 \in V$. Assume that A is not a similitude and show that there exists a unique pair of orthonormal vectors e_1 and e_2 in V such that Ae_1, Ae_2 are orthogonal in W .

- b. Use part a to prove Tissot's theorem: Let $\varphi: U_1 \subset S_1 \rightarrow S_2$ be a diffeomorphism from a neighborhood U_1 of a point p of a surface S_1 into a surface S_2 . Assume that the linear map $d\varphi$ is nowhere a similitude. Then it is possible to parametrize a neighborhood of p in S_1 by an orthogonal parametrization $\mathbf{x}_1: U \rightarrow S_1$ in such a way that $\varphi \circ \mathbf{x}_1 = \mathbf{x}_2: U \rightarrow S_2$ is also an orthogonal parametrization in a neighborhood of $\varphi(p) \in S_2$.

练习 3-4, 10* —do Carmo, Exercise 3-4, 10 Let T be the torus of Example 6 of Sec. 2-2 and define a map $\varphi: R^2 \rightarrow T$ by $\varphi(u, v) = ((r \cos u + a) \cos v, (r \cos u + a) \sin v, r \sin u)$, where u and v are the Cartesian coordinates of R^2 . Let $u = at, v = bt$ be a straight line in R^2 , passing by $(0, 0) \in R^2$, and consider the curve in T $\alpha(t) = \varphi(at, bt)$. Prove that

- φ is a local diffeomorphism.
- The curve $\alpha(t)$ is a regular curve; $\alpha(t)$ is a closed curve if and only if b/a is a rational number.
- If b/a is irrational, the curve $\alpha(t)$ is dense in T .

练习 3-4, 11 —do Carmo, Exercise 3-4, 11 Use the local uniqueness of trajectories of a vector field w in $U \subset S$ to prove the following result. Given $p \in U$, there exists a unique trajectory $\alpha: I \rightarrow U$ of w , with $\alpha(0) = p$, which is maximal in the following sense: Any other trajectory $\beta: J \rightarrow U$, with $\beta(0) = p$, is the restriction of α to J .

练习 3-4, 12 —do Carmo, Exercise 3-4, 12 Prove that if w is a differentiable vector field on a compact surface S and $\alpha(t)$ is the maximal trajectory of w with $\alpha(0) = p \in S$, then $\alpha(t)$ is defined for all $t \in R$.

练习 3-4, 13* —do Carmo, Exercise 3-4, 13 Construct a differentiable vector field on an open disk of the plane (which is not compact) such that a maximal trajectory $\alpha(t)$ is not defined for all $t \in R$ (this shows that the compactness condition of Exercise 12 is essential).

3.4 Ruled Surfaces and Minimal Surfaces

本节简要介绍两类特殊曲面：直纹面 (ruled surfaces) 和极小曲面 (minimal surfaces)。这些是曲面论中的经典研究对象。

3.4.1 Ruled Surfaces

定义 3.18 (直纹面 (Ruled surface)). 由一条可动直线 (母线) 生成的曲面称为直纹面 (ruled surface)。具体地, 若 $\alpha(t)$ 是一条空间曲线, $d(t)$ 是与 $\alpha(t)$ 正交的光滑单位向量场, 则

$$\mathbf{x}(t, v) = \alpha(t) + v d(t), \quad t \in I, v \in \mathbb{R}$$

参数化一张直纹面。

直纹面的经典例子包括：圆柱面（直母线绕一条固定轴旋转）、单叶双曲面（ $\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{z^2}{c^2} = 1$ ，可以通过两条直线族生成）。

直纹面的几何特征：至少有一个主曲率为零。这意味着直纹面在母线方向上的法曲率恒为零——沿母线方向，曲面是“平直”的。

例 3.8 (圆柱面). 圆柱面 $x^2 + y^2 = 1$ 可以视为由所有平行于 z 轴的直线（母线）绕 z 轴旋转而成。圆柱面有一个零主曲率（沿母线方向）和一个非零主曲率（沿圆周方向）。

可展曲面（developable surface）是一类特殊的直纹面，其 Gauss 曲率恒为零。平面、柱面、锥面都是可展曲面的例子。可展曲面可以在不拉伸、不撕裂的情况下平铺到平面上——这正是“可展”的含义。

例 3.9 (锥面). 锥面 $\frac{x^2}{a^2} + \frac{y^2}{b^2} = \frac{z^2}{c^2}$ 也是直纹面（从顶点出发的每条直线都是母线），且 $K \equiv 0$ 。

Gauss 映射在直纹面上的行为有特别的几何意义——直纹面上一条母线的所有点对应球面上相同的法向量（因为母线上每一点的切平面都相同）。

3.4.2 Minimal Surfaces

定义 3.19 (极小曲面 (Minimal surface)). 平均曲率恒为零的曲面称为极小曲面 (*minimal surface*):

$$H = \frac{\kappa_1 + \kappa_2}{2} \equiv 0. \quad (3-5-1)$$

极小曲面的名字来源于一个物理背景：如果将一根铁丝弯成一条封闭曲线，然后将其浸入肥皂水中，形成的肥皂膜（极小曲面）会使面积最小化——这是自然界的“能量最小化”原理。

著名的极小曲面包括：

- **Catenoid** (悬链面): $x^2 + y^2 = \cosh^2 z$ ，第一个被发现的双连通极小曲面
- **Henneberg surface**
- **Enneper's surface** (见 ??)
- **Plateau 法则**: 给定边界曲线，极小曲面是使该边界曲线所张成的面积最小的曲面

极小曲面有一个显著性质：如果 $H \equiv 0$ ，则（由 Codazzi-Mainardi 方程）有

$$\langle dN_p(w_1), dN_p(w_2) \rangle = -K(p) \langle w_1, w_2 \rangle. \quad (3-5-2)$$

这说明 $|dN_p(v)| = \sqrt{-K(p)}$ ，即 Gauss 映射的微分由 Gauss 曲率完全决定。

极小曲面是当代微分几何和复分析中的核心研究对象，其理论连接了微分几何、复分析和数学物理。

3-5 节练习

练习 3-5, 1 —do Carmo, Exercise 3-5, 1 Show that the helicoid (cf. Example 3, Sec. 2-5) is a ruled surface, its line of striction is the z axis, and its distribution parameter is constant.

解答 直观理解：螺旋面由一族直纹面组成，每条直母线绕 z 轴螺旋上升。直观上， z 轴是所有直母线的“中心线”，因为每条直母线都与 z 轴等距且等角相交。

解答：

螺旋面的参数化来自 Example 3, Sec. 2-5:

$$\mathbf{x}(t, v) = (v \cos t, v \sin t, ct), \quad (t, v) \in \mathbb{R}^2,$$

其中 t 为角度参数， v 为沿母线的坐标， $c \neq 0$ 为常数。

(1) 螺旋面是直纹面：对每个固定的 t ，曲线 $v \mapsto \mathbf{x}(t, v)$ 是一条直线（母线），方向向量为

$$\mathbf{x}_v(t, v) = (\cos t, \sin t, 0),$$

故螺旋面是直纹面。

(2) 求线限制 (line of striction)：先计算 $\alpha'(t) \wedge w(t)$ 。直母线方向 $w(t) = \mathbf{x}_v(t, 0) = (\cos t, \sin t, 0)$ ，沿 t 的方向向量取基线 $\alpha(t) = \mathbf{x}(t, 0) = (0, 0, ct)$ 。则

$$\alpha'(t) = (0, 0, c), \quad w(t) = (\cos t, \sin t, 0),$$

$$\alpha'(t) \wedge w(t) = (0, 0, c) \wedge (\cos t, \sin t, 0) = (-c \sin t, c \cos t, 0).$$

计算分布参数 (distribution parameter)

$$\lambda(t) = -\frac{\langle \alpha'(t), w(t) \wedge w'(t) \rangle}{|w'(t) \wedge w(t)|^2}.$$

先算 $w'(t) = (-\sin t, \cos t, 0)$ ，再算

$$w'(t) \wedge w(t) = (-\sin t, \cos t, 0) \wedge (\cos t, \sin t, 0) = (0, 0, -1).$$

故 $|w'(t) \wedge w(t)| = 1$ ，而

$$\langle \alpha'(t), w(t) \wedge w'(t) \rangle = (0, 0, c) \cdot (0, 0, -1) = -c.$$

于是

$$\lambda(t) = -\frac{-c}{1} = c,$$

为常数。

(3) 求中心点 (central point)：由

$$\bar{v}(t) = -\frac{\langle \alpha', w' \rangle}{\langle w', w' \rangle} = -\frac{\langle (0, 0, c), (-\sin t, \cos t, 0) \rangle}{\langle (-\sin t, \cos t, 0), (-\sin t, \cos t, 0) \rangle} = -\frac{0}{1} = 0.$$

故所有直母线的中心点为 $\alpha(t) = (0, 0, ct)$ ，即 z 轴。

结论：螺旋面是直纹面，线限制为 z 轴，分布参数恒为常数 c 。

□

练习 3-5, 2 —do Carmo, Exercise 3-5, 2 Show that on the hyperboloid of revolution $x^2 + y^2 - z^2 = 1$, the parallel of least radius is the line of striction, the rulings meet it under a constant angle, and the distribution parameter is constant.

解答 直观理解：单叶双曲面 $x^2 + y^2 - z^2 = 1$ 是直纹面（两种直母线族），绕 z 轴旋转而成。最小半径的平行圆位于 $z = 0$ 处，直观上这是所有直母线的“中心”所在。

解答：

(1) 参数化：单叶双曲面作为直纹面有标准参数化（两种直母线族之一）：

$$\mathbf{x}(u, v) = (\cos u + v \sin u, \sin u - v \cos u, v), \quad u \in \mathbb{R}, v \in \mathbb{R}.$$

验证： $x^2 + y^2 - z^2 = (\cos u + v \sin u)^2 + (\sin u - v \cos u)^2 - v^2 = \cos^2 u + 2v \cos u \sin u + v^2 \sin^2 u + \sin^2 u - 2v \sin u \cos u + v^2 \cos^2 u - v^2 = 1$ 。

这里取基线 $\alpha(u) = (\cos u, \sin u, 0)$ ，方向向量

$$\tilde{w}(u) = (\sin u, -\cos u, 1).$$

归一化得

$$w(u) = \frac{1}{\sqrt{2}}(\sin u, -\cos u, 1), \quad |w| = 1.$$

(2) 分布参数：

$$\alpha'(u) = (-\sin u, \cos u, 0), \quad w'(u) = \frac{1}{\sqrt{2}}(\cos u, \sin u, 0).$$

$$w'(u) \wedge w(u) = \frac{1}{\sqrt{2}}(\cos u, \sin u, 0) \wedge \frac{1}{\sqrt{2}}(\sin u, -\cos u, 1) = \frac{1}{2}(\sin u, -\cos u, -1).$$

故 $|w' \wedge w|^2 = \frac{1}{4}(\sin^2 u + \cos^2 u + 1) = \frac{1}{2}$ 。

$$\langle \alpha', w' \wedge w \rangle = \frac{1}{2} \langle (-\sin u, \cos u, 0), (\sin u, -\cos u, -1) \rangle = \frac{1}{2}(-\sin^2 u - \cos^2 u + 0) = -\frac{1}{2}.$$

于是分布参数

$$\lambda(u) = -\frac{\langle \alpha', w' \wedge w \rangle}{|w' \wedge w|^2} = -\frac{-1/2}{1/2} = 1,$$

为常数。

(3) 线限制：

$$\bar{v} = -\frac{\langle \alpha', w' \rangle}{\langle w', w' \rangle}.$$

$$\langle \alpha', w' \rangle = \frac{1}{\sqrt{2}} \langle (-\sin u, \cos u, 0), (\cos u, \sin u, 0) \rangle = \frac{1}{\sqrt{2}}(-\sin u \cos u + \cos u \sin u) = 0.$$

故 $\bar{v} = 0$ ，中心点在 $\alpha(u) = (\cos u, \sin u, 0)$ ，即 $z = 0$ 的单位圆——平行圆中半径最小的（ $r = 1$ ）。

结论： $z = 0$ 的平行圆（半径为 1）确是线限制；分布参数恒为 1（常数）；由 λ 为常数可知直母线与线限制的夹角为常数。□

练习 3-5, 3 —do Carmo, Exercise 3-5, 3 Let $\alpha: I \rightarrow S \subset R^3$ be a curve on a regular surface S and consider the ruled surface generated by the family $\{\alpha(t), N(t)\}$, where $N(t)$ is the normal to the surface at $\alpha(t)$. Prove that $\alpha(I) \subset S$ is a line of curvature in S if and only if this ruled surface is developable.

解答 直观理解：沿曲线 α 的法向量场 $N(t)$ 生成一个直纹面。若 α 是主曲率方向 (line of curvature), 则 $N(t)$ 沿 α 的旋转由主曲率控制, 此时该直纹面为可展曲面——即切平面沿每条母线不变。

解答：

直纹面为

$$\mathbf{x}(t, v) = \alpha(t) + vN(t).$$

由于 $|N(t)| = 1$, $N'(t)$ 与 $N(t)$ 正交。

($\Rightarrow$) 设 $\alpha(I)$ 是 S 的主曲率方向。则 $N'(t)$ 必与 $\alpha'(t)$ 共线 (因为主曲率方向是 dN 的特征向量方向, 而 $dN(\alpha') = -k_n\alpha'$, 故 N' 与 α' 平行)。于是存在函数 $\lambda(t)$ 使

$$N'(t) = \lambda(t)\alpha'(t).$$

对直纹面 $\mathbf{x}(t, v)$, 计算偏导:

$$\mathbf{x}_t = \alpha' + vN', \quad \mathbf{x}_v = N.$$

两者的叉积为

$$\mathbf{x}_t \wedge \mathbf{x}_v = (\alpha' + vN') \wedge N = \alpha' \wedge N + vN' \wedge N.$$

注意 $N' \wedge N$ 与 N 正交, 而 $\alpha' \wedge N$ 也与 N 正交, 故 $\mathbf{x}_t \wedge \mathbf{x}_v$ 总与 N 正交。由可展曲面的判别 (Theorem 3-5-1), 直纹面 $\mathbf{x}(t, v)$ 为可展当且仅当

$$\langle \mathbf{x}_t \wedge \mathbf{x}_v, \mathbf{x}_{tt} \wedge \mathbf{x}_{tv} \rangle = 0,$$

或等价地 (由直接计算), 当 $\langle N, \alpha' \wedge N' \rangle = 0$ 。因为 N' 与 α' 共线, $\alpha' \wedge N'$ 与 N 正交, 故此条件成立。

($\Leftarrow$) 设该直纹面可展。则由可展曲面的性质, 有 $\langle N, \alpha' \wedge N' \rangle = 0$ 。这意味着 $\alpha' \wedge N'$ 与 N 正交, 故 $\alpha' \wedge N'$ 落在切平面内。但 N' 本身也与 N 正交, 所以 N' 必与 α' 共线 (因为两者都在与 N 正交的平面内)。于是 $N' = \lambda\alpha'$, 说明 α' 是 dN 的特征向量方向, 即 α 是主曲率方向。

结论： α 是 S 的主曲率方向当且仅当该直纹面可展。 □

练习 3-5, 4* —do Carmo, Exercise 3-5, 4 Assume that a noncylindrical ruled surface $\mathbf{x}(t, v) = \alpha(t) + vw(t)$, $|w| = 1$, is regular. Let $w(t_1), w(t_2)$ be the directions of two rulings of $\mathbf{x}$ and let $\mathbf{x}(t_1, v_1), \mathbf{x}(t_2, v_2)$ be the feet of the common perpendicular to these two rulings. As $t_2 \rightarrow t_1$, these points tend to a point $\mathbf{x}(t_1, \bar{v})$. To determine $(t_1, \bar{v})$ prove the following:

- The unit vector of the common perpendicular converges to a unit vector tangent to the surface at $(t_1, \bar{v})$. Conclude that, at $(t_1, \bar{v})$ $\langle w' \wedge w, N \rangle = 0$.
- $\bar{v} = -(\langle \alpha', w' \rangle / \langle w', w' \rangle)$. Thus, $(t_1, \bar{v})$ is the central point of the ruling through t_1 , and this gives another interpretation of the line of striction (assumed nonsingular).

练习 3-5, 5* —do Carmo, Exercise 3-5, 5 A right conoid is a ruled surface whose rulings L_t intersect perpendicularly at fixed axis r which does not meet the directrix $\alpha: I \rightarrow R^3$.

- Find a parametrization for the right conoid and determine a condition that implies it to be noncylindrical.
- Given a noncylindrical right conoid, find the line of striction and the distribution parameter.

练习 3-5, 6 —do Carmo, Exercise 3-5, 6 Let $\mathbf{x}(t, v) = \alpha(t) + vw(t)$ be a developable surface. Prove that at a regular point we have $\langle N_v, \mathbf{x}_v \rangle = \langle N_v, \mathbf{x}_t \rangle = 0$. Conclude that the tangent plane of a developable surface is constant along (the regular points of) a fixed ruling.

练习 3-5, 7 —do Carmo, Exercise 3-5, 7 Let S be a regular surface and let $C \subset S$ be a regular curve on S , nowhere tangent to an asymptotic direction. Consider the envelope of the family of tangent planes of S along C . Prove that the direction of the ruling that passes through a point $p \in C$ is conjugate to the tangent direction of C at p .

练习 3-5, 8 —do Carmo, Exercise 3-5, 8 Show that if $C \subset S^2$ is a parallel of unit sphere S^2 , then the envelope of tangent planes of S^2 along C is either a cylinder, if C is an equator, or a cone, if C is not an equator.

练习 3-5, 9* —do Carmo, Exercise 3-5, 9 (Focal Surfaces.) Let S be a regular surface without parabolic or umbilical points. Let $\mathbf{x}: U \rightarrow S$ be a parametrization of S such that the coordinate curves are lines of curvature. The parametrized surfaces $\mathbf{y}(u, v) = \mathbf{x}(u, v) + \rho_1 N(u, v)$ and $\mathbf{z}(u, v) = \mathbf{x}(u, v) + \rho_2 N(u, v)$, where $\rho_1 = 1/k_1$, $\rho_2 = 1/k_2$, are called focal surfaces of $\mathbf{x}(U)$. Prove that

- If $(k_1)_u$ and $(k_2)_v$ are nowhere zero, then $\mathbf{y}$ and $\mathbf{z}$ are regular parametrized surfaces.
- At the regular points, the directions on a focal surface corresponding to the principal directions on $\mathbf{x}(U)$ are conjugate.
- A focal surface, say $\mathbf{y}$, can be constructed as follows: Consider the line of curvature $\mathbf{x}(u, \text{const.})$ on $\mathbf{x}(U)$, and construct the developable surface generated by the normals of $\mathbf{x}(U)$ along the curve $\mathbf{x}(u, \text{const.})$. The line of striction of such a developable lies on $\mathbf{y}(U)$, and as $\mathbf{x}(u, \text{const.})$ describes $\mathbf{x}(U)$, this line describes $\mathbf{y}(U)$.

练习 3-5, 10* —do Carmo, Exercise 3-5, 10 Example 4 can be generalized as follows. A one-parameter differentiable family of planes $\{\alpha(t), N(t)\}$ is a correspondence which assigns to each $t \in I$ a point $\alpha(t) \in R^3$ and a unit vector $N(t) \in R^3$ in such a way that both α , and N are differentiable maps. A family $\{\alpha(t), N(t)\}$, $t \in I$, is said to be a family of tangent planes if $\alpha'(t) \neq 0$, $N'(t) \neq 0$, and $\langle \alpha'(t), N(t) \rangle = 0$ for all $t \in I$.

- Give a proof that a differentiable one-parameter family of tangent planes $\{\alpha(t), N(t)\}$ determines a differentiable one-parameter family of lines $\{\alpha(t), (N \wedge N')/|N'|\}$ which generates a developable surface $\mathbf{x}(t, v) = \alpha(t) + v \frac{N \wedge N'}{|N'|}$.

- b. Prove that if $\alpha'(t) \wedge (N(t) \wedge N'(t)) \neq 0$ for all $t \in I$, then the envelope is regular in a neighborhood of $v = 0$, and the unit normal vector of $\mathbf{x}$ at $(t, 0)$ is $N(t)$.
- c. Let $\alpha = \alpha(s)$ be a curve in R^3 parametrized by arc length. Assume that the curvature $k(s)$ and the torsion $\tau(s)$ of α are nowhere zero. Prove that the family of osculating planes $\{\alpha(s), b\{s\}\}$ is a one-parameter differentiable family of tangent planes and that the envelope of this family is the tangent surface to $\alpha(s)$.

练习 3-5, 11* —do Carmo, Exercise 3-5, 11 Let $\mathbf{x} = \mathbf{x}(u, v)$ be a regular parametrized surface. A parallel surface to $\mathbf{x}$ is a parametrized surface $\mathbf{y}(u, v) = \mathbf{x}(u, v) + aN(u, v)$, where a is a constant.

- a. Prove that $y_u \wedge y_v = (1 - 2Ha + Ka^2)(\mathbf{x}_u \wedge \mathbf{x}_v)$, where K and H are the Gaussian and mean curvatures of $\mathbf{x}$, respectively.
- b. Prove that at the regular points, the Gaussian curvature of $\mathbf{y}$ is $\frac{K}{1-2Ha+Ka^2}$ and the mean curvature of $\mathbf{y}$ is $\frac{H-Ka}{1-2Ha+Ka^2}$.
- c. Let a surface $\mathbf{x}$ have constant mean curvature equal to $c \neq 0$ and consider the parallel surface to $\mathbf{x}$ at a distance $1/2c$. Prove that this parallel surface has constant Gaussian curvature equal to $4c^2$.

练习 3-5, 12* —do Carmo, Exercise 3-5, 12 Prove that there are no compact (i.e., bounded and closed in R^3) minimal surfaces.

练习 3-5, 13* —do Carmo, Exercise 3-5, 13 a. Let S be a regular surface without umbilical points. Prove that S is a minimal surface if and only if the Gauss map $N: S \rightarrow S^2$ satisfies, for all $p \in S$ and all $w_1, w_2 \in T_p(S)$, $\langle dN_p(w_1), dN_p(w_2) \rangle_{N(p)} = \lambda(p) \langle w_1, w_2 \rangle_p$, where $\lambda(p) \neq 0$ is a number which depends only on p .

- b. Let $\mathbf{x}: U \rightarrow S^2$ be a parametrization of the unit sphere S^2 by stereographic projection. Consider a neighborhood V of a point p of the minimal surface S in part a such that $N: S \rightarrow S^2$ restricted to V is a diffeomorphism. Prove that the parametrization $y = N^{-1} \circ \mathbf{x}: U \rightarrow S$ is isothermal.

练习 3-5, 14* —do Carmo, Exercise 3-5, 14 When two differentiable functions $f, g: U \subset R^2 \rightarrow R$ satisfy the Cauchy-Riemann equations $\frac{\partial f}{\partial u} = \frac{\partial g}{\partial v}$, $\frac{\partial f}{\partial v} = -\frac{\partial g}{\partial u}$, they are easily seen to be harmonic; in this situation, f and g are said to be harmonic conjugate. Let $\mathbf{x}$ and $\mathbf{y}$ be isothermal parametrizations of minimal surfaces such that their component functions are pairwise harmonic conjugate; then $\mathbf{x}$ and $\mathbf{y}$ are called conjugate minimal surfaces. Prove that

- a. The helicoid and the catenoid are conjugate minimal surfaces.
- b. Given two conjugate minimal surfaces, $\mathbf{x}$ and $\mathbf{y}$, the surface $\mathbf{z} = (\cos t)\mathbf{x} + (\sin t)\mathbf{y}$ is again minimal for all $t \in R$.
- c. All surfaces of the one-parameter family have the same fundamental form: $E = \langle \mathbf{x}_u, \mathbf{x}_u \rangle = \langle \mathbf{y}_v, \mathbf{y}_v \rangle$, $F = 0$, $G = \langle \mathbf{x}_v, \mathbf{x}_v \rangle = \langle \mathbf{y}_u, \mathbf{y}_u \rangle$.

附录：详细推导

球面的第二基本形式推导

考虑单位球面在地理坐标下的参数化：

$$\mathbf{x}(\theta, \varphi) = (\sin \theta \cos \varphi, \sin \theta \sin \varphi, \cos \theta).$$

步骤 1：计算一阶偏导数

$$\mathbf{x}_\theta = (\cos \theta \cos \varphi, \cos \theta \sin \varphi, -\sin \theta), \quad \mathbf{x}_\varphi = (-\sin \theta \sin \varphi, \sin \theta \cos \varphi, 0).$$

步骤 2：计算单位法向量

$$\mathbf{x}_\theta \times \mathbf{x}_\varphi = (\sin^2 \theta \cos \varphi, \sin^2 \theta \sin \varphi, \sin \theta \cos \theta).$$

取模长 $\|\mathbf{x}_\theta \times \mathbf{x}_\varphi\| = \sin \theta$ ，得

$$N = (\sin \theta \cos \varphi, \sin \theta \sin \varphi, \cos \theta). \quad (\text{A3-1})$$

注意这里 $N = \mathbf{x}$ ——球面上每一点的法向量恰好等于该点的位置向量。

步骤 3：计算二阶偏导数

$$\mathbf{x}_{\theta\theta} = (-\sin \theta \cos \varphi, -\sin \theta \sin \varphi, -\cos \theta), \quad \mathbf{x}_{\theta\varphi} = (-\cos \theta \sin \varphi, \cos \theta \cos \varphi, 0), \quad \mathbf{x}_{\varphi\varphi} = (-\sin \theta \cos \varphi, -\sin \theta \sin \varphi, 0).$$

步骤 4：计算第二基本形式系数

$$\begin{aligned} L = \langle \mathbf{x}_{\theta\theta}, N \rangle &= (-\sin \theta \cos \varphi)(\sin \theta \cos \varphi) + (-\sin \theta \sin \varphi)(\sin \theta \sin \varphi) + (-\cos \theta)(\cos \theta) \\ &= -\sin^2 \theta - \cos^2 \theta = -1, \end{aligned}$$

$$M = \langle \mathbf{x}_{\theta\varphi}, N \rangle = 0, \quad N_{\text{coeff}} = \langle \mathbf{x}_{\varphi\varphi}, N \rangle = -\sin^2 \theta. \quad (\text{A3-2})$$

结论：球面的第二基本形式系数为 $L = -1$, $M = 0$, $N_{\text{coeff}} = -\sin^2 \theta$ 。结合第一基本形式 $E = 1$, $F = 0$, $G = \sin^2 \theta$ ，可得 $S_p = I$ （恒等变换），因此 $\kappa_1 = \kappa_2 = 1$ 。

注意在很多教材中，球面的 Weingarten 映射定义为 $S_p = -dN_p$ ，此时 $dN_p(v) = -v$, $S_p(v) = v$ （恒等变换），与我们的定义一致。但在法曲率计算 $\kappa_n = \langle S_p(v), v \rangle$ 中，法曲率为正值。这与球面“向法向量方向弯曲”的直觉一致。

旋转抛物面的第二基本形式推导

考虑旋转抛物面 $z = x^2 + y^2$ ，参数化为

$$\mathbf{x}(u, v) = (u, v, u^2 + v^2).$$

步骤 1：一阶偏导数

$$\mathbf{x}_u = (1, 0, 2u), \quad \mathbf{x}_v = (0, 1, 2v).$$

步骤 2：单位法向量

$$\mathbf{x}_u \times \mathbf{x}_v = (-2u, -2v, 1), \quad \|\mathbf{x}_u \times \mathbf{x}_v\| = \sqrt{1 + 4u^2 + 4v^2}.$$

$$N = \frac{(-2u, -2v, 1)}{\sqrt{1 + 4u^2 + 4v^2}}. \quad (\text{A3-3})$$

步骤 3: 二阶偏导数

$$\mathbf{x}_{uu} = (0, 0, 2), \quad \mathbf{x}_{uv} = (0, 0, 0), \quad \mathbf{x}_{vv} = (0, 0, 2).$$

步骤 4: 第二基本形式系数

$$L = \langle \mathbf{x}_{uu}, N \rangle = \frac{2}{\sqrt{1 + 4u^2 + 4v^2}}, \quad M = \langle \mathbf{x}_{uv}, N \rangle = 0, \quad N_{\text{coeff}} = \langle \mathbf{x}_{vv}, N \rangle = \frac{2}{\sqrt{1 + 4u^2 + 4v^2}}. \quad (\text{A3-4})$$

在点 $(0, 0)$ 处, $L = N_{\text{coeff}} = 2$, $M = 0$ 。

此时第一基本形式为 $E = 1$, $F = 0$, $G = 1$, 因此

$$S_p = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}^{-1} \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix} = \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix} = 2I.$$

所以 $\kappa_1 = \kappa_2 = 2$, $K = 4 > 0$ (椭圆点), $H = 2$ 。

旋转抛物面在原点的 Dupin 指示线是一个椭圆, 因为 $K > 0$, 所有方向的法曲率同号。

Christoffel 符号与 Codazzi-Mainardi 方程

Christoffel 符号 Γ_{ij}^k 由第一基本形式的系数定义为:

$$\Gamma_{ij}^k = \frac{1}{2} \sum_{m=1}^2 g^{km} \left(\frac{\partial g_{im}}{\partial j} + \frac{\partial g_{jm}}{\partial i} - \frac{\partial g_{ij}}{\partial m} \right), \quad (\text{A3-5})$$

其中 $(g_{ij}) = \begin{pmatrix} E & F \\ F & G \end{pmatrix}$, (g^{ij}) 是其逆矩阵。

Codazzi-Mainardi 方程的直观含义是: 第二基本形式的”协变导数”必须对称。具体形式为:

$$\begin{aligned} \frac{\partial L}{\partial v} - \frac{\partial M}{\partial u} &= L\Gamma_{12}^1 + M(\Gamma_{12}^2 - \Gamma_{11}^1) - N_{\text{coeff}}\Gamma_{11}^2, \\ \frac{\partial N_{\text{coeff}}}{\partial u} - \frac{\partial M}{\partial v} &= L\Gamma_{22}^1 + M(\Gamma_{22}^2 - \Gamma_{21}^1) - N_{\text{coeff}}\Gamma_{21}^2. \end{aligned} \quad (\text{A3-6})$$

这些方程的深层含义是: 第一基本形式和第二基本形式不是独立的——它们必须协调一致, 才能描述一个真实嵌入 $\mathbb{R}^3$ 的曲面。这种协调性正是可积性条件的本质。

附录：问答记录

本节记录学习 Do Carmo 《曲线与曲面的微分几何》过程中的问答与思考。

待记录问题

本节将随学习进度逐步填充以下问题：

- 曲率与挠率的直观几何意义
- Frenet-Serret 公式的推导细节
- 局部标准形式的系数解释
- 全曲率定理的证明思路
- 等周不等式的证明

历史悠久的问答

曲率的直观理解

问：曲率 $k(s) = \|\alpha''(s)\|$ 的直观几何意义是什么？

答：对于单位速度曲线 $\alpha(s)$ ，速度向量 $\alpha'(s)$ 是单位切向量，其模长恒为 1。加速度向量 $\alpha''(s)$ 描述速度向量方向的变化率。 $\|\alpha''(s)\|$ 正是这种方向变化的速率——即曲线在每一点的弯曲程度。直线速度方向不变， $\alpha''(s) = 0$ ；圆周速度方向匀速变化， $\|\alpha''(s)\| = 1/a$ (a 为半径)。

挠率与扭曲

问：挠率 $\tau(s)$ 描述的是什么？

答：挠率描述曲线在空间中的“扭曲”程度。曲率只告诉我们曲线在密切平面内如何弯曲，但曲线可能“偏离”密切平面——这种偏离的速度由挠率度量。对于平面曲线， $\alpha', \alpha'', \alpha'''$ 始终共面， $\det(\alpha', \alpha'', \alpha''') = 0$ ，故挠率为零。螺旋线的挠率为常数，描述其均匀扭曲的特性。

曲率另一个表达式的推导

背景：命题 1-5, 2 给出了曲率在一般参数化下的表达式：

$$k(t) = \frac{\|\alpha'(t) \times \alpha''(t)\|}{\|\alpha'(t)\|^3}.$$

目标：从单位速度曲率的定义 $k(s) = \|\alpha''(s)\|$ 推导出一般参数下的表达式。

推导步骤：

设 $\alpha(t)$ 是正则曲线， $s = s(t)$ 是弧长参数。回忆弧长公式：

$$s(t) = \int_{t_0}^t \|\alpha'(u)\| du, \quad \frac{ds}{dt} = \|\alpha'(t)\|.$$

由链式法则， $\alpha'(s) = \frac{\alpha'(t)}{ds/dt} = \frac{\alpha'(t)}{\|\alpha'(t)\|}$ 。

进一步，

$$\alpha''(s) = \frac{d}{ds} \alpha'(s) = \frac{d}{dt} \left(\frac{\alpha'(t)}{\|\alpha'(t)\|} \right) \cdot \frac{dt}{ds}.$$

经过计算（利用向量积的模长公式 $\|u \times v\| = \|u\| \|v\| \sin \theta$ ），可得：

$$\|\alpha''(s)\| = \frac{\|\alpha'(t) \times \alpha''(t)\|}{\|\alpha'(t)\|^3}.$$

因此在参数 t 下，曲率为 $k(t) = \frac{\|\alpha'(t) \times \alpha''(t)\|}{\|\alpha'(t)\|^3}$ 。

挠率另一个表达式的推导

背景：命题 1-5, 4 给出了挠率的另一个表达式：

$$\tau(s) = -\langle \mathbf{b}'(s), \mathbf{n}(s) \rangle.$$

目标：从挠率的行列式定义 $\tau = \frac{\det(\alpha', \alpha'', \alpha''')}{k^2}$ 推导出 $\tau = -\langle \mathbf{b}', \mathbf{n} \rangle$ 。

推导步骤：

由 Frenet 标架的定义： $\mathbf{t} = \alpha'$ ， $\mathbf{n} = \frac{\alpha''}{k}$ ， $\mathbf{b} = \mathbf{t} \times \mathbf{n}$ 。

对 $\mathbf{b} = \mathbf{t} \times \mathbf{n}$ 求导：

$$\mathbf{b}' = \mathbf{t}' \times \mathbf{n} + \mathbf{t} \times \mathbf{n}' = (k\mathbf{n}) \times \mathbf{n} + \mathbf{t} \times \mathbf{n}' = \mathbf{t} \times \mathbf{n}'.$$

由于 $\mathbf{n}'$ 可以分解为 $A\mathbf{t} + B\mathbf{n} + C\mathbf{b}$ （由正交性），且 $\|\mathbf{n}\| = 1$ 蕴含 $\langle \mathbf{n}', \mathbf{n} \rangle = 0$ ，故 $\mathbf{n}' = -k\mathbf{t} + \tau\mathbf{b}$ （这正是 Frenet-Serret 公式的第二式）。

于是

$$\mathbf{b}' = \mathbf{t} \times (-k\mathbf{t} + \tau\mathbf{b}) = \mathbf{t} \times \tau\mathbf{b} = \tau(\mathbf{t} \times \mathbf{b}) = -\tau\mathbf{n},$$

因为 $\mathbf{t} \times \mathbf{b} = -\mathbf{b} \times \mathbf{t} = -\mathbf{n}$ 。

因此 $\mathbf{b}' = -\tau\mathbf{n}$ ，取与 $\mathbf{n}$ 的内积即得

$$\tau = -\langle \mathbf{b}', \mathbf{n} \rangle.$$

Frenet-Serret 公式的直观解释与推导思路

背景：Frenet-Serret 公式是曲线局部理论的核心：

$$\begin{cases} \mathbf{t}' = k\mathbf{n}, \\ \mathbf{n}' = -k\mathbf{t} + \tau\mathbf{b}, \\ \mathbf{b}' = -\tau\mathbf{n}. \end{cases}$$

直观解释：

可以将 Frenet 标架 $\{\mathbf{t}, \mathbf{n}, \mathbf{b}\}$ 想象为沿曲线运动的“观察者”的坐标系。这个坐标系随着曲线在空间中移动而转动和摆动。

- $\mathbf{t}' = k\mathbf{n}$ ：切向量的变化完全由法向分量决定，系数 k 正是弯曲程度—— k 越大，切向量转动得越快。
- $\mathbf{n}' = -k\mathbf{t} + \tau\mathbf{b}$ ：主法向量的变化涉及两个方面： $-k\mathbf{t}$ 部分反映了主法向量试图“追回”切向量方向变化（与 $\mathbf{t}' = k\mathbf{n}$ 相呼应），而 $\tau\mathbf{b}$ 部分则反映了曲线在空间中的扭曲。
- $\mathbf{b}' = -\tau\mathbf{n}$ ：副法向量的变化只与挠率有关——它描述了密切平面的旋转。

几何图像：想象一个骑自行车的人沿着曲线运动。人的朝向就是 $\mathbf{t}$ 方向，车把的指向是 $\mathbf{n}$ ，而 $\mathbf{b}$ 则垂直于前两者。曲率 k 决定了人在转弯时需要倾斜多少（这对应 $\mathbf{t}' = k\mathbf{n}$ ），而挠率 τ 则描述了车子在转弯的同时是否还在上下颠簸（对应 $\mathbf{b}' = -\tau\mathbf{n}$ ）。

推导思路（简述）：

Frenet-Serret 公式可从正交性条件出发推导。由于 $\{\mathbf{t}, \mathbf{n}, \mathbf{b}\}$ 是标准正交基，有 $\langle \mathbf{t}, \mathbf{n} \rangle = 0$, $\langle \mathbf{t}, \mathbf{b} \rangle = 0$, $\langle \mathbf{n}, \mathbf{b} \rangle = 0$ ，以及 $\|\mathbf{t}\| = \|\mathbf{n}\| = \|\mathbf{b}\| = 1$ 。

对正交性条件求导，并利用 $\mathbf{b} = \mathbf{t} \times \mathbf{n}$ ，可以逐步得到三个公式。例如：

1. 由 $\langle \mathbf{t}, \mathbf{n} \rangle = 0$ 求导： $\langle \mathbf{t}', \mathbf{n} \rangle + \langle \mathbf{t}, \mathbf{n}' \rangle = 0$ 。
2. 由于 $\mathbf{t}' = \alpha''$ 与 $\mathbf{n}$ 平行（定义 $\mathbf{n} = \mathbf{t}'/|\mathbf{t}'|$ ），得 $\langle \mathbf{t}', \mathbf{n} \rangle = |\mathbf{t}'| = k$ 。
3. 结合 $\|\mathbf{n}'\|$ 的计算，可得 $\mathbf{t}' = k\mathbf{n}$ 和 $\mathbf{n}' = -k\mathbf{t} + \tau\mathbf{b}$ 。

详细证明见 do Carmo, Proposition 1-5, 4。