

BURNS & DERMATOLOGY MEDS

1. BURN MANAGEMENT MEDICATIONS

-  IV Fluids (Lactated Ringer's)
 - Used for: fluid resuscitation in major burns (especially first 24 hrs)
 - Side effects: (fluid overload, electrolyte imbalance)
-  Silver Sulfadiazine
 - Topical antibiotic for 2nd and 3rd-degree burns
 - Side effects: (painless application, can ↓ WBCs, not for face or pregnancy)
-  Mafenide (Sulfamylon)
 - Used for: electrical burns or eschar penetration
 - Side effects: (painful application, metabolic acidosis)
-  Tetanus Vaccine
 - Given if status unknown or outdated, to prevent tetanus infection in burns
 - Side effects: (fever, soreness)

2. SKIN INFECTION & WOUND CARE MEDS

-  Mupirocin (Bactroban)
 - Used for: impetigo, minor skin infections
 - Side effects: (local irritation)
-  Bacitracin
 - Used for: small cuts and abrasions
 - Side effects: (rash, contact dermatitis)
-  Topical corticosteroids (Hydrocortisone, Triamcinolone)
 - Used for: eczema, dermatitis, allergic rashes
 - Side effects: (skin thinning with long use, irritation)
-  Antihistamines (Diphenhydramine cream or oral)
 - Used for: allergic skin reactions, itching (e.g., eczema, hives)
 - Side effects: (drowsiness if oral, dryness)

3. FUNGAL SKIN INFECTIONS

-  Clotrimazole, Miconazole (topical antifungals)
 - Used for: athlete's foot, ringworm, diaper rash (yeast)
 - Side effects: (mild irritation or burning)
-  Oral Fluconazole
 - Used for: widespread fungal infections, tinea capitis, or candidiasis
 - Side effects: (GI upset, liver toxicity)

4. PARASITIC SKIN INFECTIONS

-  Permethrin 1% or 5% Cream
 - Used for: lice (1%) or scabies (5%)
 - Side effects: (burning, itching, redness after application)

SPECIAL NOTES FOR NCLEX:

Always check for allergies to sulfa (e.g., silver sulfadiazine, mafenide)

Burn dressing changes are painful → pre-medicate if needed

Monitor for signs of infection in burns (↑ temp, odor, green drainage)