

AUTOIMMUNE, INFECTIOUS & INFLAMMATORY DISORDER MEDICATION

SYSTEMIC LUPUS ERYTHEMATOSUS (SLE)

-  Corticosteroids (Prednisone)
 - Reduce severe inflammation
 - (↑ infection risk, hyperglycemia, osteoporosis)
-  NSAIDs (Ibuprofen, Naproxen)
 - For joint pain, fever
 - (GI bleeding, kidney issues — take with food)
-  Antimalarial (Hydroxychloroquine)
 - Prevents flares long-term
 - (Eye damage — need annual eye exams)
-  Immunosuppressants (Azathioprine, Mycophenolate mofetil)
 - For severe disease; steroid-sparing
 - (↑ infection risk, avoid live vaccines)
-  Biologics (Belimumab)
 - Targets B cells → reduce inflammation
 - (Injection reaction, possible depression risk)

LYME DISEASE

-  Doxycycline
 - First-line for most cases
 - (Photosensitivity, not for pregnant women or kids <8)
-  Amoxicillin or Cefuroxime
 - Used in pregnancy or children
 - (GI upset, rash)
-  IV Ceftriaxone
 - For late Lyme with neuro/cardiac signs
 - (Pain at injection site, diarrhea)



ANTHRAX

-  Ciprofloxacin or Doxycycline
 - Post-exposure prophylaxis or active infection
 - (GI upset, tendon risk [cipro], photosensitivity [doxy])
-  Anthrax Antitoxins (for inhalation anthrax)
 - Neutralizes anthrax toxins
 - (Given IV in severe cases — side effects vary)
-  Vaccine (for prevention in high-risk groups)
 - Military, lab workers
 - (Soreness at injection site)



OTHER AUTOIMMUNE DISEASES (FROM OVERVIEW)

RHEUMATOID ARTHRITIS

- Methotrexate (DMARD) → slows disease
 - (Liver toxicity, infection risk)
- NSAIDs → pain/inflammation
 - (GI bleeding, kidney strain)
- Steroids → short-term relief
 - (Hyperglycemia, immune suppression)

MULTIPLE SCLEROSIS

- Corticosteroids → flare control
 - (Mood swings, high blood sugar)
- Immunomodulators (e.g., interferons)
 - (Flu-like symptoms, injection site issues)

MYASTHENIA GRAVIS

- Pyridostigmine → improves muscle strength
 - (GI upset, cramps)
- Plasmapheresis / IVIG → short-term relief in crisis

GRAVES' DISEASE (HYPERTHYROIDISM)

- Methimazole or PTU → block thyroid hormone
→ (Liver damage, low WBCs)
- Beta-blockers → slow heart rate
→ (Bradycardia, fatigue)

HASHIMOTO'S (HYPOTHYROIDISM)

- Levothyroxine → replaces thyroid hormone
→ (Tachycardia, insomnia if overdosed)

TYPE 1 DIABETES

- Insulin → lifelong treatment
→ (Hypoglycemia, lipodystrophy)

CELIAC DISEASE

- Gluten-free diet only
→ (No medication, just strict dietary adherence)



GUILLAIN-BARRÉ SYNDROME (GBS)

-  IVIG (Intravenous Immunoglobulin)
→ Blocks damaging antibodies
→ (Headache, chills, rare kidney injury)
-  Plasmapheresis
→ Removes harmful antibodies from blood
→ (Bleeding, low BP, infection risk)
-  Anticoagulants (e.g., heparin)
→ Prevent DVT in immobile patients
→ (Bleeding risk)



SARS (SEVERE ACUTE RESPIRATORY SYNDROME)

-  IV Fluids
→ Prevent dehydration in pneumonia
→ (Fluid overload if heart/lung issues)

-  Acetaminophen
 - For fever and pain
 - (Liver toxicity with overdose)
-  Corticosteroids
 - May be used to reduce lung inflammation
 - (Suppress immunity, ↑ blood sugar)
-  Ribavirin (experimental antiviral)
 - Not widely effective; sometimes used in severe viral cases
 - (Anemia, birth defects — NOT for pregnancy)