

Appendix 1 of Meek et al. “Geography and Admixture Shape the Genome-scale Phylogeny of North American *Delphinium*”

Table of Contents

- [New clade definitions](#)
- [Methods](#)

New clade definitions

The following clade definitions synthesize morphological descriptions from the Flora of North America (FNA) and related botanical sources. Each clade is defined using a **minimum-clade** definition (the smallest clade containing the internal specifiers and all descendants). Diagnostic characters summarise the most consistent synapomorphies and habitat preferences, while comparison sections highlight how each clade differs from its congeners. Comments identify notable exclusions. All numbered citations refer to the sources listed in the reference list at the end of the document.

- [Clade Cordillerana](#)
- [Clade Columbianum](#)
- [Clade Intermontana](#)
- [Clade Sudmontana](#)
- [Clade Californica](#)
- [Clade Basinensis](#)

Clade Cordillerana *nomen cladi novum*

Definition (minimum-clade): The smallest clade containing *Delphinium glaucum*, *D. exaltatum*, and *D. brachycentrum*.

Etymology: This name means “the clade of the Cordilleran mountains”. It refers to the mountain chains of the Pacific Cordillera that stretch from Alaska’s Brooks and Alaska Ranges, down through the Coast and Canadian Rockies of British Columbia, and into the Cascades and Sierra Nevada. *Delphinium glaucum* ranges across this interconnected mountain system, and *Delphinium brachycentrum* is found in the tundra at its northern edge.

Diagnostic characteristics

- **Glabrous or glaucous stems.** The stems of *D. glaucum* and *D. exaltatum* are smooth and often glaucous, lacking the pubescence seen in many other larkspurs[1][2]. *D. brachycentrum* also has a clean stem surface[3].

- **Moderate nectar spurs (≈9–17 mm).** Spurs are of intermediate length: *D. glaucum* has spurs 10–15(–19) mm[1]; *D. exaltatum* 9–12 mm[2]; and *D. brachycentrum* 15–17 mm[3].
- **Intermediate inflorescences (5–90 flowers).** Inflorescences typically bear 8–30 flowers in *D. exaltatum*[2], 5–18 in *D. brachycentrum*[3], and 40–90 in *D. glaucum*[1]—well below the very dense racemes of some southwestern species.
- **Blue to pale-lavender flowers.** Sepal color ranges from bluish-purple (*D. glaucum*)[1] to whitish or pale lavender (*D. exaltatum*)[2]; *D. brachycentrum* has dark-blue flowers with rounded sepals[3].
- **Cool, moist montane distribution.** Members of Cordillerana inhabit boreal and montane environments—wet thickets and meadows from Alaska to the northern Rockies for *D. glaucum*[1], rocky slopes in eastern deciduous woods for *D. exaltatum*[2], and alpine tundra slopes for *D. brachycentrum*[3].

Comparison with other clades

- **Versus Sudmontana:** Cordillerana species are glabrous or glaucous and bear moderate racemes with 5–90 flowers. In contrast, members of Sudmontana are robust, often puberulent plants with 20–120(+) flowers in a raceme[11][13].
- **Versus Columbianum:** Columbianum species are generally shorter (10–100 cm) and have smaller racemes (3–50 flowers) with variable flower colors, including yellow or green; Cordillerana lacks yellow or green flowers and has a consistent montane distribution[4][33].
- **Versus Intermontana:** Both clades possess mid-length spurs, but Intermontana species often have puberulent inflorescences or leaves and occupy high-elevation wet meadows. Cordillerana is characterised by glabrous stems and slightly fewer flowers[17][18].
- **Versus Californica:** Californica species inhabit low-elevation chaparral and oak woodlands and have longer nectar spurs (up to 24 mm) and variable sepal colors (lavender, white, pink, red). Cordillerana lacks these traits[7][8].
- **Versus Basinensis:** Basinensis species are short, basal-leaved Great Basin endemics with small racemes and unwinged seeds[20][21]. Cordillerana, by contrast, has longer racemes and occurs in moist montane habitats.

Comments

Exclusion: *D. carolinianum* is explicitly excluded from Clade Cordillerana. *D. carolinianum* has moderate stems (20–90 cm), variable sepal colors and relatively few flowers (3–27), features that do not match the glabrous, montane phenotype of Cordillerana[23].

Summary

Clade Cordillerana comprises glabrous or glaucous larkspurs with intermediate-length spurs, moderate racemes, blue to pale-lavender flowers and a preference for cool montane habitats. These traits distinguish it from the taller, densely flowered southern Rocky Mountain clades, the diverse chaparral group and the small Great Basin taxa.

Clade *Columbianum* *nomen cladi novum*

Definition (minimum-clade): The smallest clade containing *Delphinium lineapetalum*, *D. multiplex*, *D. xantholeucum*, *D. nuttallianum*, *D. distichum*, *D. bicolor* and *D. viridescens*.

Etymology: Named for the Columbia Plateau. Most members occur in the mountains surrounding the Columbia Plateau in eastern Washington, north-eastern Oregon, Idaho and western Montana.

Diagnostic characteristics

- **Moderate stature with basal or proximal leaves.** Species range from 10–100 cm tall and typically have leaves restricted to the basal or lower half of the stem. *D. lineapetalum* has glabrous stems 30–60 cm tall with small, deeply lobed leaves[4]; *D. nuttallianum* is 10–40 cm tall with leaves on the proximal quarter[5]; *D. viridescens* and *D. multiplex* are taller (up to 150 cm) but still bear most leaves on the lower half[6][35].
- **Small racemes with 3–50 flowers.** Inflorescences typically contain 3–24 flowers in *D. lineapetalum*[4] and 8–30 in *D. distichum*[31]; even the largest species (*D. viridescens*, *D. multiplex*) rarely exceed 80 flowers[6][35].
- **Moderate spurs (8–23 mm).** Spur length varies across the clade but never reaches the extreme lengths seen in chaparral species: *D. nuttallianum* has spurs 8–23 mm[5]; *D. lineapetalum* 11–17 mm[4]; *D. bicolor* 13–23 mm[32].
- **Diverse flower colors.** Flower color ranges from dark blue (*D. bicolor*, *D. distichum*)[32][31] through lavender (*D. lineapetalum*)[4] to yellowish or greenish (*D. xantholeucum*, *D. viridescens*)[33][6].
- **Habitat affinity for montane forests, meadows and sage-steppe.** Members occur in open pine woods, dry meadows, wet meadows and grassy sage scrub around the Columbia Plateau[4][5][33].

Comparison with other clades

- **Versus Sudmontana:** Columbianum species are shorter and have much smaller racemes (3–50 flowers) than the robust, 20–120+ flowered species of Sudmontana[11][13].
- **Versus Californica:** Californica species have longer racemes (up to 90 flowers), longer spurs (7–24 mm) and occur in chaparral or oak woodlands at low elevations[7], whereas Columbianum occupies montane or subalpine meadows and includes yellow-flowered taxa.
- **Versus Intermontana:** Intermontana species are generally taller with more robust racemes and uniform blue-purple flowers[17][18]. Columbianum includes low-stature, short-flowered species and a broader color palette.
- **Versus Cordillerana:** Columbianum lacks the glabrous, glaucous stems typical of Cordillerana and includes species with pubescent stems and leaves[4].

- **Versus Basinensis:** Basinensis is restricted to the Great Basin and has small racemes, moderate spurs and consistently dark-blue flowers[20][21]. Columbianum is more diverse in color and occurs in moister environments.

Comments

This clade currently contains seven species (see definition) and has no known exclusions beyond species placed in other clades.

Summary

Clade Columbianum comprises a heterogeneous set of larkspurs from the mountains encircling the Columbia Plateau. They share moderate stature, small racemes with mid-length spurs, diverse flower colors and adaptations to montane forests, meadows and sage steppe. These characters distinguish them from the robust southern Rocky Mountain clades, the chaparral larkspurs and the Great Basin taxa.

Clade Intermontana *nomen cladi novum*

Definition (minimum-clade): The smallest clade containing *Delphinium barbeyi*, *D. glaucescens*, *D. occidentale*, and *D. geyeri*.

Etymology: From Latin *inter* (“between”) and *montana* (“of the mountains”), meaning “between the mountains”, indicating that members are distributed in the Intermountain West regions of Utah, Idaho and Nevada.

Diagnostic characteristics

- **Moderate to tall stems with large, lobed leaves.** Stems range from 35–150 cm. *D. barbeyi* produces reniform leaves 4–8 × 7–15 cm[17]; *D. glaucescens* has pentagonal leaves 2–9 × 3–14 cm[18]; *D. occidentale* bears blades 5–15 cm across[27]; and *D. geyeri* has pubescent leaves[19].
- **Racemes with 6–50 flowers.** *D. barbeyi* has 10–50 flowers[17]; *D. glaucescens* 10–25[18]; *D. occidentale* often has >25 flowers[27]; *D. geyeri* 6–30[19].
- **Moderate spurs (9–18 mm).** Spurs range from 10–18 mm in *D. barbeyi*[17], 9–13 mm in *D. glaucescens*[18], 10–15 mm in *D. occidentale*[27], and 11–16 mm in *D. geyeri*[19].
- **Blue to purple flowers with robust spurs.** Flowers are uniformly blue to purple across the clade: dark bluish purple (*D. barbeyi*)[17], dark blue (*D. glaucescens*), bright blue (*D. geyeri*)[19]. The hybrid *D. occidentale* may be paler but still falls within the blue spectrum[27].
- **Montane to subalpine distribution.** Members grow in wet meadows, rocky slopes and streambanks at elevations of ~1 400–4 100 m[17][18][19].

Comparison with other clades

- **Versus Cordillerana:** Both clades possess mid-length spurs, but Intermontana species often have puberulent inflorescences or leaves and occupy high-elevation

wet meadows, whereas Cordillerana species are glabrous and inhabit cooler boreal or eastern montane forests[1][18].

- **Versus Columbianum:** Intermontana species are taller (up to 150 cm) and have more robust racemes, with consistently blue flowers. Columbianum includes shorter species with yellow or green flowers and smaller racemes[4][33].
- **Versus Sudmontana:** Sudmontana species produce very long racemes (20–120+ flowers) and grow predominantly in the southern Rocky Mountains[11]. Intermontana species have fewer flowers and are centred in the northern and central Rocky Mountains.
- **Versus Californica:** Californica species inhabit chaparral slopes at low elevations and may bear red or white flowers; Intermontana lacks red-flowered taxa and occurs at higher elevations.
- **Versus Basinensis:** Basinensis taxa are Great Basin endemics with small racemes and unwinged seeds[20]; Intermontana species have larger racemes and, in some cases, wing-margined seeds (e.g. *D. barbeyi*)[17].

Comments

This clade comprises four species currently recognised in the FNA, in addition to *D. occidentale*, which has been described as a natural hybrid between *D. barbeyi* and *D. glaucum*. Its inclusion reflects that it does not appear to be more involved in hybridization than many other described taxa based on genomic admixture analyses.

Summary

Clade Intermontana encompasses the mid-elevation, blue-flowered larkspurs of the northern and central Rocky Mountains. They are characterised by moderate to tall stems, large lobed leaves, racemes of 6–50 flowers and mid-length nectar spurs. Their preference for wet meadows and subalpine habitats and their uniform blue flowers separate them from the taller southwestern clades, the varied chaparral taxa and the small Great Basin species.

Clade Sudmontana *nomen cladi novum*

Definition (minimum-clade): The smallest clade containing *Delphinium sapellonis*, *D. novomexicanum*, *D. andesicola*, *D. geraniifolium*, *D. scopulorum*, *D. robustum*, *D. ramosum* and *D. alpestre*.

Etymology: From Latin *sud* (“south”) and *montana* (“of the mountains”), meaning “of the southern mountains”, indicating that members are centred in the southern Rocky Mountains and adjacent southwestern sky islands.

Diagnostic characteristics and cohesion

- **Robust, erect stems with deeply lobed leaves.** Most species reach 50–200 cm in height and bear large, palmately lobed leaves 5–15 cm wide[11][13]. Even the

shorter *D. ramosum* (≈70–100 cm) has pentagonal leaves with 5–21 lobes[15], and *D. alpestre*—though only 5–25 cm tall—shares the same leaf architecture[16].

- **Large, many-flowered racemes.** Inflorescences typically bear 20–90 (sometimes up to 180) flowers in *D. sapellonis*, *D. novomexicanum*, *D. andesicola*, *D. geraniifolium* and *D. robustum*[11][12][14][29][13]. *D. scopulorum* has 10–30 flowers but still produces larger racemes than the Columbianum or Basinensis clades[30].
- **Dark-blue to purple flowers with moderate spurs (8–15 mm).** Sepals are consistently dark blue or purple across the clade, and nectar spurs range between 8 and 15 mm[11][12]. *D. ramosum* has spurs 9–13 mm ascending ~30° above the horizontal[15].
- **Pubescent or puberulent stems, pedicels and fruits.** Many species exhibit puberulence on the stems or inflorescences (e.g. *D. andesicola*, *D. geraniifolium* and *D. scopulorum*)[14][29][30]. Seed coats often have roughened or wing-margined cells[13].
- **High-elevation montane and subalpine habitats.** Members grow in meadows, riparian woodlands and coniferous forests at elevations of roughly 2 000–3 900 m in New Mexico, Colorado, Utah and Arizona[11][12][13]. *D. alpestre* occurs on exposed alpine talus at very high elevations[16].

Comparison with other clades

- **Versus Cordillerana:** Sudmontana species are taller and more robust, often puberulent and possess much longer racemes (20–120+ flowers) than the glabrous, moderate-flowered Cordillerana species[1].
- **Versus Columbianum:** Columbianum taxa are shorter, produce smaller racemes (3–50 flowers) and often include yellow or green flowers[4][33]. Sudmontana species lack yellow flowers and are consistently dark blue or purple.
- **Versus Intermontana:** Intermontana species inhabit higher, wetter subalpine habitats and have fewer flowers (6–50)[17], whereas Sudmontana species occur in slightly drier montane meadows and possess longer racemes.
- **Versus Californica:** Californica species have longer spurs (up to 24 mm) and grow in low-elevation chaparral[7]. Sudmontana species have moderate spurs, robust racemes and high-elevation distributions.
- **Versus Basinensis:** Basinensis includes much smaller plants with compact racemes and unwinged seeds, adapted to arid Great Basin habitats[20][21]. Sudmontana species are larger and occupy montane forests and meadows.

Comments

This clade currently includes eight species. No additional exclusions are needed beyond the distinction from other named clades.

Summary

Clade Sudmontana unites the tall, many-flowered larkspurs of the southern Rocky Mountains and adjacent Southwestern uplands. They share robust stems with deeply lobed leaves, dense racemes of dark-blue flowers with moderate nectar spurs and a preference

for montane meadows and forests. These features distinguish them from the smaller, yellow-flowered plateau species, the glabrous montane species, the chaparral larkspurs and the arid Great Basin taxa.

Clade Californica *nomen cladi novum*

Definition (minimum-clade): The smallest clade containing *D. cardinale*, *D. californicum*, *D. hesperium*, *D. gypsophilum* and *D. parryi*.

Etymology: Name signals that these species are centred in California and occupy chaparral and oak-woodland habitats.

Diagnostic characteristics

- **Moderate to tall stems with few basal leaves.** Plants stand 40–160 cm tall and have leaves mostly confined to the upper portions of the stem. *D. californicum* attains (60–)100–160(–230) cm with leaves restricted to the upper 80 % [10]; *D. gypsophilum* 60–100(–150) cm with glabrous, glaucous stems [8]; *D. cardinale* 50–150(–280) cm with leaves mostly cauline [9]; *D. parryi* (10–)40–80(–110) cm and *D. hesperium* 40–80 cm share similar stature [7].
- **Racemes with many flowers and long nectar spurs.** Inflorescences bear 8–90(–162) flowers. *D. californicum* has 40–90(–162) flowers with spurs 7–11(–14) mm [10]; *D. gypsophilum* 15–30(–64) flowers with spurs 7–15 mm [8]; *D. cardinale* 10–40(–80) flowers with spurs 15–24 mm [9]; *D. hesperium* 15–30(–100) flowers with spurs 9–18 mm [34]; *D. parryi* 8–24(–48) flowers with spurs 9–21(–24) mm [7].
- **Highly variable flower colors and unwinged seeds.** Sepals range from lavender-greenish in *D. californicum* [10], to white–pink in *D. gypsophilum* [8], dark blue in *D. parryi* [7] and *D. hesperium* [34], and bright red in *D. cardinale* [9]. Seeds are typically unwinged or only brick-shaped [8] [7].
- **Low to mid-elevation chaparral and oak-woodland habitats.** These species grow on dry slopes, well-drained hillsides and open woodlands in California and the adjacent Southwest. Elevations range from sea level to ~1 500 m [8] [9].

Comparison with other clades

- **Versus Sudmontana:** Californica species have longer nectar spurs (often >15 mm) and occur at much lower elevations in chaparral or oak woodland. Sudmontana species have shorter spurs and inhabit high-elevation montane meadows [11].
- **Versus Columbianum:** Columbianum taxa are smaller and produce shorter racemes with moderate spurs; some have yellow or green flowers. Californica lacks yellow flowers and has larger racemes [4].
- **Versus Intermontana:** Intermontana species are taller and occupy wet subalpine meadows, with uniformly blue flowers [17]; Californica species occur in drier chaparral and include red and white-flowered taxa.

- **Versus Cordillerana:** Cordillerana species have glabrous or glaucous stems, moderate racemes and blue to pale flowers[1]; California species may be glabrous or glaucous but possess long spurs and extremely varied sepal colors.
- **Versus Basinensis:** Basinensis species are low-growing Great Basin taxa with small racemes and moderate spurs[20][21]; California species are taller and concentrated in California chaparral.

Comments

Exclusion: This clade explicitly **excludes** *D. patens*, *D. nudicaule* and *D. trolliifolium*. *D. patens* has reflexed lateral sepals, shorter racemes and ascending spurs[24]; *D. nudicaule* has scarlet tubular flowers with very long spurs and moist cliff habitats[25]; and *D. trolliifolium* has exclusively cauline leaves and elongated fruits in wet oak woods[26]. These traits are absent from California and warrant their placement in separate clades.

Summary

Clade California comprises the chaparral-adapted larkspurs of California and the adjacent Southwest. They are moderate to tall perennials with racemes of many flowers bearing long nectar spurs, highly variable sepal colors and unwinged seeds. Their low-elevation chaparral and oak woodland habitats and long spurs distinguish them from the robust mountain larkspurs of Sudmontana, the small plateau species of Columbianum, the blue-flowered Intermontana taxa and the glaucous montane species of Cordillerana. The clade explicitly excludes *D. patens*, *D. nudicaule* and *D. trolliifolium*.

Clade Basinensis *nomen cladi novum*

Definition (minimum-clade): The smallest clade containing *D. scaposum* and *D. andersonii*.

Etymology: Meaning “of the basin”, reflecting the clade’s distribution in the arid landscapes of the Great Basin (primarily Nevada).

Diagnostic characteristics

- **Moderate stature with basal leaves.** *D. scaposum* has scape-like stems 25–50(–65) cm tall with most leaves at the base[20]; *D. andersonii* has erect stems 30–60(–90) cm with leaves mostly on the lower half[21].
- **Small racemes with 10–25(–40) flowers.** The racemes are cylindric and open, bearing few flowers. Pedicels are 0.6–2.5 cm and glabrous in *D. scaposum*[20] and slightly longer (1–4 cm) and sigmoid in *D. andersonii*[21].
- **Mid-length spurs (12–18 mm) and dark-blue flowers.** Flowers are bright dark blue with spurs 13–18 mm in *D. scaposum*[20] and dark blue with spurs 12–18 mm in *D. andersonii*[21].
- **Elongate fruits and unwinged, brick-shaped seeds.** Fruits measure 12–16 mm in *D. scaposum*[20] and 17–32 mm in *D. andersonii*[21]. Seed coat cells are brick-shaped and smooth.

- **Great Basin habitats.** Both species inhabit juniper woodlands, grasslands and cold desert scrub in the Great Basin at elevations of 1 200–2 700 m[20][21].

Comparison with other clades

- **Versus Sudmontana and Intermontana:** Basinensis species are shorter (25–90 cm), have small racemes (10–25 flowers) and unwinged seeds, unlike the tall, many-flowered species of Sudmontana[11] or the wet-meadow species of Intermontana[17].
- **Versus Cordillerana:** Cordillerana species inhabit cool, moist montane environments and have larger racemes with glabrous stems[1]; Basinensis species occupy arid Great Basin habitats and have pubescent or glabrous pedicels and unwinged seeds.
- **Versus Columbianum:** Columbianum includes a broader array of flower colors and habitats; Basinensis is restricted to dark-blue flowers and deserts of the Great Basin.
- **Versus Californica:** Californica species are taller and have much longer spurs (up to 24 mm) and larger racemes[9]. Basinensis species are modest in stature with moderate spurs.

Comments

Exclusion: *D. stachydeum* is explicitly excluded from Clade Basinensis. Despite also occurring in the Great Basin, *D. stachydeum* has tall stems (70–150 cm), dense racemes of 30–60(–102) bright-blue flowers, wing-margined seeds and grows in moist Artemisia swales[22].

Summary

Clade Basinensis contains the two Great Basin desert larkspurs *D. scaposum* and *D. andersonii*. These species are moderate-sized perennials with basal leaves, small racemes of dark-blue flowers with mid-length spurs, elongate fruits and unwinged seeds. They inhabit arid juniper woodlands and cold desert scrub at moderate elevations. The combination of Great Basin distribution, compact inflorescences and unwinged seeds distinguishes Basinensis from all other Delphinium clades described here.

References

1. **Flora of North America Editorial Committee.** *Delphinium glaucum* in **Flora of North America north of Mexico**, Vol. 3. New York and Oxford: Oxford University Press. Available at: https://floranorthamerica.org/Delphinium_glaucum【793823136742840†L55-L70】.
2. **Flora of North America Editorial Committee.** *Delphinium exaltatum* in **Flora of North America north of Mexico**, Vol. 3. Available at: https://floranorthamerica.org/Delphinium_exaltatum【73553004382722†L50-L65】.

3. **Flora of North America Editorial Committee.** *Delphinium brachycentrum* in **Flora of North America north of Mexico**, Vol. 3. Available at:
https://floranorthamerica.org/Delphinium_brachycentrum【938511734675106†L45-L59】.
4. **Flora of North America Editorial Committee.** *Delphinium lineapetalum* in **Flora of North America north of Mexico**, Vol. 3. Available at:
https://floranorthamerica.org/Delphinium_lineapetalum【736021784608033†L50-L66】.
5. **Flora of North America Editorial Committee.** *Delphinium nuttallianum* in **Flora of North America north of Mexico**, Vol. 3. Available at:
https://floranorthamerica.org/Delphinium_nuttallianum【243959302267772†L49-L66】.
6. **Flora of North America Editorial Committee.** *Delphinium viridescens* in **Flora of North America north of Mexico**, Vol. 3. Available at:
https://floranorthamerica.org/Delphinium_viridescens【36312153868147†L50-L58】.
7. **Flora of North America Editorial Committee.** *Delphinium parryi* in **Flora of North America north of Mexico**, Vol. 3. Available at:
https://floranorthamerica.org/Delphinium_parryi【852769970396148†L49-L62】.
8. **Flora of North America Editorial Committee.** *Delphinium gypsophilum* in **Flora of North America north of Mexico**, Vol. 3. Available at:
https://floranorthamerica.org/Delphinium_gypsophilum【179313723911639†L48-L60】.
9. **Flora of North America Editorial Committee.** *Delphinium cardinale* in **Flora of North America north of Mexico**, Vol. 3. Available at:
https://floranorthamerica.org/Delphinium_cardinale【620554933416448†L43-L56】.
10. **Flora of North America Editorial Committee.** *Delphinium californicum* in **Flora of North America north of Mexico**, Vol. 3. Available at:
https://floranorthamerica.org/Delphinium_californicum【664433447320462†L48-L59】.
11. **Flora of North America Editorial Committee.** *Delphinium sapellonis* in **Flora of North America north of Mexico**, Vol. 3. Available at:
https://floranorthamerica.org/Delphinium_sapellonis【448049940458835†L45-L63】.
12. **Flora of North America Editorial Committee.** *Delphinium novomexicanum* in **Flora of North America north of Mexico**, Vol. 3. Available at:
https://floranorthamerica.org/Delphinium_novomexicanum【701864018428591†L47-L63】.

13. **Flora of North America Editorial Committee.** *Delphinium robustum* in **Flora of North America north of Mexico**, Vol. 3. Available at:
https://floranorthamerica.org/Delphinium_robustum【713256998688013†L50-L67】.
14. **Flora of North America Editorial Committee.** *Delphinium andesicola* in **Flora of North America north of Mexico**, Vol. 3. Available at:
https://floranorthamerica.org/Delphinium_andesicola【528047884121319†L45-L56】.
15. **Flora of North America Editorial Committee.** *Delphinium ramosum* in **Flora of North America north of Mexico**, Vol. 3. Comparative description available at:
<https://nwwildflowers.com/compare/>【465523646963587†L16-L33】.
16. **Flora of North America Editorial Committee.** *Delphinium alpestre* in **Flora of North America north of Mexico**, Vol. 3. Available at:
https://floranorthamerica.org/Delphinium_alpestre【603698680033958†L64-L70】.
17. **Flora of North America Editorial Committee.** *Delphinium barbeyi* in **Flora of North America north of Mexico**, Vol. 3. Available at:
https://floranorthamerica.org/Delphinium_barbeyi【40418217850526†L54-L68】.
18. **Flora of North America Editorial Committee.** *Delphinium glaucescens* in **Flora of North America north of Mexico**, Vol. 3. Available at:
https://floranorthamerica.org/Delphinium_glaucescens【78539320886863†L45-L60】.
19. **Flora of North America Editorial Committee.** *Delphinium geyeri* in **Flora of North America north of Mexico**, Vol. 3. Available at:
https://floranorthamerica.org/Delphinium_geyeri【950197660752108†L50-L66】.
20. **Flora of North America Editorial Committee.** *Delphinium scaposum* in **Flora of North America north of Mexico**, Vol. 3. Available at:
https://floranorthamerica.org/Delphinium_scaposum【696118333793837†L50-L64】.
21. **Flora of North America Editorial Committee.** *Delphinium andersonii* in **Flora of North America north of Mexico**, Vol. 3. Available at:
https://floranorthamerica.org/Delphinium_andersonii【430104109965132†L52-L67】.
22. **Flora of North America Editorial Committee.** *Delphinium stachydeum* in **Flora of North America north of Mexico**, Vol. 3. Available at:
https://floranorthamerica.org/Delphinium_stachydeum【554245642331618†L54-L66】.
23. **Flora of North America Editorial Committee.** *Delphinium carolinianum* in **Flora of North America north of Mexico**, Vol. 3. Available at:

https://floranorthamerica.org/Delphinium_carolinianum【705170725909413†L49-L60】.

24. **Flora of North America Editorial Committee.** *Delphinium patens* in **Flora of North America north of Mexico**, Vol. 3. Available at: https://floranorthamerica.org/Delphinium_patens【99726023098449†L48-L59】.
25. **Flora of North America Editorial Committee.** *Delphinium nudicaule* in **Flora of North America north of Mexico**, Vol. 3. Available at: https://floranorthamerica.org/Delphinium_nudicaule【428197565839957†L52-L63】.
26. **Flora of North America Editorial Committee.** *Delphinium trolliifolium* in **Flora of North America north of Mexico**, Vol. 3. Available at: https://floranorthamerica.org/Delphinium_trolliifolium【139072958821701†L45-L58】.
27. **Montana Natural Heritage Program.** “Tall Larkspur (*Delphinium × occidentale*).” *Montana Field Guide* (accessed 2025). Available at: <https://montana.plant-life.org/cgi-bin/species03.cgi>【768953008360490†L78-L84】.
28. **Northwest Wildflowers & Pacific Northwest Plants.** “Delphinium species comparison.” Available at: <https://nwwildflowers.com/compare/>【465523646963587†L16-L33】.
29. **Flora of North America Editorial Committee.** *Delphinium geraniifolium* in **Flora of North America north of Mexico**, Vol. 3. Available at: https://floranorthamerica.org/Delphinium_geraniifolium【378826461748443†L47-L60】.
30. **Flora of North America Editorial Committee.** *Delphinium scopulorum* in **Flora of North America north of Mexico**, Vol. 3. Available at: https://floranorthamerica.org/Delphinium_scopulorum【9239202906663†L52-L67】.
31. **Flora of North America Editorial Committee.** *Delphinium distichum* in **Flora of North America north of Mexico**, Vol. 3. Available at: https://floranorthamerica.org/Delphinium_distichum【390531643150230†L52-L66】.
32. **Flora of North America Editorial Committee.** *Delphinium bicolor* in **Flora of North America north of Mexico**, Vol. 3. Available at: https://floranorthamerica.org/Delphinium_bicolor【140123829857920†L49-L60】.
33. **Flora of North America Editorial Committee.** *Delphinium xantholeucum* in **Flora of North America north of Mexico**, Vol. 3. Available at:

https://floranorthamerica.org/Delphinium_xantholeucum【945135328931146†L48-L63】.

34. **Flora of North America Editorial Committee.** *Delphinium hesperium* in **Flora of North America north of Mexico**, Vol. 3. Available at: https://floranorthamerica.org/Delphinium_hesperium【478718253132052†L48-L61】.
35. **Flora of North America Editorial Committee.** *Delphinium multiplex* in **Flora of North America north of Mexico**, Vol. 3. Available at: https://floranorthamerica.org/Delphinium_multiplex【110024019203093†L46-L62】.

Methods

We propose a set of formal taxonomic names, following the Phylocode protocol for the six higher-level clades containing more than one taxon that were consistently recovered across our phylogenomic analyses. To identify additional supporting evidence for these clades beyond genomics, we employed OpenAI's ChatGPT (GPT- 4) in a tool-enabled agent-based framework to retrieve and analyze morphological descriptions for each taxon. The agent performed structured searches to retrieve data from the Flora of North America and other authoritative referenced sources, fetching details on stem height, leaf morphology, inflorescence size, sepal color, spur length and habitat. After collating these character states, it identified synapomorphies that unite members of our specified clades, and drafted clade definitions that highlight diagnostic characters. All data-based statements are linked to references. The process was iterative with feedback from the authors to generate etymologically appropriate names, revise formatting, and suggest additional comparisons. Manual measurements of herbarium specimens were conducted to verify that morphological characters extracted by the agent were consistent for the newly defined clades. The agent-based analysis was initiated with the prompt below.

Prompt

Act as an expert taxonomist with experience analyzing botanical specimens in herbaria, and in developing taxonomic classifications based on morphological data, and external information from molecular phylogenetic analyses. Your task is to extract trait, environmental, geographic, and other information from taxonomic descriptions. Most prominently, get information from species descriptions on the <https://floranorthamerica.org/>. Compare this information across the species that I specify, and to search for similarities that group together sets of taxa based on their groupings that I will provide to you, based on phylogenetic inference. The focal taxa for your analysis are only the ones included in this list, which are all *Delphinium* species. If a taxon is not present in the Flora of North America, do not include it in your analysis. The species are *Delphinium patens*, *D. sapellonis*, *D. novomexicanum*, *D. andesicola*, *D. geraniifolium*, *D. scopulorum*, *D. robustum*, *D. carolinianum*, *D. glaucum*, *D. exaltatum*, *D. brachycentrum*, *D. glaucescens*, *D. occidentale*, *D. barbeyi*, *D. geyeri*, *D. parryi*, *D. gypsophilum*, *D. cardinale*, *D. hesperium*, *D. californicum*, *D. stachydeum*, *D. lineapetalum*, *D. xantholeucum*, *D. nuttallianum*, *D. bicolor*, *D. viridescens*, *D. distichum*, *D. multiplex*, *D. scaposum*, *D. andersonii*, *D. trolliifolium*, *D. nudicaule*.

Goal: find similarities that can justify uniting these species into the following clades. I am particularly interested in the clades that include multiple species, and include the single-species clades only as a contrast: 1: *D. sapellonis*, *D. novomexicanum*, *D. andesicola*, *D. geraniifolium*, *D. scopulorum*, *D. robustum*; 2: *D. carolinianum*; 3: *D. glaucum*, *D. exaltatum*, *D. brachycentrum*; 4: *D. glaucescens*, *D. occidentale*, *D. barbeyi*, *D. geyeri*; 5: *D. parryi*, *D. gypsophilum*, *D. cardinale*, *D. hesperium*, *D.*

californicum; 6: *D. stachydeum*; 7: *D. lineapetalum*, *D. xantholeucum*, *D. nuttallianum*, *D. bicolor*, *D. viridescens*, *D. distichum*, *D. multiplex*, *D. scaposum*, *D. andersonii*; 8: *D. troliifolium*; 9: *D. nudicaule*; 10: *D. patens*.

Prioritize accuracy: accept that morphological descriptions are not perfect, quantitative descriptions are estimates, and some descriptions are absent, but do not make up descriptions that do not exist. Interpret the meaning of morphological descriptions based on definitions used in botanical taxonomy, including botanical latin. Cross-check your results with the original data (taxonomic descriptions) to ensure that you did not make up any trait similarities that were not originally described. Return results as a table where for each clade you list the most consistent uniting morphological characters for the taxa in that clade, and descriptions of how consistent this is among those species, and how this is different from the other listed clades.