

# CMPT 210: Probability and Computing

## Lecture 4

---

Sharan Vaswani

January 18, 2024

Ask about the degree of a vertex in an undirected graph where a self loop is contained.

**Number of ways of choosing size  $k$ -subsets from a size  $n$ -set:**  $\binom{n}{k}$  (E.g. Number of  $n$ -bit sequences with exactly  $k$  ones).

**Binomial Theorem:** For all  $n \in \mathbb{N}$  and  $a, b \in \mathbb{R}$ ,  $(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^{n-k} b^k$ .

# Generalization to Multinomials

We saw how to split a set into two subsets - one that contains some elements, while the other does not. Can generalize the arguments to split a set into more than two subsets.

A  $(k_1, k_2, \dots, k_m)$ -split of set  $A$  is a sequence of sets  $(A_1, A_2, \dots, A_m)$  s.t. sets  $A_i$  form a partition ( $A_1 \cup A_2 \cup \dots = A$  and for  $i \neq j$ ,  $A_i \cap A_j = \emptyset$ ) and  $|A_i| = k_i$ .

An example of a  $(2, 1, 3)$ -split of  $A = \{1, 2, 3, 4, 5, 6\}$  is  $(\{2, 4\}, \{1\}, \{3, 5, 6\})$ . Here,  $m = 3$ ,  $A_1 = \{2, 4\}$ ,  $A_2 = \{1\}$ ,  $A_3 = \{3, 5, 6\}$  s.t.  $|A_1| = 2$ ,  $|A_2| = 1$ ,  $|A_3| = 3$ ,  $A_1 \cup A_2 \cup A_3 = A$  and for  $i \neq j$ ,  $A_i \cap A_j = \emptyset$ .

*Example:* Consider strings of length 6 of  $a$ 's,  $b$ 's and  $c$ 's such that number of  $a$ 's = 2; number of  $b$ 's = 1 and number of  $c$ 's = 3. Possible strings: abaccc, ccbaac, bacacc, cbacac.

Each possible string, e.g., bacacc can be written as a  $(2, 1, 3)$ -split of  $A = \{1, 2, 3, 4, 5, 6\}$  as  $(\{2, 4\}, \{1\}, \{3, 5, 6\})$  where  $A_1$  records the positions of  $a$ ,  $A_2$  records the positions of  $b$  and  $A_3$  records the positions of  $c$ .   
 This maps to "bacacc"

Since each set is disjoint, union of all the sets = the original set

# Generalization to Multinomials

**Q:** Show that the number of ways to obtain an  $(k_1, k_2, \dots, k_m)$  split of  $A$  with  $|A| = n$  is  $\binom{n}{k_1, k_2, \dots, k_m} = \frac{n!}{k_1! k_2! \dots k_m!}$  where  $\sum_i k_i = n$ .

Can map any permutation  $(a_1, a_2, \dots, a_n)$  into a split by selecting the first  $k_1$  elements to form set  $A_1$ , next  $k_2$  to form set  $A_2$  and so on. For the same split, the order of the elements in each subset does not matter. Hence  $f$  : number of permutations  $\rightarrow$  number of splits is a  $k_1! k_2! \dots k_m!$ -to-1 function. Ask Sharan about the  $k$  to 1 function.

$$\text{Hence, } |\text{number of splits}| = \frac{|\text{number of permutations}|}{k_1! k_2! \dots k_m!} = \frac{n!}{k_1! k_2! \dots k_m!}.$$

you have  $k_1 = |A_1| = 3$

$$k_2 = |A_2| = 2$$

$$k_3 = |A_3| = 1$$

## Generalization to Multinomials - Example

**Q:** Count the number of permutations of the letters in the word BOOKKEEPER.

We want to count sequences of the form  $(1E, 1P, 2E, 1B, 1K, 1R, 2O, 1K) = EPEEBKROOK$ .

There is a bijection between such sequences and  $(1, 2, 2, 3, 1, 1)$  split of  $A = \{1, 2, \dots, 10\}$  where  $A_1$  is the set of positions of  $B$ 's,  $A_2$  is the set of positions of  $O$ 's,  $A_3$  is set of positions of  $K$  and so on.

Since there is a subscript, can we assume  $a_1$  is distinct from  $a_2$ ?  
For example, the above sequence maps to the following split:

$(\{5\}, \{8,9\}, \{6, 10\}, \{1,3,4\}, \{2\}, \{7\})$

Yes

Hence, the total number of sequences that can be formed from the letters in BOOKKEEPER = number of  $(1, 2, 2, 3, 1, 1)$  splits of  $A = [10] = \{1, 2, \dots, 10\} = \frac{10!}{1! 2! 2! 3! 1! 1!}$ .

**Q:** Count the number of permutations of the letters in the word (i) ABBA (ii)  $A_1 B B A_2$  and (iii)  $A_1 B_1 B_2 A_2$ ?

i:  $4!/2!2!$

ii:  $4!/2!$

Text

iii:  $4!$

## Generalization to Multinomials - Example

Each walk can be written as a (5, 5, 5, 5) split of the set.

Q: Suppose we are planning a 20 km walk, which should include 5 northward km, 5 eastward km, 5 southward km, and 5 westward km. We can move in steps of 1 km in any direction. For example, a valid walk is (*NENWSNSSSENSWWESWEENW*) that corresponds to 1 km north followed by 1 km east and so on. How many different walks are possible?

$$20!/(5!)^4$$

# Multinomial Theorem

For all  $m, n \in \mathbb{N}$  and  $z_1, z_2, \dots, z_m \in \mathbb{R}$ , **multinomial coefficient.**

$$(z_1 + z_2 + \dots + z_m)^n = \sum_{\substack{k_1, k_2, \dots, k_m \\ k_1 + k_2 + \dots + k_m = n}} \binom{n}{k_1, k_2, \dots, k_m} z_1^{k_1} z_2^{k_2} \dots z_m^{k_m}$$

**$k_i$  represents the coefficient of the  $z_i$  term.**

where  $\binom{n}{k_1, k_2, \dots, k_m} = \frac{n!}{k_1! k_2! \dots k_m!}$ . **Sum of all  $k_i = n$**

*Example 1:* If  $m = 2$ ,  $k_1 = k$ ,  $k_2 = n - k$  and  $z_1 = a$ ,  $z_2 = b$ , recover the Binomial theorem.

*Example 2:* If  $n = 4$ ,  $m = 3$ , then the coefficient of  $abc^2$  in  $(a + b + c)^4$  is  $\binom{4}{1, 1, 2} = \frac{4!}{1! 1! 2!}$ .

**the coefficient of  $ab$  in  $(a + b + c)(a + b + c)$  is**  
**(**

**Should I not be dividing by another factor of 2!?**

Questions?



# Inclusion-Exclusion Principle

Recall that if  $A, B, C$  are disjoint subsets, then,  $|A \cup B \cup C| = |A| + |B| + |C|$  (this is the Sum rule from Lecture 2).

For two general sets  $A, B$ ,  $|A \cup B| = |A| + |B| - |A \cap B|$ . The last term fixes the “double counting”.

Similarly,  $|A \cup B \cup C| = |A| + |B| + |C| - |A \cap B| - |B \cap C| - |A \cap C| + |A \cap B \cap C|$ .

In general,

$$|\cup_{i=1,2,\dots,n} A_i| = \sum_i |A_i| - \sum_{i,j \text{ s.t. } 1 \leq i < j \leq n} |A_i \cap A_j| + \sum_{i,j,k \text{ s.t. } 1 \leq i < j < k \leq n} |A_i \cap A_j \cap A_k| \\ + \dots + (-1)^{n-1} |\cap_{i=1,2,\dots,n} A_i|$$

union on n sets

Intersection of n sets

$$|\bigcup_{i=1}^3 A_i| = |A_1| + |A_2| + |A_3|$$


## Inclusion-Exclusion Principle - Example

**Q:** Suppose there are 60 math majors, 200 EECS majors, and 40 physics majors. A student is allowed to double or even triple major. There are 4 math-EECS double majors, 3 math-physics double majors, 11 EECS-physics double majors and 2-triple majors. What is the total number of students across these three departments?

If  $M, E, P$  are the sets of students majoring in math, EECS and physics respectively, then we wish to compute  $|M \cup E \cup P| = |M| + |E| + |P| - |M \cap E| - |M \cap P| - |E \cap P| + |M \cap E \cap P|$   
 $= 300 - |M \cap E| - |M \cap P| - |E \cap P| + |M \cap E \cap P|.$

$$|M \cap E| = 4 + 2 = 6, |M \cap P| = 3 + 2 = 5, |E \cap P| = 11 + 2 = 13. |M \cap E \cap P| = 2$$

$$|M \cup E \cup P| = 300 - 6 - 5 - 13 + 2 = 278.$$



## Inclusion-Exclusion Principle - Example

**Q:** In how many permutations of the set  $\{0, 1, 2, \dots, 9\}$  do either 4 and 2, 0 and 4, or 6 and 0 appear consecutively? For example, in the following permutation 42067891235, 4 and 2 appear consecutively, but 6 and 0 do not (the order matters).

Treat 42 as one number and keep the rest of the numbers. There are 8 remaining numbers and 42 where you need to reorder. Therefore, there are 9! permutations.

Let  $P_{42}$  be the set of sequences such that 4 and 2 appear consecutively. Similarly, we define  $P_{60}$  and  $P_{04}$ . So we want to compute

$$|P_{42} \cup P_{60} \cup P_{04}| = |P_{42}| + |P_{60}| + |P_{04}| - |P_{42} \cap P_{60}| - |P_{42} \cap P_{04}| - |P_{60} \cap P_{04}| + |P_{42} \cap P_{60} \cap P_{04}|.$$

Let us first compute  $|P_{42}| = 9!$ . Similarly,  $|P_{60}| = |P_{04}| = 9!$ .

What about intersections?  $|P_{42} \cap P_{60}| =$  Number of sequences of the form  $(42, 60, 1, 3, 5, 7, 8, 9) = 8!$ . Similarly,  $|P_{60} \cap P_{04}| = |P_{42} \cap P_{04}| = 8!$ .

$|P_{42} \cap P_{60} \cap P_{04}| =$  Number of sequences of the form  $(6042, 1, 3, 5, 7, 8, 9) = 7!$ .

By the inclusion-exclusion principle,  $|P_{42} \cup P_{60} \cup P_{04}| = 3 \times 9! - 3 \times 8! + 7!$ .

# Combinatorial Proofs

What are the number of ways of choosing  $k$  students if we have to include a specific student?

Recall that if we have to choose  $k$  elements out of a size  $n$  set. Number of ways to do this is  $\binom{n}{k}$ . But this is equivalent to saying, we want to find the number of ways to throw away  $n - k$

elements =  $\binom{n}{n-k}$ . Hence  $\binom{n}{k} = \binom{n}{n-k}$ . Can prove algebraic statements using combinatorial arguments.

What are the number of ways of choosing  $k$  students if we do not include the specific student?

**Q:** Prove Pascal's identity using a combinatorial proof:  $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$

Consider  $n$  students in this class. What is the number of ways of selecting  $k$  students?  $\binom{n}{k}$ .

What is the number of ways of selecting  $k$  students if we have to ensure to include a particular student?  $\binom{n-1}{k-1}$ .

What is the number of ways of selecting  $k$  students if we have to ensure to NOT include a particular student?  $\binom{n-1}{k}$ .

Number of ways to select  $k$  students = number of ways of selecting  $k$  students to include a particular student + number of ways of selecting  $k$  students to NOT include a particular student.


Hence,  $\binom{n}{k} = \binom{n-1}{k-1} + \binom{n-1}{k}$ .

Not including a specific student

# Counting Practice

Q: In how many ways can we place (i) two identical black rooks (ii) a black rook and a white rook such that they do not share the same row or column?

You have sequence of (row, column)  
After first choice, you now have 7 options left for the remaining columns and rows.



**Figure 15.2** Two ways to place 2 rooks (♖♜) on a chessboard. The configuration in (b) is invalid because the rooks are in the same column.

64 options for the first position, and then 49 possible positions for the next rook

There are  $86 * 49$  possible ways to place the rooks.

Questions?

# Pigeonhole principle

**Q:** A drawer in a dark room contains red socks, green socks, and blue socks. How many socks must you withdraw to be sure that you have a matching pair?

Such problems can be tackled using the Pigeonhole principle.

**Pigeonhole Principle:** If there are more pigeons than holes they occupy, then there must be at least two pigeons in the same hole.

Formally, if  $|A| > |B|$ , then for every total function (one that has an assignment for every element in  $A$ ),  $f : A \rightarrow B$ , there exist two different elements of  $A$  that are mapped by  $f$  to the same element of  $B$ .

For the above problem,  $A$  = set of socks we picked = pigeons,  $B$  = set of colors {red, blue, green} = pigeonholes.  $|A|$  = number of socks we picked.  $|B| = 3$ .  $f : A \rightarrow B$  s.t.  $f(\text{sock we picked}) = \text{it's color}$ .

If there are more pigeons than holes (picked socks than colors), then at least two pigeons will be in the same hole (two of the picked socks will have the same color, and we get a matching pair). Hence, to ensure a matching pair, we need to pick 4 socks.

## Pigeonhole principle - Example

1: A student can have a birthday in one of the 52 weeks of the year.

Let us assign a week for every one of the 54 students.

Since we have more students than weeks in the year, this means that there will be two students who have a birthday in the same week.

Let  $A$  be the numbers from 1 to 100.

Let  $B$  be the remainders after doing  $n \bmod 41$ .  $\{0, \dots, 40\}$   
Q: A class has 54 students. Prove that there exist at least 2 students with their birthday in the same week.  
 $f: A \rightarrow B = (n \bmod 41)$

Since  $|A| > |B|$ , this means there are two numbers with the same remainder.  
Q: In the set of integers  $\{1, 2, \dots, 100\}$ , use the pigeonhole principle to prove that there exist two numbers whose difference is a multiple of 41.

$$\text{Let } x = 41m_1 + r_1$$

$$\text{Let } y = 41m_2 + r_1$$

$$x - y = 41(m_1 - m_2)$$

$$x - y \text{ is divisible by } 41$$



# Pigeonhole principle - Example

A kind of problem that arises in cryptography is to find different subsets of numbers with the same sum. For example, in this list of 25-digit numbers, find a subset of numbers that have the same sum. For example, maybe the sum of the last ten numbers in the first column is equal to the sum of the first eleven numbers in the second column.

Breaking a password is very hard

0020480135385502964448038	3171004832173501394113017
5763257331083479647409308	8247331000042995311646021
0489445991866915676240992	3208234421597368647019265
5800949123548989122628663	8496243997123475922766310
1082662032430379651370981	3437254656355157864869113
6042900801199280218026001	8518399140676002660747477
1178480894769706178994993	3574883393058653923711365
6116171789137737896701405	8543691283470191452333763
1253127351683239693851327	3644909946040480189969149
6144868973001582369723512	8675309258374137092461352
1301505129234077811069011	3790044132737084094417246
6247314593851169234746152	86943211123639086729665
1311567111143866433882194	3870332127437971355322815
6814428944266874963488274	8772321203608477245851154
1470029452721203587686214	4080505804577801451363100
6870852945543886849147881	8791422161722582546341091
1578271047286257499433886	4167283461025702348124920
6914955508120950093732397	9062628024592126283973285
1638243921852176243192354	423599683112377788211249
6949632451365987152423541	9137845566925526349897794
1763580219131985963102365	4670939445749439042111220
7128211143613619828415650	9153762966803189291934419
183622775601842231029694	4815379351865384279613427
7173920083651862307925394	9270880194077636406984249
1843971862675102037201420	4837052948212922604442190

This is a hard problem which is why it is used in cryptography. The first step to figure out is whether there even exists such a subset of numbers. We can do this using the pigeonhole principle!

## Pigeonhole principle - Example

**Q:** More generally, in a list of  $n$   $b$ -digit numbers, are there two different subsets of numbers that have the same sum?

Let  $A$  = set of all subsets of the  $n$  numbers. For example, if  $b = 3$ , an element of  $A$  is  $\{113, 221, 42\}$ .  $|A| = 2^n$


Let  $B$  be the set of possible sums of such subsets.  $f$  is a function that maps each subset to its corresponding sum. For example, if  $b = 3$ ,  $f(\{113, 221\}) = 334$ .

Let us compute  $|B|$ . For any list of  $n$  numbers, Minimum possible sum = 0. Max possible sum  $< 10^b \times n$ . For example, if  $b = 3$  and  $n = 5$ , then the maximum possible sum =  $999 \times 5 < 1000 \times 5$ . Hence,  $|B| < 10^b \times n$ . **Upper bound**

By the pigeonhole principle, there exist different subsets with the same sum if  $|A| > |B|$  i.e. if  $2^n > 10^b \times n$ .

For  $b = 3$ , this is possible if  $2^n > 1000n$ , meaning this is possible if  $n \log(2) > 3 + \log(n)$  (since  $\log$  is a monotonic function) Let's plot.

# Pigeonhole - Example



Hence, it is possible when  $n > 15$ . Similarly, for a general  $b$ , there exist different subsets with the same sum if  $n \log(2) > b + \log(n)$ .

Questions?