

CMPT 210: Probability and Computing

Lecture 11

Sharan Vaswani

February 13, 2024

Verifying Matrix Multiplication

As an example, let us focus on A, B being binary 2×2 matrices.

Example: $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, B = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$ then $C = AB = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$

Verifying Matrix Multiplication

As an example, let us focus on A, B being binary 2×2 matrices.

Example: $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, B = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$ then $C = AB = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$

Objective: Verify whether a matrix multiplication operation is correct.

Verifying Matrix Multiplication

As an example, let us focus on A, B being binary 2×2 matrices.

Example: $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, B = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$ then $C = AB = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$

Objective: Verify whether a matrix multiplication operation is correct.

Trivial way: Do the matrix multiplication ourselves, and verify it using $O(n^3)$ (or $O(n^{2.373})$) operations.

Verifying Matrix Multiplication

As an example, let us focus on A, B being binary 2×2 matrices.

Example: $A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, B = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}$ then $C = AB = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$

Objective: Verify whether a matrix multiplication operation is correct.

Trivial way: Do the matrix multiplication ourselves, and verify it using $O(n^3)$ (or $O(n^{2.373})$) operations.

Frievald's Algorithm: Randomized algorithm to verify matrix multiplication with high probability in $O(n^2)$ time.

(Basic) Freivald's Algorithm

Q: For $n \times n$ matrices A , B and D , is $D = AB$?

Algorithm:

1. Generate a random n -bit vector x , by making each bit x_i either 0 or 1 *independently* with probability $\frac{1}{2}$. E.g, for $n = 2$, toss a fair coin independently twice with the scheme – H is 0 and T is 1). If we get HT , then set $x = [0; 1]$.

(Basic) Freivald's Algorithm

Q: For $n \times n$ matrices A , B and D , is $D = AB$?

Algorithm:

1. Generate a random n -bit vector x , by making each bit x_i either 0 or 1 *independently* with probability $\frac{1}{2}$. E.g, for $n = 2$, toss a fair coin independently twice with the scheme – H is 0 and T is 1). If we get HT , then set $x = [0; 1]$.
2. Compute $t = Bx$ and $y = At = A(Bx)$ and $z = Dx$.

Since we are only doing matrix vector multiplication, it is $O(n^2)$

(Basic) Freivald's Algorithm

Q: For $n \times n$ matrices A , B and D , is $D = AB$?

Algorithm:

1. Generate a random n -bit vector x , by making each bit x_i either 0 or 1 *independently* with probability $\frac{1}{2}$. E.g, for $n = 2$, toss a fair coin independently twice with the scheme – H is 0 and T is 1). If we get HT , then set $x = [0; 1]$.
2. Compute $t = Bx$ and $y = At = A(Bx)$ and $z = Dx$.
3. Output “yes” if $y = z$ (all entries need to be equal), else output “no”.

(Basic) Freivald's Algorithm

Q: For $n \times n$ matrices A , B and D , is $D = AB$?

Algorithm:

1. Generate a random n -bit vector x , by making each bit x_i either 0 or 1 *independently* with probability $\frac{1}{2}$. E.g, for $n = 2$, toss a fair coin independently twice with the scheme – H is 0 and T is 1). If we get HT , then set $x = [0; 1]$.
2. Compute $t = Bx$ and $y = At = A(Bx)$ and $z = Dx$.
3. Output “yes” if $y = z$ (all entries need to be equal), else output “no”.

Computational complexity: Step 1 can be done in $O(n)$ time. Step 2 requires 3 matrix vector multiplications and can be done in $O(n^2)$ time. Step 3 requires comparing two n -dimensional vectors and can be done in $O(n)$ time. Hence, the total computational complexity is $O(n^2)$.

(Basic) Frievald's Algorithm

Let us run the algorithm on an example. Suppose we have generated $x = [1; 0]$

$$A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad ; \quad B = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \quad ; \quad D = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$
$$Bx = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad ; \quad y = A(Bx) = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad ; \quad z = Dx = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

(Basic) Frievald's Algorithm

Let us run the algorithm on an example. Suppose we have generated $x = [1; 0]$

$$A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad ; \quad B = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \quad ; \quad D = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$
$$Bx = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad ; \quad y = A(Bx) = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad ; \quad z = Dx = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

Hence the algorithm will correctly output “no” since $D \neq AB$.

(Basic) Frievald's Algorithm

Let us run the algorithm on an example. Suppose we have generated $x = [1; 0]$

$$A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad ; \quad B = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \quad ; \quad D = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$
$$Bx = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad ; \quad y = A(Bx) = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad ; \quad z = Dx = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

Hence the algorithm will correctly output “no” since $D \neq AB$.

Q: Suppose we have generated $x = [0; 0]$. What is y and z ?

(Basic) Frievald's Algorithm

Let us run the algorithm on an example. Suppose we have generated $x = [1; 0]$

$$A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad ; \quad B = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \quad ; \quad D = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$
$$Bx = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad ; \quad y = A(Bx) = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad ; \quad z = Dx = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

Hence the algorithm will correctly output “no” since $D \neq AB$.

Q: Suppose we have generated $x = [0; 0]$. What is y and z ?

In this case, $y = z$ and the algorithm will incorrectly output “yes” even though $D \neq AB$.

Correctness of algorithm depends on vector x

(Basic) Frievald's Algorithm

Let us run the algorithm on an example. Suppose we have generated $x = [1; 0]$.

$$A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad ; \quad B = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \quad ; \quad C = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$$
$$Bx = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad ; \quad y = A(Bx) = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad ; \quad z = Cx = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

(Basic) Frievald's Algorithm

Let us run the algorithm on an example. Suppose we have generated $x = [1; 0]$.

$$A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad ; \quad B = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \quad ; \quad C = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$$
$$Bx = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad ; \quad y = A(Bx) = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad ; \quad z = Cx = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

Hence the algorithm will correctly output “yes” since $C = AB$.

(Basic) Frievald's Algorithm

Let us run the algorithm on an example. Suppose we have generated $x = [1; 0]$.

$$A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad ; \quad B = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \quad ; \quad C = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$$
$$Bx = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad ; \quad y = A(Bx) = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad ; \quad z = Cx = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

Hence the algorithm will correctly output “yes” since $C = AB$.

Q: Suppose we have generated $x = [0; 1]$. What is y and z ?

(Basic) Frievald's Algorithm

Let us run the algorithm on an example. Suppose we have generated $x = [1; 0]$.

$$A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} \quad ; \quad B = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix} \quad ; \quad C = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$$
$$Bx = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad ; \quad y = A(Bx) = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \quad ; \quad z = Cx = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

Hence the algorithm will correctly output “yes” since $C = AB$.

Q: Suppose we have generated $x = [0; 1]$. What is y and z ?

In this case again, $y = z$ and the algorithm will correctly output “yes”.

(Basic) Freivald's Algorithm

Let us analyze the algorithm for general matrix multiplication.

Case (i): If $D = AB$, does the algorithm always output “yes”?

(Basic) Frievald's Algorithm

Let us analyze the algorithm for general matrix multiplication.

Case (i): If $D = AB$, does the algorithm always output “yes”? Yes! Since $D = AB$, for any vector x , $Dx = ABx$.

(Basic) Freivald's Algorithm

Let us analyze the algorithm for general matrix multiplication.

Case (i): If $D = AB$, does the algorithm always output “yes”? Yes! Since $D = AB$, for any vector x , $Dx = ABx$.

Case (ii) If $D \neq AB$, does the algorithm always output “no”?
Some probability of a mistake here

(Basic) Freivald's Algorithm

Let us analyze the algorithm for general matrix multiplication.

Case (i): If $D = AB$, does the algorithm always output “yes”? Yes! Since $D = AB$, for any vector x , $Dx = ABx$.

Case (ii) If $D \neq AB$, does the algorithm always output “no”?

Claim: For any input matrices A, B, D if $D \neq AB$, then the (Basic) Freivald's algorithm will output “no” with probability $\geq \frac{1}{2}$. What is the non-basic version?

(Basic) Freivald's Algorithm

Let us analyze the algorithm for general matrix multiplication.

Case (i): If $D = AB$, does the algorithm always output “yes”? Yes! Since $D = AB$, for any vector x , $Dx = ABx$.

Case (ii) If $D \neq AB$, does the algorithm always output “no”?

Claim: For any input matrices A, B, D if $D \neq AB$, then the (Basic) Freivald's algorithm will output “no” with probability $\geq \frac{1}{2}$.

	Yes	No	
$D = AB$	1	0	No mistake here
$D \neq AB$	$< \frac{1}{2}$	$\geq \frac{1}{2}$	Some mistake here

Table 1: Probabilities for Basic Freivalds Algorithm

One sided in the sense that an error only occurs if $D \neq AB$

(Basic) Freivald's Algorithm

Proof: If $D \neq AB$, we wish to compute the probability that algorithm outputs “yes” and prove that it less than $\frac{1}{2}$.

Let $E = AB - D$

If $D \neq AB$, there is at least an (i,j) such that $e_{ij} \neq 0$.

If E only has zeros, $AB = D$

$$Ex = ABx - Dx = (AB - D)x$$

(Basic) Frievald's Algorithm

Proof: If $D \neq AB$, we wish to compute the probability that algorithm outputs “yes” and prove that it less than $\frac{1}{2}$.

Define $E := (AB - D)$ and $r := Ex = (AB - D)x = y - z$. If $D \neq AB$, then $\exists(i, j)$ s.t. $E_{i,j} \neq 0$.

(Basic) Frievald's Algorithm

Proof: If $D \neq AB$, we wish to compute the probability that algorithm outputs “yes” and prove that it less than $\frac{1}{2}$.

Define $E := (AB - D)$ and $r := Ex = (AB - D)x = y - z$. If $D \neq AB$, then $\exists(i, j)$ s.t. $E_{i,j} \neq 0$.

$$\Pr(y = z \mid D \neq AB)$$

$$\Pr(R = 0 \mid D \neq AB)$$

$$\begin{aligned}\Pr[\text{Algorithm outputs “yes”}] &= \Pr[y = z] = \Pr[r = \mathbf{0}] \\ &= \Pr[(r_1 = 0) \cap (r_2 = 0) \cap \dots \cap (r_i = 0) \cap \dots]\end{aligned}$$

$$\Pr(R = 0) = \Pr(r_1 = 0 \ \& \ r_2 = 0 \ \& \ \dots \ \& \ r_n = 0 \mid D \neq AB)$$

(Basic) Frievald's Algorithm

Proof: If $D \neq AB$, we wish to compute the probability that algorithm outputs “yes” and prove that it less than $\frac{1}{2}$.

Define $E := (AB - D)$ and $r := Ex = (AB - D)x = y - z$. If $D \neq AB$, then $\exists(i, j)$ s.t. $E_{i,j} \neq 0$.

$$\begin{aligned}\Pr[\text{Algorithm outputs “yes”}] &= \Pr[y = z] = \Pr[r = \mathbf{0}] \\ &= \Pr[(r_1 = 0) \cap (r_2 = 0) \cap \dots \cap (r_i = 0) \cap \dots] \\ &= \Pr[(r_i = 0)] \Pr[(r_1 = 0) \cap (r_2 = 0) \cap \dots \cap (r_n = 0) | r_i = 0] \\ &\hspace{15em} (\text{By def. of conditional probability})\end{aligned}$$

(Basic) Frievald's Algorithm

Proof: If $D \neq AB$, we wish to compute the probability that algorithm outputs “yes” and prove that it less than $\frac{1}{2}$.

Define $E := (AB - D)$ and $r := Ex = (AB - D)x = y - z$. If $D \neq AB$, then $\exists(i, j)$ s.t. $E_{i,j} \neq 0$.

$$\begin{aligned}\Pr[\text{Algorithm outputs “yes”}] &= \Pr[y = z] = \Pr[r = \mathbf{0}] \\ &= \Pr[(r_1 = 0) \cap (r_2 = 0) \cap \dots \cap (r_i = 0) \cap \dots] \\ &= \Pr[(r_i = 0)] \underbrace{\Pr[(r_1 = 0) \cap (r_2 = 0) \cap \dots \cap (r_n = 0) | r_i = 0]}_{\text{(By def. of conditional probability)}}\end{aligned}$$

$$\implies \Pr[\text{Algorithm outputs “yes”}] \leq \Pr[r_i = 0] \quad (\text{Probabilities are in } [0, 1])$$

(Basic) Frievald's Algorithm

Proof: If $D \neq AB$, we wish to compute the probability that algorithm outputs “yes” and prove that it less than $\frac{1}{2}$.

Define $E := (AB - D)$ and $r := Ex = (AB - D)x = y - z$. If $D \neq AB$, then $\exists(i, j)$ s.t. $E_{i,j} \neq 0$.

$$\begin{aligned}\Pr[\text{Algorithm outputs “yes”}] &= \Pr[y = z] = \Pr[r = \mathbf{0}] \\ &= \Pr[(r_1 = 0) \cap (r_2 = 0) \cap \dots \cap (r_i = 0) \cap \dots] \\ &= \Pr[(r_i = 0)] \Pr[(r_1 = 0) \cap (r_2 = 0) \cap \dots \cap (r_n = 0) | r_i = 0] \\ &\hspace{15em} (\text{By def. of conditional probability})\end{aligned}$$

$$\implies \Pr[\text{Algorithm outputs “yes”}] \leq \Pr[r_i = 0] \hspace{10em} (\text{Probabilities are in } [0, 1])$$

To complete the proof, on the next slide, we will prove that $\Pr[r_i = 0] \leq \frac{1}{2}$.

(Basic) Frievald's Algorithm

Row i is the columns of matrix E

$$r_i = \sum_{k=1} E_{i,k} x_k = E_{i,j} x_j + \sum_{k \neq j} E_{i,k} x_k = E_{i,j} x_j + \omega \quad (\omega := \sum_{k \neq j} E_{i,k} x_k)$$

~~X~~ Goes through row of matrix E

$\begin{pmatrix} \vdots \\ i \end{pmatrix} \rightarrow$

$\begin{pmatrix} \vdots \\ j \end{pmatrix}$

If we are wanting to obtain the i th component, is it not the i th row multiplied by the vector X ? How does this formula come about?

(Basic) Frievald's Algorithm

$$r_i = \sum_{k=1}^n E_{i,k} x_k = E_{i,j} x_j + \sum_{k \neq j} E_{i,k} x_k = E_{i,j} x_j + \omega \quad (\omega := \sum_{k \neq j} E_{i,k} x_k)$$

$$\Pr[r_i = 0] = \Pr[r_i = 0 | \omega = 0] \Pr[\omega = 0] + \Pr[r_i = 0 | \omega \neq 0] \Pr[\omega \neq 0]$$

(By the law of total probability)

(Basic) Frievald's Algorithm

$$r_i = \sum_{k=1}^n E_{i,k} x_k = E_{i,j} x_j + \sum_{k \neq j} E_{i,k} x_k = E_{i,j} x_j + \omega \quad (\omega := \sum_{k \neq j} E_{i,k} x_k)$$

$$\Pr[r_i = 0] = \Pr[r_i = 0 | \omega = 0] \Pr[\omega = 0] + \Pr[r_i = 0 | \omega \neq 0] \Pr[\omega \neq 0]$$

(By the law of total probability)

$$\Pr[r_i = 0 | \omega = 0] = \Pr[x_j = 0] = \frac{1}{2} \quad (\text{Since } E_{i,j} \neq 0 \text{ and } \Pr[x_j = 1] = \frac{1}{2})$$

if ω is 0, then E_{ij} must be zero.

(Basic) Friedvald's Algorithm

$$r_i = \sum_{k=1}^n E_{i,k} x_k = E_{i,j} x_j + \sum_{k \neq j} E_{i,k} x_k = E_{i,j} x_j + \omega \quad (\omega := \sum_{k \neq j} E_{i,k} x_k)$$

$$\Pr[r_i = 0] = \Pr[r_i = 0 | \omega = 0] \Pr[\omega = 0] + \Pr[r_i = 0 | \omega \neq 0] \Pr[\omega \neq 0]$$

(By the law of total probability)

$$\Pr[r_i = 0 | \omega = 0] = \Pr[x_j = 0] = \frac{1}{2}$$

(Since $E_{i,j} \neq 0$ and $\Pr[x_j = 1] = \frac{1}{2}$)

Where did -w come from?

$$\Pr[r_i = 0 | \omega \neq 0] = \Pr[(x_j = 1) \cap E_{i,j} = -\omega] = \Pr[(x_j = 1)] \Pr[E_{i,j} = -\omega | x_j = 1]$$

(By def. of conditional probability)

$$\implies \Pr[r_i = 0 | \omega \neq 0] \leq \Pr[(x_j = 1)] = \frac{1}{2}$$

(Probabilities are in $[0, 1]$, $\Pr[x_j = 1] = \frac{1}{2}$)

(Basic) Frievald's Algorithm

$$r_i = \sum_{k=1}^n E_{i,k} x_k = E_{i,j} x_j + \sum_{k \neq j} E_{i,k} x_k = E_{i,j} x_j + \omega \quad (\omega := \sum_{k \neq j} E_{i,k} x_k)$$

$$\Pr[r_i = 0] = \Pr[r_i = 0 | \omega = 0] \Pr[\omega = 0] + \Pr[r_i = 0 | \omega \neq 0] \Pr[\omega \neq 0]$$

(By the law of total probability)

$$\Pr[r_i = 0 | \omega = 0] = \Pr[x_j = 0] = \frac{1}{2} \quad (\text{Since } E_{i,j} \neq 0 \text{ and } \Pr[x_j = 1] = \frac{1}{2})$$

$$\Pr[r_i = 0 | \omega \neq 0] = \Pr[(x_j = 1) \cap E_{i,j} = -\omega] = \Pr[(x_j = 1)] \Pr[E_{i,j} = -\omega | x_j = 1]$$

(By def. of conditional probability)

$$\implies \Pr[r_i = 0 | \omega \neq 0] \leq \Pr[(x_j = 1)] = \frac{1}{2} \quad (\text{Probabilities are in } [0, 1], \Pr[x_j = 1] = \frac{1}{2})$$

$$\implies \Pr[r_i = 0] \leq \frac{1}{2} \Pr[\omega = 0] + \frac{1}{2} \Pr[\omega \neq 0] = \frac{1}{2} \Pr[\omega = 0] + \frac{1}{2} [1 - \Pr[\omega = 0]] = \frac{1}{2}$$

($\Pr[E^c] = 1 - \Pr[E]$)

(Basic) Freivald's Algorithm

$$r_i = \sum_{k=1}^n E_{i,k} x_k = E_{i,j} x_j + \sum_{k \neq j} E_{i,k} x_k = E_{i,j} x_j + \omega \quad (\omega := \sum_{k \neq j} E_{i,k} x_k)$$

$$\Pr[r_i = 0] = \Pr[r_i = 0 | \omega = 0] \Pr[\omega = 0] + \Pr[r_i = 0 | \omega \neq 0] \Pr[\omega \neq 0]$$

(By the law of total probability)

$$\Pr[r_i = 0 | \omega = 0] = \Pr[x_j = 0] = \frac{1}{2} \quad (\text{Since } E_{i,j} \neq 0 \text{ and } \Pr[x_j = 1] = \frac{1}{2})$$

$$\Pr[r_i = 0 | \omega \neq 0] = \Pr[(x_j = 1) \cap E_{i,j} = -\omega] = \Pr[(x_j = 1)] \Pr[E_{i,j} = -\omega | x_j = 1]$$

(By def. of conditional probability)

$$\implies \Pr[r_i = 0 | \omega \neq 0] \leq \Pr[(x_j = 1)] = \frac{1}{2} \quad (\text{Probabilities are in } [0, 1], \Pr[x_j = 1] = \frac{1}{2})$$

$$\implies \Pr[r_i = 0] \leq \frac{1}{2} \Pr[\omega = 0] + \frac{1}{2} \Pr[\omega \neq 0] = \frac{1}{2} \Pr[\omega = 0] + \frac{1}{2} [1 - \Pr[\omega = 0]] = \frac{1}{2}$$

($\Pr[E^c] = 1 - \Pr[E]$)

$$\implies \Pr[\text{Algorithm outputs "yes"}] \leq \Pr[r_i = 0] \leq \frac{1}{2}.$$

(Basic) Freivald's Algorithm

$$r_i = \sum_{k=1}^n E_{i,k} x_k = E_{i,j} x_j + \sum_{k \neq j} E_{i,k} x_k = E_{i,j} x_j + \omega \quad (\omega := \sum_{k \neq j} E_{i,k} x_k)$$

$$\Pr[r_i = 0] = \Pr[r_i = 0 | \omega = 0] \Pr[\omega = 0] + \Pr[r_i = 0 | \omega \neq 0] \Pr[\omega \neq 0]$$

(By the law of total probability)

$$\Pr[r_i = 0 | \omega = 0] = \Pr[x_j = 0] = \frac{1}{2} \quad (\text{Since } E_{i,j} \neq 0 \text{ and } \Pr[x_j = 1] = \frac{1}{2})$$

$$\Pr[r_i = 0 | \omega \neq 0] = \Pr[(x_j = 1) \cap E_{i,j} = -\omega] = \Pr[(x_j = 1)] \Pr[E_{i,j} = -\omega | x_j = 1]$$

(By def. of conditional probability)

$$\implies \Pr[r_i = 0 | \omega \neq 0] \leq \Pr[(x_j = 1)] = \frac{1}{2} \quad (\text{Probabilities are in } [0, 1], \Pr[x_j = 1] = \frac{1}{2})$$

$$\implies \Pr[r_i = 0] \leq \frac{1}{2} \Pr[\omega = 0] + \frac{1}{2} \Pr[\omega \neq 0] = \frac{1}{2} \Pr[\omega = 0] + \frac{1}{2} [1 - \Pr[\omega = 0]] = \frac{1}{2}$$

($\Pr[E^c] = 1 - \Pr[E]$)

$$\implies \Pr[\text{Algorithm outputs "yes"}] \leq \Pr[r_i = 0] \leq \frac{1}{2}.$$

(Basic) Freivald's Algorithm

Hence, if $D \neq AB$, the Algorithm outputs “yes” with probability $\leq \frac{1}{2} \implies$ the Algorithm outputs “no” with probability $\geq \frac{1}{2}$.

In the worst case, the algorithm can be incorrect half the time! We promised the algorithm would return the correct answer with “high” probability close to 1.

(Basic) Freivald's Algorithm

Hence, if $D \neq AB$, the Algorithm outputs “yes” with probability $\leq \frac{1}{2} \implies$ the Algorithm outputs “no” with probability $\geq \frac{1}{2}$.

In the worst case, the algorithm can be incorrect half the time! We promised the algorithm would return the correct answer with “high” probability close to 1.

A common trick in randomized algorithms is to have m independent trials of an algorithm and aggregate the answer in some way, reducing the probability of error, thus *amplifying the probability of success*.

Questions?

Frievald's Algorithm

By repeating the *Basic Frievald's Algorithm* m times, we will amplify the probability of success. The resulting complete Frievald's Algorithm is given by:

- 1 Run the Basic Frievald's Algorithm for m independent runs.

Frievald's Algorithm

By repeating the *Basic Frievald's Algorithm* m times, we will amplify the probability of success. The resulting complete Frievald's Algorithm is given by:

- 1 Run the Basic Frievald's Algorithm for m independent runs.
- 2 If *any* run of the Basic Frievald's Algorithm outputs “no”, output “no”.
- 3 If *all* runs of the Basic Frievald's Algorithm output “yes”, output “yes”.

Frievald's Algorithm

By repeating the *Basic Frievald's Algorithm* m times, we will amplify the probability of success. The resulting complete Frievald's Algorithm is given by:

- 1 Run the Basic Frievald's Algorithm for m independent runs.
- 2 If *any* run of the Basic Frievald's Algorithm outputs “no”, output “no”.
- 3 If *all* runs of the Basic Frievald's Algorithm output “yes”, output “yes”.

	Yes	No
$D = AB$	1	0
$D \neq AB$	$< \frac{1}{2^m}$	$\geq 1 - \frac{1}{2^m}$

Table 2: Probabilities for Frievald's Algorithm

Frievald's Algorithm

By repeating the *Basic Frievald's Algorithm* m times, we will amplify the probability of success. The resulting complete Frievald's Algorithm is given by:

- 1 Run the Basic Frievald's Algorithm for m independent runs.
- 2 If *any* run of the Basic Frievald's Algorithm outputs "no", output "no".
- 3 If *all* runs of the Basic Frievald's Algorithm output "yes", output "yes".

	Yes	No
$D = AB$	1	0
$D \neq AB$	$< \frac{1}{2^m}$	$\geq 1 - \frac{1}{2^m}$

Table 2: Probabilities for Frievald's Algorithm

If $m = 20$, then Frievald's algorithm will make mistake with probability $1/2^{20} \approx 10^{-6}$.

Computational Complexity: $O(mn^2)$

Probability Amplification

Consider a randomized algorithm \mathcal{A} that is supposed to solve a binary decision problem i.e. it is supposed to answer either Yes or No. It has a one-sided error – (i) if the true answer is Yes, then the algorithm \mathcal{A} correctly outputs Yes with probability 1, but (ii) if the true answer is No, the algorithm \mathcal{A} incorrectly outputs Yes with probability $\leq \frac{1}{2}$.

Probability Amplification

Consider a randomized algorithm \mathcal{A} that is supposed to solve a binary decision problem i.e. it is supposed to answer either Yes or No. It has a one-sided error – (i) if the true answer is Yes, then the algorithm \mathcal{A} correctly outputs Yes with probability 1, but (ii) if the true answer is No, the algorithm \mathcal{A} incorrectly outputs Yes with probability $\leq \frac{1}{2}$.

Let us define a new algorithm \mathcal{B} that runs algorithm \mathcal{A} m times, and if *any* run of \mathcal{A} outputs No, algorithm \mathcal{B} outputs No. If *all* runs of \mathcal{A} output Yes, algorithm \mathcal{B} outputs Yes.

Probability Amplification

Consider a randomized algorithm \mathcal{A} that is supposed to solve a binary decision problem i.e. it is supposed to answer either Yes or No. It has a one-sided error – (i) if the true answer is Yes, then the algorithm \mathcal{A} correctly outputs Yes with probability 1, but (ii) if the true answer is No, the algorithm \mathcal{A} incorrectly outputs Yes with probability $\leq \frac{1}{2}$.

Let us define a new algorithm \mathcal{B} that runs algorithm \mathcal{A} m times, and if *any* run of \mathcal{A} outputs No, algorithm \mathcal{B} outputs No. If *all* runs of \mathcal{A} output Yes, algorithm \mathcal{B} outputs Yes.

Q: What is the probability that algorithm \mathcal{B} correctly outputs Yes if the true answer is Yes, and correctly outputs No if the true answer is No?

Probability Amplification - Analysis

If A_i denotes run i of Algorithm \mathcal{A} , then

$$\Pr[\mathcal{B} \text{ outputs Yes} \mid \text{true answer is Yes}]$$

$$= \Pr[\mathcal{A}_1 \text{ outputs Yes} \cap \mathcal{A}_2 \text{ outputs Yes} \cap \dots \cap \mathcal{A}_m \text{ outputs Yes} \mid \text{true answer is Yes}]$$

$$= \prod_{i=1}^m \Pr[\mathcal{A}_i \text{ outputs Yes} \mid \text{true answer is Yes}] = 1 \quad (\text{Independence of runs})$$

Probability Amplification - Analysis

If A_i denotes run i of Algorithm \mathcal{A} , then

$$\Pr[\mathcal{B} \text{ outputs Yes} \mid \text{true answer is Yes}]$$

$$= \Pr[\mathcal{A}_1 \text{ outputs Yes} \cap \mathcal{A}_2 \text{ outputs Yes} \cap \dots \cap \mathcal{A}_m \text{ outputs Yes} \mid \text{true answer is Yes}]$$

$$= \prod_{i=1}^m \Pr[\mathcal{A}_i \text{ outputs Yes} \mid \text{true answer is Yes}] = 1 \quad (\text{Independence of runs})$$

$$\Pr[\mathcal{B} \text{ outputs No} \mid \text{true answer is No}]$$

$$= 1 - \Pr[\mathcal{B} \text{ outputs Yes} \mid \text{true answer is No}]$$

$$= 1 - \Pr[\mathcal{A}_1 \text{ outputs Yes} \cap \mathcal{A}_2 \text{ outputs Yes} \cap \dots \cap \mathcal{A}_m \text{ outputs Yes} \mid \text{true answer is No}]$$

$$= 1 - \prod_{i=1}^m \Pr[\mathcal{A}_i \text{ outputs Yes} \mid \text{true answer is No}] \geq 1 - \frac{1}{2^m}.$$

Probability Amplification - Analysis

If A_i denotes run i of Algorithm \mathcal{A} , then

$$\begin{aligned} & \Pr[\mathcal{B} \text{ outputs Yes} \mid \text{true answer is Yes}] \\ &= \Pr[\mathcal{A}_1 \text{ outputs Yes} \cap \mathcal{A}_2 \text{ outputs Yes} \cap \dots \cap \mathcal{A}_m \text{ outputs Yes} \mid \text{true answer is Yes}] \\ &= \prod_{i=1}^m \Pr[\mathcal{A}_i \text{ outputs Yes} \mid \text{true answer is Yes}] = 1 \end{aligned} \quad \text{(Independence of runs)}$$

$$\begin{aligned} & \Pr[\mathcal{B} \text{ outputs No} \mid \text{true answer is No}] \\ &= 1 - \Pr[\mathcal{B} \text{ outputs Yes} \mid \text{true answer is No}] \\ &= 1 - \Pr[\mathcal{A}_1 \text{ outputs Yes} \cap \mathcal{A}_2 \text{ outputs Yes} \cap \dots \cap \mathcal{A}_m \text{ outputs Yes} \mid \text{true answer is No}] \\ &= 1 - \prod_{i=1}^m \Pr[\mathcal{A}_i \text{ outputs Yes} \mid \text{true answer is No}] \geq 1 - \frac{1}{2^m}. \end{aligned}$$

When the true answer is Yes, both \mathcal{B} and \mathcal{A} correctly output Yes. When the true answer is No, \mathcal{A} incorrectly outputs Yes with probability $< \frac{1}{2}$, but \mathcal{B} incorrectly outputs Yes with probability $< \frac{1}{2^m} \ll \frac{1}{2}$. By repeating the experiment, we have “amplified” the probability of success.

Questions?