

CMPT 210: Probability and Computing

Lecture 2

Sharan Vaswani

January 11, 2024

Functions

We can also define a function with a set as the argument. For a set $S \in D$,
 $f(S) := \{x \mid \forall s \in S, x = f(s)\}$.

$A = \{a, b, c, \dots, z\}$, $B = \{1, 2, 3, \dots, 26\}$. $f : A \rightarrow B$ such that $f(a) = 1$, $f(b) = 2, \dots$
 $f(\{e, f, z\}) = \{5, 6, 26\}$.

If D is the domain of f , then $\text{range}(f) := f(D) = f(\text{domain}(f))$.

Q: If $f : \mathbb{N} \rightarrow \mathbb{R}$, and $f(x) = x^2$. What is the domain and codomain of f ? What is the range?

Q: Consider $f : \{0, 1\}^5 \rightarrow \mathbb{N}$ s.t. $f(x)$ counts the length of a left to right search of the bits in the binary string x until a 1 appears. $f(01000) = 2$.

What is $f(00001)$, $f(00000)$? Is f a total function?

Surjective Functions

Surjective functions: $f : A \rightarrow B$ is a surjective function iff for every $b \in B$, there exists an $a \in A$ s.t. $f(a) = b$. $f : \mathbb{R} \rightarrow \mathbb{R}$ such that $f(x) = x + 1$ is a surjective function.

For surjective functions, $|\#arrows| \geq |B|$.

Since each element of A is assigned at most one value, and some need not be assigned a value at all, $|\#arrows| \leq |A|$.

Hence, if f is a surjective function, then $|A| \geq |B|$.

$A = \{a, b, c, \dots, z, \alpha, \beta, \gamma, \dots\}$, $B = \{1, 2, 3, \dots, 26\}$. $f : A \rightarrow B$ such that $f(a) = 1$, $f(b) = 2, \dots$. f does not assign any value to the Greek letters. For every number in B , there is a letter in A . Hence, f is surjective, and $|A| > |B|$.

Injective & Bijective Functions

Injective functions: $f : A \rightarrow B$ is an injective function iff $\forall a \in A$, there is a *unique* $b \in B$ s.t. $f(a) = b$. If f is injective and $f(a) = f(b)$, then it implies that $a = b$.

Hence, $|\#\text{arrows}| = |A| \leq |B|$. Hence, if f is a injective function, then $|A| \leq |B|$.

$A = \{a, b, c, \dots z\}$, $B = \{1, 2, 3, \dots 26, 27, \dots 100\}$. $f : A \rightarrow B$ such that $f(a) = 1$, $f(b) = 2, \dots$. No element in A is assigned values $27, 28, \dots$, and for every letter in A , there is a unique number in B . Hence, f is injective, and $|A| < |B|$.

Bijective functions: f is a bijective function iff it is both surjective and injective, implying that $|A| = |B|$.

$A = \{a, b, c, \dots z\}$, $B = \{1, 2, 3, \dots 26\}$. $f : A \rightarrow B$ such that $f(a) = 1$, $f(b) = 2, \dots$. Every element in A is assigned a unique value in B and for every element in B , there is a value in A that is mapped to it. f is bijective, and $|A| = |B|$.

Converse of the previous statements is also true.

- If $|A| \geq |B|$, then it's always possible to define a surjective function $f : A \rightarrow B$.
- If $|A| \leq |B|$, then it's always possible to define an injective function $f : A \rightarrow B$.
- If $|A| = |B|$, then it's always possible to define a bijective function $f : A \rightarrow B$.

Q: Recall that the Cartesian product of two sets $S = \{s_1, s_2, \dots, s_m\}$, $T = \{t_1, t_2, \dots, t_n\}$ is $S \times T := \{(s, t) | s \in S, t \in T\}$. Construct a bijective function $f : (S \times T) \rightarrow \{1, \dots, nm\}$, and prove that $|S \times T| = nm$.

Examples: (a, b, a) , $(1,3,4)$, $(4,3,1)$

An element can appear twice. E.g. $(a, a, b) \neq (a, b)$.

The order of the elements does matter. E.g. $(a, b) \neq (b, a)$.

Q: What is the size of $(1, 2, 2, 3)$? What is the size of $\{1, 2, 2, 3\}$?

Sets and Sequences: The Cartesian product of sets $S \times T \times U$ is a set consisting of all sequences where the first component is drawn from S , the second component is drawn from T and the third from U . $S \times T \times U = \{(s, t, u) | s \in S, t \in T, u \in U\}$.

Q: For set $S = \{0, 1\}$, $S^3 = S \times S \times S$. Enumerate S^3 . What is $|S^3|$?


Questions?

Counting Sets – using a bijection

Q: Suppose we want to buy 10 donuts. There are 5 donut varieties – chocolate, lemon-filled, sugar, glazed, plain. What is the number of ways to select the 10 donuts?

Let A be the set of ways to select the 10 donuts. Each element of A is a potential selection. For example, 4 chocolate, 3 lemon, 0 sugar, 2 glazed and 1 plain.

Let's map each way to a string as follows:

0000	000		00	0
				
chocolate	lemon	sugar	glazed	plain

Lets fix the ordering – chocolate, lemon, sugar, glazed and plain, and abstract this out further to get the sequence: 00001000110010. Hence, each way of choosing donuts is mapped to a binary sequence of length 14 with exactly 4 ones. Now, let B be all 14-bit sequences with exactly 4 ones. An element of B is 11110000000000.

Q: The above sequence corresponds to what donut order?

For every way to select donuts, we have an equivalent sequence in B . And every sequence in B implies a unique way to select donuts. Hence, the mapping from $A \rightarrow B$ is a bijective function.

Counting Sets – using a bijection

Hence, $|A| = |B|$, meaning that we have reduced the problem of counting the number of ways to select donuts to counting the number of 14-bit sequences with exactly 4 ones.

General result: The number of ways to choose n elements with k available varieties is equal to the number of $n + k - 1$ -bit binary sequences with exactly $k - 1$ ones.

Q: There are 2 donut varieties – chocolate and lemon-filled. Suppose we want to buy only 2 donuts. Use the above result to count the number of ways in which we can select the donuts? What are these ways? **There are three ways to buy the donuts**

Q: In the above example, I want at least one chocolate donut. What are the types of acceptable 3-bit sequences with this criterion? How many ways can we do this?

0 | 0 | 0 |

two ways of doing this

0, 0, 1

0, 1, 0

Counting Sets – using the sum rule

Union

Q: Let R be the set of rainy days, S be the set of snowy days and H be the set of really hot days in 2023. A bad day can be either rainy, snowy or really hot. What is the number of good days?

Let B be the set of bad days. $B = R \cup S \cup H$, and we want to estimate $|\bar{B}|$. $|D| = 365$.

$$|\bar{B}| = |D| - |B| = 365 - |B| = 365 - |R \cup S \cup H|.$$

Since the sets R , S and H are disjoint, $|R \cup S \cup H| = |R| + |S| + |H|$, and hence the number of good days = $365 - |R| - |S| - |H|$.

Sum rule: If $A_1, A_2 \dots A_m$ are disjoint sets, then, $|A_1 \cup A_2 \cup \dots \cup A_m| = \sum_{i=1}^m |A_i|$.

Counting Sequences – using the product rule

Either or: union

Pick one from each: sequences

Q: Suppose the university offers Math courses (denoted by the set M), CS courses (set C) and Statistics courses (set S). We need to pick one course from each subject – Math, CS and Statistics. What is the number of ways we can select we can select the 3 courses?

The above problem is equivalent to counting the number of sequences of the form (m, c, s) that maps to choose the Math course m , CS course c and Stats course s .

Recall that the product of sets $M \times C \times S$ is a set consisting of all sequences where the first component is drawn from M , the second component is drawn from C and the third from S , i.e. $M \times C \times S = \{(m, c, s) | m \in M, c \in C, s \in S\}$. Hence, counting the number of sequences is equivalent to computing $|M \times C \times S|$.

It is equivalent to counting sequences since you need one class from each class type, giving you an implicit order.

Product Rule: $|M \times C \times S| = |M| \times |C| \times |S|$.

Using the above equivalence, the number of sequences and hence, the number of ways to select the 3 courses is $|M| \times |C| \times |S|$.

x is a character of the alphabet}
of length n passwords = $|A|^n$

Q: What is the number of length n -passwords that can be generated if each character in the password is allowed to be lower-case letter?

26^n since each letter can have 26 possible values

Counting – Example

Q: What is the number of passwords that can be generated if the (i) first character is only allowed to be a lower-case letter, (ii) each subsequent character in the password is allowed to be lower-case letter or digit (0 – 9) and (iii) the length of the password is required to be between 6-8 characters?

Let $L = \{a, b, \dots, z\}$ and $D = \{0, 1, 2, \dots\}$. Using the equivalence between sequences and products of sets, the set of passwords of length 6 is given by $P_6 = L \times (L \cup D)^5$. Using the product rule, $|P_6| = |L| \times (|L \cup D|)^5 = |L| \times (|L| + |D|)^5$.

Since the total set of passwords are $P = P_6 \cup P_7 \cup P_8$, and a password can be either of length 6, 7 or 8, sets P_6 , P_7 and P_8 are disjoint. Using the sum rule, $|P| = |P_6| + |P_7| + |P_8| = |L| \times [(|L| + |D|)^5(1 + (|L| + |D|) + (|L| + |D|)^2)] = 26 \times 36^5 \times [1 + 36 + 1296]$.

In tests, show what rule you are using
Do it systematically

Counting sequences – using the generalized product rule

Q: Suppose we have p prizes to be handed amongst the set A of n students. What are the number of ways in which we can distribute the prizes?

Q: Suppose we have p prizes to be handed amongst the set A of n students. What are the number of ways in which we can distribute the prizes such that each prize goes to a different student i.e. no student receives more than one prize?

Other way:

(2, 7, 4, 5) represents the sequence of the ids of the students who won prizes.

You want to count the number of possible sequences

Consider sequences of length p . The first entry can be chosen in n ways (the first prize can be given to one of the n students). After the first entry is chosen, since the same student cannot receive the prize, the second entry can be chosen in $n - 1$ ways, and so on. Hence, the total number of ways to distribute the prizes $= n \times (n - 1) \times \dots \times (n - (p - 1))$.

Generalized product rule: If S is the set of length k sequences such that the first entry can be selected in n_1 ways, after the first entry is chosen, the second one can be chosen in n_2 ways, and so on, then $|S| = n_1 \times n_2 \times \dots \times n_k$. If $n_1 = n_2 = \dots = n_k$, we recover the product rule.

n^p ways since a student can win multiple prizes

Counting - Example

Q: A dollar bill is defective if some digit appears more than once in the 8-digit serial number. What is the fraction of non-defective bills?

In order to compute the fraction of non-defective bills, we need to compute the quantity

$$\frac{|\text{serial numbers with all different digits}|}{|\text{possible serial numbers}|}.$$

For computing $|\text{possible serial numbers}|$, each digit can be one of 10 numbers. Hence, using the product rule, $|\text{possible serial numbers}| = 10 \times 10 \dots = 10^8$.

For computing $|\text{serial numbers with all different digits}|$, the first digit can be one of 10 numbers. Once the first digit is chosen, the second one can be chosen in 9 ways, and so on. By the generalized product rule, $|\text{serial numbers with all different digits}| = 10 \times 9 \times \dots \times 3 = 1,814,400$.

$$\text{Fraction of non-defective bills} = \frac{1,814,400}{10^8} = 1.8144\%$$

For each combination of digits, you need to compute the permutations to generate each serial number

Permutations

A permutation of a set S is a sequence of length $|S|$ that contains every element of S exactly once. Permutations of $\{a, b, c\}$ are $(a, b, c), (a, c, b), (b, c, a), (b, a, c), (c, a, b), (c, b, a)$.

Q: Given a set of size n , what is the total number of permutations?

Considering sequences of length n , the first entry can be chosen in n ways. Since each element can be chosen only once, the second entry can be chosen in $n - 1$ ways, and so on. **N! many permutations**

By the generalized product rule, the number of permutations = $n \times (n - 1) \times \dots \times 1$.
Text
Is this supposed to be +2 or -2?

Factorial: $n! := n \times (n - 1) \times \dots \times 1$. By convention: $0! = 1$.

How big is $n!$? **Stirling approximation:** $n! \approx \sqrt{2\pi n} \left(\frac{n}{e}\right)^n$.

Q: Which is bigger? $n!$ vs $n(n - 1)(n + 2)(n - 3)!$?

Q: In how many ways can we arrange n people in a line?

Counting – Division rule

k -to-1 function: Maps exactly k elements of the domain to every element of the codomain.

If $f : A \rightarrow B$ is a k -to-1 function, then, $|A| = k|B|$.

Example: E is the set of ears in this room, and P is the set of people. Then f mapping the ears to people is a 2-to-1 function. Hence, $|E| = 2|P|$.

Q: If $f : A \rightarrow B$ is a k -to-1 function, and $g : B \rightarrow C$ is a m -to-1 function, then what is $|A|/|C|$?

Q: If $f : A \rightarrow B$ is a k -to-1 function, and $g : C \rightarrow B$ is a m -to-1 function, then what is $|A|/|C|$?

$$A = kB, B = Cm,$$

$$C = B / m$$

$$A / B = kB * m / B = mk$$

$$A = kB, C = Bm,$$

$$A / C = kB / Bm = k / m$$

Counting – Example

er of seatings: all possible ways to seat people
ngements: all unique ways of seating people

Q: In how many ways can we arrange n people around a round table? Two seatings are considered to be the same *arrangement* if each person sits with the same person on their left in both seatings.

Starting from the head of the table, and going clockwise, each seating has an equivalent sequence. $|\text{seatings}| = \text{number of permutations} = n!$.

n different seatings can result in the same arrangement (by clockwise rotation).

Hence, $f : \text{seatings} \rightarrow \text{arrangements}$ is an n -to-1 function. Hence, the $|\text{seatings}| = n |\text{arrangements}|$, meaning that the $|\text{arrangements}| = (n - 1)!$.

Questions?

Counting subsets

Q: How many size- k subsets of a size- n set are there?

Example: How many ways can we select 5 books from 100?

Let us form a permutation of the n elements, and pick the first k elements to form the subset. Every size k subset can be generated this way. There are $n!$ total such permutations.

The order of the first k elements in the permutation does not matter in forming the subset, and neither does the order of the remaining $n - k$ elements.

The first k elements can be ordered in $k!$ ways and the remaining $n - k$ elements can be ordered in $(n - k)!$ ways. Using the product rule, $k! \times (n - k)!$ permutations map to the same size k subset.

Hence, the function $f : \text{permutations} \rightarrow \text{size } k \text{ subsets}$ is a $k! \times (n - k)!$ -to-1 function. By the division rule, $|\text{permutations}| = k! \times (n - k)! |\text{size } k \text{ subsets}|$. Hence, the total number of size k subsets $= \frac{n!}{k! \times (n - k)!}$.

$$n \text{ choose } k = \binom{n}{k} := \frac{n!}{k! \times (n - k)!}.$$

Q: Prove that $\binom{n}{k} = \binom{n}{n-k}$ - both algebraically (using the formula for $\binom{n}{k}$) and combinatorially (without using the formula)

Q: Which is bigger? $\binom{8}{4}$ vs $\binom{8}{5}$?

$$\begin{aligned} 8 * 7 * 6 * 5 / 4 * 3 * 2 \\ = 7 * 6 * 5 / 3 \\ = 7 * 3 * 5 = 105 \end{aligned}$$

$$\begin{aligned} 8 * 7 * 6 / 3 * 2 \\ = 8 * 7 = 56 \end{aligned}$$

Counting subsets – Example

Q: How many m -bit binary sequences contain exactly k ones?

Consider set $A = \{1, \dots, m\}$ and selecting S , a subset of size k . For example, say $m = 10, k = 3$ and $S = \{1, 7, 10\}$. S records the positions of the 1's, and can be mapped to the sequence 0010001001. Similarly, every m -bit sequence with exactly k ones can be mapped to a subset S of size k . Hence, there is a bijection:

$f : m\text{-bit sequence with exactly } k \text{ ones} \rightarrow \text{subsets of size } k \text{ from size } m\text{-set, and}$
 $|m\text{-bit sequence with exactly } k \text{ ones}| = |\text{subsets of size } k| = \binom{m}{k}.$

Q: Suppose we want to buy 10 donuts. There are 5 donut varieties – chocolate, lemon-filled, sugar, glazed, plain. What is the number of ways to select the 10 donuts?

Recall that the number of ways of selecting 10 donuts with 5 varieties = number of 14-bit sequences with exactly 4 ones = $\binom{14}{4} = 1001$.

Q: What is the number of ways of choosing n things with k varieties?

Counting subsets – Example

sum of $\binom{n}{i}$, where i ranges from k to n

Q: What is the number of n -bit binary sequences with at least k ones?

Q: What is the number of n -bit binary sequences with less than k ones?

sum of $\binom{n}{i}$, where i ranges from 0 to $k - 1$

Q: What is the total number of n -bit binary sequences?

sum of $\binom{n}{i}$, where i ranges from 0 to n , which is 2^n

Questions?