

Wilson lines on Loewner-generated contours

Defect supersymmetry, reflection, and endpoint positivity

Drafted for Edward Baker

19 September 2026 – Version 0.1

Abstract

We examine a construction in which Loewner evolution supplies a contour and a supersymmetric gauge theory supplies an open Wilson observable on it. The motivation is a positivity program for the Weil quadratic form, but the results concern the geometric and field-theoretic prerequisites of that program. We give an exact constant-driver description of the inherited semicircles and prove their rigidity for one fixed scalar coupling and bulk supercharge. A smooth transport variation identifies the insertions produced by a regular inverse Loewner map. Allowing a tangent-dependent scalar coupling restores contour freedom: a full chiral D3–D5 projector calculation gives a common complex Poincaré charge, including one charge for all defect directions. Within this constant-map ansatz, however, the scalar endpoints killed by the same charge have zero mutual contraction and a charged, vanishing vacuum pairing. Ordinary time reflection instead gives a nonzero free Gram kernel, with a different scalar map on the reflected branch. We construct a smooth reference junction and compute a finite, nonzero direct bulk response to a polynomial contour deformation. This is one perturbative sector, not the complete interacting response. The paper supplies the arithmetic target, relevant background, proofs, controls, and reproducible diagnostics. Interacting reflection positivity, a protected nonzero pairing, and the identification with the full Weil form remain open.

Status. Working research draft. Classical and free-field statements are separated from conditional quantum statements throughout. No independent mathematical review has been completed. Model assistance and source provenance are disclosed in Appendix B.

Contents

1	Introduction and relation to earlier work	2
1.1	The relevant literature	3
1.2	What survives from the parent investigations	4
1.3	Results and scope	4
2	The arithmetic positivity target	5
2.1	The compactly supported Weil form	5
2.2	The positive kinetic term and the indispensable local term	5
2.3	Shifted transfers and a sufficient contraction route	6
2.4	The exact positive part, and what it leaves unresolved	7
3	Why defect matter reproduces the archimedean exponents	7
3.1	An auxiliary Gaussian realization of the gamma ratio	9

4	Semicircles, constant drivers, and a restricted rigidity theorem	9
4.1	Fixed scalar coupling and fixed special charge	10
5	Smooth Wilson variation and the Loewner insertion	11
5.1	Conventions and the exact variation	11
5.2	Regular inverse-map deformations	12
5.3	A precise obstruction to geometry-only closure	12
5.4	The tangent connection as a one-form	13
6	Tangent coupling with the full chiral defect projector	13
6.1	Field content and conventions	13
6.2	A normal plane	13
6.3	One charge across all defect directions	14
6.4	The remaining H weight	15
7	Endpoint polarization, vanishing, and the adjoint	15
7.1	The same-charge endpoint calculation	15
7.2	The full bilinear has nonzero R charge	16
7.3	Why the original semicircle is a nonzero control	16
7.4	The adjoint changes the scalar map	16
8	Physical reflection and the reference color junction	16
8.1	Ordinary time reflection	16
8.2	The actual paired operator and the conditional Gram statement	17
8.3	An unconditional free kernel	18
8.4	The two branches preserve different constant charges	18
8.5	Internal rotations do not automatically preserve positivity	18
8.6	A free test of an odd-H normal reflection	19
9	A smooth reference join and a direct bulk response	19
9.1	A contour family that meets the reflection plane smoothly	19
9.2	The specified Gaussian perturbative sector	20
9.3	Why a positive norm need not be protected	21
10	Status and the remaining route to a proof	22
10.1	Statement ledger	22
10.2	The next discriminating calculation	22
10.3	What would turn the construction into an RH route	23
A	Reproducible diagnostics and controls	23
B	Provenance and contribution disclosure	24

1 Introduction and relation to earlier work

There are two distinct uses of Loewner theory in a positivity construction. One may try to identify a conformal-map observable directly with an arithmetic transfer function. Alternatively, one may use the map to generate a curve, evaluate a field-dependent observable on that curve, and only then

seek a transfer law. This paper develops the second option. The additional field and endpoint data are essential: the conformal map alone does not determine Wilson transport or its variation.

The proposed application is the Riemann Hypothesis (RH). In the normalization used here, RH is equivalent to positivity of an explicit quadratic form on every compact interval. Its archimedean part contains the kernel

$$n_{\Gamma}(u) = \frac{e^{-u/2}}{1 - e^{-2u}} = \sum_{n=0}^{\infty} e^{-(2n+1/2)u}, \quad u > 0. \quad (1.1)$$

A free three-dimensional defect scalar, after a logarithmic radial change of variables and even angular smearing, supplies this sequence of exponents. That observation motivates the use of defect matter. It does not supply the local subtraction, the pole terms, or the prime-power terms of the Weil form. Section 2 states that target explicitly, and Section 3 explains exactly what the free construction provides.

1.1 The relevant literature

The scalar-coupled Wilson operator in four-dimensional $\mathcal{N} = 4$ supersymmetric Yang–Mills theory is part of the Wilson-loop framework introduced in the AdS/CFT setting by Maldacena [1]. Zarembo [2] constructed families whose internal scalar direction follows the spacetime tangent through a constant isometry. The resulting linear Clifford projectors allow arbitrary contour shapes while retaining common supersymmetry. Our tangent prescription is an application of that construction. The additional questions here are its compatibility with the chiral defect projector, physical endpoint matter, and reflection.

DeWolfe, Freedman and Ooguri [3] described a four-dimensional $\mathcal{N} = 4$ bulk theory coupled to a three-dimensional fundamental hypermultiplet at a D3–D5 intersection. The defect preserves an $SU(2)_H \times SU(2)_V$ R symmetry. Gaiotto and Witten [4] give the associated brane supersymmetry projectors in their study of supersymmetric boundary conditions. We use the unfluxed intersection projector, not their additional conditions for D3-branes terminating on a D5-brane. Baker [5] constructed supersymmetric open Wilson lines in this defect theory, including a semicircle obtained from a ray by a conformal transformation, and analyzed their scalar endpoints. These examples are controls for our endpoint test with a constant Poincare charge; that test does not exclude Baker’s semicircle.

Loewner’s equation describes the conformal maps of growing planar hulls. We use the chordal normalization in Lawler’s lectures [6]. The relation between stochastic Loewner evolution and conformal-field-theory martingales developed by Bauer and Bernard [7] is relevant motivation. A map-level stochastic differential equation, however, does not by itself construct a renormalized Wilson observable on a rough trace. We work with smooth curves and regular pullbacks; the one stochastic formula below is explicitly limited to the inverse map.

Osterwalder–Schrader reflection positivity [8] explains how Euclidean reflected pairings can define Hilbert-space norms. For scalar-coupled Wilson transport, reflection must account for both complex conjugation and path ordering. Dorn and Pershin [9] analyzed this distinction and separated ordinary reflection inequalities from a conjectural inequality involving an internal rotation in a special large-rectangle limit. Our internal-rotation counterexample concerns the full defect scalar doublet and does not address that restricted conjecture.

On the arithmetic side, Weil’s positivity criterion provides the logical target. Suzuki’s recent treatment [10] places the localized Weil form, the screw function and associated operator constructions in a common framework, with references to the earlier work of Weil, Yoshida, Bombieri, and Connes

and collaborators. We use the compact-support criterion, not a conjectural limiting spectral construction. The present field-theoretic calculations establish no new equivalence with RH.

1.2 What survives from the parent investigations

The preceding Wilson-lines, Loewner, and fractional-dimension investigations [11, 13, 14] are working research documents in the same program. Their relevant identities are reproduced here, so they are not prerequisites for reading this paper. In particular, the shifted transfer has an explicit positive Beta-kernel/prime-comb part and an essential rational correction. The earlier direct Loewner realization tests did not identify a bare map observable producing this entire arithmetic object. The fractional-dimension calculation explains the gamma ratio by a dimensionally continued integral, but a natural continued lattice count has negative coefficients. These are restrictions on particular realizations, not an exclusion of field-dependent observables on Loewner-generated curves.

The combined construction studied here therefore has four separate ingredients: a contour law; a gauge and scalar connection on that contour; endpoint and reference states with a specified adjoint; and an eventual arithmetic identity. We establish parts of the first three. We do not identify the fourth.

1.3 Results and scope

The argument proceeds through the following results.

1. Inversion sends the inherited common-origin semicircles to vertical slits of constant-driver chordal evolution. Under the fixed scalar and special bulk charge specified in Section 4, those curves are rigid.
2. The exact smooth variation of an open Wilson transport contains field-strength, scalar-gradient, arclength, and endpoint terms. An abelian counterexample rules out a universal geometry-only scalar closure of the transport variation (Section 5).
3. A real tangent isometry compatible with chirality and the D3–D5 projector admits a two-dimensional complex charge space in a normal plane, and a one-dimensional space across all defect directions. Its orientation matters (Section 6).
4. Ordinary scalar endpoints annihilated by the same constant charge form a zero-contraction pair. The full bilinear has nonzero residual R charge and zero expectation in an invariant vacuum. Physical adjoints evade the zero contraction but change the supersymmetry question (Section 7).
5. Ordinary time reflection gives a different scalar map on the reflected branch. The resulting free kernel is nonzero and positive. Positivity of the interacting network is stated conditionally, with its color and regulator requirements (Section 8).
6. A family with a smooth reference join has a finite direct Gaussian bulk response with leading coefficient $3/2 - 2\log 2 > 0$. Endpoint, defect and renormalization contributions have not been computed (Section 9).

All supersymmetry ranks are complex ranks in a specified Euclidean Clifford module. All transport variation theorems are statements about smooth finite matrix fields. A regulated quantum use of those identities requires additional analytic and renormalization input. The finite diagnostics in Appendix A check signs, ranks and integrals; they neither establish a continuum quantum field theory nor prove an inequality for the Weil form.

2 The arithmetic positivity target

This section fixes the target independently of any field-theory model. The formulas also explain why positivity of a single covariance kernel would be insufficient.

2.1 The compactly supported Weil form

Let $I_L = (-L/2, L/2)$, with $L > 0$, and let $f \in C_c^\infty(I_L; \mathbb{C})$. Write F for its zero extension to $\mathbb{R}$ and use

$$\widehat{F}(\tau) = \int_{\mathbb{R}} e^{-i\tau x} F(x) dx, \quad \langle f, g \rangle = \int_{I_L} \overline{f(x)} g(x) dx. \quad (2.1)$$

For $a > 0$, define the compressed shift $T_a f(x) = F(x - a)$ on I_L . Let $\psi = \Gamma'/\Gamma$ be the digamma function, and let $\Lambda(n) = \log p$ when $n = p^k$, p prime and $k \geq 1$, and zero otherwise. Finally put

$$C(f) = \int_{I_L} f(x) \cosh(x/2) dx, \quad S(f) = \int_{I_L} f(x) \sinh(x/2) dx. \quad (2.2)$$

The normalization of the program is

$$\begin{aligned} Q_{0,L}[f] &= \frac{1}{2\pi} \int_{\mathbb{R}} [\operatorname{Re} \psi(\tfrac{1}{4} + \tfrac{i\tau}{2}) - \log \pi] |\widehat{F}(\tau)|^2 d\tau + 2|C(f)|^2 - 2|S(f)|^2 \\ &\quad - \sum_{\substack{n \geq 2 \\ \log n < L}} \frac{\Lambda(n)}{\sqrt{n}} \langle f, (T_{\log n} + T_{\log n}^*) f \rangle. \end{aligned} \quad (2.3)$$

The prime-power sum is finite on a fixed interval. The pole term is indefinite, and the archimedean multiplier includes its local constant. Neither may be dropped in a positivity argument.

For comparison with the usual explicit formula, set $h = F * \tilde{F}$, where $\tilde{F}(x) = \overline{F(-x)}$. The two prime sums in the Weil functional give the last line of (2.3); its $e^{x/2} + e^{-x/2}$ term gives $2|C|^2 - 2|S|^2$. Its archimedean distribution has Fourier multiplier $\operatorname{Re} \psi(1/4 + i\tau/2) - \log \pi$. Thus $Q_{0,L}[f] = W(F * \tilde{F})$ in the convention of [10, p. 1]. In particular the weight is $\Lambda(n)/\sqrt{n}$.

Theorem 2.1 (Weil positivity criterion, used as background). *RH holds if and only if $Q_{0,L}[f] \geq 0$ for every $L > 0$ and every $f \in C_c^\infty(I_L; \mathbb{C})$.*

The equivalence is the classical criterion, not a result of the present construction. Every compact support lies in one such interval, so an unbounded sequence of interval lengths suffices. A finite collection of lengths or numerical matrices does not.

2.2 The positive kinetic term and the indispensable local term

Put $a_n = 2n + 1/2$, for $n \geq 0$, and

$$b(\tau^2) = \operatorname{Re} \psi(\tfrac{1}{4} + \tfrac{i\tau}{2}) - \psi(\tfrac{1}{4}) = 2 \sum_{n=0}^{\infty} \frac{\tau^2}{a_n(a_n^2 + \tau^2)} \geq 0, \quad w_0 = \psi(\tfrac{1}{4}) - \log \pi < 0. \quad (2.4)$$

The series follows by subtracting the convergent digamma partial fractions. Each exponential in (1.1) satisfies $\int_0^\infty e^{-au} (1 - \cos \tau u) du = \tau^2/[a(a^2 + \tau^2)]$. Plancherel therefore gives

$$\mathcal{E}_\Gamma[F] := \int_0^\infty n_\Gamma(u) \|F(\cdot + u) - F\|_2^2 du = \frac{1}{2\pi} \int_{\mathbb{R}} b(\tau^2) |\widehat{F}(\tau)|^2 d\tau. \quad (2.5)$$

Consequently the first line of (2.3), before its pole term, is $\mathcal{E}_\Gamma[F] + w_0 \|f\|_2^2$. The positive difference energy is only one part of the desired form.

The natural form domain on a fixed interval is given by zero extensions with $\int \log(2 + |\tau|)|\widehat{F}(\tau)|^2 d\tau < \infty$. Indeed $b(\tau^2)$ grows logarithmically at infinity, whereas the pole and finite translation terms are bounded forms on $L^2(I_L)$. This explains the logarithmic form domain and semiboundedness of the localized form. For the logical proof target in Theorem 2.1, the smooth compactly supported class is enough.

2.3 Shifted transfers and a sufficient contraction route

Define the entire completed zeta function and its centered version by

$$\xi(s) = \frac{1}{2}s(s-1)\pi^{-s/2}\Gamma(s/2)\zeta(s), \quad \Xi(p) = \xi(\frac{1}{2} + p). \quad (2.6)$$

For $0 < \omega \leq 1/2$, the shifted transfer is

$$K_\omega(p) = \frac{\Xi(p - \omega)}{\Xi(p + \omega)}. \quad (2.7)$$

The functional equation and reality imply $|K_\omega(i\tau)| = 1$ where the ratio is defined. This boundary identity is unconditional; it is not a proof that an unweighted convolution operator is a contraction. In particular poles away from that boundary cannot be ignored.

The causal interpretation uses inverse Laplace transform on a line $\operatorname{Re} p = \sigma > 1$. Let k_ω be that causal distribution and $V_{\omega,L} = P_L(k_\omega *) P_L$, where P_L denotes restriction to I_L together with zero extension when used as an input. The explicit kernel below is locally integrable for $\omega > 0$, so these finite-window operators are bounded. At $\omega = 0$, $k_0 = \delta_0$ and $V_{0,L} = I$. Logarithmic differentiation gives

$$\partial_\omega K_\omega(p) = -a_\omega(p)K_\omega(p), \quad a_\omega(p) = \frac{\Xi'(p - \omega)}{\Xi(p - \omega)} + \frac{\Xi'(p + \omega)}{\Xi(p + \omega)}. \quad (2.8)$$

At zero shift the causal generator has Laplace symbol

$$a_0(p) = \frac{2}{p + 1/2} + \frac{2}{p - 1/2} - \log \pi + \psi(\frac{1}{4} + \frac{p}{2}) - 2 \sum_{n \geq 2} \frac{\Lambda(n)}{\sqrt{n}} e^{-p \log n}. \quad (2.9)$$

The identity

$$\psi(\frac{1}{4} + \frac{p}{2}) - \psi(\frac{1}{4}) = 2 \int_0^\infty n_\Gamma(u)(1 - e^{-pu}) du \quad (2.10)$$

shows that the real quadratic form of this causal generator is exactly (2.3). The two rational terms give the symmetric kernel $2 \cosh((x - y)/2)$; its form is $2|C|^2 - 2|S|^2$. On smooth test functions the first variation is therefore

$$\langle f, (I - V_{\omega,L}^* V_{\omega,L}) f \rangle = 2\omega Q_{0,L}[f] + o(\omega), \quad \omega \downarrow 0. \quad (2.11)$$

One obtains this expansion by differentiating the causal kernel at $\omega = 0$ using (2.9); smooth zero extensions make the subtracted difference integral convergent in L^2 on a finite window. Thus an independent proof of $\|V_{\omega,L}\| \leq 1$ for every L and shifts tending to zero would imply RH. We need only this sufficient implication here; no global unweighted realization of K_ω is assumed.

2.4 The exact positive part, and what it leaves unresolved

Write $a = 1/2 - \omega$, $b = 1/2 + \omega$, $s = 1/2 + p$, and $\mathcal{L}(s) = \pi^{-s/2} \Gamma(s/2) \zeta(s)$, distinguished from the von Mangoldt function. Direct factorization yields

$$K_\omega(p) = \frac{(p+a)(p-b)}{(p+b)(p-a)} \widetilde{K}_\omega(p), \quad \widetilde{K}_\omega(p) = \frac{\mathcal{L}(s-\omega)}{\mathcal{L}(s+\omega)}. \quad (2.12)$$

The beta integral and Euler product, for $\operatorname{Re} p > b$, give

$$g_\omega(t) = \frac{2\pi^\omega}{\Gamma(\omega)} e^{-at} (1 - e^{-2t})^{\omega-1} \mathbf{1}_{t>0}, \quad (2.13)$$

$$c_n(\omega) = n^{\omega-1/2} \prod_{\ell|n, \ell \text{ prime}} (1 - \ell^{-2\omega}), \quad c_1 = 1, \quad (2.14)$$

$$\widetilde{k}_\omega = g_\omega * \sum_{n \geq 1} c_n(\omega) \delta_{\log n} \geq 0. \quad (2.15)$$

For example, the substitution $r = e^{-2t}$ in the Laplace transform of (2.13) gives $\pi^\omega \Gamma((p+a)/2) / \Gamma((p+b)/2)$. For the zeta quotient the coefficient at a prime power ℓ^k is $\ell^{k\omega} (1 - \ell^{-2\omega})$ before multiplication by $n^{-1/2}$, proving (2.14) by multiplicativity. Only finitely many shifts contribute on a fixed window. These formulas define positive locally finite measures, not normalized probability laws; the endpoint $a = 0$ in particular precludes an unqualified Beta($a/2, \omega$) probability interpretation.

Absorb one rational factor into the positive part:

$$\begin{aligned} \widehat{k}_\omega &= \widetilde{k}_\omega + 2a(e^a \cdot \mathbf{1}_{>0}) * \widetilde{k}_\omega \geq 0, \\ k_\omega &= \widehat{k}_\omega - 2b(e^{-b} \cdot \mathbf{1}_{>0}) * \widehat{k}_\omega. \end{aligned} \quad (2.16)$$

Thus the full kernel is a positive part minus an exponential average. This exact reduction was central to the parent Loewner investigation [13]. It neither makes k_ω positive nor bounds the norm of its finite-window convolution. The required cancellation is still the arithmetic problem.

The fractional-dimension interpretation follows from the radial beta integral

$$\int_{\mathbb{R}^d} (1 + |u|^2)^{-(p+b)/2} d^d u = \pi^{d/2} \frac{\Gamma((p+b-d)/2)}{\Gamma((p+b)/2)}. \quad (2.17)$$

For integer d this is an ordinary integral in its convergence region; its radial continuation at $d = 2\omega$ reproduces the gamma factor. It is not a continuously variable lattice dimension. To see a concrete obstruction to one natural lattice continuation, put $\vartheta(q) = \sum_{k \in \mathbb{Z}} q^{k^2}$ and $m = 1 + d$. The coefficient of q^3 in $\vartheta(q)^m$ is

$$8 \binom{m}{3} = \frac{4}{3} m(m-1)(m-2) = \frac{4}{3} (d^3 - d) < 0 \quad (0 < d < 1). \quad (2.18)$$

It cannot be a positive point count. This excludes that interpolation, not every positive kernel or analytic continuation. Neither (2.17) nor a bare Loewner map supplies the prime comb and rational subtraction in (2.15)–(2.16).

3 Why defect matter reproduces the archimedean exponents

The geometric reason for using three-dimensional defect matter can be seen without supersymmetry. Normalize a free massless scalar covariance on the defect to $|X - Y|^{-1}$; coupling and color constants

will be restored when needed. Write $X = e^x \Omega$, $Y = e^y \Omega'$ with $\Omega, \Omega' \in S^2$, and rescale the field by $e^{x/2}$. With $u = x - y$ and $v = \Omega \cdot \Omega'$, the covariance becomes

$$C(u, v) = \frac{1}{\sqrt{2(\cosh u - v)}} = \sum_{\ell=0}^{\infty} e^{-(\ell+1/2)|u|} P_{\ell}(v), \quad u \neq 0. \quad (3.1)$$

This is the Legendre generating function with parameter $e^{-|u|}$.

Fix an axis $e \in S^2$ and $0 \leq \rho < 1$. The Poisson density on S^2 and its even part are

$$P_{\rho}(v) = \frac{1 - \rho^2}{4\pi(1 - 2\rho v + \rho^2)^{3/2}}, \quad h_{\rho}(\Omega) = \frac{1}{2}\{P_{\rho}(e \cdot \Omega) + P_{\rho}(-e \cdot \Omega)\}. \quad (3.2)$$

Integrating (3.1) against $h_{\rho}(\Omega)h_{\rho}(\Omega')$ projects onto even angular degrees. Orthogonality of the Legendre polynomials gives

$$C_{\rho}(u) = \sum_{n=0}^{\infty} \rho^{4n} e^{-a_n|u|} = \frac{e^{-|u|/2}}{1 - \rho^4 e^{-2|u|}}, \quad a_n = 2n + \frac{1}{2}. \quad (3.3)$$

For $\rho < 1$ this is a bounded continuous positive-definite covariance. For $u \neq 0$ it tends to $n_{\Gamma}(|u|)$ as $\rho \uparrow 1$. The limit is singular at $u = 0$, so positivity of the regulated covariance cannot simply be passed to a finite contact-subtracted covariance.

Proposition 3.1 (The sign of the subtraction). *Let T_{ρ} be convolution by C_{ρ} on $\mathbb{R}$, and put $M_{\rho} = \widehat{C}_{\rho}(0)$. Then $M_{\rho}I - T_{\rho}$ is positive, and its quadratic form tends to $\mathcal{E}_{\Gamma}$ on smooth compactly supported functions. The opposite subtraction $T_{\rho} - M_{\rho}I$ tends to the negative of that form. Adding any finite constant multiple of the identity to the latter limit does not give a positive form on all smooth compactly supported functions.*

Proof. Fourier transformation gives

$$\begin{aligned} \widehat{C}_{\rho}(\tau) &= \sum_{n \geq 0} \rho^{4n} \frac{2a_n}{a_n^2 + \tau^2}, \\ M_{\rho} - \widehat{C}_{\rho}(\tau) &= 2 \sum_{n \geq 0} \rho^{4n} \frac{\tau^2}{a_n(a_n^2 + \tau^2)} \uparrow b(\tau^2). \end{aligned} \quad (3.4)$$

Monotone convergence and (2.5) prove the first claims. Moreover $M_{\rho} = \sum_{n \geq 0} \rho^{4n}/(n + 1/4)$ diverges logarithmically. Since $b(\tau^2) \rightarrow \infty$, any multiplier $c - b(\tau^2)$ is negative at high frequencies. Modulating a fixed nonzero compactly supported smooth function to arbitrarily high frequency gives a negative form value, also on any fixed nonempty interval. $\square$

There is a second useful decomposition. For $u \neq 0$ define the same-ray and opposite-ray functions

$$G_s(u) = \frac{1}{2 \sinh(|u|/2)}, \quad G_o(u) = \frac{1}{2 \cosh(u/2)}. \quad (3.5)$$

Then $n_{\Gamma}(|u|) = \frac{1}{2}(G_s(u) + G_o(u))$. The reflected free network constructed in Section 8 gives G_o . That is a positive and finite kernel, but it is not the even tower by itself. Conversely, the full angular smearing above is an ordinary covariance construction with support on all of S^2 . It is not automatically a state supported in a positive-time half-space, as required for a time-reflection argument. Restricting the angular profile to a hemisphere changes its spectral weights.

The useful conclusion is therefore precise: free defect matter supplies the desired exponents and a regulated positive covariance. The RH target requires the associated difference energy, its fixed local term, its pole term and its arithmetic translations. These are distinct operations on the free data.

3.1 An auxiliary Gaussian realization of the gamma ratio

The parent endpoint work also gave a useful fixed-space alternative to fractional dimension [12]. It can be understood from elementary Gaussian integration. Retain one complex mode for each $a_n = 2n + 1/2$, and for real p with $a_0 + p > \omega$ define

$$Z_N(p) = \prod_{n=0}^{N-1} \int_{\mathbb{C}} \frac{d^2 z_n}{\pi} e^{-(a_n+p)|z_n|^2} = \prod_{n=0}^{N-1} (a_n + p)^{-1}. \quad (3.6)$$

For a half-line field $\phi_n(t)$ with action $\int_0^\infty (|\phi'_n|^2 + a_n^2 |\phi_n|^2) dt$ and boundary value z_n , the decaying classical solution is $\phi_n(t) = z_n e^{-a_n t}$. Its action is $a_n |z_n|^2$. Adding the boundary term $p|z_n|^2$ gives the same Gaussian; varying the boundary value gives the Robin condition $(-\partial_t + p)\phi_n(0) = 0$ for the full variational problem. Constants common to the two partition functions cancel in the ratio.

Set $x = (p + 1/2 - \omega)/2$ and $y = x + \omega$. The finite product gives

$$\frac{Z_N(p - \omega)}{Z_N(p + \omega)} = \prod_{n=0}^{N-1} \frac{n + y}{n + x} = \frac{\Gamma(N + y)\Gamma(x)}{\Gamma(y)\Gamma(N + x)}. \quad (3.7)$$

Since $\Gamma(N + y)/\Gamma(N + x) \sim N^\omega$,

$$\pi^\omega \lim_{N \rightarrow \infty} N^{-\omega} \frac{Z_N(p - \omega)}{Z_N(p + \omega)} = \pi^\omega \frac{\Gamma((p + a)/2)}{\Gamma((p + b)/2)}. \quad (3.8)$$

Equivalently one can divide by the same finite product at a reference p_0 to cancel the normalization. One real Gaussian mode would give the square root of the ratio, so statistics and multiplicities matter.

This is an auxiliary partition-function model, not a localization of the Wilson network. Angular smearing samples a channel; it does not remove the other angular modes from a partition function. The full scalar sphere has multiplicity $2\ell + 1$, whereas the model retains one copy of each even axial mode. A physical projection or cancellation of the extra modes has not been derived. Here p is a boundary mass, and identifying it with the arithmetic Laplace frequency requires a dynamical dictionary. The Robin response $(a_n + p)^{-1}$, the partition ratio, and a causal transfer are different objects. The supplied factor π^ω , the prime comb, and the rational pole correction remain outside this Gaussian mechanism.

4 Semicircles, constant drivers, and a restricted rigidity theorem

Choose a plane containing a direction tangent to the defect and its normal, and write $z = x + iy$ with $y > 0$. The common-reference semicircle from $r > 0$ to zero is

$$z_r(\theta) = \frac{r}{2}(1 + \cos \theta + i \sin \theta), \quad 0 \leq \theta < \pi. \quad (4.1)$$

Inversion preserves the upper half-plane and gives

$$w_r(\theta) = -\frac{1}{z_r(\theta)} = -\frac{1}{r} + \frac{i}{r} \tan(\theta/2). \quad (4.2)$$

This is a vertical slit. In the chordal convention

$$\partial_t g_t(w) = \frac{2}{g_t(w) - U_t}, \quad g_0(w) = w, \quad (4.3)$$

the constant driver $U_t = a = -1/r$ has solution

$$g_t(w) = a + \sqrt{(w-a)^2 + 4t}, \quad \gamma_t = a + 2i\sqrt{t}, \quad z_t = -\frac{1}{a + 2i\sqrt{t}}. \quad (4.4)$$

The square-root branch is fixed by $g_t(w) \sim w$ at infinity. Its expansion is $g_t(w) = w + 2t/w + O(w^{-2})$, so the half-plane capacity is $2t$, consistent with [6, Lecture 2, Proposition 2.17]. Comparing the traces gives the exact dictionary

$$t = \frac{\tan^2(\theta/2)}{4r^2}. \quad (4.5)$$

The complete semicircle reaches the reference only as $t \rightarrow \infty$. At fixed angle, $r \mapsto \lambda r$ sends $t \mapsto \lambda^{-2}t$. Thus this inverted capacity is neither the coordinate $\log r$ nor the arithmetic shift ω . The formulas identify contours; they do not assert equality of renormalized expectations through an inversion singular at an endpoint.

4.1 Fixed scalar coupling and fixed special charge

Let the Euclidean Clifford generators satisfy $\{\Gamma_A, \Gamma_B\} = 2\delta_{AB}$. Take one internal direction I and fix a nonzero complex spinor ϵ_c obeying

$$\epsilon_s = 0, \quad J\epsilon_c = -\epsilon_c, \quad J = i\Gamma_I\Gamma_3. \quad (4.6)$$

Here $\epsilon_s + X^\mu \Gamma_\mu \epsilon_c$ is the relevant conformal Killing spinor, with its constant part set to zero. These are the inherited common bulk conditions. For a regular real curve $X = (x, y) \neq 0$ in the $(1, 3)$ -plane and constant positive scalar arclength coupling, its local line condition is

$$(i\dot{x}\Gamma_1 + i\dot{y}\Gamma_3 + v\Gamma_I)(x\Gamma_1 + y\Gamma_3)\epsilon_c = 0, \quad v = |\dot{X}| > 0. \quad (4.7)$$

Proposition 4.1 (Rigidity under (4.6)). *Equation (4.7) holds if and only if*

$$\frac{\dot{z}}{|\dot{z}|} = \frac{iz^2}{|z|^2}. \quad (4.8)$$

Consequently $-1/z$ has constant real part and increasing imaginary part. Connected solutions in the upper half-plane are subarcs of circles through zero with center on the real axis, or the degenerate imaginary-axis line. A solution from a nonzero real boundary point to zero is the corresponding oriented upper semicircle.

Proof. The projector implies $\Gamma_I\epsilon_c = -i\Gamma_3\epsilon_c$. Clifford multiplication reduces (4.7) to $(A + B\Gamma_1\Gamma_3)\epsilon_c = 0$, where

$$A = x\dot{x} + y\dot{y} + vy, \quad B = y\dot{x} - x\dot{y} + vx. \quad (4.9)$$

Since $(\Gamma_1\Gamma_3)^2 = -1$, multiplication by $A - B\Gamma_1\Gamma_3$ gives $(A^2 + B^2)\epsilon_c = 0$. The coefficients are real and the spinor is nonzero, so $A = B = 0$. Solving gives

$$\frac{\dot{x}}{v} = -\frac{2xy}{x^2 + y^2}, \quad \frac{\dot{y}}{v} = \frac{x^2 - y^2}{x^2 + y^2}, \quad (4.10)$$

which is (4.8). Conversely it makes $A = B = 0$. Moreover $d(-1/z)/ds = iv/|z|^2$. The level set $\text{Re}(-1/z) = a \neq 0$ is $x^2 + y^2 = -x/a$, and $a = 0$ gives the imaginary axis. $\square$

For instance, an inverted deformation $w(s) = a + \varepsilon h(s)$ preserves these conditions only if $\varepsilon h'(s) = 0$. A constant h changes the slit location; a nontrivial fixed-endpoint deformation fails. A regular capacity-parametrized Loewner trace satisfying the same conditions therefore has constant driving.

This is a bulk theorem for one plane, one charge and one scalar prescription. It is not a classification of supersymmetric contours with varying scalar directions, other conformal Killing spinors, additional endpoint degrees of freedom, or nonplanar embeddings. Section 6 supplies a constructive control showing that a changed scalar prescription allows arbitrary tangents.

5 Smooth Wilson variation and the Loewner insertion

5.1 Conventions and the exact variation

Work with smooth finite matrix fields and a regular C^2 curve $X : [0, 1] \rightarrow \mathbb{R}^4$. Let A_μ and the scalar fields X_I be Hermitian, and let $n^I(s)$ be a unit internal vector. Define

$$\mathcal{M}(s) = iA_\mu(X(s))\dot{X}^\mu(s) + \Phi(X(s), s)v(s), \quad \Phi = n^I X_I, \quad v = |\dot{X}|. \quad (5.1)$$

The ordered transport $U(b, a)$ solves $\partial_b U(b, a) = \mathcal{M}(b)U(b, a)$, $U(a, a) = I$; larger parameters stand to the left. Our curvature and covariant derivative are

$$F_{\nu\mu} = \partial_\nu A_\mu - \partial_\mu A_\nu - i[A_\nu, A_\mu], \quad D_\nu \Phi = \partial_\nu \Phi - i[A_\nu, \Phi]. \quad (5.2)$$

Vary the curve by $\eta = \delta X$ and the internal vector by δn , holding the spacetime fields fixed.

Proposition 5.1 (Ordered transport variation). *With $C_s = iA_\nu(X(s))\eta^\nu(s)$,*

$$\delta U(1, 0) = C_1 U(1, 0) - U(1, 0) C_0 + \int_0^1 U(1, s) \mathcal{I}(s) U(s, 0) ds, \quad (5.3)$$

$$\mathcal{I} = iF_{\nu\mu}\eta^\nu \dot{X}^\mu + v D_\nu \Phi \eta^\nu + \Phi \frac{\dot{X} \cdot \dot{\eta}}{v} + v X_I \delta n^I. \quad (5.4)$$

Proof. Differentiating the defining ODE gives $\delta U = \int U \delta \mathcal{M} U$. Direct differentiation of $\mathcal{M}$ yields $\delta \mathcal{M} = \dot{C} + [C, \mathcal{M}] + \mathcal{I}$. Meanwhile

$$\frac{d}{ds} \{U(1, s) C_s U(s, 0)\} = U(1, s) (\dot{C} + [C, \mathcal{M}]) U(s, 0). \quad (5.5)$$

Integration proves the formula, including its ordering and boundary signs. $\square$

For constant n and fixed endpoints, integrate the arclength term by parts. Writing $T = dX/d\ell$, $k = dT/d\ell$, one obtains

$$\begin{aligned} \delta U &= \int_C U(1, s) \eta^\nu \mathcal{D}_\nu U(s, 0) d\ell, \\ \mathcal{D}_\nu &= iF_{\nu\mu} T^\mu + (\delta_\nu^\mu - T_\nu T^\mu) D_\mu \Phi - \Phi k_\nu. \end{aligned} \quad (5.6)$$

In particular $T^\nu \mathcal{D}_\nu = 0$, as required for a fixed-endpoint reparametrization. The curvature term is necessary because the scalar couples to arclength. If n varies, one must retain its additional terms in (5.4).

Let the endpoints be $x = X(0)$ and $y = X(1)$, both on the defect, and consider $\mathcal{O} = \bar{q}(y)U(1,0)q(x)$. The boundary terms in (5.3) combine with endpoint differentiation to give

$$\begin{aligned} \delta\mathcal{O} &= \eta_y^\nu(\partial_\nu\bar{q} + i\bar{q}A_\nu)(y)Uq(x) + \bar{q}(y)U\eta_x^\nu(D_\nu q)(x) \\ &\quad + \int_0^1 \bar{q}(y)U(1,s)\mathcal{I}(s)U(s,0)q(x) ds. \end{aligned} \quad (5.7)$$

Here $D_\nu q = \partial_\nu q - iA_\nu q$, and endpoint displacements are tangent to the defect. Changing endpoint R polarizations contributes further terms. A defect field q is not defined at a bulk tip.

5.2 Regular inverse-map deformations

Let $f_t = g_t^{-1}$ for (4.3), and choose a fixed smooth reference contour $\zeta(s)$ in a region where f_t is regular and $\zeta(s) \neq U_t$. Differentiating $g_t(f_t(w)) = w$ gives

$$X_t(s) = f_t(\zeta(s)), \quad \eta_t(s) = \partial_t X_t(s) = -\frac{2f_t'(\zeta(s))}{\zeta(s) - U_t}. \quad (5.8)$$

Insert its real and imaginary components into (5.4) and (5.7). This is a concrete field-dependent evolution of the transport on a pulled-back test curve. It is not a variation formula for the singular growing tip. Regular real boundary segments remain on the defect in this planar interpretation. A growing prefix instead carries an uncontracted color index at its interior endpoint; it requires a specified tip state or a conditional completion of the curve.

If $dU_t = \sqrt{\kappa}dB_t$ for standard Brownian motion B_t , then at fixed w the inverse-map equation is of finite variation until a singularity is reached. In centered coordinates $\mathfrak{f}_t(w) = f_t(w + U_t)$, stopped smooth Ito calculus gives

$$d\mathfrak{f}_t(w) = \left[-\frac{2\mathfrak{f}_t'(w)}{w} + \frac{\kappa}{2}\mathfrak{f}_t''(w) \right] dt + \sqrt{\kappa}\mathfrak{f}_t'(w)dB_t. \quad (5.9)$$

This is only a map identity. No scalar arclength integral or quantum Wilson transport on an unregulated stochastic trace is defined by it.

5.3 A precise obstruction to geometry-only closure

Proposition 5.2. *There is no universal multiplicative shape generator for a single Wilson transport whose coefficient depends only on the curve and its deformation, independently of the background fields.*

Proof. In an abelian theory take $A_x = cy$, $A_y = \Phi = 0$ and the straight path $X(s) = (s, 0)$, $0 \leq s \leq 1$. Then $U = 1$ for every real c . For $\eta = (0, h(s))$ with $h(0) = h(1) = 0$,

$$\delta U = ic \int_0^1 h(s) ds. \quad (5.10)$$

The same curve, deformation and original transport thus have different derivatives. As a positive control, a pure gauge $A = d\chi$ has fixed-endpoint transport independent of the contour. $\square$

The proposition does not exclude closure of an averaged correlation function after field equations, Ward identities or a protected reduction are supplied. It identifies the extra information such a closure must use. At leading free endpoint order $U = I$, fixed-endpoint interior deformations do not change (3.3); the nontrivial information must enter at higher order or through another specified sector.

5.4 The tangent connection as a one-form

For a constant real isometry $M : \mathbb{R}^4 \rightarrow \mathbb{R}^6$, set $n = MT$, so that $\Phi|dX| = \Phi_\mu dX^\mu$ with $\Phi_\mu = M^A{}_\mu X_A$. Define $\mathcal{B}_\mu = iA_\mu + \Phi_\mu$. Then the fixed-endpoint variation takes the ordinary one-form form

$$\begin{aligned} \delta U_M &= \int U_M(1, s) \mathcal{F}_{\nu\mu} \eta^\nu \dot{X}^\mu U_M(s, 0) ds, \\ \mathcal{F}_{\nu\mu} &= \partial_\nu \mathcal{B}_\mu - \partial_\mu \mathcal{B}_\nu - [\mathcal{B}_\nu, \mathcal{B}_\mu] \\ &= iF_{\nu\mu} + D_\nu \Phi_\mu - D_\mu \Phi_\nu - [\Phi_\nu, \Phi_\mu]. \end{aligned} \quad (5.11)$$

For moving endpoints the boundary insertion is $\mathcal{B}_\nu \eta^\nu$. These signs follow from the same ordered ODE. Supersymmetry of the line is not an assertion that this complex curvature vanishes on arbitrary field configurations.

6 Tangent coupling with the full chiral defect projector

6.1 Field content and conventions

The defect is $x^3 = 0$ in Euclidean spacetime (x^0, x^1, x^2, x^3) . The bulk has a gauge field A_μ , six adjoint scalar fields X_A , $A = 4, \dots, 9$, and their supersymmetric fermion partners. The triplets (X_4, X_5, X_6) and (X_7, X_8, X_9) transform under $SU(2)_H$ and $SU(2)_V$, respectively. Defect matter includes fundamental scalars q_m , $m = 1, 2$, forming an H doublet, and fermions forming a V doublet; $\bar{q}_m$ are the physical antifundamental conjugates. These are the bulk/defect assignments of [3, 5].

Work in a 32-dimensional complex Euclidean Clifford module, with

$$\chi = -i\Gamma_0\Gamma_1 \cdots \Gamma_9, \quad D = \Gamma_3\Gamma_4\Gamma_5\Gamma_6, \quad \chi\epsilon = \epsilon, \quad D\epsilon = \epsilon. \quad (6.1)$$

The second equation is the unfluxed D3–D5 intersection condition. It is the projector appearing in [4, eqs. (2.7), (4.1)–(4.3), (4.9)]; the terminating-brane boundary conditions and Nahm pole are not part of our assumptions. The displayed conventions fix the chirality and orientation signs. Complex ranks below are not counts of real Euclidean Majorana–Weyl supercharges.

6.2 A normal plane

For the (x^1, x^3) -plane take the connection

$$\mathcal{B} = iA_1 dx^1 + iA_3 dx^3 + X_8 dx^1 + X_6 dx^3. \quad (6.2)$$

This is the tangent prescription $n = (\dot{x}^1 e_8 + \dot{x}^3 e_6)/|\dot{x}|$, so the scalar direction changes with orientation. In the usual local supersymmetry variation of scalar-coupled transport, a constant Poincare parameter is annihilated by every tangent if

$$[\dot{x}^1(i\Gamma_1 + \Gamma_8) + \dot{x}^3(i\Gamma_3 + \Gamma_6)]\epsilon = 0. \quad (6.3)$$

This is the tangent construction of [2], now with (6.1) imposed as well. Define

$$K_1 = i\Gamma_8\Gamma_1, \quad K_n = i\Gamma_6\Gamma_3. \quad (6.4)$$

Proposition 6.1 (Planar charge space). *The common subspace $\chi = D = +1$, $K_1 = K_n = -1$ has complex dimension two. Every parameter in it obeys (6.3) for every smooth planar tangent.*

Proof. The four displayed matrices are commuting Hermitian involutions. Their orthogonal projector is

$$P_{\text{plane}} = \frac{1}{16}(1 + \chi)(1 + D)(1 - K_1)(1 - K_n). \quad (6.5)$$

Every nonidentity Clifford word in its expansion has zero trace, and no nonempty product of these four generators is scalar. Hence $\text{tr } P_{\text{plane}} = 32/16 = 2$. Multiplication of $K_1\epsilon = -\epsilon$ by Γ_8 , and of $K_n\epsilon = -\epsilon$ by Γ_6 , gives the two linear equations in (6.3) separately. $\square$

The triplet assignments are necessary within the stated real, constant-map ansatz. The defect projector D commutes with a defect tangent gamma and with a V gamma; it anticommutes with the normal gamma and with an H gamma. Pairing a tangent with H, or the normal with V, therefore gives an involution anticommuting with D and no simultaneous eigenspinor. More generally, split a proposed real scalar vector into H and V components. For an independently variable defect tangent its wrong H component lies in the opposite D -eigenspace from the remaining line equation, so it must annihilate ϵ separately. Its Clifford square is its squared real length, making it invertible unless that component is zero. The normal argument exchanges H and V. Complex null internal vectors and couplings restricted to one constrained tangent are outside this argument. Proposition 6.1 is the nonzero control.

6.3 One charge across all defect directions

To accommodate an angular family, introduce three tangent conditions

$$K_{\mu}^{(s)} = is_{\mu}\Gamma_{7+\mu}\Gamma_{\mu}, \quad \mu = 0, 1, 2, \quad s_{\mu} \in \{+1, -1\}. \quad (6.6)$$

The corresponding oriented tangent isometry has determinant $s_0s_1s_2$. Direct Clifford multiplication gives

$$K_0^{(s)} K_1^{(s)} K_2^{(s)} = (s_0s_1s_2)\chi D. \quad (6.7)$$

Proposition 6.2 (Orientation and the all-direction charge). *For chirality $\chi = c$ and defect sign $D = d$, all three tangent eigenvalues can be -1 , together with $K_n = -1$, if and only if the tangent isometry has determinant $-cd$. With that sign the common space has complex dimension one; with the opposite sign it is zero.*

Proof. Taking eigenvalues in (6.7) proves necessity. For the allowed sign, one tangent condition follows from the other two and χ, D . The remaining five commuting conditions $\chi = c$, $D = d$, $K_n = -1$, $K_0^{(s)} = K_1^{(s)} = -1$ are independent. Their projector has trace $32/2^5 = 1$, by the same nonidentity-word trace argument as above. Orthogonal changes of the oriented tangent and internal frames give the general isometry statement. $\square$

For $c = d = +1$, choose once and for all

$$Me_0 = -e_7, \quad Me_1 = e_8, \quad Me_2 = e_9, \quad Me_3 = e_6. \quad (6.8)$$

Then $M^T M = I$ and the connection is

$$\mathcal{B} = iA_{\mu} dx^{\mu} - X_7 dx^0 + X_8 dx^1 + X_9 dx^2 + X_6 dx^3, \quad \mu = 0, 1, 2, 3. \quad (6.9)$$

The same nonzero constant complex parameter works for every Euclidean four-dimensional tangent, hence for all normal planes indexed by S^2 . This is not a statement about arbitrary real Lorentzian trajectories with real internal couplings.

6.4 The remaining H weight

Set $T_H = i\Gamma_4\Gamma_5$. The exact identity

$$D = K_n T_H \quad (6.10)$$

implies $T_H \epsilon = -\epsilon$ for our signs. The tangent projectors reduce the spacetime/V factors but do not introduce the other H weight. Thus a surviving constant charge has the same one-dimensional H factor at both endpoints. Using Baker's generator convention, the plus-normal combination $iA_3 + X_6$ is labeled by preserved generators Q_{i2} ; the conjugate convention exchanges 1 and 2. Spinor-parameter and generator indices are related by charge conjugation, so the endpoint labels must be fixed from the physical transformations, not read off from a gamma eigenvalue alone.

7 Endpoint polarization, vanishing, and the adjoint

7.1 The same-charge endpoint calculation

The physical scalar variations may be written $\delta q_m = 2\bar{\zeta}_{im}\Psi_i$ and $\delta \bar{q}_m = 2\bar{\Psi}_i\zeta_{im}$, with the related physical generator conjugation; see [5, Appendix C]. Complexification does not make the actions on q and $\bar{q}$ independently adjustable. For the plus-normal choice the annihilated scalar components are q_2 and $\bar{q}_1$.

Here is a basis-independent polarization version of that statement. Suppress the nonzero spacetime/V factor of the charge, let its H factor be $a \in \mathbb{C}^2 \setminus \{0\}$, and write

$$q(v) = \sum_m v_m q_m, \quad \bar{q}(w) = \sum_m w_m \bar{q}_m, \quad E = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}. \quad (7.1)$$

In the generator convention just specified, the invariant tensors give

$$Q(a)q(v) \propto (a^T E v)\Psi, \quad Q(a)\bar{q}(w) \propto (a^T w)\tilde{\Psi}. \quad (7.2)$$

This follows by rotating the representative conditions $Q_{i2}q_2 = Q_{i2}\bar{q}_1 = 0$, with E the fundamental antisymmetric tensor and the second contraction the fundamental/dual evaluation. The suppressed fermion components are arbitrary, so an operator annihilator requires the scalar coefficient to vanish.

Proposition 7.1 (Zero same-charge scalar pairing). *For the constant fixed-H-factor charge above, nonzero ordinary fundamental and antifundamental scalar endpoints separately killed by that charge have zero mutual free contraction. They are not a physical adjoint pair.*

Proof. The two nullspaces in (7.2) are

$$v = \lambda a, \quad w = \mu E a. \quad (7.3)$$

The H tensor of the free propagator is δ_{mn} , so its polarization factor is $v^T w = \lambda \mu a^T E a = 0$. Physical conjugation instead requires $w = v^*$ and gives $v^T v^* = \|v\|^2 > 0$. If $v = \lambda a \neq 0$, then $a^T v^* = \lambda^* \|a\|^2 \neq 0$, so the physical conjugate is not killed by that same charge. $\square$

In the representative basis, q_2 pairs nontrivially with $\bar{q}_2$, whereas its same-charge partner is $\bar{q}_1$. The physical adjoint pair is a useful nonzero control, but its dressed expectation still requires an independent positivity argument. At distinct endpoints the two supersymmetry variations contain independent fermion fields, so they cannot cancel each other as bare operator identities. In this ansatz the bulk line variation already vanishes separately. Junction matter or a fermionic superconnection would change the operator and require a new calculation.

7.2 The full bilinear has nonzero R charge

The fields in (6.2) and (6.9) are neutral under the H rotation fixing X_6 . Assign q_1, q_2 weights $+1/2, -1/2$. Then $\bar{q}_1$ also has weight $-1/2$ and

$$\mathcal{O}_C = \bar{q}_1(b)U_M(C)q_2(a) \quad (7.4)$$

has charge -1 . If the vacuum and the definition of the operator preserve this symmetry, its expectation obeys

$$\langle \mathcal{O}_C \rangle = e^{-i\alpha} \langle \mathcal{O}_C \rangle \quad \text{for every } \alpha, \quad \langle \mathcal{O}_C \rangle = 0. \quad (7.5)$$

This selection rule extends beyond the free contraction. It does not construct an ultraviolet-finite line operator. A compensating charged insertion, a different endpoint sector or a symmetry-breaking vacuum changes its assumptions. A fixed-endpoint deformation of the same neutral connection carries no compensating charge.

7.3 Why the original semicircle is a nonzero control

The restriction is to a constant Poincare charge and the displayed tangent map. On (4.1), that map has scalar direction $n_8 = -\sin \theta$, $n_6 = \cos \theta$: the scalar normal direction changes sign along with the tangent. The common Poincare charge therefore keeps its H weight fixed.

The original semicircle with constant scalar arclength coupling instead uses a position-dependent conformal Killing spinor. Its local endpoint conditions can select different H factors at the two ends. Baker's same-polarization semicircle has a nonzero leading endpoint contraction, proportional to $g^2/(4\pi|a-b|)$ in that paper's normalization [5, Section 3.2]. It is consequently a successful control outside Proposition 7.1, rather than a counterexample to it. This distinction is also why the rigidity theorem and tangent-map construction must not be conflated.

7.4 The adjoint changes the scalar map

For a tangent-coupled line and Hermitian matrix fields,

$$U_M(C)^\dagger = U_{-M}(C^{-1}), \quad U_M(C^{-1}) = U_M(C)^{-1}. \quad (7.6)$$

To prove the first identity, reverse the product order in the discretized ordered exponential and conjugate each exponent. The gauge term changes sign under conjugation; after reversing the contour it is again $iA_\mu dx^\mu$, while the scalar one-form has acquired $-M$. The second identity is ordinary inverse transport. Thus contour reversal with unchanged tangent map gives the inverse, not the Hermitian adjoint. A reflected norm must implement the former identity with the appropriate spacetime pullback. This is the purpose of the next section.

8 Physical reflection and the reference color junction

8.1 Ordinary time reflection

Reflect a time direction tangent to the defect:

$$\vartheta x = Rx, \quad R = \text{diag}(-1, 1, 1, 1), \quad b = 0 \in \{x^0 = 0\} \cap \{x^3 = 0\}. \quad (8.1)$$

The positive half-space is $x^0 > 0$. In the bosonic adjoint convention, A transforms as a one-form, all six Hermitian scalars are even under geometric time reflection, and conjugation exchanges q_m with

$\bar{q}_m$. Fermionic reflection and an interacting regulator are additional requirements, not consequences of these bosonic rules.

Let $C : a \rightarrow b$ lie in the positive half-space except at b , with a on the defect. Use M from (6.8). Pullback of the fields first gives $U_{MR}(\vartheta C)$; applying (7.6) then gives

$$\Theta_0 U_M(C) = U_N(\vartheta C^{-1}), \quad N = -MR. \quad (8.2)$$

Explicitly,

$$Ne_0 = -e_7, \quad Ne_1 = -e_8, \quad Ne_2 = -e_9, \quad Ne_3 = -e_6. \quad (8.3)$$

The time component retains its scalar map and the spatial components reverse it. This ordering calculation agrees with the distinction between geometric reflection and scalar-coupled adjoint transport in [9, eqs. (4)–(7)].

8.2 The actual paired operator and the conditional Gram statement

Define a covariant color vector at the reference by

$$F_i^\alpha = [U_M(C_i)]^\alpha{}_\beta q_2^\beta(a_i). \quad (8.4)$$

Its reflected row is $\bar{q}_{2,\gamma}(\vartheta a_i)[U_N(\vartheta C_i^{-1})]^\gamma{}_\alpha$. The proposed kernel is

$$K_{ij} = \left\langle \bar{q}_{2,\gamma}(\vartheta a_i)[U_N(\vartheta C_i^{-1})]^\gamma{}_\alpha [U_M(C_j)]^\alpha{}_\beta q_2^\beta(a_j) \right\rangle. \quad (8.5)$$

There is an identity color contraction at b , not another defect field there. Under a gauge transformation, $F_i \mapsto g(b)F_i$ and its reflected row transforms by $g(b)^{-1}$, so the paired observable is gauge invariant. The individual vector is not a gauge-invariant state without a specified reference charge sector.

Assumption 8.1. A regulator and reference color sector admit Osterwalder–Schrader positivity for these half-space functionals. The regulated endpoint and junction operators belong to that domain, and their limiting renormalization preserves the inequality whenever a limit is claimed.

Proposition 8.2 (Conditional positivity). *Under the regulated part of Assumption 8.1, the matrix in (8.5) is positive semidefinite for every finite collection of admissible contours and endpoints.*

Proof. Apply the assumed OS inequality to $\sum_i c_i F_i$, including its reference color fiber, to obtain $\sum_{ij} \bar{c}_i K_{ij} c_j \geq 0$. For a concrete bosonic sufficient model, suppose the two half-integrals factor over boundary data B with positive measure ν . They give vector wavefunctions $\psi_i^\alpha(B)$ and

$$K_{ij} = \int \sum_\alpha \overline{\psi_i^\alpha(B)} \psi_j^\alpha(B) d\nu(B), \quad \nu \geq 0. \quad (8.6)$$

The claimed inequality is then the integral of a squared norm. Boundary gauge transformations act unitarily on the displayed fiber. $\square$

This conditional statement locates the missing physical input. A positive bosonic factorization is not asserted for the full fermionic $\mathcal{N} = 4$ defect measure. Gauge invariance of (8.5) alone proves no reflection inequality. Gauss law, the reference charge sector, the regulator, and junction counterterms must all be addressed in an interacting realization.

8.3 An unconditional free kernel

At leading free endpoint order $U = I$. Strip the positive color and coupling constants and write $a_i = (\tau_i, \mathbf{z}_i, 0)$ with $\tau_i > 0$, $\mathbf{z}_i \in \mathbb{R}^2$. Then

$$K^{(0)}(a_i, a_j) = \frac{1}{\sqrt{(\tau_i + \tau_j)^2 + |\mathbf{z}_i - \mathbf{z}_j|^2}}. \quad (8.7)$$

Its representation

$$K^{(0)}(a_i, a_j) = 4\pi \int_{\mathbb{R}^2} \frac{d^2k}{(2\pi)^2} \frac{e^{-|k|(\tau_i + \tau_j)} e^{ik \cdot (\mathbf{z}_j - \mathbf{z}_i)}}{2|k|} \quad (8.8)$$

is an integral of products $\overline{\psi_i(k)} \psi_j(k)$ with positive measure. This proves free positivity independently of Assumption 8.1. On the time ray $a_r = (r, 0, 0, 0)$ it reduces to

$$K^{(0)}(r, s) = \frac{1}{r + s} = \int_0^\infty e^{-\lambda r} e^{-\lambda s} d\lambda, \quad \sqrt{rs} K^{(0)}(r, s) = G_o(\log r - \log s). \quad (8.9)$$

The diagonal is finite for $r > 0$. The kernel is Hankel in r, s , and translation invariant after the radial weights and logarithmic coordinates. It is the opposite-ray kernel of (3.5), not the full even archimedean tower. The full-sphere profile in Section 3 cannot be used as the same half-space state without changing its support and recomputing its weights.

8.4 The two branches preserve different constant charges

Let $\widetilde{K}_\mu = i\Gamma_{Me_\mu} \Gamma_\mu$ for $\mu = 0, 1, 2$, and $K_n = i\Gamma_6 \Gamma_3$. With $\chi = D = +1$, the two branches require the following eigenvalues:

Involution	Ket with M	Reflected bra with N
$\widetilde{K}_0$	-1	-1
$\widetilde{K}_1, \widetilde{K}_2$	-1	+1
K_n	-1	+1

Each all-direction family has one complex compatible charge; in the (x^0, x^3) -plane each has two. Their common subspace is zero when time and normal tangents are independently required, because the normal conditions are opposite. The normal projectors also select opposite H weights: the ket uses q_2 and the physical bra uses $\bar{q}_2$ with the conjugate-weight charge. The identity tensor at b cannot compensate a nonzero variation on a branch.

As a bulk control, a straight time-directed contour needs only $\widetilde{K}_0 = -1$. That condition is common to the maps and gives a four-dimensional complex chiral defect solution space. It is a bulk control, not a statement that a constant charge annihilates the physical adjoint pair of endpoint scalars.

8.5 Internal rotations do not automatically preserve positivity

The proper rotations by π about H axis 4 and V axis 7 are

$$S_H = \text{diag}(1, -1, -1), \quad S_V = \text{diag}(1, -1, -1). \quad (8.10)$$

For $S = S_H \oplus S_V$, one checks $SN = M$. Thus an internally rotated reflection aligns the two bulk scalar maps. But S_H acts on the endpoint doublet by $\pm i\sigma_1$, up to basis convention. It takes $\bar{q}_2$ to a phase times $\bar{q}_1$, restoring the zero diagonal contraction of Proposition 7.1.

On the full free endpoint doublet the inserted metric is not positive. Without a phase adjustment its eigenvalues are $\pm i$; with a phase making it Hermitian they are $+1, -1$. For instance

$$(1, -1)\sigma_1 \begin{pmatrix} 1 \\ -1 \end{pmatrix} / 2 = -1. \quad (8.11)$$

No overall phase makes both eigenvalues positive. The identity metric from ordinary adjoint pairing is the control. More generally, if a unitary S on a positive Hilbert space satisfies $\langle \psi, S\psi \rangle \geq 0$ for every ψ , then S is positive self-adjoint; unitarity forces its spectrum to be $\{1\}$ and $S = I$. Projection to one rotation eigenspace is a different operation and here mixes the fixed H endpoint weights.

This is a counterexample for the displayed doublet state space. It does not contradict the special large-rectangle rotated-reflection conjecture distinguished from ordinary reflection in [9, eqs. (11)–(16)].

8.6 A free test of an odd-H normal reflection

A different proposal in the parent investigation reflected across the defect and assigned odd parity to the H scalars. For the canonical free scalar algebra, a real mode of frequency $\nu > 0$ at distance $a > 0$ from a reflection plane has ordinary norm $e^{-2\nu a}/(2\nu) > 0$; assigning it odd reflection parity changes the sign. The failure is also visible in a gauge-invariant observable. For a non-Abelian group with $\text{tr}(T_a T_b) = \delta_{ab}/2$, take the Hermitian Wick polynomial

$$\mathcal{O} = i \text{tr}(H[V_1, V_2]), \quad (8.12)$$

where H is one H scalar and V_1, V_2 are distinct V scalars. Writing $[T_b, T_c] = i f_{bca} T_a$, its coefficient is $-f_{bca}/2$. Distinct species have only their matching free contractions, hence at separated reflected points

$$\langle \Theta_{\text{even}} \mathcal{O}(x) \mathcal{O}(x) \rangle = \frac{1}{4} \sum_{abc} f_{abc}^2 D_0(\vartheta x - x)^3 > 0, \quad (8.13)$$

with D_0 the positive scalar covariance. Odd H and even V parity give the negative of this value. This excludes that reflection on the unrestricted canonical free bulk observable algebra. It does not classify all interacting normal-reflection actions or restricted observable sectors. The even assignment supplies the positive control.

9 A smooth reference join and a direct bulk response

9.1 A contour family that meets the reflection plane smoothly

Let v be the incoming unit ket tangent at b . Its reflected reverse has outgoing tangent $-Rv$. A smooth join requires

$$v = -Rv, \quad (9.1)$$

so v is parallel to the time direction e_0 . The internal scalar directions then agree as well, since $N(-Rv) = Mv$. For a kernel involving many contours, every ket should have the same reference tangent. The old semicircles arrive along $-e_3$ and return along $+e_3$ under time reflection, producing a backtracking join instead.

Contour freedom provides a simple replacement. In the (x^0, x^3) -plane set

$$X_{T,\eta}(t) = (Tt, \eta T f(t)), \quad f(t) = t^2(1-t), \quad 0 \leq t \leq 1, \quad T > 0, \quad \eta \geq 0. \quad (9.2)$$

The ket traverses the curve from $t = 1$ to zero. The endpoints are on the defect, its interior is in positive time and, for $\eta > 0$, strictly above the defect. Since $f'(0) = 0$, its incoming unit tangent is $-e_0$. The reflected return is regular with a continuous scalar direction even when the two branches have different T or η . Curvature need not agree.




Figure 1: A polynomial bulge and its ordinary reflected adjoint. The ket runs from a to b , and the bra from b to ϑa . Both branches lie above the defect and meet with tangent $-e_0$. The scalar maps differ away from the join. Vertical scale is enlarged.

This construction removes a fixed-angle geometric and scalar-direction cusp in the direct bulk term. It does not prove absence of defect contact or junction counterterms. No driver or capacity law for (9.2) is computed here; it is a smooth contour control for the field-theory problem, not an identification with ω .

9.2 The specified Gaussian perturbative sector

Use the free Feynman-gauge propagators

$$\langle A_\mu^a(x) A_\nu^b(y) \rangle_0 = \frac{g^2 \delta^{ab} \delta_{\mu\nu}}{4\pi^2 |x - y|^2}, \quad \langle X_A^a(x) X_B^b(y) \rangle_0 = \frac{g^2 \delta^{ab} \delta_{AB}}{4\pi^2 |x - y|^2}. \quad (9.3)$$

This defines the direct exchange of a bulk field between the two line branches. Endpoint self-energy, endpoint-to-line interactions, defect vertices and counterterms are excluded from this sector. Only their consistent sum could be a gauge-independent interacting answer.

On a single branch the gauge/scalar numerator cancels by $M^T M = I$ or $N^T N = I$. If v_i, v_j denote oriented original ket tangents, the cross-branch numerator instead is

$$\begin{aligned} -(-Rv_i) \cdot v_j + [N(-Rv_i)] \cdot Mv_j &= v_i \cdot (R + I)v_j \\ &= 2v_{i,\text{sp}} \cdot v_{j,\text{sp}}. \end{aligned} \quad (9.4)$$

Here sp consists of the three directions unchanged by time reflection, including the defect normal. Near a regular reference join satisfying (9.1), the spatial tangents vanish linearly in arclength. The numerator is $O(st)$ and the denominator is comparable to $(s + t)^2$. The local double integral is finite. This local estimate concerns (9.3), not all possible defect graphs.

For the self-pairing of (9.2), T cancels by scale invariance and the stripped cross term is

$$I(\eta) = 2\eta^2 \int_0^1 \int_0^1 \frac{f'(s)f'(t)}{(s+t)^2 + \eta^2[f(s) - f(t)]^2} ds dt. \quad (9.5)$$

If $\sum_a T_a T_a = C_F I$ in the endpoint representation, its contribution relative to the free endpoint propagator is $g^2 C_F I(\eta)/(4\pi^2)$. There is no factor 1/2 for an exchange between two distinct ordered branches.

Proposition 9.1 (The direct bulk bulge coefficient). *For $f(t) = t^2(1 - t)$,*

$$I(\eta) = \left(\frac{3}{2} - 2 \log 2\right) \eta^2 + O(\eta^4), \quad \frac{3}{2} - 2 \log 2 = 0.1137056388801094 \dots > 0. \quad (9.6)$$

One may bound the absolute remainder by $2\eta^4$ for real η .

Proof. The quadratic coefficient is $2A$, where

$$A = \int_0^1 \int_0^1 \frac{(2s - 3s^2)(2t - 3t^2)}{(s + t)^2} ds dt. \quad (9.7)$$

Put $r = s/(s + t)$ and $u = s + t$. The Jacobian is u . By symmetry, integrate $0 \leq r \leq 1/2$ and $0 \leq u \leq 1/(1 - r)$ and multiply by two. The integrand including the Jacobian is $r(1 - r)[4 - 6u + 9r(1 - r)u^2]u$. Its u integral, including that factor two, simplifies to $r^2/[2(1 - r)^2]$. Hence

$$A = \frac{1}{2} \int_0^{1/2} \frac{r^2}{(1 - r)^2} dr = \frac{3}{4} - \log 2. \quad (9.8)$$

For the remainder, $|f'| \leq 1$, $|f'(s)| \leq 2s$ and $|f(s) - f(t)| \leq |s - t| \leq s + t$. Subtracting the denominator at $\eta = 0$ bounds the absolute integrand difference by

$$2\eta^4 \frac{|f'(s)f'(t)||f(s) - f(t)|^2}{(s + t)^4} \leq 8\eta^4 \frac{st}{(s + t)^2} \leq 2\eta^4. \quad (9.9)$$

The integration square has area one. This justifies the expansion uniformly even at the apparent singular corner. $\square$

Numerical controls give $I(0) = 0$, $I(0.3) \simeq 0.0102220294386856$ and $I(0.7) \simeq 0.0553790171218514$. These check the integral and its small-bulge coefficient; they are not estimates of the complete interacting response.

There is also a positive Gaussian interpretation of this cross term. The reflected four-dimensional scalar covariance has representation

$$D_0(\vartheta x - y) = \int_{\mathbb{R}^3} \frac{d^3k}{(2\pi)^3} \frac{e^{-|k|(x^0 + y^0)} e^{ik \cdot (y - x)}}{2|k|}. \quad (9.10)$$

Smear it against the three spatial tangent currents of each ket and sum the components as in (9.4). This is a Gram matrix. The vanishing of those currents at b permits the finite join limit just estimated. This proves positivity of the displayed Gaussian current sector, without extending it to the omitted diagrams.

9.3 Why a positive norm need not be protected

For an actual reflected pairing a smooth deformation has the structure

$$\delta K_{ij} = \langle \Theta_0(\delta F_i) F_j \rangle + \langle \Theta_0 F_i \delta F_j \rangle, \quad (9.11)$$

with the insertions of Section 5 and any renormalized contact terms. Even if a deformation of a ket is Q -exact, $\delta\psi = Q\lambda$, and $Q\psi = 0$, its physical norm obeys

$$\delta \|\psi\|^2 = 2 \operatorname{Re} \langle \psi, Q\lambda \rangle = 2 \operatorname{Re} \langle Q^\dagger \psi, \lambda \rangle. \quad (9.12)$$

The first condition does not imply $Q^\dagger \psi = 0$. For example, on a Hilbert space with orthonormal e_1, e_2 , let $Qe_2 = e_1$, $Qe_1 = 0$, $\psi = e_1$, $\lambda = e_2$. Then $Q^2 = 0$, $Q\psi = 0$, but (9.12) equals two. Adding a vector e_3 killed by both Q and $Q^\dagger$ supplies a zero-first-variation control. Thus a common-charge Ward identity and a physical positive norm are different requirements.

10 Status and the remaining route to a proof

The combined investigation remains open as a field-theoretic research problem. The original fixed-scalar family has an exact Loewner description but little contour freedom at fixed charge. Tangent coupling removes that geometric restriction and admits the full chiral defect projectors. Its ordinary same-charge scalar pairing vanishes. Physical reflection produces a different, nonzero free pairing and a smooth reference join, but it changes the branch charges and has no established protection theorem. These conclusions can coexist: they concern different pairings of the same basic fields.

10.1 Statement ledger

Statement	Status in this draft
Weil positivity criterion	Classical background theorem; target normalization given in (2.3).
Beta/comb and rational factorization	Exact identities reproduced with their convergence region; no contraction conclusion.
Free even angular kernel	Positive for $\rho < 1$; its difference energy has a positive limit. The opposite contact subtraction does not.
Auxiliary gamma partition ratio	Exact finite Gaussian product and renormalized limit; no physical mode selection or transfer identified.
Constant-driver dictionary and rigidity	Exact contour identity and proof under the specified fixed-scalar bulk charge.
Smooth transport and inverse-map variation	Exact matrix-field ODE identities away from singularities; no rough-trace quantum construction.
Tangent-map defect supersymmetry	Complex chiral projector ranks proved for the real constant-map ansatz.
Same-charge scalar endpoint pairing	Zero free contraction; vanishing expectation conditional on an invariant vacuum and operator definition.
Ordinary reflected network	Exact classical map and color gluing; free endpoint positivity proved. Interacting positivity is conditional.
Rotation and odd-H tests	Explicit failures on the stated free algebras, with ordinary-adjoint positive controls.
Smooth join and bulge response	Direct Feynman-gauge Gaussian bulk sector computed; full interacting response remains open.
Arithmetic identification	No prime-weight mechanism, complete transfer, or proof of positivity of $Q_{0,L}$ has been obtained.

10.2 The next discriminating calculation

The immediate continuation is the complete first interaction-order shape response of (8.5), beginning with a positive bulge $\eta_0 > 0$ in (9.2). Fix the gauge, endpoint normalization, reference color sector and renormalization prescription before comparing deformations. Alongside the direct bulk exchange, include endpoint-to-line gauge exchange, the defect H-scalar derivative coupling, endpoint self-energy, and all defect and junction counterterms at that order. The interaction vertices must be taken from

a consistent defect action, such as the conventions used in [3, 5]. Endpoint renormalization may cancel in a specified ratio, but that cancellation must be shown.

The flat limit needs separate analysis. At $\eta = 0$ the entire contour lies on the defect; for $\eta > 0$ only its endpoints do. Defect contact terms can therefore make the full small-height expansion nonuniform. Analyticity of the bulk integral in η^2 does not justify analyticity of the complete answer: terms linear in height or involving logarithms have not been excluded. A complete calculation at positive height would first determine whether the nonzero bulk response survives, cancels, or requires a different operator definition.

10.3 What would turn the construction into an RH route

A precise sufficient endpoint would be a Hilbert space $\mathcal{H}_L$ and a linear map J_L defined for every compactly supported smooth f such that

$$Q_{0,L}[f] = \|J_L f\|_{\mathcal{H}_L}^2 \quad \text{for every } L > 0. \quad (10.1)$$

Alternatively, an independently defined positive system could realize the causal transfer (2.7) with $\|V_{\omega,L}\| \leq 1$ for every L and shifts tending to zero. Equation (2.11) would then give the same positivity conclusion. Either identity must be derived, rather than built into an assumed positive metric.

For the present proposal this entails the following concrete steps.

1. Define a nonzero observable and its physical adjoint, including the reference sector, all endpoint and junction data, a regulator, and a limit preserving the required positivity.
2. Derive an evolution law for the relevant averaged observable. The insertions in Section 5 must close in a specified sector or be controlled by a proved identity. A common bulk charge alone does not establish this.
3. Identify the contour or boundary parameter with the arithmetic shift and frequency. The capacity relation (4.5) and the auxiliary Gaussian ratio (3.8) are partial dictionaries with different meanings; neither makes this identification.
4. Produce the full arithmetic form: the local normalization w_0 , the pole contribution, and each $\Lambda(n)/\sqrt{n}$ translation with its sign. Equivalently produce the full transfer, including the prime comb and rational correction. A free spectral resemblance or a positive kernel cannot replace these exact terms.
5. Prove the equality and bounds for all test functions and an unbounded range of interval lengths, with the limits needed in (10.1) or (2.11). Finite Gram matrices and perturbative coefficients are diagnostics for this task, not substitutes for it.

The present work supplies explicit objects on which to test the first two steps and exposes where protection and positivity separate. Its value for the RH program depends on making one of those tests succeed and then deriving the missing arithmetic structure.

A Reproducible diagnostics and controls

The accompanying package contains three Python standard-library programs and their preserved JSON results. They register 268 finite cases. Each case records its name, numerical value, comparison

direction, threshold and pass status. Replay requires identical parameters, case identities and thresholds and a passing result; small platform-dependent changes in floating values are permitted. No case uses zeta zeros or infers an arithmetic positivity theorem.

Program in numerics/	Cases	Main comparisons
<code>check_smooth_variation.py</code>	102	Inverse-map geometry, ordered ODE variation, endpoint motion, reparametrization, abelian and scalar-coupling controls.
<code>check_defect_endpoints.py</code>	99	Chiral projectors, all orientation signs, endpoint nullspaces, R charge and noncommuting adjoint transport.
<code>check_reflected_junction.py</code>	67	Reflected maps and color gluing, free Gram matrices, twist failures, smooth joins and independent bulk quadrature.

For transparency the Clifford matrices may be generated by five tensor factors. With $j = 0, \dots, 4$ and Pauli matrices σ_i ,

$$\Gamma_{2j} = \sigma_3^{\otimes j} \otimes \sigma_1 \otimes I^{\otimes(4-j)}, \quad \Gamma_{2j+1} = \sigma_3^{\otimes j} \otimes \sigma_2 \otimes I^{\otimes(4-j)}. \quad (\text{A.1})$$

They satisfy the stated Clifford relations. Projector traces are compared with independent row reduction. Correct triplet assignments and the allowed orientation supply nonzero controls for the excluded choices. The adjoint check uses noncommuting Hermitian matrix segments, so a mistaken reversal of matrix order cannot pass merely because an abelian example commutes.

The transport checks compare direct variation of an ordered ODE solution with the integrated insertion formula on smooth matrix backgrounds, including mesh refinement. The reflected response checks compare the analytic coefficient with a separately integrated double integral and check finite Gaussian current Gram matrices. The flat and ordinary-adjoint cases are positive or zero controls for the claimed failures. These comparisons support the written derivations; they do not certify physical supersymmetry transformations, quantum operator existence, continuum reflection positivity, all-order Ward identities or a stochastic contour limit.

The standalone manuscript package can be checked with

```
python3 validation/drafts.py check -replay
```

The live investigation additionally has `validation/check_package.py`, which inventories the research notes and their inherited repository sources. That broader inventory is separate from the portable manuscript snapshot.

The auxiliary free-kernel background in Sections 2 and 3 is derived explicitly in the text. The 268 count refers to this investigation's three programs, not to every calculation in its parent investigations. Hash and build records establish file identity and reproducibility, not mathematical correctness.

B Provenance and contribution disclosure

This first manuscript version was assembled on 19 September 2026 by OpenAI GPT-6 (Codex), for Edward Baker, from the Wilson–Loewner investigation and its parent research. It is a working synthesis, not a submitted or independently refereed paper. The title-page designation records for whom it was drafted and does not assert human verification of every derivation. No independent review of this package has yet been completed.

The principal local source notes, all dated 19 September 2026, are

1. *Smooth Wilson variation and the constant Loewner driver*, supplying Sections 4 and 5;
2. *Defect-compatible contour freedom and the endpoint pairing*, supplying Sections 6 and 7;
3. *Physical reflection, the reference junction, and a first bulk response*, supplying Sections 8 and 9.

They are retained with the earlier handoffs in the live investigation’s `notes/` directory. The parent Wilson-lines continuation supplies the endpoint motivation, free angular regulator and auxiliary boundary Gaussian; the Loewner and fractional-dimension manuscripts supply the arithmetic factorization and the scoped realization obstructions. All formulas needed for the present arguments have been included here. The manuscript has no shared-background TeX dependency.

The derivations use published conventions where stated. The printed equations for the Loewner normalization, tangent-coupled projectors, D3–D5 projector, Baker endpoint transformations and semicircle control, Dorn–Pershin reflection rules, and the Weil-form weights were checked against the corresponding PDFs. Bibliographic metadata and primary links were checked during preparation. This is a focused background review, not an exhaustive literature or priority survey. No novelty is claimed for standard Wilson variation, tangent-coupled supersymmetry, the ray endpoint selection rule, or the classical Weil criterion.

The machine-readable `MANUSCRIPT_SOURCES.json` identifies the local research inputs and their hashes at assembly. The `BUILD_RECORD.json` binds the reviewed PDF to its TeX sources and supporting checks; `SNAPSHOT.json` binds a dated archive. Neither record is a proof certificate. Historical notes and snapshots are preserved so that a later correction can be distinguished from the claims made in this version.

References

- [1] J. M. Maldacena, *Wilson loops in large N field theories*, Physical Review Letters **80** (1998), 4859–4862, [arXiv:hep-th/9803002](#).
- [2] K. Zarembo, *Supersymmetric Wilson loops*, Nuclear Physics B **643** (2002), 157–171, [arXiv:hep-th/0205160](#).
- [3] O. DeWolfe, D. Z. Freedman and H. Ooguri, *Holography and defect conformal field theories*, Physical Review D **66** (2002), 025009, [arXiv:hep-th/0111135](#).
- [4] D. Gaiotto and E. Witten, *Supersymmetric boundary conditions in $\mathcal{N} = 4$ super Yang–Mills theory*, Journal of Statistical Physics **135** (2009), 789–855, [arXiv:0804.2902](#).
- [5] E. B. Baker III, *Supersymmetric open Wilson lines*, Journal of High Energy Physics **03** (2011), 140, [arXiv:1102.4948](#).
- [6] G. F. Lawler, *Schramm–Loewner evolution (SLE)*, lecture notes, especially Lecture 2, Proposition 2.17 and equation (2.8), [author’s PDF](#), accessed 19 September 2026.
- [7] M. Bauer and D. Bernard, *Conformal field theories of stochastic Loewner evolutions*, Communications in Mathematical Physics **239** (2003), 493–521, [arXiv:hep-th/0210015](#).
- [8] K. Osterwalder and R. Schrader, *Axioms for Euclidean Green’s functions*, Communications in Mathematical Physics **31** (1973), 83–112, [doi:10.1007/BF01645738](#).

- [9] H. Dorn and V. D. Pershin, *Concavity of the $Q\bar{Q}$ potential in $\mathcal{N} = 4$ super Yang–Mills gauge theory and AdS/CFT duality*, Physics Letters B **461** (1999), 338–344, [arXiv:hep-th/9906073v3](#).
- [10] M. Suzuki, *Weil’s quadratic form via the screw function*, preprint (2026), version 2, [arXiv:2606.09096v2](#). Used for the classical compact-support Weil criterion and conventions; its proposed limiting spectral construction is not assumed here.
- [11] Riemann Hypothesis research program, *Deformation flows for the shifted Weil family: the transfer as an all-pass filter*, working draft 0.8, 17 September 2026, and endpoint continuation notes, 18–19 September 2026. Repository investigation [papers/susy-positivity/investigations/wilson-lines/](#). Internal research, with model-assisted contributions disclosed in the source documents.
- [12] OpenAI GPT-6 (Codex), for Edward Baker, *Angular smearing, the finite difference form, and a Robin model of the shift*, research note, 19 September 2026, in the Wilson-lines investigation, [notes/ANGULAR_SMEARING_AND_ROBIN_MODEL_20260919.md](#).
- [13] Riemann Hypothesis research program, *The Markov part of the shifted Weil transfer*, working draft 0.3, 18 September 2026. Repository investigation [papers/susy-positivity/investigations/loewner/](#). Internal research; its direct realization proposal was closed within the tested scope.
- [14] Riemann Hypothesis research program, *A scattering matrix without a space*, working draft 0.1, 18 September 2026. Repository investigation [papers/susy-positivity/investigations/fractional-dimension/](#). Internal research; the continued lattice count and its limitations are discussed in its Section 6.